

The complete mitochondrial genome of the Antarctic marbled rockcod, *Notothenia rossii* (Perciformes, Nototheniidae)

Euna Jo^{a,b,*} , Yll Hwan Cho^{a,*}, Seung Jae Lee^a , Eunkyung Choi^a, Jeong-Hoon Kim^c, Young Min Chi^a, Jin-Hyoung Kim^b and Hyun Park^a 

^aDivision of Biotechnology, College of Life Sciences and Biotechnology, Korea University, Seoul, Korea; ^bUnit of Research for Practical Application, Korea Polar Research Institute (KOPRI), Incheon, Korea; ^cDivision of Polar Life Science, Korea Polar Research Institute, Incheon, Korea

ABSTRACT

The complete mitochondrial genome of *Notothenia rossii* was obtained using PacBio Sequel long-read sequencing platform. The mitogenome of *N. rossii* was circular form and 18,274 bp long, which consists of 13 protein-coding genes, 24 tRNAs, 2 rRNAs, and non-coding control region. Particularly, we found duplicated tRNA^{Thr} and tRNA^{Pro} in addition to the typical 22 tRNAs. The phylogenetic tree revealed that *N. rossii* was most closely related to *N. coriiceps* among species in the Nototheniidae clade within the suborder Notothenioidei.

ARTICLE HISTORY

Received 6 April 2020
Accepted 24 May 2020

KEYWORDS





Notothenia rossii;
mitochondrial genome;
Antarctica;
Notothenioidei; PacBio

Notothenia rossii Richardson, 1844 is a species belonging to the family Nototheniidae, also called the common name marbled rockcod. Like other notothenid fish, *N. rossii* widely distributes in the Southern Ocean and around Antarctica, although the population has been decreased by overfishing due to commercial importance in the fisheries (Barrera-Oro and Marschoff 2007). There are 7 recognized species in the genus *Notothenia* according to the FishBase (Froese and Pauly 2019). Up to now, mitochondrial and whole genome sequences in the genus *Notothenia* have been reported for only one species of *N. coriiceps* (Shin et al. 2014; Oh et al. 2016). Here, we provide the complete mitochondrial genome of *N. rossii* (GenBank accession No. MT192936) and the results of the phylogenetic analysis with Antarctic fish species.

The sample of *N. rossii* was collected from the sea near Barton Peninsula, King George Island, West Antarctica (62°14'S, 58°47'W). The specimen was deposited at the Earth Biocollection in the Division of Biotechnology, Korea University with accession number KAN0007030. The genomic DNA was extracted by phenol/chloroform method and then g-TUBE (Covaris, CA, USA) shearing device and BluePippin system (Sage Science, MA, USA) were used for preparing 20 kb size-selected templates. The SMRTbell library preparation and sequencing were performed using PacBio Sequel platform according to the manufacturer's protocol (Pacific Biosciences, CA, USA). The quality and quantity for library were checked with Fragment analyzer (Agilent Technologies,

CA, USA) and Qubit 2.0 Fluorometer (Invitrogen, Life Technologies, CA, USA), respectively. PacBio subreads for mitochondrial genome were filtered out using partial sequences of 16S rRNA and COX1 genes. De novo assembly was conducted by CANU assembler (Koren et al. 2017) and the assembled genome was annotated via MITOS web server (Bernt et al. 2013).

The complete mitochondrial genome of *N. rossii* (GenBank number: MT192936) was 18,274 bp in length, including 13 protein-coding genes, 24 transfer RNA genes (tRNAs), 2 ribosomal RNA genes (rRNAs) and non-coding control region. In particular, duplicated tRNA^{Thr} and tRNA^{Pro} are present in addition to the 22 tRNAs commonly found in fishes. The base composition of A+T (54.10%) was higher than the G+C (45.90%) in the mitogenome. Most of the protein coding genes have typical start codon (ATG) except for COX1, ATP8, and ATP6 genes (GTG). Three types of stop codons (TAA, TAG, and T(AA)) were identified. The phylogenetic relationships of *N. rossii* were analyzed with 10 Antarctic species in the suborder Notothenioidei using 13 protein-coding genes (Figure 1). Maximum Likelihood (ML) tree was built with 500 bootstrap replications and JTT matrix-based model (Jones et al. 1992) by MEGA X software (Kumar et al. 2018). The tree showed that *N. rossii* was first clustered with *N. coriiceps*, and they were grouped with species belonging to Nototheniidae (*Dissostichus eleginoides*, *D. mawsoni*, *Pagothenia borchgrevinki* and *Pleuragramma antarcticum*). This result provides

CONTACT Jin-Hyoung Kim  kimjh@kopri.re.kr  Unit of Research for Practical Application, Korea Polar Research Institute (KOPRI), Incheon 21990, Korea; Hyun Park  hpark@korea.ac.kr  Division of Biotechnology, College of Life Sciences and Biotechnology, Korea University, Seoul 02841, Korea

*These authors have contributed equally to this work.

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Maximum likelihood (ML) tree for *Notothenia rossii* within the suborder Notothenioidei. The scientific name and GenBank accession number were shown for each species.

fundamental data to further our understanding of the evolutionary relationship of Antarctic fishes in the Southern ocean.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Data availability

The data that support the findings of this study are openly available in NCBI under the accession MT192936 (<https://www.ncbi.nlm.nih.gov/nucleotide/MT192936.1/>).

Funding

This research was supported by Post-Polar Genomics Project: Functional genomic study for securing of polar useful genes [PE20040] and a Korea University Grant to H. P.

ORCID

Euna Jo  <http://orcid.org/0000-0003-2666-6743>

Seung Jae Lee  <http://orcid.org/0000-0002-2404-3107>

Hyun Park  <http://orcid.org/0000-0002-8055-2010>

References

- Barrera-Oro ER, Marschoff ER. 2007. Information on the status of fjord *Notothenia rossii*, *Gobionotothen gibberifrons* and *Notothenia coriiceps* in the lower South Shetland Islands, derived from the 2000–2006 monitoring program at Potter Cove. *CCAMLR Science*. 14:83–87.
- Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsche G, Pütz J, Middendorf M, Stadler PF. 2013. MITOS: improved de novo metazoan mitochondrial genome annotation. *Mol Phylogenet Evol*. 69(2): 313–319.
- Froese, R. & Pauly, D., Editors. 2019. FishBase. World Wide Web electronic publication. www.fishbase.org. version (12/2019).
- Jones DT, Taylor WR, Thornton JM. 1992. The rapid generation of mutation data matrices from protein sequences. *Bioinformatics*. 8(3):275–282.
- Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res*. 27(5):722–736.
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol*. 35(6):1547–1549.
- Oh JS, Ahn DH, Lee J, Choi J, Chi YM, Park H. 2016. Complete mitochondrial genome of the Antarctic bullhead notothen, *Notothenia coriiceps* (Perciformes, Nototheniidae). *Mitochondrial DNA Part A*. 27(2): 1407–1408.
- Shin SC, Ahn DH, Kim SJ, Pyo CW, Lee H, Kim M-K, Lee J, Lee JE, Detrich HW, Postlethwait JH, et al. 2014. The genome sequence of the Antarctic bullhead notothen reveals evolutionary adaptations to a cold environment. *Genome Biol*. 15(9):468.