

## Editorial



# Exploring the Role of Vitamin D Deficiency Correction in Heart Failure Management: Insights and Prospects

Hyung Yoon Kim , MD, PhD

Department of Cardiovascular Medicine, Gachon University Gil Medical Center, Incheon, Korea



► See the article “Vitamin D Deficiency in Patients Hospitalized for Heart Failure Living in the Tropics” in volume 6 on page 84.

**Received:** Apr 10, 2024

**Accepted:** Apr 15, 2024

**Published online:** Apr 17, 2024

### Correspondence to

**Hyung Yoon Kim, MD, PhD**

Department of Cardiovascular Medicine,  
Gachon University Gil Medical Center, 21  
Namdong-daero 774beon-gil, Namdong-gu,  
Incheon 21565, Korea.  
Email: medoc7@gmail.com

© 2024. Korean Society of Heart Failure

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

This issue of the *International Journal of Heart Failure* focuses on the prevalence of vitamin D deficiency (VDD) and its association with cardiometabolic parameters in heart failure (HF) patients living in the city of Recife (latitude 8° South).

HF remains a tremendous challenge in current healthcare, with its prevalence steadily increasing worldwide.<sup>1)</sup> Among the variable factors contributing to HF pathogenesis and progression, research has begun to illuminate the potential role of VDD as a modifiable risk factor.<sup>2-5)</sup> A study conducted through a bidirectional Mendelian randomization approach found that higher genetically predicted levels of 25-hydroxyvitamin D were associated with a reduced risk of HF. This suggested a potential link where correcting VDD might have a role in HF management, although the causal mechanisms and clinical implications still require further investigation.<sup>6)</sup>

In this issue of the journal, de Oliveira et al.<sup>7)</sup> present compelling findings from their cross-sectional study investigating the prevalence of VDD in HF patients residing in tropical regions and its association with cardiometabolic parameters. This study not only adds to our understanding of the complex interplay between vitamin D status and cardiovascular health but also highlights the importance of tailored management strategies in diverse patient populations. Vitamin D, long revered for its classical role in calcium homeostasis and bone health, has increasingly been recognized for its pleiotropic effects on various organ systems, including the cardiovascular system.<sup>3,5)</sup>

VDD is prevalent globally, with the lowest levels observed in the Middle East and South Asia, predominantly due to factors including elderly female demographics, higher latitudes, seasonal variations, limited sunlight exposure, skin pigmentation, dietary habits, and insufficient vitamin D fortification.<sup>8,10)</sup> Despite abundant sunlight in tropical regions, which facilitates endogenous vitamin D synthesis, the study findings reveal a high prevalence of VDD among HF patients living in tropical regions, challenging conventional assumptions regarding vitamin D sufficiency in such geographical areas. It is crucial to note that even with sunlight exposure facilitate endogenous Vitamin D synthesis, VDD persists as a global concern, affecting people across various geographic locations, including those with year-round sun exposure.<sup>8,11)</sup> This suggests that factors beyond geographic location and sunlight exposure influence Vitamin D status. This raises intriguing questions about the underlying mechanisms behind VDD in this population.<sup>12,13)</sup>

Further investigation into environmental, genetic, and lifestyle factors contributing to suboptimal vitamin D levels is warranted.

The association between VDD and adverse metabolic parameters in HF patients uncovered by this study provides valuable insights into potential pathophysiological pathways linking VDD to cardiovascular risk. Particularly interesting is the observed like between VDD and conditions like diabetes mellitus and dyslipidemia, suggesting a complex relationship with metabolic health.<sup>13,14</sup> These findings underscore the importance of holistic metabolic evaluation in HF patients, focusing on identifying and addressing modifiable risk factors to optimize clinical outcomes.

Moreover, the study highlights the need for tailored interventions to address VDD in HF patients, taking into account individual patient characteristics and environmental factors. While sunlight exposure remains a primary source of vitamin D, factors such as skin pigmentation, dietary habits, and comorbidities may influence vitamin D synthesis and metabolism. Thus, strategies aimed at optimizing vitamin D status should be individualized and integrated into comprehensive HF management plans, with the goal of mitigating cardiovascular risk and improving patient outcomes.

In conclusion, de Oliveira et al.'s study<sup>7</sup>) represents a significant contribution to the growing body of evidence exploring the relationship between VDD and HF. By elucidating the prevalence of VDD and its association with cardiometabolic parameters in HF patients residing in tropical regions, this study emphasizes the importance of considering vitamin D status in the management of cardiovascular disease. Moving forward, further research is warranted to elucidate the mechanistic links between VDD and adverse cardiovascular outcomes, paving the way for personalized therapeutic strategies aimed at optimizing vitamin D status and improving clinical outcomes in HF patients.

#### ORCID iDs

Hyung Yoon Kim   
<https://orcid.org/0000-0002-0692-5640>

#### Conflict of Interest

The author has no financial conflicts of interest.

## REFERENCES

1. Park JJ, Lee CJ, Park SJ, et al. Heart failure statistics in Korea, 2020: a report from the Korean Society of Heart Failure. *Int J Heart Fail* 2021;3:224-36. [PUBMED](#) | [CROSSREF](#)
2. D'Amore C, Marsico F, Parente A, et al. Vitamin D deficiency and clinical outcome in patients with chronic heart failure: a review. *Nutr Metab Cardiovasc Dis* 2017;27:837-49. [PUBMED](#) | [CROSSREF](#)
3. Cosentino N, Campodonico J, Milazzo V, et al. Vitamin D and cardiovascular disease: current evidence and future perspectives. *Nutrients* 2021;13:3603. [PUBMED](#) | [CROSSREF](#)
4. Luo Q, Yan W, Nie Q, Han W. Vitamin D and heart failure: a two-sample mendelian randomization study. *Nutr Metab Cardiovasc Dis* 2022;32:2612-20. [PUBMED](#) | [CROSSREF](#)
5. Latic N, Erben RG. Vitamin D and cardiovascular disease, with emphasis on hypertension, atherosclerosis, and heart failure. *Int J Mol Sci* 2020;21:6483. [PUBMED](#) | [CROSSREF](#)
6. Gao N, Li X, Kong M, et al. Associations between vitamin D levels and risk of heart failure: a bidirectional Mendelian randomization study. *Front Nutr* 2022;9:910949. [PUBMED](#) | [CROSSREF](#)
7. de Oliveira LB, de Figueiredo Martins Siqueira MA, de Macedo Gadêlha RB, Garcia J, Bandeira F. Vitamin D deficiency in patients hospitalized for heart failure living in the tropics. *Int J Heart Fail* 2024;6:84-90. [CROSSREF](#)
8. Mendes MM, Darling AL, Hart KH, Morse S, Murphy RJ, Lanham-New SA. Impact of high latitude, urban living and ethnicity on 25-hydroxyvitamin D status: a need for multidisciplinary action? *J Steroid Biochem Mol Biol* 2019;188:95-102. [PUBMED](#) | [CROSSREF](#)
9. Darling AL. Vitamin D deficiency in western dwelling South Asian populations: an unrecognised epidemic. *Proc Nutr Soc* 2020;79:259-71. [PUBMED](#) | [CROSSREF](#)
10. Porto CM, Silva VL, da Luz JS, Filho BM, da Silveira VM. Association between vitamin D deficiency and heart failure risk in the elderly. *ESC Heart Fail* 2018;5:63-74. [PUBMED](#) | [CROSSREF](#)
11. Luo Y, Qu C, Zhang R, Zhang J, Han D, Na L. Geographic location and ethnicity comprehensively influenced vitamin D status in college students: a cross-section study from China. *J Health Popul Nutr* 2023;42:145. [PUBMED](#) | [CROSSREF](#)
12. Lee JH, O'Keefe JH, Bell D, Hensrud DD, Holick MF. Vitamin D deficiency an important, common, and easily treatable cardiovascular risk factor? *J Am Coll Cardiol* 2008;52:1949-56. [PUBMED](#) | [CROSSREF](#)
13. Patel R, Rizvi AA. Vitamin D deficiency in patients with congestive heart failure: mechanisms, manifestations, and management. *South Med J* 2011;104:325-30. [PUBMED](#) | [CROSSREF](#)
14. Nolte K, Herrmann-Lingen C, Platschek L, et al. Vitamin D deficiency in patients with diastolic dysfunction or heart failure with preserved ejection fraction. *ESC Heart Fail* 2019;6:262-70. [PUBMED](#) | [CROSSREF](#)