


## Endoscopic ultrasound-guided drainage of a liver abscess with a self-expandable metal stent as rescue therapy after plastic stent misdeployment


OPEN  
ACCESS



► **Fig. 1** Computed tomography scan showing a 5-cm abscess in liver segment II.



► **Fig. 2** Radiological view of a plastic biliary stent into the right duct and an endoscopic ultrasound (EUS)-guided hepaticogastrostomy into the left duct. Also, the liver abscess is outlined after contrast injection.




► **Fig. 3** Radiological view of the self-expandable metal stent being deployed from the liver abscess containing the double-pigtail stent. The previously placed EUS-guided hepaticogastrostomy can be observed in the lower right quadrant.

Endoscopic ultrasound (EUS) is an alternative to percutaneous drainage of abdominal abscesses [1]. Percutaneous abscess drainage may be challenging in poorly accessible locations [2].

A 53-year-old woman underwent palliative biliary drainage for a Bismuth IIIb hilar cholangiocarcinoma. A transpapillary plastic biliary stent was placed by endoscopic retrograde cholangiopancreatography (ERCP) into the right hepatic duct and EUS-guided hepaticogastrostomy performed with a metal stent into the left hepatic duct. Four weeks later, a 5-cm subphrenic abscess was noted in liver segment II (► **Fig. 1**).

The abscess location was deemed unfavorable for percutaneous drainage. An EUS-guided approach was suggested instead. The abscess was imaged under linear EUS and punctured with a 19-G needle from the distal esophagus. Serial dilation with a 6F cystotome and 4-mm balloon dilation was performed (► **Fig. 2**). A 7-Fr 5-cm double-pigtail stent (DPS) was then inadvertently deployed fully within the abscess (► **Fig. 3**). A covered biliary self-expandable metal stent (SEMS) was placed across the tract from the gastroesophageal junction just below the Z line into the abscess, balloon-dilated to 10 mm, and anchored to the esophageal wall with a hemostatic clip (► **Fig. 4**, ► **Fig. 5**). A 0.035-inch guidewire was coiled within the abscess. The echoendoscope was removed over the wire. An ultra-slim gastroscope was carefully advanced over the wire through the SEMS into the abscess. The DPS was grasped with a 5-F tripod forceps under endoscopic view and repositioned into the stomach under gentle traction (► **Video 1**).

A computed tomography (CT) scan performed 2 weeks later confirmed abscess resolution with in-situ SEMS and coaxial DPS. Both stents were removed 1 week later using a standard gastroscope.




► **Fig. 4** Endoscopic ultrasound view of the misdeployed double-pigtail stent inside the liver abscess. Note the close proximity to the left ventricle.



► **Fig. 5** Endoscopic view of the self-expandable metal stent in the distal esophagus clipped to the mucosa.

Drainage of high-grade hilar cholangiocarcinoma remains challenging. ERCP with transpapillary biliary stenting combined with left-sided EUS-guided hepaticogastrostomy appears promising [3–4]. Misdeployment of a DPS within an acute collection is a potentially serious adverse event [5]. As in other related scenarios, placement of a fully covered SEMS bridged the puncture tract. This allowed transluminal access into the abscess similar to that provided by natural orifice transluminal endoscopic surgery (NOTES), and eventually DPS reposition-



**Video 1** Radiological view of the insertion through the self-expandable metal stent of an ultra-thin gastroscope with a tripod grasping forceps repositioning the previously mis-deployed double-pigtail stent.


ing and successful transluminal abscess drainage.

Endoscopy\_UCTN\_Code\_TTT\_1AS\_2AD

### Competing interests

Dr. Manuel Perez-Miranda is a consultant for Boston Scientific, Olympus, Medtronic and M.I.Tech.

### The authors

**Esteban Fuentes-Valenzuela, Beatriz Burgueño Gomez, Carlos Chavarria, Ramon Sanchez-Ocana, Carlos de la Serna-Higuera, Manuel Perez-Miranda** 

Gastroenterology Department, Hospital Universitario Rio Hortega, Valladolid, Spain

### Corresponding author

**Manuel Perez-Miranda, MD, PhD**

Gastroenterology Department, Hospital Universitario Rio Hortega, Calle Dulzaina 2, 47012 Valladolid, Spain  
mperezmiranda@saludcastillayleon.es

### References

- [1] Carbajo AY, Brunie Vegas FJ, García-Alonso FJ et al. Retrospective cohort study comparing endoscopic ultrasound-guided and percutaneous drainage of upper abdominal abscesses. *Dig Endosc* 2019; 31: 431–438. doi:10.1111/DEN.13342
- [2] Chin YK, Asokkumar R. Endoscopic ultrasound-guided drainage of difficult-to-access liver abscesses. *SAGE Open Med* 2020; 8: 205031212092127. doi:10.1177/2050312120921273
- [3] Sundaram S, Dhir V. EUS-guided biliary drainage for malignant hilar biliary obstruction: A concise review. *Endosc Ultrasound* 2021; 10: 154–160. doi:10.4103/EUS-D-21-00004
- [4] Kongkam P, Orprayoon T, Boonmee C et al. ERCP plus endoscopic ultrasound-guided biliary drainage versus percutaneous transhepatic biliary drainage for malignant hilar biliary obstruction: A multicenter observational open-label study. *Endoscopy* 2021; 53: 55–62. doi:10.1055/A-1195-8197
- [5] Vinay Chandrasekhara A, Barthet M, Devière J et al. Safety and efficacy of lumen-apposing metal stents versus plastic stents to treat walled-off pancreatic necrosis: Systematic review and meta-analysis. *Endosc Int Open* 2020; 8: E1639–E1653. doi:10.1055/a-1243-0092

### Bibliography

*Endoscopy* 2023; 55: E328–E329

DOI 10.1055/a-1974-9558

ISSN 0013-726X

published online 14.12.2022

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



### ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is an open access online section, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>