



Olfactory Dysfunction is Associated with More Severe Clinical Course in COVID-19

Azin Tabari¹ · Goli Golpayegani¹ · Azadeh Tabari² · Babak Saedi¹ · Aydin Mahdkhah¹ · Amin Amali¹ · Saber Jazinizadeh¹ · Leyla Sahebi³ · Negin Saffarzadeh⁴ · Seyed Mousa Sadrehosseini¹

Received: 21 October 2020 / Accepted: 2 March 2021
© Association of Otolaryngologists of India 2021

Abstract To perform a quantitative olfactory test in positive COVID19 RT-PCR admitted patients and asymptomatic ones, to evaluate the association between hyposmia and disease severity. This is a Cross sectional study. Ninety-one patients including 68 inpatients and 23 asymptomatic healthcare workers with positive COVID-19 RT-PCRs. Methods: Demographics and clinical characteristics were collected. Iran Smell Identification Test (IR-SIT), a highly accurate 6-odorant test was used to evaluate

the reliability of self-reported hyposmia and determine the correlation of the measured olfactory dysfunction with disease severity. Twenty-two of 91 patients (24%) reported hyposmia, while 41/91 (45%) patients had measurable olfactory dysfunction (IR-SIT score 1–4, $p < 0.05$). Mean age of the 68 inpatients and 23 asymptomatic patients were 43.97 ± 16.13 years; M:F 43:25, and 43.87 ± 12.76 years; M:F 8:15 respectively. Of 68 patients, 20 were graded as severe, and 48/68 had mild course of disease. IR-

✉ Goli Golpayegani
golpayegani.goli@gmail.com

Azin Tabari
atabari@sina.tums.ac.ir

Azadeh Tabari
atabari@mgh.harvard.edu

Babak Saedi
saedi@tums.ac.ir

Aydin Mahdkhah
aydinmahdkhah@gmail.com

Amin Amali
aminamali@yahoo.com

Saber Jazinizadeh
Saber.jazini@gmail.com

Leyla Sahebi
sahebileila@yahoo.com

Negin Saffarzadeh
neginsaffarzade68@gmail.com

Seyed Mousa Sadrehosseini
sadrehosseini@gmail.com

² Department of Radiology, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA

³ Family Health Research Institute, Maternal-Fetal and Neonatal Research Center, Tehran University of Medical Science, Tehran, Iran

⁴ Department of Medical Genetics, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

¹ Department of Otolaryngology-Head and Neck Surgery, Otorhinolaryngology Research Center, Imam Khomeini Hospital Complex, Tehran University of Medical Sciences, Keshavarz Blvd, Tehran, Iran

SIT detected hyposmia in 80% of patients with severe disease, and 50% with mild disease, respectively. The risk of disease severity was significantly increased for patients with olfactory dysfunction and was detected 4 times higher when compared to patients with mild disease (OR 4, 95% CI: 1.166–13.728, $p = 0.028$). Olfactory Dysfunction was present in 80% of patients with severe course. The risk of disease severity is significantly increased with olfactory dysfunction in admitted patients.

Keywords COVID-19 · SARS-CoV-2 · Hyposmia · Olfactory dysfunction · Olfactory test

Introduction

There is rapidly accumulating anecdotal evidence that anosmia (loss of smell) is frequently reported symptom associated with the COVID-19 pandemic. Previously described human strains of coronavirus have also been demonstrated to invade the central nervous system and propagate from within the olfactory bulb [1–3]. It is therefore perhaps no surprise that alterations in sense of smell would be reported by patients with COVID-19. Multiple reports are surfacing from countries around the world including Italy, Iran and Spain showing the sudden surge in the number of patients presenting with anosmia [4–10]. COVID-19 is now recognized to be highly heterogeneous in severity—ranging from asymptomatic (or symptoms too mild to be noticed) to severe acute respiratory distress syndrome and death—and in symptomatology, which extends to include otolaryngologic symptoms. Patients with the well-known and characteristic symptoms of COVID-19 may be readily identified (by themselves or others). There have been reports that olfactory dysfunction can be the primary or the only symptom, therefore, patients experiencing nonclassical symptoms of COVID-19 may be missed [9–12]. In this study we have investigated the reliability of subjective reports of smell loss and the correlation of olfactory dysfunction with severity of COVID-19.

Methods

The study protocol was approved by the local ethics committee and the National Ministry of Health (license number IR.TUMS.VCR.REC.1399.305). Detailed information about the study was given to the participants, and an informed written consent was obtained from each participant.

Subjects

This study included 91 individuals from June 2020 to July 2020. All the subjects were divided into 2 groups: 1-patients who had confirmed COVID-19 with RT-PCR of nasopharyngeal and throat, done based on the World Health Organization (WHO) recommendation [13], and were admitted to a tertiary COVID referral hospital ($n = 68$), 2- asymptomatic health care workers who, were screened as part of hospital protocol, and were positive for COVID-19 RT-PCR ($n = 23$). The inclusion criteria comprised: hemodynamic stability, age > 18 years, understand Farsi and ability to follow the testing instructions. Patients with history of head trauma, skull base fracture, neoplasms of nose, sinus or brain, exposure to irritant chemical cleansers, prior olfactory dysfunction, prior neurological disorders that can affect olfactory function, nasal polyposis and acute sinusitis were excluded.

Clinical Severity Classification of COVID-19

Patients

COVID-19 detection protocol of National Ministry of Health was used which includes CT imaging (64 slice high speed CT, GH Healthcare) with positive findings. Moreover, the diagnosis of COVID-19 disease was confirmed by quantitative detection of SARS-CoV-2 RNA using RT-PCR (viral nucleic acid extraction kit, RBCBioscience Inc., Brussels, Belgium) of the patients' nasopharyngeal wash/aspirate or nasal aspirate. The severity of COVID-19 was categorized as mild ($SpO_2 > 90\%$ with or without risk factors), and severe (patients with $SPO_2 < 90\%$ at initial presentation or progressive disease course during hospital stay; drop in SPO_2 , ICU admission, intubation, acute respiratory distress syndrome). Age > 55 years old, pre-existing pulmonary disease, chronic kidney disease, diabetes with $A1c > 7.6\%$, history of hypertension or cardiovascular disease or transplant or immunosuppression or HIV were considered as epidemiological risk factors. Critical vital signs were respiratory rate > 24 breaths/min, heart rate > 125 beats/min and $SpO_2 < 90\%$ on ambient air. In lab findings, D-dimer > 1000 ng/mL, CPK > twice upper limit of normal, CRP > 100, LDH > 245 U/L, elevated troponin, admission absolute lymphocyte count < 0.8 and ferritin > 300 ug/L.

Subjective Evaluation of Loss of Sense of Smell

All subjects were asked to fill out a questionnaire including medical conditions, medications, prior olfactory problems, smoking habit, suspicious recent close contacts, current symptoms, duration of symptoms and contact information.

Objective Evaluation of Olfactory Function

The olfactory function was assessed using the quick IR-SIT [14]. IR-SIT is the Iranian verified version of UPSIT, using six odors including banana, rosewater, cinnamon, garlic, mint, and melon, with high accuracy in differentiating anosmia, hyposmia and normosmia. Six odors were presented to the patient as a sticker. Stickers were scratched by the physician and patients were asked to smell them from 2-cm distance. Patients had to select the correct odor name from the 4-alternative test sheet. Patients with a total score of 5 and 6 (number of odorants diagnosed correctly) were considered normosmic, while hyposmic patients were scored 1–4. Anosmia was defined as 0.

Assessment of Olfactory Dysfunction and Severity of the Disease

All patients were divided into two groups with normal sense of smell and hyposmia using IR-SIT testing. We then compared the 2 groups in terms of demographic, comorbidity variables and severity of the disease.

Statistical Analysis

Chi-square test was used for comparison of categorical variables. Independent sample t-test was used for comparison of means. The association of COVID-19 severity and hyposmia was determined using logistic regression. The agreement between subjective hyposmia and objective test was performed by Phi test. *P* values less than 0.05 were considered statistically significant. Statistical analyses were performed using SPSS 24.

Results

This cohort included 91 individuals (51 males, 40 females; mean age, 43.95 ± 15.28 years old; age range, 15–82 years old) from June 2020 to July 2020. Subjects' characteristics and demographics are summarized in Table 1. The COVID-19 symptomatic patients' most common presenting symptoms were cough ($n = 45$), dyspnea ($n = 40$), fever ($n = 37$), myalgia ($n = 33$) and sore throat ($n = 17$). As shown in Table 1, inpatients tend to have more comorbidities when compared to asymptomatic individuals ($p < 0.001$). Inpatients were significantly more hypertensive comparing to asymptomatic individuals ($p = 0.03$). In addition, COVID-19 positive patients with negative smoking history, no diabetes mellites and no hypertension were more likely to remain asymptomatic when compared to the inpatient group (Table 1).

10/23(43.5%) of asymptomatic individuals being positive screened for COVID-19 has reported hyposmia in history taking but their objective test results were normal. Interestingly, only 1/23 (4%) of asymptomatic subjects were considered hyposmic when performed the objective olfactory test.

All 68 inpatients were divided into two groups with normal sense of smell ($n = 28$) and hyposmia ($n = 40$) using objective olfactory testing (Table 2). Fifty-eight percent ($n = 40$) of inpatients with COVID-19 had measured olfactory dysfunction and were scored 1–4 (hyposmia) on IR-SIT test; 10(25.0%) had severe smell dysfunction (IR-SIT score = 1–2) and 30(75.0%) patients had mild to moderate olfactory dysfunction (IR-SIT score = 3–4), while there were no pure anosmic subjects. The IR-SIT testing revealed that the admitted patients exhibited marked olfactory dysfunction compared to asymptomatic individuals (40/68 and 1/23 respectively, $p = 0.001$).

Generally, 34 cases of inpatients (60.7%) who did not report hyposmia, were classified as some degrees of olfactory dysfunction recognized by IR-SIT test. Inpatients with hyposmia under-reported their loss of sense of smell by 70% (12/68 self-report vs. 40/68 with objective olfactory testing). Therefore, there is no significant agreement between subjective hyposmia and objective test results among inpatients (Phi Test value = -0.083, p -value = 0.434).

Of 68 inpatients, 20 and 48 patients had severe and mild disease, based on the criteria such as SPO₂, vital signs, risk factors and their respiratory progressive disease course during their hospital stay. It is important to point out that 80% (16/20) of patients with more progressive disease had objective olfactory dysfunction whereas, only 50% (24/48) of mild patients had objective hyposmia. The risk of disease severity was significantly increased in patients with olfactory dysfunction and was detected at 4 times higher rate, when compared to patients with mild disease (OR 4, 95% CI: 1.166–13.728, $p = 0.028$).

Discussion

This study quantitatively evaluated olfaction in a sizable cohort of patients diagnosed with the SARS-CoV-2 virus infection. By employing a well-validated 6-item smell test [14], COVID-19 patients were able to be classified into distinct categories of olfactory dysfunction, with 40 of 68 (60%) admitted patients and 1 of 23 (1.7%) asymptomatic healthcare workers with confirmed COVID-19, exhibiting hyposmia. In the present study, only (13/91) 14% of the subjects were aware of their olfactory deficit before testing, which is a lower percentage compared to previous reports

Table 1 Characteristics and demographics of the 2 groups

	Inpatients (68)	Asymptomatic individuals (23)	<i>P</i> value
Mean Age (years) ± SD	43.97 ± 16.13	43.87 ± 12.76	0.978
Sex; n (%)			
Female (40)	25(62.5%)	15(37.5%)	0.011
Male (51)	43(84.3%)	8(15.6%)	
Smoking history; n (%)			
Yes	8(88.8%)	1(11.2%)	0.440*
No	60(73.1%)	22(26.8%)	
Smell complaints	12(54.5%)	10(45.4%)	
Smell dysfunction(IR-SIT)	40(97.6%)	1(2.4%)	< 0.001
Comorbidities	41(93.1%)	3(6.9%)	< 0.001
Diabetes	13(92.8%)	1(7.2%)	0.108*
Hypertension	12(100%)	0	0033*
Cancer	2(100%)	0	1*
Chronic kidney disease	1(100%)	0	1*
Chronic pulmonary disease	3(75%)	1(25%)	1*

**P* value based on fisher exact test

with interviews from inpatients or online surveys (34% and 60%) [15, 16]. Since the first published studies about olfactory dysfunction surge during COVID-19 pandemic, different studies have investigated its prevalence through population based online surveys (60%) [16], self-report or questionnaire in outpatient (47%) [17] or inpatients (5.1%, 33.9%, 85.6%) [15, 18, 19] and validated smell test (98%) [20]. One of the main reasons resulting in the gap between self-report rate and quantified smell assessment is that in general, anosmia is more noticeable compared to hyposmia. Therefore, many confirmed patients may not report hyposmia. The difference between self-report rate and quantified smell assessment proved the importance of performing a quantitative olfactory dysfunction test for all the patients regardless of their complaint.

Ninety one percent of admitted patients did not present with nasal obstruction or rhinorrhea. This leads to suspecting another reason for hyposmia than mechanical nasal obstruction. SARS-COV-2 may enter the CNS through retrograde neuronal route that is supported by the fact that some patients develop hyposmia [21, 22]. This phenomenon may also be indicative of the immunosuppression in COVID-19 patients with CNS symptoms, especially in the severe subgroup [22]. Hyposmia showed significant correlation with severity of the disease in the inpatient group. We observed that 80% of patients with more severe course of the disease presented with olfactory dysfunction. This is consistent with a previous study that pointed a correlation between neurologic symptoms such as anosmia and COVID-19 disease severity in inpatient setting

[21, 23–27]. Occurrence of olfactory dysfunction during other viral infections has strong correlation with disease severity [18, 19, 25–28]. Hyposmia as a neurologic symptom may need for further close follow-up in inpatients. In addition, neuro-invasion may increase the risk of multiple sclerosis and Parkinson in future [22].

The strengths of this study were using quantitative test to evaluate olfactory function which is more sensitive instead of relying on self-reports or subjective questionnaires. Moreover, we included both symptomatic and asymptomatic patients which makes it a unique study population. To our knowledge, this is the first study designed to evaluate if loss of smell has a prognostic significance on the disease severity.

This study has some limitations. First, the low number of subjects due to the unknown nature of COVID-19 in the first months of pandemic. The low number of cases were directly the result of safety issue concerns in critical care. We included the data of the admitted patients, therefore the symptomatic ambulatory patients who showed milder disease were excluded since they mostly preferred to stay at home during quarantine. Second, examiners were not blinded to the diagnosis when performing the olfactory test. Lastly, we were unable to perform the objective test on the more severely ill such as ICU patients, due to safety precautions.

Based on our findings, objective smell testing enables confirmatory diagnosis in the absence of other symptoms and helps to predict the clinical course of COVID-19 in terms of progression and severity. We would also like to

Table 2 Baseline characteristics and Objective test results of the admitted patient group

	Normal sense of smell (28)	Hyposmia (40)	P-value
Age; Mean (SD)	43.32 ± 14.74	44.43 ± 17.20	0.784
Sex (F:M)	13:15	12:28	0.167
Positive smoking history (%)	2(25%)	6(75%)	0.455**
Comorbidities	15(36.5%)	26(63.4%)	0.343
Diabetes	4(30.8%)	9(69.2%)	0.397
Hypertension	2(16.7%)	10(83.3%)	0.084**
Symptoms			
Cough	18(40%)	27(60%)	0.873
Dyspnea	17(42.5%)	23(57.5%)	0.791
Fever	18(48.6%)	19(51.4%)	0.171
Myalgia	16(48.5%)	17(51.5%)	0.187
Sore throat	7(41.2%)	10(58.8%)	1
Dysgeusia	8(61.5%)	5(38.5%)	0.097
Hyposmia	6(50%)	6(50%)	0.494
Sneezing	1(20%)	4(80%)	0.642**
Rhinorrhea	1(25%)	3(75%)	0.498
Nasal obstruction	0	6(100%)	0.032
Diarrhea	10(47.6%)	11(52.4%)	0.471
Nausea/vomiting	11(55%)	9(45%)	0.135
Signs			
SpO ₂			0.077
< 90%	8(28.6%)	20(71.4%)	
90%	20(50%)	20(50%)	
CRP			0.961
100	7(41.2%)	10(58.8%)	
< 100	18(41.9%)	25(58.1%)	
Lymph count			0.155
< 0.8	9(31%)	20(69%)	
0.8 <	17(48.6%)	18(51.4%)	

*p-value based on t-test

**p-value based on fisher exact test

SpO₂ = Oxygen saturation; CRP = C-reactive protein; Lymph count = absolute lymphocyte count

draw attention to the importance of objective assessment of olfactory dysfunction. It is important to perform the testing in early stages, regardless of patient complaint of loss of sense of smell. It is time for public health workers to recognize the link between COVID-19 severity and smell disturbances.

Key messages

COVID-19-related olfactory dysfunction is a known description in the medical literature. Overall, 60% of the patients with COVID-19 present with hyposmia. Olfactory dysfunction is significantly associated with more

progressive course of the disease in current study. The outcome is consistent in 80% of patients with severe disease. This notion may need to be communicated to the medical community.

Funding This research received no specific grant from funding agency in the public, commercial, or non-for-profit sectors.

References

1. Dubé M, Le Coupanec A, Wong AH, Rini JM, Desforges M, Talbot PJ (2018) Axonal transport enables neuron-to-neuron propagation of human coronavirus OC43. *J Virol* 92 (17)

2. Desforges M, Le Coupanec A, Stodola JK, Meessen-Pinard M, Talbot PJ (2014) Human coronaviruses: viral and cellular factors involved in neuroinvasiveness and neuropathogenesis. *Virus Res* 194:145–158
3. Morris M, Zohrabian V (2020) Neuroradiologists, be mindful of the neuroinvasive potential of COVID-19. *Am J Neuroradiol* 41:E37–E39. <https://doi.org/10.3174/ajnr.A6551>
4. Cheng Q, Yang Y, Gao J (2020) Infectivity of human coronavirus in the brain. *Biomedicine* 56:102799
5. Hopkins C, Surda P, Kumar N (2020) Presentation of new onset anosmia during the COVID-19 pandemic. *Rhinology* 58(3):295–298
6. Acter T, Uddin N, Das J, Akhter A, Choudhury TR, Kim S (2020) Evolution of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) as coronavirus disease 2019 (COVID-19) pandemic: a global health emergency. *Sci Total Environ* 730:138996
7. Heidari F, Karimi E, Firouzifar M, Khamushian P, Ansari R, Ardehali MM, Heidari F (2020) Anosmia as a prominent symptom of COVID-19 infection. *Rhinology* 58(3):302–303
8. Krajewska J, Krajewski W, Zub K, Zatoński T (2020) COVID-19 in otolaryngologist practice: a review of current knowledge. *Eur Arch Oto-Rhino-Laryngol* 277(7):1885–1897
9. Gane SB, Kelly C, Hopkins C (2020) Isolated sudden onset anosmia in COVID-19 infection. A novel syndrome. *Rhinology* 58(3):299–301
10. Cao Y-C, Deng Q-X, Dai S-X (2020) Remdesivir for severe acute respiratory syndrome coronavirus 2 causing COVID-19: An evaluation of the evidence. *Travel Med Infect Dis* 35:101647
11. Nicola M, O'Neill N, Sohrabi C, Khan M, Agha M, Agha R (2020) Evidence based management guideline for the COVID-19 pandemic-review article. *Int J Surg* 77:206–216. <https://doi.org/10.1016/j.ijsu.2020.04.001>
12. Gilani S, Roditi R, Naraghi M (2020) COVID-19 and anosmia in Tehran Iran. *Med Hypotheses* 141:109757
13. WHO (2020) Laboratory testing for coronavirus disease (COVID-19) in suspected human cases: interim guidance, 19 March 2020. World Health Organization
14. Kamrava SK, Jalessi M, Ghalehbaghi S, Amini E, Alizadeh R, Rafiei F, Moosa S, Farhadi M (2020) Validity and reliability of persian smell identification test. *Iranian J Otorhinolaryngol* 32(109):65
15. Giacomelli A, Pezzati L, Conti F, Bernacchia D, Siano M, Oreni L, Rusconi S, Gervasoni C, Ridolfo AL, Rizzardini G (2020) Self-reported olfactory and taste disorders in patients with severe acute respiratory coronavirus 2 infection: a cross-sectional study. *Clin Infect Dis* 71:889–890. <https://doi.org/10.1093/cid/ciaa330>
16. Bagheri SH, Asghari A, Farhadi M, Shamshiri AR, Kabir A, Kamrava SK, Jalessi M, Mohebbi A, Alizadeh R, Honarmand AA (2020) Coincidence of COVID-19 epidemic and olfactory dysfunction outbreak in Iran. *Med J Islamic Repub Iran* 34:62
17. Spinato G, Fabbris C, Polesel J, Cazzador D, Borsetto D, Hopkins C, Boscolo-Rizzo P (2020) Alterations in smell or taste in mildly symptomatic outpatients with SARS-CoV-2 infection. *JAMA* 323:2089. <https://doi.org/10.1001/jama.2020.6771>
18. Besser G, Liu DT, Renner B, Hummel T, Mueller CA (2020) Reversible obstruction of the olfactory cleft: impact on olfactory perception and nasal patency. In: *International forum of allergy & rhinology*. Wiley Online Library 6:713–718
19. Lechien JR, Chiesa-Estomba CM, De Siati DR, Horoi M, Le Bon SD, Rodriguez A, Dequanter D, Blecic S, El Afia F, Distinguin L (2020) Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch Oto-Rhino-Laryngol* 277:1–11
20. Moein ST, Hashemian SM, Mansourafshar B, Khorram-Tousi A, Tabarsi P, Doty RL (2020) Smell dysfunction: a biomarker for COVID-19. In: *International forum of allergy & rhinology*. Wiley Online Library 8:944–950
21. Mao L, Jin H, Wang M, Hu Y, Chen S, He Q, Chang J, Hong C, Zhou Y, Wang D (2020) Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurol* 77(6):683–690
22. Das G, Mukherjee N, Ghosh S (2020) Neurological insights of COVID-19 pandemic. *ACS Chem Neurosci* 11(9):1206–1209
23. Montalvan V, Lee J, Bueso T, De Toledo J, Rivas K (2020) Neurological manifestations of COVID-19 and other coronavirus infections: a systematic review. *Clinical Neurol Neurosurg* 194:105921
24. Román GC, Spencer PS, Reis J, Buguet A, Faris MEA, Katrak SM, Láinez M, Medina MT, Meshram C, Mizusawa H (2020) The neurology of COVID-19 revisited: a proposal from the Environmental Neurology Specialty Group of the World Federation of Neurology to implement international neurological registries. *J Neurol Sci* 414:116884
25. Yan CH, Faraji F, Prajapati DP, Ostrander BT, DeConde AS (2020) Self-reported olfactory loss associates with outpatient clinical course in COVID-19. In: *International forum of allergy & rhinology*. Wiley Online Library 7:821–831
26. Kotfis K, Williams Roberson S, Wilson JE, Dabrowski W, Pun BT, Ely EW (2020) COVID-19: ICU delirium management during SARS-CoV-2 pandemic. *Crit Care* 24:1–9
27. Munhoz RP, Pedrosa JL, Nascimento FA, Almeida SMD, Barsottini OGP, Cardoso FEC, Teive HAG (2020) Neurological complications in patients with SARS-CoV-2 infection: a systematic review. *Arq Neuropsiquiatr* 78(5):290–300
28. Trotier D, Bensimon JL, Herman P, Tran Ba Huy P, Døving KB, Eloit C (2007) Inflammatory obstruction of the olfactory clefts and olfactory loss in humans: a new syndrome? *Chem Senses* 32(3):285–292

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.