



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Behavioral considerations and impact on personal protective equipment use: Early lessons from the coronavirus (COVID-19) pandemic



Jonathan Kantor, MD, MSCE

*Philadelphia, Pennsylvania and Saint Augustine, Florida*

In this issue of the JAAD, Lan et al<sup>1</sup> provide preliminary data on the prevalence of cutaneous irritation associated with using personal protective equipment (PPE) by health care workers (HCWs) during the early months of the coronavirus 2019 (COVID-19) outbreak in Wuhan, China. The authors highlight the high prevalence of cutaneous irritation associated with N95 mask and goggle use—echoing previous reports highlighting cutaneous adverse events associated with PPE use for severe acute respiratory syndrome<sup>2</sup>—and their data suggest that increased duration of PPE use may be associated with an increased risk of cutaneous irritation.

Given that most cases of PPE-associated skin irritation are clinically mild, why should dermatologists and frontline HCWs worry about mild, self-limited, pressure- and abrasion-induced injuries?

Although HCWs and the general public often focus on direct droplet spread to mucosa as a means of infection, research has highlighted the role of behaviors such as face touching in the spread of viral disease.<sup>3-5</sup> Indeed, the role of face touching and surface contact has been recognized as a driver of viral transmission for decades,<sup>6</sup> and a meta-analysis demonstrated that hand washing was associated with a 24% reduction in viral transmission, suggesting that hand-to-face contact may play a substantial role in infection.

The presence of even mild abrasions on the central face may increase the likelihood of face touching while not using PPE or inadvertent PPE protocol breaches, such as mask touching or adjustment, in an unconscious effort to relieve a source of irritation.

The importance of adherence to strict PPE protocols is paramount, as the infection of 2 HCWs during the Ebola outbreak was tied to possible PPE protocol adherence breaches.<sup>7</sup> Inappropriate doffing of PPE in HCWs is common: 1 study found that 26% of HCWs inappropriately touched the front of their mask while doffing, and approximately one-half touched a potentially contaminated PPE surface with an ungloved hand.<sup>7</sup>

What can be done to limit these risks? First, educating HCWs to expect some mild skin irritation may be helpful.

Second, if topical agents are used to reduce irritation, this could be considered a high-risk activity, and introducing topical ointment to the skin should be done with great care; for example, using a sterile cotton-tipped applicator and a single-use petroleum jelly pack before possible exposure.

Third, those with a history of sensitivity may wish to explore other options aside from an N95 respirator and goggles, such as a full-face respirator or using a full-face shield rather than goggles. Given their added cost and decreased availability, however, these may not be feasible options.

Finally, Lan et al mention the possibility of prophylactic dressing use to mitigate the risk of skin-related complications of PPE use, but this approach has not been studied, and—more importantly—the potential impact of such dressings on PPE efficacy is unexplored. Given the very high stakes associated with adequate and reliable PPE functioning, future studies exploring approaches to mitigate the risk of PPE-induced irritation and potential improvements in PPE design are warranted.

From the Department of Dermatology and Center for Global Health, University of Pennsylvania Perelman School of Medicine, and Florida Center for Dermatology, P.A.

Funding sources: None.

Conflicts of interest: None disclosed.

IRB approval status: Not applicable.

Accepted for publication March 4, 2020.

Reprints not available from the author.

Correspondence to: Jonathan Kantor, MD, MSCE, Florida Center for Dermatology, P.A., PO Box 3044, Saint Augustine, FL 32085.

E-mail: [jonkantor@gmail.com](mailto:jonkantor@gmail.com).

J Am Acad Dermatol 2020;82:1087-8.

0190-9622/\$36.00

© 2020 by the American Academy of Dermatology, Inc.

<https://doi.org/10.1016/j.jaad.2020.03.013>

## REFERENCES

1. Lan J, Song Z, Miao X, et al. Skin damage among health care workers managing coronavirus disease-2019. *J Am Acad Dermatol*. 2020;82:1215-1216.
2. Foo CCI, Goon ATJ, Leow YH, Goh CL. Adverse skin reactions to personal protective equipment against severe acute respiratory syndrome—a descriptive study in Singapore. *Contact Dermatitis*. 2006;55(5):291-294.
3. Kwok YLA, Gralton J, McLaws M-L. Face touching: a frequent habit that has implications for hand hygiene. *Am J Infect Control*. 2015;43(2):112-114.
4. Bertsch RA. Avoiding upper respiratory tract infections by not touching the face. *Arch Intern Med*. 2010;170(9):833-834.
5. Nicas M, Best D. A study quantifying the hand-to-face contact rate and its potential application to predicting respiratory tract infection. *J Occup Environ Hyg*. 2008;5(6):347-352.
6. Hendley JO, Wenzel RP, Gwaltney JM. Transmission of rhinovirus colds by self-inoculation. *N Engl J Med*. 1973;288(26):1361.
7. Phan LT, Maita D, Mortiz DC, et al. Personal protective equipment doffing practices of healthcare workers. *J Occup Environ Hyg*. 2019;16(8):575-581.