


## CORRECTION

# Correction: PID1 alters the antilipolytic action of insulin and increases lipolysis via inhibition of AKT/PKA pathway activation

Chunyan Yin, Wei hua Liu, Yuesheng Liu, Li Wang, Yanfeng Xiao

[Fig 4](#) is incorrect. Please see the correct [Fig 4](#) here.




## OPEN ACCESS

**Citation:** Yin C, Liu Wh, Liu Y, Wang L, Xiao Y (2019) Correction: PID1 alters the antilipolytic action of insulin and increases lipolysis via inhibition of AKT/PKA pathway activation. PLoS ONE 14(6): e0218721. <https://doi.org/10.1371/journal.pone.0218721>

**Published:** June 17, 2019

**Copyright:** © 2019 Yin et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



**Fig 4. Effects of PID1 expression on lipolysis and the phosphorylation of AKT/PDE3B/PKA/HSL signaling molecules and ATGL.** Preadipocytes were subjected to PID1 knockout or upregulation and allowed to differentiate into 3T3-L1 adipocytes; these cells were treated with 1  $\mu$ M isoproterenol and 100 nM insulin for 24 h (in triplicate). (a-b) Immunofluorescence analysis was performed to assess the expression of the PID1 gene in empty vector cells, PID1-overexpressing cells, and control cells. (c) RT-PCR analyses of the mRNA expression of PID1 in empty vector cells, PID1-overexpressing cells, and control cells. (d) Glycerol released into the medium after the upregulation of PID1. (e-f) Protein expression of AKT, PDE3B, PKA, HSL and ATGL in empty vector cells, PID1-overexpressing cells, and control cells. Phosphorylated AKT (p-AKT), phosphorylated PKA (p-PKA), phosphorylated HSL (p-HSL) and phosphorylated ATGL (p-ATGL) expression was normalized to their total protein level as a loading control. (g) The phosphorylated protein/total protein ratios for AKT, PKA, HSL, and ATGL in 3T3-L1 adipocytes after transfection with the PID1 overexpression plasmid. (h) The expression of PDE3B was determined by Western blot after the PID1 overexpression plasmid. (i-j) Protein expression of AKT, PDE3B, PKA, HSL and ATGL after transfection with PID1 shRNA. Phosphorylated AKT (p-AKT), phosphorylated PKA (p-PKA), phosphorylated HSL (p-HSL) and phosphorylated ATGL (p-ATGL) expression was normalized to their total protein level as a loading control. (k) RT-PCR analyses of mRNA after transfection with PID1 shRNA. (l) Glycerol was released into the medium after knockdown of PID1. (m) Phosphorylated protein/total protein ratios for AKT, PKA, HSL, and ATGL after PID1 knockdown. (n) The expression of PDE3B was determined by Western blot after transfection with PID1 shRNA.

<https://doi.org/10.1371/journal.pone.0218721.g001>

## Reference

1. Yin C, Liu Wh, Liu Y, Wang L, Xiao Y (2019) PID1 alters the antilipolytic action of insulin and increases lipolysis via inhibition of AKT/PKA pathway activation. *PLoS ONE* 14(4): e0214606. <https://doi.org/10.1371/journal.pone.0214606> PMID: 30990811