


Article

Effects of Pre-Processing Hot-Water Treatment on Aroma Relevant VOCs of Fresh-Cut Apple Slices Stored in Sugar Syrup

Guido Rux ^{1,2,*}, Efecan Efe ^{1,2}, Christian Ulrichs ², Susanne Huyskens-Keil ², Karin Hassenberg ¹ and Werner B. Herppich ¹ 

¹ Department of Horticultural Engineering, Leibniz Institute for Agricultural Engineering and Bioeconomy (ATB), 14469 Potsdam, Germany; efeefeca@cms.hu-berlin.de (E.E.); khassenberg@atb-potsdam.de (K.H.); wherppich@atb-potsdam.de (W.B.H.)

² Research Group Quality Dynamics/Postharvest Physiology, Division Urban Plant Ecophysiology, Humboldt-Universität zu Berlin, 14195 Berlin, Germany; christian.ulrichs@hu-berlin.de (C.U.); susanne.huyskens@hu-berlin.de (S.H.-K.)

* Correspondence: grux@atb-potsdam.de; Tel.: +49-331-5599-920

Received: 16 December 2019; Accepted: 8 January 2020; Published: 10 January 2020



Abstract: In practice, fresh-cut fruit and fruit salads are currently stored submerged in sugar syrup (approx. 20%) to prevent browning, to slow down physiological processes and to extend shelf life. To minimize browning and microbial spoilage, slices may also be dipped in a citric acid/ascorbic acid solution for 5 min before storage in sugar syrup. To prevent the use of chemicals in organic production, short-term (30 s) hot-water treatment (sHWT) may be an alternative for gentle sanitation. Currently, profound knowledge on the impact of both sugar solution and sHWT on aroma and physiological properties of immersed fresh-cuts is lacking. Aroma is a very important aspect of fruit quality and generated by a great variety of volatile organic compounds (VOCs). Thus, potential interactive effects of sHWT and sugar syrup storage on quality of fresh-cut apple slices were evaluated, focusing on processing-induced changes in VOCs profiles. Intact ‘Braeburn’ apples were sHW-treated at 55 °C and 65 °C for 30 s, sliced, partially treated with a commercial ascorbic/citric acid solution and slices stored in sugar syrup at 4 °C up to 13 days. Volatile emission, respiration and ethylene release were measured on storage days 5, 10 and 13. The impact of sHWT on VOCs was low while immersion and storage in sugar syrup had a much higher influence on aroma. sHWT did not negatively affect aroma quality of products and may replace acid dipping.

Keywords: minimal processing; sugar syrup immersion; volatile organic compounds; chemical prevention; ready-to-eat fruit salads

1. Introduction

Fresh-cut processing induces a catena of physiological responses [1,2] finally resulting in the loss of quality and aroma and pronouncedly shortens the storage life of fresh-cut produce [3]. In current practice, fresh-cut fruit for fruit salads are often stored in sugar syrup (ca. 20%), especially for use by bulk purchasers [4]. This storage method may extend product shelf life by preventing enzymatic and oxidative browning and transpiration, slows down respiration, ethylene metabolism and other physiological processes [5–7]. In this context, however, microbial spoilage is the main factor limiting shelf life [7]. It is therefore very important to remove the microorganisms adherent to the fruit skin [8] before processing.

In practice, e.g., apple slices are dipped in a mixture of citric and ascorbic acid solutions for sanitation purposes. To prevent the consumption of these chemicals especially in organic production, gentle physical sanitation methods are demanded.

As such, hot-water treatments (HWT) in the temperature range of 40–80 °C were shown to effectively reduce microbial contamination, and are relatively inexpensive and easy to use [9]. In addition, HWT maintains storage quality of fruits [10–12]. Since they are chemical-free, particularly short-term (15–60 s) hot-water treatments (sHWT) are suitable for organic production [13,14]. Besides earlier studies on the impacts of HWT on structure and function of fruit epidermal tissue [15,16], only recently, the effects of sHWT on surface tissue, heat transfer dynamics and suitability for pre-processing of intact apples for fresh-cut salads were investigated in detail [12,17]. Furthermore, the implications of sHWT on important quality parameters such as tissue browning, tissue strength and on microbial loads of apple slices immersed in sugar syrup have been studied under semi-practical conditions [18]. On the effects of sugar syrup storage on fresh-cut fruit quality attributes only very few studies are available [7,19].

In contrast to analyses of sHWT impacts on visual and internal quality attributes of fresh-cut fruit [12,17,18], studies on potential variations in fruit aroma, a very important aspect of fruit quality sensation [2,20], are completely lacking. Aroma is generated by a great variety of permanent or secondary VOCs [21], synthesized via numerous biosynthetic pathways, which, in turn, are regulated by a great variety of enzymes and substrates [2,22,23]. Moreover, both respiration and ethylene biosynthesis are also involved in VOCs evolution [24]. In addition, microbial growth may, directly or indirectly, negatively affect product aroma [25,26]. Short-term-HWT and sugar syrup immersions may influence all of these processes. Knowledge on the impact of both sugar solution and sHWT on aroma development and physiological properties of immersed fresh-cuts is lacking.

Thus, the present study focused on the evaluation of the potential effects of sHWT (30 s, at 55 °C or 65 °C) on the aroma-related quality of 'Braeburn' apple slices stored in sugar syrup under strictly simulated practical condition. Some parts of the samples were also pre-treated with a commercial ascorbic/citric acid solution to test the potential synergistic effects of this treatment on aroma. During a 13 days-storage at 4 °C, respiration, ethylene emission and, for the first time, the processing-induced direct and indirect changes in VOC profiles were measured at defined intervals. This will enable comprehension of the respective quality-related physiological processes, to effectively select the optimal process conditions and to verify whether sHWT can safely replace the use of chemicals in processing of ecologically produced fresh-cut fruit salads.

2. Materials and Methods

2.1. Material

Fresh mature 'Braeburn' apples (*Malus domestica* Borkh.) were obtained from a commercial fresh-cut salad producer. At the Department of Horticultural Engineering (Leibniz Institute for Agricultural Engineering and Bioeconomy, Potsdam, Germany), the apples were stored at 4 °C and 95% relative humidity for up to 3 days until the start of the experiments. Undamaged apples of uniform size (mean fresh mass: 150.7 ± 5.1 g and mean dry matter content: 177 ± 11 g kg⁻¹) were selected.

2.2. Pre-Processing Short-Term Hot-Water Treatment

Before cutting, apples were divided in five batches of 16 fruit each. Apples from the control batch were water-washed at approx. 20 °C. The other samples were hot-water-treated in a GFL 1086 water bath (Gesellschaft für Labortechnik mbH, Burgwedel, Germany) by submerging four apples each for 30 s. Samples of two batches each were hot-water-treated at 55 °C or 65 °C, respectively, according to Kabelitz and Hassenberg [12] and Kabelitz et al. [19]. These authors indicated that HWT at 55 °C for 30 s effectively reduced microbial loads without negatively affecting the external quality of samples, while 65 °C showed to be a negative control treatment in terms of quality maintenance.

2.3. Fresh-Cut Preparation and Sampling

After washing and hot-water treatments, all apples were cut under semi-practical hygienic conditions in a cooling room at 4 °C. For this, apples were at first halved equatorially and then each half-segment was cut into 16 pieces by a Parti apple cutter/corer (Gefu GmbH, Eslohe, Germany). The controls and slices of one of each 55 °C and 65 °C sHWT batches were additionally chemically treated (= chemical prevention, cp) by immersing in ascorbic/citric acid solution (40 g ascorbic and 20 g citric acid solved in 1 L deionized water) for 5 min (Figure 1). Slices of each batch were randomly filled in 12 commercial 840 mL plastic pails. Closely following practice, each pail containing approx. 48 apple slices (approx. 228 g) was filled up with 450 mL sugar syrup (200 g L⁻¹ invert sugar syrup, 72.7%; Hanseatische Zuckerraffinerie GmbH & Co. KG, Hamburg, Germany plus 10 g L⁻¹ OBSTSERVAL HC-2 browning inhibitor (Konserval, Pharmacon Lebensmittelzusätze GmbH, Trittau, Germany). The browning inhibitor contained ascorbic acid, sodium ascorbate and citric acid. All pails were tightly closed and stored at 4 °C for up to 13 days. Three pails (replicates) of each batch were opened on days 5, 10 (the common maximum shelf-life) and on day 13. For further analyses, 26 apple slices of each pail were removed.




Figure 1. Scheme of the various pre- (short-term hot-water, sHWT, vs. washed) and post-cutting (chemical prevention, cp, vs. no chemical prevention, non-cp) treatments of fresh-cut apples slices stored in sugar syrup at 4 °C for up to 13 days.

2.4. Sampling of Volatile Organic Compounds (VOCs) and Ethylene

To determine VOCs emission, ethylene evolution and respiration rate (RR), 20 apple slices were filled in a glass jar (1 L), which was hermetically closed by a glass lid. VOCs emitted by samples accumulated at 4 °C for 1 h. Then VOCs were 10 min-extracted with 50/30 µm divinylbenzene/carboxen/polydimethylsiloxane (DVB/CAR/PDMS) SPME fiber (stableFlex/SS, Supelco, Bellefonte, PA, USA) via septum in glass lid. After VOCs extraction, 0.5 mL sample air was taken from headspace with a 0.5 mL A-2 Series syringe (VICI Precision Sampling, Baton Rouge, LA, USA) via the same septum to measure ethylene and CO₂ concentrations.

2.5. Analysis of Volatile Organic Compounds

VOCs were analyzed with a GCMS-QP2010 SE gas chromatograph-mass spectrometer (GC-MS; Shimadzu Europa GmbH, Duisburg, Germany) on a DB-WAX (Agilent Technologies, Palo Alto, CA, USA) column (30 m × 0.25 mm × 0.25 µm). SPME fiber was introduced into the GC injector (set to 250 °C and fitted with a SPME liner (0.75 mm × 5.0 mm × 95 mm); Shimadzu GCs; Restek, Bellefonte, PA, USA) for 30 min for VOC desorption and conditioning before next extraction. Helium was used as carrier gas (flow rate: 0.8 mL min⁻¹). The following temperature program was used: 35 °C (hold 1 min), ramp to 110 °C at 5 °C min⁻¹ and ramp to 230 °C at 10 °C min⁻¹ (hold 3 min). The mass spectrometer (MS) detector worked in full scan mode (mass-to-charge ratio: 35–250 m/z) and operated in electron impact mode at 70 eV. Measured chromatogram peaks were integrated based on quantifier and qualifier ions of target compound (Table 1) using MSD Chem-Station E.02.01.1177 (Agilent Technologies, Palo Alto, CA, USA). Specific VOCs were identified (1) by NIST v2.0f library (NIST, Gaithersburg, MD, USA) and (2) by retention index (RI) calculated for each compound according to van Den Dool and Kratz [27].

RIs were determined based on an alkane standard (C7–C30; 1000 $\mu\text{g mL}^{-1}$; Supelco, Bellefonte, PA, USA). For this, 5 μL of alkane standard was injected in a 3.9 L glass jar, extracted with SPME after equilibrium time of 1 d and analyzed as described above. The RIs were compared with those given by van Den Dool and Kratz [27] by using online NIST Chemistry WebBook [28]. The total integrated peak area of each VOC was used to calculate a semi-quantitative VOC concentration expressed in nL L^{-1} based on the external standard 2-methylbutyl 3-methylbutanoate (CAS-number: 2245-77-4). A standard curve of 2-methylbutyl-3-methylbutanoate was created in the concentration range from 5 to 1000 nL L^{-1} .

Table 1. All identified volatile organic compounds (VOCs) emitted by intact, freshly cut and variously treated (sHWT/chemical prevention) fresh-cut apple slices as characterized by their CAS registry numbers, calculated retention indices (RI), previously reported retention indexes ($\text{RI}_{\text{literature}}$), and quantifier and qualifier ions used for peak integration.

Chemical Group	VOCs	CAS-Number	RI	$\text{RI}_{\text{literature}}$	Quantifier (Qualifier) Ions (m z^{-1})
esters (21)	Ethyl acetate	141-78-6	903	863–908	43 (45–61)
	Ethyl propionate	105-37-3	963	939–976	57 (74–75)
	Ethyl 2-methylpropanoate	97-62-1	968	957–969	43 (41–71)
	Propyl acetate	109-60-4	977	952–996	43 (61–73)
	Methyl butyrate	623-42-7	988	969–993	74 (43–71)
	Methyl 2-methylbutyrate	868-57-5	1011	1000–1010	57 (41–88)
	Isobutyl acetate	110-19-0	1014	1000–1031	43 (56–73)
	Ethyl butyrate	105-54-4	1035	1000–1073	71 (43–88)
	Ethyl 2-methylbutanoate	7452-79-1	1050	1022–1073	57 (41–102)
	Butyl acetate	123-86-4	1069	1049–1105	43 (56–73)
	2-methylbutyl acetate	624-41-9	1116	1111–1125	43 (55–70)
	Ethyl valerate	539-82-2	1129	1131–1139	88 (57–85)
	Ethyl 2-butenoate	10544-63-5	1154	1158–1158	69 (41–99)
	Pentyl acetate	628-63-7	1165	1175–1181	43 (55–70)
	Methyl hexanoate	106-70-7	1178	1176–1189	74 (43–99)
	Ethyl hexanoate	123-66-0	1222	1196–1245	71 (43–89)
	Hexyl acetate	142-92-7	1258	1251–1311	43 (56–61)
	2-Hexen-1-yl acetate	10094-40-3	1317	-	
	Hexyl butyrate	2639-63-6	1397	1393–1410	71 (43–89)
	Hexyl 2-methylbutanoate	10032-15-2	1414	1415–1416	
	Hexyl hexanoate	6378-65-0	1610	1596–1599	43 (56–117)
ketones (2)	2-Butanone	78-93-3	908	875–926	43 (57–72)
	1-Penten-3-one	1629-58-9	1020	1019–1056	
alcohols (6)	Ethanol	64-17-5	943	900–955	45 (43–46)
	2-Methyl-1-propanol	78-83-1	1096	1092–1114	43 (41–42)
	1-Butanol	71-36-3	1143	1116–1166	
	2-Methyl-1-butanol	1565-80-6	1200	-	57 (41–56)
	1-Hexanol	111-27-3	1341	1339–1396	
	2-(2-Ethoxyethoxy)-ethanol	111-90-0	1628	1615–1619	45 (59–72)
aldehydes	Hexanal	66-25-1	1077	1048–1120	44 (41–56)
terpenes (2)	D-Limonene	5989-27-5	1186	1176–1238	68 (67–93)
	α -Farnesene	502-61-4	1747	1720–1764	93 (41–69)
benzenic derivatives	Estragole	140-67-0	1724	1624–1661	148 (117–147)

2.6. Quantification of Ethylene and CO_2 Evolution

A small volume of 0.5 mL air was taken from canning glass jar and analyzed with a GC17A gas chromatograph (Shimadzu) to measure ethylene and CO_2 concentrations. The GC was equipped with both FID and TCD, 80/100 Porapak N (Supelco, Bellefonte, PA, USA) column (1.8 m \times 1/8 in \times 2.1 mm) and a 5605PC mole sieve (Alltech GmbH, Unterhaching, Germany). As carrier gas, helium was used (constant flow rate: 1.4 mL min^{-1}), oven temperature was kept constant at 60 $^\circ\text{C}$. Measured peaks were integrated using Class-VP chromatography data system software v4.2 (Shimadzu Europa GmbH, Duisburg, Germany). Rates of ethylene release were expressed in $\mu\text{g kg}^{-1} \text{h}^{-1}$, those of CO_2 production in $\text{mg kg}^{-1} \text{h}^{-1}$.

2.7. Statistical Analysis

For each treatment, three pails were used as replicates for each sample day. VOCs and ethylene emissions were each determined on 20 apple slices from each pail ($n = 3$), while respiration was analyzed on a total of 60 apple slices, i.e., 20 from each pail. Statistical analyses (ANOVA) were carried out with WinSTAT (R. Fitch Software, Staufen, Germany) and results were presented as mean \pm standard deviation (SD). The significance of differences between the calculated means was analyzed by Duncan's multiple range test ($p < 0.05$). The relationships between VOC evolution and sHWT were additionally analyzed by principal components analysis (PCA) using Latentix Ver. 2.00 (Latent5, Copenhagen, Denmark).

3. Results

3.1. Volatile Organic Compounds

In total, 33 different VOCs were identified in fresh-cut apple slices, with esters (21) forming the major group of compounds (Table 1). The emission of individual VOCs was significantly affected by cutting, sHWT and/or storage time (Table 2). In fresh, intact 'Braeburn' apples, 2-methylbutyl acetate (29.0%) and α -Farnesene (15.9%) were the most abundant VOCs. Cutting immediately and significantly increased the emissions of 17 VOCs, while that of only three VOCs significantly decreased. The cumulative VOCs concentration increased sevenfold, particularly concentrations of acetate esters (es2–es8) increased most.

Table 2. Semi-quantitative concentrations (nL L^{-1}) of VOCs, emitted by intact, freshly cut (cut) and variously treated (C: control with chemical prevention; 55: 55 °C sHWT without chemical prevention; 55_cp: 55 °C sHWT with chemical prevention; 65: 65 °C sHWT without chemical prevention; 65_cp: 65 °C sHWT with chemical prevention) fresh-cut apple slices on days 5, 10 and 13 of storage in sugar syrup at 4 °C. VOCs concentrations given resulted from the emissions of 20 apple slices hermetically enclosed in a 1 L-glass jar at 4 °C for 1 h. Given are means ($n = 3$). Different letters indicate significant differences between means ($p < 0.05$). The different colors help to indicate increasing (green) or decreasing (red) emissions of VOCs compared to intact apples (yellow).

VOC	Time	Initial	Time	C	55	55_cp	65	65_cp
Cumulative VOC concentration	intact	87 a	5 d	445 b–d	515 c–e	388 b	415 b–c	425 b–c
	cut	618 e	10 d	364 b	493 c–e	320 b	379 b	354 b
			13 d	385 b	502 d–e	305 b	317 b	369 b
Ethyl acetate es1	intact	0.86 a	5 d	109 c–d	152 e	80.9 b–c	93.3 b–d	83.5 b–c
	cut	1.98 a	10 d	104 b–d	180 f	75.8 b	121 d	83.2 b–c
			13 d	121 d	196 f	82 b–c	106 b–d	105 b–d
Propyl acetate es2	intact	nd	5 d	3.79 g	2.53 d–e	2.45 c–e	1.78 b–c	2.59 d–e
	cut	2.91 e–f	10 d	3.35 f–g	2.27 b–e	2.31 b–e	1.71 b	2.39 b–e
			13 d	3.74 g	2.21 b–e	2.11 b–d	0.39 a	nd
Butyl acetate * es3	intact	2.53 a	5 d	27.6 b	25.3 b	24.4 b	26.3 b	24.7 b
	cut	56.0 c	10 d	20.0 b	18.7 b	18.4 b	17.7 b	17.8 b
			13 d	19.0 b	17.3 b	16.9 b	14.9 b	17.2 b
Isobutyl acetate es4	intact	1.52 a	5 d	21.9 b	18.1 b	17.9 b	22.2 b	17.9 b
	cut	64.1 c	10 d	16.7 a–b	15.1 a–b	13.7 a–b	16.1 a–b	13.7 a–b
			13 d	15.9 a–b	15.8 a–b	12.3 a–b	13.9 a–b	14.2 a–b
2-methylbutyl acetate * es5	intact	25.2 a	5 d	139 d	127 c–d	110 b–d	114 b–d	128 c–d
	cut	323 e	10 d	94.1 b–d	94.7 b–d	79.5 a–c	74.8 a–c	91.1 b–d
			13 d	87.3 b–d	89.9 b–d	66.5 a–b	62 a–b	87.8 b–d
Pentyl acetate * es6	intact	0.78 a	5 d	4.36 b	3.34 a–b	3.36 a–b	4.04 b	3.16 a–b
	cut	13.1 c	10 d	2.64 a–b	2.44 a–b	2.02 a–b	2.37 a–b	1.91 a–b
			13 d	2.38 a–b	1.83 a–b	1.61 a–b	1.80 a–b	1.43 a–b
Hexyl acetate * es7	intact	3.15 a	5 d	5.98 a	4.64 a	4.45 a	5.19 a	3.73 a
	cut	59.8 b	10 d	3.48 a	3.56 a	2.31 a	2.85 a	2.14 a
			13 d	3.15 a	2.56 a	1.73 a	2.13 a	1.88 a

Table 2. Cont.

VOC	Time	Initial	Time	C	55	55_cp	65	65_cp
2-Hexen-1-yl acetate es8	intact	nd	5 d	nd	nd	nd	nd	nd
	cut	22.9 b	10 d	nd	nd	nd	nd	nd
			13 d	nd	nd	nd	nd	nd
Ethyl propionate es9	intact	1.46 b-c	5 d	3.17 e	3.16 e	2.03 c-d	1.22 b	1.65 b-d
	cut	nd	10 d	3.27 e	4.23 f	2.34 d	1.63 b-d	1.79 b-d
			13 d	5.01 g	4.49 f-g	2.33 d	1.34 b-c	1.82 b-d
Ethyl 2-methylpropanoate es10	intact	nd	5 d	1.06 b-c	2.88 e	1.32 b-c	1.64 c-d	1.24 b-c
	cut	nd	10 d	0.97 b-c	3.94 f	1.18 b-c	2.20 d-e	1.31 b-c
			13 d	1.35 b-c	4.76 g	0.63 a-b	0.82 a-c	0.89 a-c
Methyl butyrate es11	intact	0.63 a	5 d	1.37 a-b	1.06 a-b	1.39 a-b	0.77 a-b	1.79 b
	cut	4.20 c	10 d	1.13 a-b	0.93 a-b	1.36 a-b	0.63 a	1.48 a-b
			13 d	1.06 a-b	0.93 a-b	1.14 a-b	0.57 a	1.5 a-b
Methyl 2-methylbutyrate * es12	intact	nd	5 d	2.27 c-e	1.60 b-c	2.78 d-g	1.09 b	2.90 e-g
	cut	nd	10 d	1.97 b-d	1.62 b-c	3.48 g	1.09 b	3.27 f-g
			13 d	2.44 c-f	1.80 b-c	2.92 e-g	1.06 b	2.94 e-g
Ethyl butyrate * es13	intact	nd	5 d	12.1 b-c	16.8 c-d	12.4 b-c	13.4 b-c	12.5 b-c
	cut	1.64 a	10 d	10.9 b	18.4 d	11.9 b	13.3 b-c	11.6 b
			13 d	12.9 b-c	18.2 d	11.0 b	10.2 b	12.3 b-c
Ethyl 2-butenolate es14	intact	nd	5 d	0.54 b-d	0.62 b-e	0.51 b-c	0.91 e-f	0.93 e-f
	cut	nd	10 d	0.70 b-f	0.82 c-f	0.55 b-d	0.98 f	0.78 b-f
			13 d	0.84 d-f	0.98 f	0.49 b	0.79 b-f	nd
Ethyl 2-methylbutanoate * es15	intact	nd	5 d	25.1 b-c	37.9 d-e	30.4 b-d	23.3 b	31.9 b-d
	cut	0.98 a	10 d	25.1 b-c	43.3 e-f	29.6 b-d	33.6 c-d	32.8 b-d
			13 d	33.7 c-d	51.0 f	27.4 b-c	24.8 b-c	33.4 c-d
Hexyl butyrate * es16	intact	5.24 a	5 d	1.55 b-d	1.92 b-d	2.45 b-c	2.66 b	2.51 b-c
	cut	5.39 a	10 d	1.00 d-e	1.16 c-e	1.44 b-e	1.38 b-e	1.65 b-d
			13 d	0.63 d-e	nd	nd	nd	1.28 c-e
Hexyl 2-methylbutanoate * es17	intact	5.56 a	5 d	11 b-e	12.9 e	12.6 d-e	11.3 c-e	11.2 c-e
	cut	8.20 b	10 d	11.2 c-e	11.1 b-e	10.2 b-e	9.76 b-d	12.3 d-e
			13 d	11.4 c-e	9.96 b-d	9.80 b-d	8.67 b-c	12.1 d-e
Ethyl valerate es18	intact	nd	5 d	0.37 d-e	0.47 f-g	nd	0.39 e-f	nd
	cut	nd	10 d	0.34 c-e	0.54 g	0.26 b-c	0.28 b-d	nd
			13 d	0.38 e	0.48 g	0.21 b	nd	0.3 b-e
Methyl hexanoate es19	intact	nd	5 d	0.58 b-d	1.80 e-f	0.54 a-d	1.44 e	1.01 d
	cut	2.09 f	10 d	0.41 a-b	1.89 e-f	0.39 a-b	0.98 c-d	0.44 a-b
			13 d	0.49 a-c	2.17 f	nd	nd	nd
Ethyl hexanoate es20	intact	nd	5 d	13.3 i-j	15.7 j	9.95 g-h	8.04 f-g	7.67 e-g
	cut	1.56 a-b	10 d	7.15 d-g	11.6 h-i	4.88 c-e	4.74 b-e	4.28 b-d
			13 d	6.34 d-f	7.72 e-g	2.86 a-c	2.74 a-c	3.22 a-c
Hexyl hexanoate * es21	intact	2.11 a-b	5 d	0.98 d-g	1.28 c-g	1.57 b-e	2.48 a	1.75 a-d
	cut	1.92 a-c	10 d	0.71 f-g	1.07 d-g	0.82 e-g	1.27 c-g	1.43 b-f
			13 d	0.56 g	0.71 f-g	0.73 f-g	0.92 e-g	1.14 c-g
Ethanol al1	intact	9.72 b	5 d	13.1 c	27.7 f	14.9 c	22.0 d-e	20.5 d
	cut	2.21 a	10 d	13.1 c	27.8 f	14.0 c	20.9 d-e	15.6 c
			13 d	15.0 c	32.4 g	19.7 d	23.8 e	20.1 d
2-(2-Ethoxyethoxy)ethanol al2	intact	0.65 a	5 d	0.35 b-d	0.45 b	0.43 b	0.37 b-d	0.38 b-c
	cut	0.62 a	10 d	0.30 c-e	0.25 d-e	0.22 e	0.29 c-e	0.22 e
			13 d	0.26 c-e	0.25 d-e	0.28 c-e	0.27 c-e	0.29 c-e
2-Methyl-1-propanol al3	intact	0.86 a	5 d	1.02 a-b	0.96 a	0.94 a	1.21 a-c	1.13 a-b
	cut	2.99 e	10 d	0.92 a	1.01 a-b	0.88 a	1.07 a-b	0.93 a
			13 d	1.07 a-b	1.68 d	1.50 c-d	1.36 b-d	1.09 a-b
1-Butanol * al4	intact	1.51 a	5 d	1.56 a	2.01 b-c	1.72 a-b	2.13 c	1.99 b-c
	cut	2.82 d	10 d	1.59 a	1.82 a-c	1.72 a-b	1.79 a-c	1.66 a-b
			13 d	1.46 a	1.67 a-b	1.67 a-b	1.67 a-b	1.64 a-b
2-Methyl-1-butanol al5	intact	3.24 a	5 d	4.25 a-b	5.32 b-d	4.44 a-c	5.14 b-d	5.89 c-d
	cut	9.02 e	10 d	4.63 a-c	5.58 b-d	4.67 b-c	4.60 a-c	5.21 b-d
			13 d	4.51 a-c	6.3 d	4.77 b-c	4.67 b-c	5.55 b-d

Table 2. Cont.

VOC	Time	Initial	Time	C	55	55_cp	65	65_cp
1-Hexanol *	intact	1.44 a	5 d	2.90 b	3.99 b-c	3.07 b	4.54 c	3.60 b-c
	cut	3.94 b-c	10 d	3.06 b	4.21 b-c	3.08 b	3.39 b-c	2.84 b
			13 d	2.88 b	3.93 b-c	3.27 b-c	3.52 b-c	3.11 b
2-Butanone	intact	2.67 a-e	5 d	0.86 a-b	4.71 e-f	3.62 c-e	8.89 g	6.75 f-g
	cut	0.33 a	10 d	0.73 a-b	7.32 g	1.96 a-d	4.68 e-f	4.22 d-e
			13 d	1.32 a-c	7.12 g	2.44 a-e	3.30 b-e	3.76 c-e
1-Penten-3-one	intact	nd	5 d	1.35 b	2.29 b-d	1.77 b-c	3.35 e-f	1.98 b-c
	cut	nd	10 d	3.71 e-f	3.98 f	3.21 d-f	5.16 g	3.72 e-f
			13 d	3.52 e-f	2.78 c-e	4.18 f	4.06 f	3.76 e-f
Hexanal *	intact	0.50 a	5 d	3.23 b-d	6.47 f-g	4.66 d-f	7.14 g	5.25 d-g
	cut	1.92 a-b	10 d	4.49 d-f	5.08 d-f	3.72 b-d	6.26 e-g	4.43 d-e
			13 d	4.19 c-d	2.30 a-c	3.46 b-d	3.77 b-d	3.81 b-d
Estragole	intact	0.30 b-e	5 d	0.61 g	0.42 c-f	0.45 d-g	0.35 b-f	0.47 e-g
	cut	0.41 b-f	10 d	0.52 f-g	nd	nd	0.29 b-d	0.37 b-f
			13 d	0.51 f-g	nd	0.26 b-c	0.24 b	0.36 b-f
D-Limonene	intact	3.27 a	5 d	6.26 a-d	10.4 e	9.77 e	7.54 b-e	8.45 d-e
	cut	4.93 a-c	10 d	4.74 a-b	4.54 a	9.54 e	7.65 c-e	8.67 d-e
			13 d	4.70 a-b	4.40 a	9.06 d-e	5.23 a-c	9.28 e
α -Farnesene	intact	13.8 a-c	5 d	24.0 d-e	18.8 a-d	21.3 c-e	16.8 a-d	28.0 e
	cut	19.1 a-d	10 d	17.4 a-d	14.4 a-c	14.9 a-c	15.0 a-c	20.5 b-e
			13 d	16.5 a-d	10.6 a	12.1 a-b	12.5 a-b	17.7 a-d

* marked VOCs described as generally important “character impact” compounds in apples [29]. nd = not detected/below detection limit; dark green = $\geq 500\%$ increase; light green = $\geq 33\%$ increase; yellow = $\leq 33\%$ increase/decrease; light red = $\geq 80\%$ decrease; dark red = $\geq 500\%$ decrease.

During further storage, the emissions of the majority of VOCs were significantly reduced compared to those observed immediately after cutting. Cumulative VOCs concentrations decreased by 37.7% for control and by 18.8%–50.7% for sHWT samples, but were nevertheless 3.5–5.8 times higher than for intact apples (Table 2, 1st position). While cutting immediately and strongly increased the emission of acetate esters, the concentrations of these compounds decreased or, at least, remained constant during storage. Only the concentration of ethyl-acetate continued to increase after 3 days of storage as did the concentrations of all other ethyl esters, methyl 2-methylbutyrate, ethanol, hexanal and both ketones.

Responses of sHWT samples depended on both the treatment temperatures and the post-processing acid treatment (Table 3). Head space concentration of ethanol was generally higher and that of estragole and propyl-acetate lower in sHW-treated samples than in controls; propyl acetate concentration was more reduced at 65 °C than at 55 °C. Furthermore, sHWT at 65 °C significantly reduced emissions of ethyl-propionate, ethyl hexanoate and methyl hexanoate. These VOCs were also reduced at sHWT of 55 °C combined with acid treatment, i.e. chemical prevention. The combination of sHWT and acid application also resulted in increased D-Limonene and decreased ethyl 2-butenate emissions, while the enhancement of the ethanol emission was attenuated. At a sHWT of 55 °C, the combination with organic acid resulted in lower ethyl acetate emission. In contrast, only the sole sHWT at 55 °C (i.e., without cp) let the emissions of nearly all ethyl esters increase especially that of ethyl-acetate, methyl hexanoate and 2-butanone, compared to controls.

Table 3. Treatment-specific qualitative changes in the emissions of relevant VOCs and corresponding aroma threshold values of fresh-cut apple slices stored in sugar syrup at 4 °C for up to 13 days.

Treatment	Reaction	VOCs	Aroma Threshold Values (nL L ⁻¹)
all sHWT	↑	Ethanol	8–900 ^a / $>1 \times 10^{5,b}$
	↓	Propyl acetate	2000–11,000 ^{a,b}
		Estragole	n/a
sHWT without cp	↑	Ethanol	8–900 ^a / $>1 \times 10^{5,b}$
sHWT combined with cp	↑	D-limonene	4–229 ^a
	↓	Ethyl 2-butenate	n/a
only 55 °C sHWT	↑	2-methyl-1-propanol	360–3300 ^a
		Ethyl acetate	5–13,500 ^b
		Ethyl 2-methylpropanoate	0.01–1 ^a
		Ethyl butyrate *	0.1–18 ^a
		Ethyl 2-methylbutanoate *	0.006–0.1 ^{a,b}
only 55 °C sHWT without cp	↑	Ethyl valerate	1.5–5 ^{a,b}
		Methyl hexanoate	10–87 ^a
		2-butanone	n/a
		Estragole	n/a
		1-penten-3-one	400 ^a
only 55 °C sHWT combined with cp	↓	Ethyl acetate	5–13,500 ^{a,b}
only 65 °C sHWT	↓	Propyl acetate	2000–11,000 ^{a,b}
only sHWT at 65 °C or combined with cp	↓	Ethyl propionate	9–45 ^a
		Ethyl hexanoate	0.3–5 ^a
		Methyl hexanoate	10–87 ^a

sHWT: short-term hot water treatment; cp: chemical prevention by applying organic acid dipping of fresh-cut slices; 55 °C/65 °C: sHWT at the specified temperature; green ↑: increase; red ↓: decrease. Given are aroma threshold values according to ^a Burdock [30] and ^b Dixon and Hewett [29]. * marked VOCs described as generally important “character impact” compounds in apples [29].

The PCA provides two principal components (PC), which together represent 98.2% of the total variance in VOCs profiles. The effects of cutting, sHWT temperature and duration of storage can be visualized by a scatter point plot (Figure 2A). Decreasing values of PC1 (representing 65.3% of total variance in VOCs profiles) can be associated with the effects of cutting, while increasing values are associated with storage time. Decreasing values of PC2 (representing 33.0% of total variance in VOCs profiles) can also be associated with cutting, and, beyond that, with the distinct effects of the various treatments. Interestingly, sHWT at 55 °C induced the highest alteration of the VOCs profile from that of intact apples (c.f. colored bars in Figure 2A).

For both PCs, the impact of each single VOC on the total variance in the VOCs profiles can be identified as visualized in Figure 2B. Mostly, ethyl acetate (es1) and, to a much smaller extent, ethyl 2-methylbutanoate (es15) and ethanol (al1) are related to effects of the sHWT (PC2), but also to changes during storage (PC1). Cutting-induced changes in the VOCs profiles (both PC decreased) are mainly associated with 2-methylbutyl acetate (es5), but also with isobutyl acetate (es4), butyl acetate (es3) and hexyl acetate (es7).




Figure 2. (A) Scatterplot of principal components analysis (PCA)-results of the variability of VOCs profiles during storage of treated and untreated fresh-cut apple slices. Decreasing scores of PC1 and PC2 can be associated with the effects of cutting; scores values of PC1 are associated with storage time, marked by colored areas (blue = 3 days and red = 10 days); decreasing scores of PC2 additional associated with distinct effects of the various treatments, marked by colored bars. (B) Loading plot of PCA-results of single-VOC impact on the total variance in VOCs profiles. VOCs with negative loadings of both PCs can be associated with the effects of cutting; VOCs with positive PC1 and negative PC2 loadings are related during storage and distinct effects of the treatments.

3.2. Ethylene Evolution

Cutting immediately and significantly intensified ethylene release of untreated apple slices by approx. 30%; however, ethylene emission also rapidly, within 4 h, declined to rates below that of intact apples (Figure 3). Storage in sugar syrup generally reduced ethylene release as measured after removing the samples from the solution. Compared to controls, sHWT at 55 °C initially (i.e., at day 5 of storage) intensified ethylene emissions of apples slices, while that at 65 °C pronouncedly lowered them. Irrespective of treatments, ethylene emission of all apple slices continuously declined during further storage (Figure 3; days 10 and 13).




Figure 3. Ethylene release rates of short-term hot-water treated and untreated fresh-cut apple slices at days 5, 10 and 13 of storage in sugar syrup at 4 °C, compared to intact apples and untreated fresh-cut apple slices (initial). Given are means \pm standard deviation ($n = 3$). Different letters indicate significant differences between means ($p < 0.05$). 5 d = 5 days, 10 d = 10 days, 13 d = 13 days.

3.3. Respiration

CO₂-based respiration rates of untreated 'Braeburn' apples increased 3.4 times immediately after cutting and then slightly declined again within 4 h (Figure 4). During storage in sugar syrup, CO₂ release of all samples was higher than that of intact apples. In addition, respiration of sHWT-samples treated at 55 °C without acid treatment was significantly higher than that of the other apple slices at all sampling days.




Figure 4. Respiration rates of short-term hot-water treated and untreated fresh-cut apple slices at days 5, 10 and 13 of storage in sugar syrup at 4 °C, compared to intact apples and untreated fresh-cut apple slices (initial). Given are means ± standard deviation ($n = 3$). Different letters indicate significant differences between means ($p < 0.05$). 5 d = 5 days, 10 d = 10 days, 13 d = 13 days.

4. Discussion

In the presented experiment, several overlaying and potentially interactive effects were responsible for the development of respective VOCs profiles during storage of apple slices, i.e., cutting, sHW-treatment at two temperatures and storage in sugar syrup with its pronounced effect on O₂ availability. The most pronouncedly emitted VOCs were esters. Similarly, Paillard [31] reported that esters of acetic, butanoic and hexanoic acids with ethyl, butyl and hexyl alcohols are the most frequent VOCs detected in the headspace of intact apples. The respective composition of the above esters determines the typical fruity (i.e., 'apple-like') cultivar-specific aroma of apples [32]. In the presented study, 10 of the identified esters were described as generally important "character impact" VOCs [29]. Although the overall composition of the VOCs may clearly vary among different apple cultivars [33], 2-methylbutyl acetate and α -Farnesene are most often the major components in 'Braeburn' apples [34,35]. This was confirmed by the present results.

4.1. Effect of Cutting on the Release of VOCs, Ethylene and CO₂

The cutting-related increase in acetates probably impacted the aroma of fresh-cut apples, as 2-methylbutyl acetate, butyl-acetate and hexyl acetate are characterized as "character impact compounds" [29]. These VOCs are predominantly associated with a characteristic sweet, cox-like apple aroma [36]. The presented results confirmed earlier findings that cutting inevitably induces physiological wounding-stress responses characterized by temporarily increased ethylene synthesis [37,38], respiration activity [7] and VOCs release [39], which all return to pre-processing levels within 24 h [40]. Emissions of acetate-esters may strongly increase, due to increased lipoxygenase activity [41] resulting in enhanced esterification of membrane lipids [42,43]. Furthermore, a largely accelerated glycolysis increases the acetyl-CoA availability, and thus also boosts amino-acids production, which serve as precursors to many VOCs [43,44]. During storage, however, the concentrations of the aroma-relevant acetates decreased, which may indicate that the cutting induced stress response was reduced or no longer existed. Therefore, the generation of precursors relevant for VOC synthesis was reduced resulting in a decline of emissions. However, the emissions were still higher than those from intact apples. Nevertheless, the typical aroma, appearing immediately after cutting the apples,

got continuously lost during storage. This, however, is generally monitored during storage of fresh-cut apples [44].

4.2. Impact of Hot-Water Treatment on the Release of VOCs, Ethylene and CO₂

Even the sHWT at 65 °C did not alter the emission of most VOCs or pronouncedly impacted VOCs profiles but only marginally reduced that of few esters. In contrast to cutting, sHWT did obviously also not affect the synthesis of acetate-esters. This is surprising because a high number of VOCs emanate from the epidermal layers of apples, while only a lower proportion derived from the pulp tissue [45]. However, dipping intact apples for 30 s in hot-water at 55 °C exclusively heated the epidermis and few hypodermal cell layers [17]. Furthermore, although treatments at higher temperatures (i.e., >55 °C) resulted in faster increase in tissue temperature of deeper cell layers, only 70 °C pronouncedly damaged the epidermis [17]. Thus, sHWT, in the temperature range used, did not decrease the emission of important “character impact” compounds [29] or increase that of VOCs associated with off-odor. Therefore, sHWT did not reduce the aroma quality. Similarly, sHWT (of up to 65 °C) did not adversely affect color attributes, tissue strength or other important quality parameters of fresh-cut apple slices [18].

Although the increased ethanol emission may be assumed as critically, the aroma-relevant threshold for this compound is much higher than for other important VOCs (Table 3). In addition, the ethanol concentrations in total were very low. The observed increase in ethanol emission, therefore, did not negatively affect the aroma quality at all. It may, nevertheless, indicate some sHWT-induced heat stress because ethanol and ethyl-acetate emissions are known to increase in response to and may be used as indicators of this stressor [46,47]. Heat stress was reported to alter the glycolytic pathway by disturbing the mitochondrial electron transport, which, similar to reduced O₂ availability, results in increased ethanol formation [48]. This may also be reflected by the marginally higher respiratory activity observed at the end of storage. Heat treatment-enhanced respiration was also reported earlier for intact ‘Granny Smith’ and ‘Anna’ apples [49]. In addition, the increased D-limonene emissions of acid-treated sHWT samples may be attributed to major perturbations of the cellular metabolism that result in the expression of multiple genes and, finally, in enhanced terpene emissions [50,51].

Omitting the post-processing acid treatment increased the ethanol emission, which, however, did not negatively affect the aroma (see above). Hot-water treatments without post-processing acid application only slightly enhanced the alteration of the VOCs profile typical for intact apples in comparison to those with additional acid dipping. Interestingly, sHWT at 55 °C without additional acid treatment clearly intensified the storage-induced physiological responses of fresh-cut apple slices, i.e., it slightly enhanced their respiration activity and ethylene release. This may indicate a direct response of the treated apples to moderate but not to excessive heat or to acid pretreatment [52,53]. Additionally, the emission of ethyl esters is further intensified in sHW-treated (55 °C) apple slices. This is especially obvious for ethyl acetate and ethanol, which, however, both increased in all samples during storage (see Section 4.3). As ethylene may directly regulate the synthesis of VOCs [44,54,55], the increase in ethyl 2-methylpropanoate, ethyl butyrate and ethyl 2-methylbutanoate may be directly related to the enhanced ethylene tissue concentrations. Ethyl acetate correlates with pronounced off-odor [56], while the “character impact compounds” [29] ethyl-butyrates and ethyl 2-methylbutanoate are generally associated with fruity and apple-like aroma [57]. The increase in the head space concentrations of the latter compounds, which own aroma thresholds much lower than ethyl acetate [29,30], may thus even improve the overall aroma of the apple samples (Table 3).

4.3. Effects of Storage in Sugar Syrup on the Release of VOCs, Ethylene and CO₂

Storing fresh-cut apple slices in sugar syrup pronouncedly affected the development of their VOCs profiles, as indicated by the distinct increase in ethyl ester emissions. Following sHWT, head space concentration of ethyl acetate of samples increased 0.5–1.6-fold compared to controls; it, however, increased approx. 100 times due to the sugar syrup.

The reduction of ethylene and CO₂ release observed after immersion in the syrup, corroborate the findings of Rux et al. [7]. These authors stored apple slices in sugar syrups of different concentrations and attributed major changes in respiration of samples to the reduced O₂ availability due to impaired gas exchange between tissues and ambient air [7,58]. Therefore, the fast decrease in O₂ and increase in CO₂ concentrations within the tissue increasingly inhibits physiological processes [59]. The increased emissions of ethanol and other alcohols probably indicate semi-anaerobic conditions [60].

Likewise, the increased emissions of ethyl esters may be explained by the increased ethanol formation under these conditions. Alcohol acetyl-CoA transferase (AAT) catalyzes the esterification of carboxylic acids and alcohols [46,61], and enhanced ethanol concentrations may enhance the formation of ethyl esters, especially ethyl acetate [62]. In this context, Cortellino et al. [62] measured a progressive increase of ethyl acetate in fresh-cut apple tissue within 11 days of storage in modified atmosphere, most pronounced at an O₂ concentration of 1%.

However, in this study, the influence of oxygen availability on VOC synthesis during storage in sugar syrup submerged apple pieces cannot be sufficiently clarified. Further studies focusing on oxygen availability are necessary.

5. Conclusions

In a semi-practical approach, this study comprehensively evaluated the potential food quality-related effects of sHWT (at 55 °C and 65 °C), cutting and/or post-processing acid treatment and their interactions on the development of aroma-relevant VOCs as well as on the physiological indicators ethylene synthesis and respiration activity in fresh-cut apple slices stored in sugar syrup. Cutting temporarily enhanced ethylene synthesis, respiration activity and VOCs production. In this context, 2-methylbutyl acetate, butyl acetate and hexyl acetate were identified as most relevant cutting-enhanced VOCs, but their emissions declined again during syrup-storage. Syrup-storage increased ethanol and ethyl-esters, and in particular ethyl-acetate, which might be attributed to reduced O₂ availability under this condition. Solely applying sHWT at both 55 °C and 65 °C enhanced the respiration of apples slices, which potentially indicated marginal heat stress. This, however, only slightly intensified ethylene release but increased ethanol emission of samples, indicating alterations in respiratory pathways. Additionally, sHWT at 65 °C resulted only in minor declines of few esters. Nevertheless, the general impact of sHWT on VOCs profiles of apples slices was low, especially in comparison to storage in sugar syrup and did not negatively affect aroma. The omission of pre-processing acid application did not negatively affect the aroma to a crucial degree. Consequently, sHWT might replace the post-processing acid treatments for fresh-cut products.

Author Contributions: Conceptualization, K.H., W.B.H. and G.R.; data curation, W.B.H.; formal analysis, G.R. and E.E.; funding acquisition, K.H.; investigation, G.R. and E.E.; methodology, K.H., W.B.H. and G.R.; project administration, K.H.; resources, S.H.-K.; supervision, S.H.-K. and C.U.; validation, K.H., W.B.H., G.R. and S.H.-K.; visualization, W.B.H., G.R. and S.H.-K.; writing—original draft, G.R. and W.B.H.; writing—review and editing, W.B.H., K.H., S.H.-K. and C.U. All authors have read and agreed to the published version of the manuscript.

Funding: The current research was funded by the German Federation of Industrial Research Associations—“Otto von Guericke” e.V. (AiF)/German Federal Ministry for Economic Affairs and Energy (BMWi) (NO:F4001902MD5), within the research program Zentrales Innovationsprogramm Mittelstand (ZIM). The publication of this article was funded by the Open Access Fund of the Leibniz Association.

Acknowledgments: The authors thank Kathrin Ilte for perfect technical assistance at ATB.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Watada, A.E.; Qi, L. Quality of fresh-cut produce. *Postharvest Biol. Technol.* **1999**, *15*, 201–205. [[CrossRef](#)]
2. Beaulieu, J. Factors affecting sensory quality of fresh-cut produce. In *Advances in Fresh-Cut Fruits and Vegetables*; Martin-Belloso, O., Soliva-Fortuny, R., Eds.; CRC Press: Boca Raton, FL, USA, 2010; pp. 115–143.

3. Toivonen, P.M.; DeEll, J.R. Physiology of Fresh-Cut Fruits and Vegetables. In *Fresh-Cut Fruits and Vegetables*; Lamikanra, O., Ed.; CRC Press: Boca Raton, FL, USA, 2002; pp. 91–123.
4. Alzamora, S.M.; Tapia, M.S.; Argaiz, A.; Welli, J. Application of combined methods technology in minimally processed fruits. *Food Res. Int.* **1993**, *26*, 125–130. [[CrossRef](#)]
5. Nishikawa, F.; Iwama, T.; Kato, M.; Hyodo, H.; Ikoma, Y.; Yano, M. Effect of sugars on ethylene synthesis and responsiveness in harvested broccoli florets. *Postharvest Biol. Technol.* **2005**, *36*, 157–165. [[CrossRef](#)]
6. Coupe, S.A.; Sinclair, B.K.; Watson, L.M.; Heyes, J.A.; Eason, J.R. Identification of dehydration-responsive cysteine proteases during post-harvest senescence of broccoli florets. *J. Exp. Bot.* **2003**, *54*, 1045–1056. [[CrossRef](#)]
7. Rux, G.; Caleb, O.J.; Fröhling, A.; Herppich, W.B.; Mahajan, P.V. Respiration and storage quality of fresh-cut apple slices immersed in sugar syrup and orange juice. *Food Bioprocess Technol.* **2017**, *10*, 2081–2091. [[CrossRef](#)]
8. Pietrysiak, E.; Ganjyal, G.M. Apple peel morphology and attachment of *Listeria innocua* through aqueous environment as shown by scanning electron microscopy. *Food Control* **2018**, *92*, 362–369. [[CrossRef](#)]
9. Lurie, S. Postharvest heat treatments. *Postharvest Biol. Technol.* **1998**, *14*, 257–269. [[CrossRef](#)]
10. Shao, X.F.; Tu, K.; Zhao, Y.Z.; Chen, L.; Chen, Y.Y.; Wang, H. Effects of prestorage heat treatment on fruit ripening and decay development in different apple cultivars. *J. Hort. Sci. Biotech.* **2007**, *82*, 297–303. [[CrossRef](#)]
11. Spadoni, A.; Guidarelli, M.; Phillips, J.; Mari, M.; Wisniewski, M. Transcriptional profiling of apple fruit in response to heat treatment: Involvement of a defense response during *Penicillium expansum* infection. *Postharvest Biol. Technol.* **2015**, *101*, 37–48. [[CrossRef](#)]
12. Kabelitz, T.; Hassenberg, K. Control of apple surface microflora for fresh-cut produce by post-harvest hot-water treatment. *LWT Food Sci. Technol.* **2018**, *98*, 492–499. [[CrossRef](#)]
13. Fallik, E. Prestorage hot water treatments (immersion, rinsing and brushing). *Postharvest Biol. Technol.* **2004**, *32*, 125–134. [[CrossRef](#)]
14. Maxin, P.; Williams, M.; Weber, R.W. Control of fungal storage rots of apples by hot-water treatments: A Northern European perspective. *Erwerbs-Obstbau* **2014**, *56*, 25–34. [[CrossRef](#)]
15. Roy, S.; Conway, W.S.; Watada, A.E.; Sams, C.E.; Erbe, E.F.; Wergin, W.P. Heat treatment affects epicuticular wax structure and postharvest calcium uptake in Golden Delicious apples. *HortScience* **1994**, *29*, 1056–1058. [[CrossRef](#)]
16. Lurie, S.; Fallik, E.; Klein, J.D. The effect of heat treatment on apple epicuticular wax and calcium uptake. *Postharvest Biol. Technol.* **1996**, *8*, 271–277. [[CrossRef](#)]
17. Kabelitz, T.; Schmidt, B.; Herppich, W.B.; Hassenberg, K. Effects of hot water dipping on apple heat transfer and post-harvest fruit quality. *LWT Food Sci. Technol.* **2019**, *108*, 416–420. [[CrossRef](#)]
18. Rux, G.; Efe, E.; Ulrichs, C.; Huyskens-Keil, S.; Hassenberg, K.; Herppich, W.B. Effects of pre-processing short-term hot-water treatments on quality and shelf life of fresh-cut apple slices. *Foods* **2019**, *8*, 653. [[CrossRef](#)]
19. Biegańska-Marecik, R.; Czapski, J. The effect of selected compounds as inhibitors of enzymatic browning and softening of minimally processed apples. *Acta Sci. Pol. Technol. Aliment.* **2007**, *6*, 37–49.
20. Barrett, D.M.; Beaulieu, J.C.; Shewfelt, R.L. Color, flavor, texture and nutritional quality of fresh-cut fruits and vegetables: Desirable levels, instrumental and sensory measurement, and effects of processing. *Crit. Rev. Food Sci. Nutr.* **2010**, *50*, 369–389. [[CrossRef](#)]
21. Schwab, W.; Davidovich-Rikanati, R.; Lewinsohn, E. Biosynthesis of plant-derived flavor compounds. *Plant J.* **2008**, *54*, 712–732. [[CrossRef](#)]
22. Baldwin, E.A.; Scott, J.W.; Shewmaker, C.K.; Schuch, W. Flavor trivia and tomato aroma: Biochemistry and possible mechanisms for control of important aroma components. *HortScience* **2000**, *35*, 1013–1022. [[CrossRef](#)]
23. Espino-Díaz, M.; Sepúlveda, D.R.; González-Aguilar, G.; Olivás, G.I. Biochemistry of apple aroma: A review. *Food Technol. Biotech.* **2016**, *54*, 375–394. [[CrossRef](#)] [[PubMed](#)]
24. Schaffer, R.J.; Friel, E.N.; Souleyre, E.J.; Bolitho, K.; Thodey, K.; Ledger, S.; Bowen, J.H.; Ma, J.; Nain, B.; Cohan, D.; et al. A genomics approach reveals that aroma production in apple is controlled by ethylene predominantly at the final step in each biosynthetic pathway. *Plant Physiol.* **2007**, *144*, 1899–1912. [[CrossRef](#)] [[PubMed](#)]
25. Lamikanra, O.; Chen, J.C.; Banks, D.; Hunter, P.A. Biochemical and microbial changes during the storage of minimally processed cantaloupe. *J. Agric. Food Chem.* **2000**, *48*, 5955–5961. [[CrossRef](#)] [[PubMed](#)]

26. Rux, G.; Luca, A.; Mahajan, P.V. Changes in volatile organic compounds in the headspace of modified atmosphere packed and unpacked white sausages. *Food Packag. Shelf Life* **2019**, *19*, 167–173. [CrossRef]
27. Van den Dool, H.; Kratz, P.D. A generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography. *J. Chromatogr. A* **1963**, *11*, 463–471. [CrossRef]
28. NIST. *Chemistry WebBook: NIST Standard Reference Database Number 69*; Linstrom, P.J., Mallard, W.G., Eds.; National Institute of Standards and Technology: Gaithersburg, MD, USA, 2012. Available online: <https://webbook.nist.gov/chemistry/> (accessed on 25 August 2019). [CrossRef]
29. Dixon, J.; Hewett, E.W. Factors affecting apple aroma/flavour volatile concentration: A review. *N. Z. J. Crop Hortic. Sci.* **2000**, *28*, 155–173. [CrossRef]
30. Burdock, G.A. *Fenaroli's Handbook of Flavor Ingredients*, 6th ed.; CRC Press: Boca Raton, FL, USA, 2016.
31. Paillard, N.M.M. The flavour of apples, pears and quinces. In *Food Flavours. Part C: The Flavour of Fruits*; Morton, I.D., MacLeod, A.J., Eds.; Elsevier Science Publishers: Amsterdam, The Netherlands, 1990; pp. 1–41, ISBN 0-444-87362-7.
32. Fellman, J.K.; Mattinson, D.S.; Bostick, B.C.; Mattheis, J.P.; Patterson, M.E. Ester biosynthesis in “Rome” apples subjected to low-oxygen atmospheres. *Postharvest Biol. Technol.* **1993**, *3*, 201–214. [CrossRef]
33. Poll, L. Evaluation of 18 apple varieties for their suitability for juice production. *J. Sci. Food Agric.* **1981**, *32*, 1081–1090. [CrossRef]
34. Aaby, K.; Haffner, K.; Skrede, G. Aroma quality of Gravenstein apples influenced by regular and controlled atmosphere storage. *LWT Food Sci. Technol.* **2002**, *35*, 254–259. [CrossRef]
35. Baiamonte, I.; Raffo, A.; Nardo, N.; Moneta, E.; Peparario, M.; D’Aloise, A.; Kelderer, M.; Casera, C.; Paoletti, F. Effect of the use of anti-hail nets on codling moth (*Cydia pomonella*) and organoleptic quality of apple (cv. Braeburn) grown in Alto Adige Region (northern Italy). *J. Sci. Food Agric.* **2016**, *96*, 2025–2032. [CrossRef]
36. Young, H.; Gilbert, J.M.; Murray, S.H.; Ball, R.D. Causal effects of aroma compounds on royal gala apple flavours. *J. Sci. Food Agric.* **1996**, *71*, 329–336. [CrossRef]
37. Soliva-Fortuny, R.C.; Ricart-Coll, M.; Martín-Belloso, O. Sensory quality and internal atmosphere of fresh-cut golden delicious apples. *Int. J. Food Sci. Tech.* **2005**, *40*, 369–375. [CrossRef]
38. Soliva-Fortuny, R.C.; Oms-Oliu, G.; Martín-Belloso, O. Effects of ripeness stages on the storage atmosphere, color, and textural properties of minimally processed apple slices. *J. Food Sci.* **2002**, *67*, 1958–1963. [CrossRef]
39. Bett, K.L.; Ingram, D.A.; Grimm, C.C.; Lloyd, S.W.; Spanier, A.M.; Miller, J.M.; Gross, E.A.; Baldwin, B.T.; Vinyard, B.T. Flavor of fresh-cut gala apples in barrier film packaging as affected by storage time. *J. Food Qual.* **2001**, *24*, 141–156. [CrossRef]
40. Finnegan, E.; Mahajan, P.V.; O’Connell, M.; Francis, G.A.; O’Beirne, D. Modelling respiration in fresh-cut pineapple and prediction of gas permeability needs for optimal modified atmosphere packaging. *Postharvest Biol. Technol.* **2013**, *79*, 47–53. [CrossRef]
41. Schreier, P. *Chromatographic Studies of Biogenesis of Plant Volatiles*; Alfred Hüthig Verlag GmbH: Berlin/Heidelberg, Germany, 1984.
42. De Pooter, H.L.; Montens, J.P.; Willaert, G.A.; Dirinck, P.J.; Schamp, N.M. Treatment of Golden Delicious apples with aldehydes and carboxylic acids: Effect on the headspace composition. *J. Agric. Food Chem.* **1983**, *31*, 813–818. [CrossRef]
43. Rowan, D.D.; Allen, J.M.; Fielder, S.; Hunt, M.B. Biosynthesis of straight-chain ester volatiles in Red Delicious and Granny Smith apples using deuterium-labeled precursors. *J. Agric. Food Chem.* **1999**, *47*, 2553–2562. [CrossRef]
44. Beaulieu, J.C. Effect of cutting and storage on acetate and nonacetate esters in convenient, ready-to-eat fresh-cut melons and apples. *HortScience* **2006**, *41*, 65–73. [CrossRef]
45. Ferreira, L.; Perestrelo, R.; Caldeira, M.; Câmara, J.S. Characterization of volatile substances in apples from rosaceae family by headspace solid-phase microextraction followed by GC-qMS. *J. Sep. Sci.* **2009**, *32*, 1875–1888. [CrossRef]
46. Forney, C.F.; Kalt, W.; Jordan, M.A. The composition of strawberry aroma is influenced by cultivar, maturity, and storage. *HortScience* **2000**, *35*, 1022–1026. [CrossRef]
47. Song, J.; Fan, L.; Forney, C.F.; Jordan, M.A. Using volatile emissions and chlorophyll fluorescence as indicators of heat injury in apples. *J. Am. Soc. Hortic. Sci.* **2001**, *126*, 771–777. [CrossRef]

48. Fan, L.; Song, J.; Forney, C.F.; Jordan, M.A. Ethanol production and chlorophyll fluorescence predict breakdown of heat-stressed apple fruit during cold storage. *J. Am. Soc. Hortic. Sci.* **2005**, *130*, 237–243. [[CrossRef](#)]
49. Klein, J.D. Ethylene biosynthesis in heat-treated apples. In *Biochemical and Physiological Aspects of Ethylene Production in Lower and Higher Plants*; Clijsters, H., de Proft, M., Marcelle, R., van Poucke, M., Eds.; Springer: Dordrecht, The Netherlands, 1989; pp. 181–189.
50. Singh, M.; Johnson-Flanagan, A. Co-ordination of photosynthetic gene expression during low-temperature acclimation and development in *Brassica napus* cv. jet neuf leaves. *Plant Sci.* **1998**, *135*, 171–181. [[CrossRef](#)]
51. Pateraki, I.; Kanellis, A.K. Stress and developmental responses of terpenoid biosynthetic genes in *Cistus creticus* subsp. *creticus*. *Plant Cell Rep.* **2010**, *29*, 629–641. [[CrossRef](#)]
52. Charron, C.S.; Cantliffe, D.J.; Heath, R.R. Volatile emissions from plants. *Hort. Rev.* **1995**, *17*, 43–72.
53. Woodstock, L.W.; Taylorson, R.B. Ethanol and acetaldehyde in imbibing soybean seeds in relation to deterioration. *Plant Physiol.* **1981**, *67*, 424–428. [[CrossRef](#)]
54. Bauchot, A.D.; Mottram, D.S.; Dodson, A.T.; John, P. Effect of aminocyclopropane-1-carboxylic acid oxidase antisense gene on the formation of volatile esters in cantaloupe Charentais melon (cv. Vedrandais). *J. Agric. Food Chem.* **1998**, *46*, 4787–4792. [[CrossRef](#)]
55. Obando-Ulloa, J.M.; Nicolai, B.; Lammertyn, J.; Bueso, M.C.; Monforte, A.J.; Fernández-Trujillo, J.P. Aroma volatiles associated with the senescence of climacteric or non-climacteric melon fruit. *Postharvest Biol. Technol.* **2009**, *52*, 146–155. [[CrossRef](#)]
56. Larsen, M.; Watkins, C.B. Firmness and concentrations of acetaldehyde, ethyl acetate and ethanol in strawberries stored in controlled and modified atmospheres. *Postharvest Biol. Technol.* **1995**, *5*, 39–50. [[CrossRef](#)]
57. Rizzolo, A.; Polesello, A.; Teleky-Vámosy, G. CGC/Sensory analysis of volatile compounds developed from ripening apple fruit. *J. High Resolut. Chromatogr.* **1989**, *12*, 824–827. [[CrossRef](#)]
58. Von Willert, D.J.; Matyssek, R.; Herppich, W.B. *Experimentelle Pflanzenökologie: Grundlagen und Anwendungen*; Georg Thieme Verlag: Stuttgart, Germany, 1995; p. 344, ISBN 3-13-134401-6.
59. Rocculi, P.; Del Nobile, M.A.; Romani, S.; Baiano, A.; Dalla Rosa, M. Use of a simple mathematical model to evaluate dipping and MAP effects on aerobic respiration of minimally processed apples. *J. Food Eng.* **2006**, *76*, 334–340. [[CrossRef](#)]
60. Lara, I.; Graell, J.; López, M.L.; Echeverría, G. Multivariate analysis of modifications in biosynthesis of volatile compounds after CA storage of 'Fuji' apples. *Postharvest Biol. Technol.* **2006**, *39*, 19–28. [[CrossRef](#)]
61. Fellman, J.K.; Mattheis, J.P.; Matthinson, D.S.; Bostick, B.C. Assay of acetyl-CoA alcohol transferase in delicious apples. *HortScience* **1991**, *27*, 773–776.
62. Cortellino, G.; Gobbi, S.; Bianchi, G.; Rizzolo, A. Modified atmosphere packaging for shelf life extension of fresh-cut apples. *Trends Food Sci. Technol.* **2015**, *46*, 320–330. [[CrossRef](#)]

