

Pentacarbonyl-2 κ^5 C-chlorido-1 κ Cl-bis-[1(η^5)-cyclopentadienyl][μ -oxido(phenyl)-methylene-1:2 κ^2 O:C]hafnium(IV)-tungsten(0)

Catharine Esterhuysen,* I. B. Jacques Nel† and Stephanie Cronje

Department of Chemistry and Polymer Science, University of Stellenbosch, Private Bag X1, Matieland 7602, South Africa
Correspondence e-mail: ce@sun.ac.za


Received 4 August 2008; accepted 5 August 2008

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å;
 R factor = 0.033; wR factor = 0.080; data-to-parameter ratio = 18.2.

The title compound, $[\text{HfW}(\text{C}_5\text{H}_5)_2(\text{C}_7\text{H}_5\text{O})\text{Cl}(\text{CO})_5]$ or $[\text{W}(\text{CO})_5(\text{C}_7\text{H}_5\text{O})\{\text{Hf}(\text{C}_5\text{H}_5)_2\text{Cl}\}]$, contains two metal centres, with a (tungstenpentacarbonyl)oxyphenylcarbene unit coordinated to a hafnocene chloride. The Hf–O–C angle is nearly linear, and the C=O distance is slightly shorter than for equivalent alkoxy carbene. One of the cyclopentadienyl (Cp) rings undergoes an offset face-to-face π – π interaction [3.495 (7) Å] with the symmetry-related Cp ring of a neighbouring molecule.

Related literature

For related literature regarding anionic Fischer-type carbenes, see: Barluenga & Fañanás (2000); Brüll *et al.* (2001). For comparable structures, see: Berlekamp *et al.* (1993); Erker *et al.* (1989, 1991). For comparable bond lengths, see: Orpen *et al.* (1989).



Experimental

Crystal data

$[\text{HfW}(\text{C}_5\text{H}_5)_2(\text{C}_7\text{H}_5\text{O})\text{Cl}(\text{CO})_5]$
 $M_r = 773.13$
Monoclinic, $P2_1/c$
 $a = 8.5422$ (2) Å
 $b = 12.5546$ (3) Å
 $c = 21.0237$ (7) Å
 $\beta = 96.152$ (1)°

$V = 2241.68$ (11) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 9.91$ mm⁻¹
 $T = 173$ (2) K
 $0.33 \times 0.27 \times 0.25$ mm

Data collection

Nonius KappaCCD diffractometer
Absorption correction: multi-scan (*DENZO-SMN*; Otwinowski & Minor, 1997)
 $T_{\min} = 0.056$, $T_{\max} = 0.089$
(expected range = 0.053–0.084)

12410 measured reflections
5106 independent reflections
4234 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.079$
 $S = 1.01$
5106 reflections

280 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 2.61$ e Å⁻³
 $\Delta\rho_{\min} = -1.76$ e Å⁻³

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *DENZO-SMN* (Otwinowski & Minor, 1997); data reduction: *DENZO-SMN*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001; Atwood & Barbour, 2003); software used to prepare material for publication: *publCIF* (Westrip, 2008).

We thank the NRF and the University of Stellenbosch for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2291).

References

- Atwood, J. L. & Barbour, L. J. (2003). *Cryst. Growth Des.* **3**, 3–8.
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Barluenga, J. & Fañanás, F. J. (2000). *Tetrahedron*, **56**, 4597–4628.
Berlekamp, M., Erker, G. & Petersen, J. L. (1993). *J. Organomet. Chem.* **458**, 97–103.
Brüll, R., Kgosane, D., Neveling, A., Pasch, H., Raubenheimer, H. G., Sanderson, R. & Wahner, U. M. (2001). *Macromol. Symp.* **165**, 11–18.
Erker, G., Dorf, U., Lecht, R., Ashby, M. T., Aulbach, M., Schlund, R., Krüger, C. & Mynott, R. (1989). *Organometallics*, **8**, 2037–2044.
Erker, G., Pfaff, R., Krüger, C. & Werner, S. (1991). *Organometallics*, **10**, 3559–3568.
Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
Orpen, A. G., Brammer, L., Allen, F. H., Kennard, O., Watson, D. G. & Taylor, R. (1989). *J. Chem. Soc. Dalton Trans.* pp. S1–S3.
Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Westrip, S. P. (2008). *publCIF*. In preparation.

‡ Currently at Indus Consulting, PO Box 67283, Centurion, 0169, South Africa.

supplementary materials

Acta Cryst. (2008). E64, m1150 [doi:10.1107/S1600536808025245]

Pentacarbonyl-2 κ^5 C-chlorido-1 κ Cl-bis[1(η^5)-cyclopentadienyl][μ -oxido(phenyl)methylene-1:2 κ^2 O:C]hafnium(IV)tungsten(0)

C. Esterhuysen, I. B. J. Nel and S. Cronje

Comment

Anionic Fischer-type carbene ligands, prepared by the standard addition of organolithium compounds to metal carbonyls, act as monodentate ligands towards transition metals like Ti and Zr (Barluenga and Fañanás, 2000). When the second metal unit is a zirconocene moiety, Cp₂ZrCl, then such complexes have been proven to catalyse the oligomerization of 1-pentene in the presence of methylaluminoxane, MAO (Brüll *et al.*, 2001). Herein, we report the Hf equivalent of these zirconocene alkoxy carbene complexes.

In the title compound, (I, Fig. 1), the W=C_{carbene} and C_{carbene}—C distances are similar to those found in alkoxy carbene complexes, whereas the C—O distance is shorter [2.16 (1), 1.50 (3) and 1.47 (2) Å, respectively; Orpen *et al.*, 1989]. The Hf—O distance is also shorter than those in the metallocyclic compounds C₂₆H₂₇HfO₅V [2.063 (3) Å; Erker *et al.*, 1991] and C₂₈H₂₉HfO₅V [2.066 (3) Å; Berlekamp *et al.*, 1993]. The Hf—O—C angle is nearly linear, with a larger value [171.4 (3)°] than the equivalent Hf—O—C angles of 163.6 (3) and 169.0 (3)° in C₂₆H₂₇HfO₅V (Erker *et al.*, 1991) and C₂₈H₂₉HfO₅V (Berlekamp *et al.*, 1993), respectively, as well as the Zr—O—C angle of 166.1 (5)° in W(CO)₅C(C₆H₅)OZr(C₅H₅)₂OC₆H₅ (Erker *et al.*, 1989).

The C21/C22/C23/C24/C25 Cp ring [with centroid Cg(1)] undergoes offset face-to-face π – π interactions with the symmetry related Cp ring on a neighbouring molecule [$Cg(1)\cdots Cg(1)^i = 3.495$ (7) Å; Symmetry code: (i) 1 - x , 2 - y , 1 - z].

Experimental

A solution of LiCH₃ (31 ml, 1.6M) in diethylether (50 ml) was added to a well stirred suspension of W(CO)₆ (17.802 g) in diethylether (100 ml). After solvent removal, dissolution of the residue in cold water (150 ml) and filtration, a solution of Et₄NCl (8.721 g) in cold water (50 ml) was added to the filtrate. Upon further filtration 0.740 g of the product {[W(CO)₅C(C₆H₅)O][NEt₄]}) was dissolved in dichloromethane (70 ml) and added to a solution of Cp₂HfCl₂ (0.505 g) in dichloromethane (40 ml). After stirring for 30 min at -40°C AgBF₄ (0.261 g) was added. The solvent was removed and the residue extracted in 5 portions of 10 ml toluene. The extract was cooled to -40°C and filtered. The filtrate was dried over anhydrous MgSO₄, concentrated to saturation, and kept at -6°C, whereupon red crystals of the title compound suitable for X-ray diffraction analysis were obtained in 19% yield.

Refinement

H atoms were positioned geometrically with C—H = 0.95 Å and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The maximum and minimum residual electron density peaks were located 0.93 and 0.83 Å, respectively from the Hf1 and W1 atoms.

supplementary materials

Figures




Fig. 1. The molecular structure of I showing the atomic labelling scheme and displacement ellipsoids drawn at the 50% probability level.

Pentacarbonyl-2 κ^5 C-chlorido-1 κ Cl-bis[1(η^5)-cyclopentadienyl][μ -oxido(phenyl)methylene-1:2 κ^2 O:C]hafnium(IV)tungsten(0)

Crystal data

[HfW(C ₅ H ₅) ₂ (C ₇ H ₅ O)Cl(CO) ₅]	$F_{000} = 1432$
$M_r = 773.13$	$D_x = 2.291 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 8.5422 (2) \text{ \AA}$	Cell parameters from 12410 reflections
$b = 12.5546 (3) \text{ \AA}$	$\theta = 1.9\text{--}27.5^\circ$
$c = 21.0237 (7) \text{ \AA}$	$\mu = 9.91 \text{ mm}^{-1}$
$\beta = 96.152 (1)^\circ$	$T = 173 (2) \text{ K}$
$V = 2241.68 (11) \text{ \AA}^3$	Prism, red
$Z = 4$	$0.33 \times 0.27 \times 0.25 \text{ mm}$

Data collection

Nonius KappaCCD diffractometer	5106 independent reflections
Radiation source: fine-focus sealed tube	4234 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.049$
$T = 173(2) \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
φ and ω scans to fill Ewald sphere	$\theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan (DENZO-SMN; Otwinowski & Minor, 1997)	$h = -11 \rightarrow 8$
$T_{\text{min}} = 0.056$, $T_{\text{max}} = 0.089$	$k = -16 \rightarrow 15$
12410 measured reflections	$l = -26 \rightarrow 27$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.032$	H-atom parameters constrained

$wR(F^2) = 0.079$	$w = 1/[\sigma^2(F_o^2) + (0.0436P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.01$	$(\Delta/\sigma)_{\max} = 0.002$
5106 reflections	$\Delta\rho_{\max} = 2.61 \text{ e \AA}^{-3}$
280 parameters	$\Delta\rho_{\min} = -1.76 \text{ e \AA}^{-3}$
Primary atom site location: heavy-atom method	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Hf1	0.80934 (3)	0.779762 (17)	0.542399 (9)	0.01909 (8)
W1	0.46096 (3)	0.614746 (17)	0.676791 (9)	0.02237 (8)
C11	1.06982 (17)	0.83902 (12)	0.58304 (7)	0.0346 (3)
O1	0.3696 (5)	0.6436 (4)	0.52755 (18)	0.0401 (11)
O2	0.1151 (5)	0.5226 (4)	0.6774 (2)	0.0515 (12)
O3	0.4999 (6)	0.6018 (4)	0.82830 (19)	0.0530 (13)
O4	0.3679 (6)	0.8575 (4)	0.6959 (2)	0.0525 (13)
O5	0.5903 (7)	0.3811 (4)	0.6553 (2)	0.0590 (15)
O6	0.7432 (5)	0.7258 (3)	0.62527 (15)	0.0235 (8)
C1	0.4097 (6)	0.6334 (4)	0.5806 (3)	0.0250 (12)
C2	0.2402 (7)	0.5579 (5)	0.6782 (3)	0.0324 (13)
C3	0.4897 (7)	0.6036 (5)	0.7740 (3)	0.0328 (14)
C4	0.3955 (7)	0.7690 (5)	0.6888 (3)	0.0305 (14)
C5	0.5421 (7)	0.4640 (5)	0.6637 (3)	0.0321 (14)
C6	0.6979 (7)	0.6775 (4)	0.6743 (2)	0.0228 (11)
C7	0.8265 (6)	0.6792 (4)	0.7293 (2)	0.0218 (11)
C8	0.9276 (7)	0.7663 (5)	0.7384 (3)	0.0298 (13)
H8	0.9138	0.8259	0.7105	0.036*
C9	1.0477 (8)	0.7664 (5)	0.7879 (3)	0.0405 (16)
H9	1.1132	0.8273	0.7950	0.049*
C10	1.0734 (7)	0.6781 (6)	0.8273 (2)	0.0384 (16)
H10	1.1578	0.6776	0.8606	0.046*
C11	0.9740 (8)	0.5901 (5)	0.8175 (3)	0.0368 (15)
H11	0.9918	0.5289	0.8439	0.044*
C12	0.8507 (7)	0.5916 (5)	0.7701 (2)	0.0299 (13)
H12	0.7812	0.5324	0.7649	0.036*

supplementary materials

C13	0.7818 (7)	0.9777 (4)	0.5270 (3)	0.0306 (13)
H13	0.8715	1.0229	0.5316	0.037*
C14	0.6886 (7)	0.9488 (5)	0.5760 (3)	0.0320 (13)
H14	0.7068	0.9681	0.6199	0.038*
C15	0.5638 (7)	0.8859 (4)	0.5475 (3)	0.0308 (13)
H15	0.4799	0.8578	0.5685	0.037*
C16	0.5844 (7)	0.8717 (5)	0.4830 (3)	0.0326 (14)
H16	0.5187	0.8310	0.4528	0.039*
C17	0.7188 (7)	0.9281 (5)	0.4710 (3)	0.0333 (14)
H17	0.7605	0.9320	0.4309	0.040*
C18	0.7079 (8)	0.6358 (5)	0.4693 (3)	0.0393 (16)
H18	0.5976	0.6332	0.4569	0.047*
C19	0.8207 (8)	0.6914 (5)	0.4373 (3)	0.0395 (16)
H19	0.8003	0.7322	0.3992	0.047*
C20	0.9681 (8)	0.6753 (6)	0.4722 (3)	0.0413 (16)
H20	1.0653	0.7037	0.4618	0.050*
C21	0.9484 (9)	0.6108 (5)	0.5246 (3)	0.0459 (18)
H21	1.0292	0.5881	0.5563	0.055*
C22	0.7886 (9)	0.5853 (5)	0.5224 (3)	0.0439 (17)
H22	0.7425	0.5411	0.5520	0.053*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Hf1	0.01979 (13)	0.02021 (13)	0.01715 (11)	0.00217 (9)	0.00150 (8)	0.00140 (8)
W1	0.02045 (13)	0.02513 (14)	0.02122 (12)	-0.00133 (9)	0.00071 (9)	0.00187 (8)
Cl1	0.0253 (8)	0.0370 (8)	0.0403 (7)	-0.0042 (7)	-0.0014 (6)	0.0026 (6)
O1	0.040 (3)	0.053 (3)	0.025 (2)	-0.007 (2)	-0.0049 (18)	0.0019 (19)
O2	0.025 (2)	0.066 (3)	0.065 (3)	-0.013 (2)	0.012 (2)	-0.007 (3)
O3	0.053 (3)	0.084 (4)	0.024 (2)	0.002 (3)	0.008 (2)	0.013 (2)
O4	0.067 (4)	0.036 (3)	0.052 (3)	0.016 (3)	-0.003 (2)	-0.011 (2)
O5	0.074 (4)	0.040 (3)	0.057 (3)	0.024 (3)	-0.020 (3)	-0.009 (2)
O6	0.028 (2)	0.024 (2)	0.0179 (16)	-0.0010 (17)	0.0020 (15)	0.0039 (14)
C1	0.017 (3)	0.023 (3)	0.034 (3)	-0.007 (2)	0.002 (2)	0.000 (2)
C2	0.033 (4)	0.033 (3)	0.031 (3)	0.000 (3)	0.006 (2)	-0.002 (3)
C3	0.028 (3)	0.040 (4)	0.030 (3)	-0.007 (3)	0.003 (2)	0.004 (3)
C4	0.028 (3)	0.038 (4)	0.025 (3)	0.001 (3)	0.001 (2)	-0.005 (3)
C5	0.026 (3)	0.039 (4)	0.029 (3)	0.003 (3)	-0.010 (2)	-0.001 (3)
C6	0.031 (3)	0.016 (3)	0.021 (2)	0.003 (2)	0.003 (2)	0.005 (2)
C7	0.017 (3)	0.027 (3)	0.022 (2)	0.005 (2)	0.002 (2)	-0.001 (2)
C8	0.023 (3)	0.034 (3)	0.032 (3)	-0.007 (3)	0.004 (2)	0.004 (3)
C9	0.036 (4)	0.051 (4)	0.034 (3)	-0.013 (3)	0.000 (3)	-0.005 (3)
C10	0.027 (3)	0.065 (5)	0.021 (3)	0.011 (3)	-0.004 (2)	-0.005 (3)
C11	0.032 (3)	0.050 (4)	0.027 (3)	0.007 (3)	0.000 (2)	0.005 (3)
C12	0.035 (3)	0.030 (3)	0.025 (3)	0.001 (3)	0.005 (2)	0.008 (2)
C13	0.028 (3)	0.014 (3)	0.051 (3)	-0.002 (2)	0.008 (3)	0.010 (2)
C14	0.037 (4)	0.026 (3)	0.033 (3)	0.005 (3)	0.005 (3)	0.003 (3)
C15	0.021 (3)	0.028 (3)	0.045 (3)	0.010 (2)	0.010 (3)	0.014 (3)

C16	0.029 (3)	0.030 (3)	0.036 (3)	0.006 (3)	-0.011 (3)	0.005 (2)
C17	0.029 (3)	0.038 (3)	0.033 (3)	0.006 (3)	0.005 (2)	0.019 (3)
C18	0.037 (4)	0.047 (4)	0.033 (3)	-0.009 (3)	0.004 (3)	-0.021 (3)
C19	0.047 (4)	0.049 (4)	0.023 (3)	0.004 (3)	0.005 (3)	-0.006 (3)
C20	0.030 (4)	0.050 (4)	0.046 (3)	0.010 (3)	0.012 (3)	-0.011 (3)
C21	0.055 (5)	0.038 (4)	0.043 (4)	0.018 (4)	0.002 (3)	-0.007 (3)
C22	0.071 (5)	0.022 (3)	0.041 (3)	-0.007 (3)	0.017 (3)	-0.013 (3)

Geometric parameters (Å, °)

Hf1—O6	2.006 (3)	C8—H8	0.9500
Hf1—Cl1	2.4139 (14)	C9—C10	1.387 (9)
Hf1—C16	2.464 (5)	C9—H9	0.9500
Hf1—C17	2.465 (5)	C10—C11	1.395 (9)
Hf1—C18	2.469 (6)	C10—H10	0.9500
Hf1—C21	2.479 (6)	C11—C12	1.371 (8)
Hf1—C22	2.480 (6)	C11—H11	0.9500
Hf1—C20	2.483 (6)	C12—H12	0.9500
Hf1—C19	2.484 (5)	C13—C17	1.389 (8)
Hf1—C14	2.494 (6)	C13—C14	1.415 (8)
Hf1—C15	2.496 (5)	C13—H13	0.9500
Hf1—C13	2.514 (5)	C14—C15	1.408 (8)
W1—C2	2.019 (6)	C14—H14	0.9500
W1—C1	2.037 (6)	C15—C16	1.397 (8)
W1—C3	2.038 (6)	C15—H15	0.9500
W1—C4	2.040 (6)	C16—C17	1.395 (8)
W1—C5	2.044 (6)	C16—H16	0.9500
W1—C6	2.177 (6)	C17—H17	0.9500
O1—C1	1.137 (6)	C18—C22	1.401 (9)
O2—C2	1.156 (7)	C18—C19	1.417 (9)
O3—C3	1.135 (7)	C18—H18	0.9500
O4—C4	1.149 (7)	C19—C20	1.403 (9)
O5—C5	1.139 (7)	C19—H19	0.9500
O6—C6	1.291 (6)	C20—C21	1.393 (9)
C6—C7	1.508 (7)	C20—H20	0.9500
C7—C8	1.394 (8)	C21—C22	1.398 (10)
C7—C12	1.397 (7)	C21—H21	0.9500
C8—C9	1.381 (8)	C22—H22	0.9500
O6—Hf1—Cl1	97.55 (11)	O3—C3—W1	176.2 (6)
O6—Hf1—C16	108.77 (18)	O4—C4—W1	175.9 (6)
Cl1—Hf1—C16	132.40 (15)	O5—C5—W1	178.1 (6)
O6—Hf1—C17	133.27 (18)	O6—C6—C7	110.4 (5)
Cl1—Hf1—C17	101.53 (15)	O6—C6—W1	123.3 (4)
C16—Hf1—C17	32.89 (19)	C7—C6—W1	126.1 (4)
O6—Hf1—C18	100.41 (18)	C8—C7—C12	119.0 (5)
Cl1—Hf1—C18	134.01 (16)	C8—C7—C6	120.5 (5)
C16—Hf1—C18	79.9 (2)	C12—C7—C6	120.5 (5)
C17—Hf1—C18	96.2 (2)	C9—C8—C7	120.3 (6)
O6—Hf1—C21	91.47 (19)	C9—C8—H8	119.9

supplementary materials

Cl1—Hf1—C21	83.03 (18)	C7—C8—H8	119.9
C16—Hf1—C21	133.2 (2)	C8—C9—C10	120.4 (6)
C17—Hf1—C21	132.8 (2)	C8—C9—H9	119.8
C18—Hf1—C21	54.7 (2)	C10—C9—H9	119.8
O6—Hf1—C22	78.04 (18)	C9—C10—C11	119.4 (5)
Cl1—Hf1—C22	114.19 (19)	C9—C10—H10	120.3
C16—Hf1—C22	109.7 (2)	C11—C10—H10	120.3
C17—Hf1—C22	129.0 (2)	C12—C11—C10	120.3 (6)
C18—Hf1—C22	32.9 (2)	C12—C11—H11	119.9
C21—Hf1—C22	32.7 (2)	C10—C11—H11	119.9
O6—Hf1—C20	124.08 (19)	C11—C12—C7	120.6 (6)
Cl1—Hf1—C20	80.31 (16)	C11—C12—H12	119.7
C16—Hf1—C20	113.1 (2)	C7—C12—H12	119.7
C17—Hf1—C20	101.1 (2)	C17—C13—C14	107.8 (5)
C18—Hf1—C20	54.6 (2)	C17—C13—Hf1	71.9 (3)
C21—Hf1—C20	32.6 (2)	C14—C13—Hf1	72.8 (3)
C22—Hf1—C20	54.1 (2)	C17—C13—H13	126.1
O6—Hf1—C19	131.56 (19)	C14—C13—H13	126.1
Cl1—Hf1—C19	109.27 (17)	Hf1—C13—H13	121.0
C16—Hf1—C19	82.0 (2)	C15—C14—C13	107.0 (5)
C17—Hf1—C19	80.5 (2)	C15—C14—Hf1	73.7 (3)
C18—Hf1—C19	33.3 (2)	C13—C14—Hf1	74.3 (3)
C21—Hf1—C19	54.5 (2)	C15—C14—H14	126.5
C22—Hf1—C19	54.5 (2)	C13—C14—H14	126.5
C20—Hf1—C19	32.8 (2)	Hf1—C14—H14	117.6
O6—Hf1—C14	82.99 (16)	C16—C15—C14	108.5 (5)
Cl1—Hf1—C14	91.77 (15)	C16—C15—Hf1	72.4 (3)
C16—Hf1—C14	54.65 (19)	C14—C15—Hf1	73.5 (3)
C17—Hf1—C14	54.37 (19)	C16—C15—H15	125.8
C18—Hf1—C14	132.1 (2)	C14—C15—H15	125.8
C21—Hf1—C14	171.9 (2)	Hf1—C15—H15	120.1
C22—Hf1—C14	149.4 (2)	C17—C16—C15	107.6 (5)
C20—Hf1—C14	152.4 (2)	C17—C16—Hf1	73.6 (3)
C19—Hf1—C14	133.5 (2)	C15—C16—Hf1	74.9 (3)
O6—Hf1—C15	80.05 (17)	C17—C16—H16	126.2
Cl1—Hf1—C15	124.55 (15)	C15—C16—H16	126.2
C16—Hf1—C15	32.70 (19)	Hf1—C16—H16	117.3
C17—Hf1—C15	54.02 (19)	C13—C17—C16	109.0 (5)
C18—Hf1—C15	100.2 (2)	C13—C17—Hf1	75.7 (3)
C21—Hf1—C15	151.8 (2)	C16—C17—Hf1	73.5 (3)
C22—Hf1—C15	119.2 (2)	C13—C17—H17	125.5
C20—Hf1—C15	145.3 (2)	C16—C17—H17	125.5
C19—Hf1—C15	112.9 (2)	Hf1—C17—H17	117.1
C14—Hf1—C15	32.78 (19)	C22—C18—C19	107.5 (6)
O6—Hf1—C13	114.49 (16)	C22—C18—Hf1	74.0 (3)
Cl1—Hf1—C13	79.09 (14)	C19—C18—Hf1	73.9 (3)
C16—Hf1—C13	54.2 (2)	C22—C18—H18	126.3
C17—Hf1—C13	32.39 (19)	C19—C18—H18	126.3
C18—Hf1—C13	128.5 (2)	Hf1—C18—H18	117.8

supplementary materials

C21—Hf1—C13	150.0 (2)	C20—C19—C18	107.4 (6)
C22—Hf1—C13	161.3 (2)	C20—C19—Hf1	73.6 (3)
C20—Hf1—C13	119.7 (2)	C18—C19—Hf1	72.8 (3)
C19—Hf1—C13	109.8 (2)	C20—C19—H19	126.3
C14—Hf1—C13	32.82 (18)	C18—C19—H19	126.3
C15—Hf1—C13	53.89 (19)	Hf1—C19—H19	119.2
C2—W1—C1	87.3 (2)	C21—C20—C19	108.7 (6)
C2—W1—C3	88.5 (2)	C21—C20—Hf1	73.5 (4)
C1—W1—C3	173.9 (2)	C19—C20—Hf1	73.6 (3)
C2—W1—C4	93.7 (2)	C21—C20—H20	125.7
C1—W1—C4	88.9 (2)	C19—C20—H20	125.7
C3—W1—C4	86.9 (2)	Hf1—C20—H20	119.0
C2—W1—C5	90.3 (2)	C20—C21—C22	107.9 (6)
C1—W1—C5	90.7 (2)	C20—C21—Hf1	73.9 (4)
C3—W1—C5	93.8 (2)	C22—C21—Hf1	73.7 (4)
C4—W1—C5	175.9 (2)	C20—C21—H21	126.1
C2—W1—C6	179.2 (2)	C22—C21—H21	126.1
C1—W1—C6	92.1 (2)	Hf1—C21—H21	118.4
C3—W1—C6	92.1 (2)	C21—C22—C18	108.6 (6)
C4—W1—C6	85.8 (2)	C21—C22—Hf1	73.6 (4)
C5—W1—C6	90.1 (2)	C18—C22—Hf1	73.1 (4)
C6—O6—Hf1	171.4 (3)	C21—C22—H22	125.7
O1—C1—W1	174.9 (5)	C18—C22—H22	125.7
O2—C2—W1	177.5 (5)	Hf1—C22—H22	119.4

supplementary materials

Fig. 1

