




RESEARCH ARTICLE

Cognitive enhancement: Effects of methylphenidate, modafinil, and caffeine on latent memory and resting state functional connectivity in healthy adults

Maxi Becker^{1,2}  | Dimitris Repantis³ | Martin Dresler⁴  | Simone Kühn^{2,5} ¹Department of Psychology, Humboldt-University Berlin, Berlin, Germany²Department of Psychiatry and Psychotherapy, University Medical Center Hamburg-Eppendorf, Hamburg, Germany³Department of Psychiatry and Psychotherapy, Charité – Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin, Humboldt-Universität zu Berlin, Berlin, Germany⁴Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Center Nijmegen, Nijmegen, The Netherlands⁵Lise Meitner Group for Environmental Neuroscience, Max Planck Institute for Human Development, Berlin, Germany**Correspondence**

Maxi Becker, Department of Psychology, Humboldt-University Berlin, Berlin, Germany.
Email: maxi_becker@gmx.net

Dimitris Repantis, Department of Psychiatry and Psychotherapy, Charité – Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin, Humboldt-Universität zu Berlin, Campus Benjamin Franklin, Berlin, Germany.
Email: dimitris.repantis@charite.de

Funding information

Einstein Foundation Berlin, Grant/Award Number: EPP-2017-423; European Research Council, Grant/Award Number: ERC-2016-StG-Self-Control-677804; German Science Foundation, Grant/Award Number: DFG SFB 936/C7; Jakobs Foundation, Grant/Award Number: JRF 2016-2018; Volkswagen Foundation, Grant/Award Number: 85648

Abstract

Stimulants like methylphenidate, modafinil, and caffeine have repeatedly shown to enhance cognitive processes such as attention and memory. However, brain-functional mechanisms underlying such cognitive enhancing effects of stimulants are still poorly characterized. Here, we utilized behavioral and resting-state fMRI data from a double-blind randomized placebocontrolled study of methylphenidate, modafinil, and caffeine in 48 healthy male adults. The results show that performance in different memory tasks is enhanced, and functional connectivity (FC) specifically between the frontoparietal network (FPN) and default mode network (DMN) is modulated by the stimulants in comparison to placebo. Decreased negative connectivity between right prefrontal and medial parietal but also between medial temporal lobe and visual brain regions predicted stimulant-induced latent memory enhancement. We discuss dopamine's role in attention and memory as well as its ability to modulate FC between large-scale neural networks (e.g., FPN and DMN) as a potential cognitive enhancement mechanism.

KEYWORDS

caffeine, memory, methylphenidate, modafinil, neuroenhancement, neuroimaging, resting state functional connectivity

1 | INTRODUCTION

The umbrella term *cognitive enhancement* refers to interventions by which healthy individuals attempt to improve their cognitive

functions, for example, attention, cognitive control, or memory (Repantis et al., 2010). Next to cognitive training, the use of psychoactive substances has become a popular way to obtain cognitive enhancement (Franke & Lieb, 2010; Talbot, 2009). Caffeine (CAF) in

Maxi Becker and Dimitris Repantis contributed equally to this study.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2022 The Authors. *Human Brain Mapping* published by Wiley Periodicals LLC.

the form of caffeine-containing beverages is probably one of the most widely used prescription-free psychoactive substance worldwide (Ferré, 2008). But also prescription-stimulants such as methylphenidate (MPH, Ritalin®) and modafinil (MOD, Vigil®) are increasingly used by healthy individuals in the attempt to enhance cognition (Battleday & Brem, 2015; Compton et al., 2018). The mode of action as well as different cognitive enhancement effects of all three stimulants have been subject to several studies.

The natural stimulant CAF acts as a nonselective antagonist by blocking adenosine A₁ and A₂ receptors increasing energy metabolism (Koppelstaetter et al., 2008; Nehlig, 2010). Caffeine is most often used and shows strong effects as countermeasure to prolonged wakefulness (Walsh et al., 1990; for a review see Irwin et al., 2020). However, beneficial effects of CAF have been also reported on sustained attention, processing speed, vigilance, and memory (Borota et al., 2014; Koppelstaetter et al., 2008; Nehlig, 2010; Ullrich et al., 2015; for a critical review on the cognitive effects of caffeine, see James, 2014; James & Keane, 2007).

MPH functions as a catecholamine reuptake inhibitor increasing extracellular dopamine in prefrontal, striatal, and hippocampal regions and noradrenaline specifically in frontal brain regions by blocking their respective transporters due to binding to it (Berridge et al., 2006; Kuczenski & Segal, 1997; Spencer et al., 2015; Volkow et al., 2001). There is evidence that MPH enhances working memory and memory consolidation while evidence for processing speed, cognitive control, and attention is rather mixed (Caviola & Faber, 2015; Linszen et al., 2014; Repantis et al., 2010 for a review).

Similar to CAF, MOD is a wakefulness-promoting agent, meaning it is most often used to stave off the effects of sleep deprivation or excessive daytime sleepiness (Bastoji & Jouvet, 1988; Boivin et al., 1993; Pigeau et al., 1995; Repantis et al., 2010). Its precise mechanism is not entirely known up to date. Similar to MPH, MOD also elevates extracellular levels of catecholamines through inhibition of dopamine and noradrenaline transporters (Franke et al., 2017). However, MOD is believed to additionally affect other neurotransmitter systems such as promoting glutamate, serotonin, histamine pathways (Minzenberg & Carter, 2008; Repantis et al., 2010). In general, MOD affects frontal lobe areas (Scoriels et al., 2013 for a review) and has been associated with improved attention, vigilance as well as memory and learning (Battleday & Brem, 2015; Linszen et al., 2014; Müller et al., 2013; Randall et al., 2005; Repantis et al., 2010).

Although the cognitive enhancement effect of all three stimulants is heavily dose-dependent (Wood et al., 2014), there seems to be converging evidence that single-dose intake can improve memory function in a healthy population (for different meta-analyses, see Ilieva et al., 2015; Linszen et al., 2014; Repantis et al., 2010). Despite the different primary modes of action of the stimulants, this poses the question whether there is a shared cognitive enhancement effect related to memory. One possible common neurobiological denominator could be that the stimulants directly or indirectly increase the extracellular levels of catecholamines in the brain. Although CAF does not exert its primary actions on the dopamine and noradrenaline systems, unlike MPH and MOD, it is assumed to indirectly modulate

them via its antagonistic adenosine A₁ receptors (Manalo & Medina, 2018; Nehlig, 2010; Volkow et al., 2015). For example, in vitro studies in animals have shown that CAF affects the local release of catecholamines, especially dopamine (Nehlig, 2010).

Although midbrain dopaminergic neurons mostly project to the basal ganglia and prefrontal cortex, evidence suggests that dopamine transmission plays an essential role in shaping large-scale neural networks underlying cognitive functions (Bentivoglio & Morelli, 2005; Birn et al., 2019; Rosenberg et al., 2016). For example, previous work found widespread effects of MPH, MOD and CAF on functional connectivity (FC). Single-dose MPH intake has been found to modulate functional network connectivity related to enhanced attention (Rosenberg et al., 2016) and modulate coupling between prefrontal regions with inhibitory and attention networks related to improved response inhibition (Pauls et al., 2012). Another study found that MPH reduced connectivity between default mode and dorsal attention, ventral attention, and visual networks in healthy adults (Sripada et al., 2013). MOD increased negative coupling between executive and the default mode network (DMN) which was associated with improved cognitive control in alcohol-dependent patients (Schmaal et al., 2013). Single-dose MOD intake has also been found to increase FC between prefrontal and striatal areas (Cera et al., 2014). Finally, single-dose CAF intake was associated with increased frontoparietal network (FPN) activation and FC in different working memory tasks (Haller et al., 2013; Koppelstaetter et al., 2008). It is important to note that the relationship between CAF and BOLD activity is not trivial because CAF acts not only as an excitatory neurostimulant but also as a vasoconstrictor reducing cerebral blood flow (Fredholm et al., 1999). This may explain why several studies also find CAF-induced widespread decreases in FC (Rack-Gomer et al., 2009; Tal et al., 2013; Wong et al., 2012).

However, up to date there are no studies investigating the cognitive enhancement effect of different stimulants in neither memory nor FC. To fill this gap, we used resting state fMRI and memory related behavioral data from a double blind randomized placebocontrolled study (Repantis et al., 2021). In this study, 48 male participants were randomized into three groups receiving a stimulant (CAF, MOD, and MPH) or a placebo first in two separate sessions. Next to fMRI data acquisition, the participants performed a series of cognitive tasks including different memory tasks (see Section 2 and Repantis et al., 2021).

Given prior evidence, we assumed enhanced performance in memory tests as a function of all three stimulants. Due to the reported widespread effects of MPH, MOD, and CAF on FC, we chose an agnostic, data-driven approach employing whole brain FC analyses on resting state fMRI data. However, since all three stimulants have repeatedly shown to affect prefrontal areas, we assumed that stimulant-induced changes in large-scale neural networks should involve prefrontal cortices. Additionally, stimulant-induced FC changes specific to memory should involve medial temporal lobe structures including the hippocampus, due to its consistent association with successful memory formation (Ranganath et al., 2005; for a meta-analysis, see Grady, 2020; Wais, 2008).

2 | METHODS

2.1 | Participants

The study included 48 healthy, right-handed male participants ($M = 26.27$, $SD = 3.47$, range: 21–36). The participants were recruited via internet recruitment and screened for presence of psychiatric or medical disorders as determined by a physician and using the Mini-International Neuropsychiatric Interview (Sheehan et al., 1998). Only male participants were included due to prior evidence showing an interaction of the female hormone cycle with resting state FC as well as performance in memory tasks (Lisofsky et al., 2015). Further exclusion criteria were: regular or excessive consumption of caffeinated drinks (regularly drinking >4 cups per day) within the last 6 months,¹ current usage of any medications, lifetime consumption of prescription stimulants (modafinil, methylphenidate) or illicit *hard* substances (e.g., cocaine, crack, and heroine), consumption of other illicit substances (e.g., speed, amphetamine, THC, ecstasy, and MDMA) within the past year, regular or irregular heavy smoking within the last 5 years (two subjects reported to smoke irregularly and little) as well as current irregular day–night rhythm (e.g., shift-work).

To further control for irregular sleeping patterns, bad sleep quality as well as strong differences in circadian rhythm between the participants, we administered the German translated versions of the Pittsburgh Sleep quality index (PSQI, Buysse et al., 1989) and the Morning-Eveningness Questionnaires (D-MEQ, Griefahn et al., 2001; Horne & Östberg, 1976). According to the PSQI, none of the participants self-reported an unhealthy sleeping pattern ($M = 2.46$, $SD = 1.65$; max = 5; note, subjects with values between 0 and 5 are considered healthy sleepers). Furthermore, according to the D-MEQ, the participants exhibited as similar circadian rhythm ($M = 2.52$, $SD = 0.46$; max = 3.47, min = 1.63; note, the D-MEQ ranges from 0 = clearly “evening person” to 6 = clearly “morning person”).

Written informed consent was obtained from all subjects and they were financially compensated for their time of participation. The study was approved by the Berlin State Ethics Committee (LAGeSo Berlin, Germany; 13/0138-EK12) and conducted according to the Declaration of Helsinki (1964).

2.2 | Study design and procedure

We used a placebocontrolled, randomized, double-blind within-subject study design with three arms in which a stimulant was tested against placebo. Each participant received only one stimulant (and placebo) in the form of a white capsule for oral ingestion and was tested at two different occasions at the same time of the day in the early afternoon. The sessions were separated by about 1 week. The stimulants that were used were immediate-release MPH (20 mg), MOD

(200 mg), and CAF (200 mg). The doses were chosen based on known equipotency in clinical practice and prior trials as well as (for MPH and MOD) similar occupancy of the dopamine transporter (Franke et al., 2017; Repantis et al., 2010; Theunissen et al., 2009; Volkow et al., 1998, 2009).

The white capsules containing one of the stimulants or the placebo (=microcrystalline cellulose) had been prepared by a pharmacy of the Charité (Berlin) in such a manner that they looked identical and weighed the same. Participants encoded the memory task with visual material in the MRI scanner and completed the rest of the tasks (including retrieval of this memory task) afterwards outside of the scanner (see Section 2.3). The start of the fMRI procedure was 90 min after oral substance ingestion. This timeframe was chosen so that all three stimulants had reached approximately their peak concentration in blood during testing. Two resting state scans were acquired directly before and after the encoding of the memory task with visual material in the MRI scanner, hence, 90 and 120 min after oral substance ingestion. The reason why we acquired two instead of one resting state scan was to account for general within-subject resting state variability (Bijsterbosch et al., 2017) and potential within-subject variation in elimination half-life of the stimulants (Wagner, 1973).

2.3 | Behavioral measures of memory

In the original study two different memory tasks were administered—memory of visual and audio material (see Repantis et al., 2021). For better comparability, we used the same memory outcome measures as reported in Repantis et al. (2021).

2.3.1 | Memory task of visual material

In a declarative memory task, participants were exposed to 72 common nouns. The encoding difficulty was matched between lists and pretested. Based on previous research, word lists counting about 70 words prevent ceiling effects (Riedel & Blokland, 2015). The words were presented in 12 blocks (six words each) inside the scanner. Each word was presented for 2000 ms with a jittered inter stimulus interval of 2–5 s. Blocks were interspersed with fixation periods. An early recall was performed outside the scanner ~20 min after learning phase.

Memory task of auditory material

Implicit and explicit verbal memory was measured with a false memory test outside of the scanner (Roediger & McDermott, 1995). Participants heard five sets of 15 words each (75 words in total). Every set contained semantically similar words that could be associated with one critical lure which was not presented itself. As an example, the presented words “drive,” “street,” “key,” “garage,” and so on are all associated to the lure “car.” After the presentation an early recall was performed outside the scanner. Afterwards, participants underwent a recognition test in which a list of 40 words was presented. These

¹Participants were instructed to refrain from drinking any caffeinated drinks throughout the entire length of the study. However, this affected only three participants who reported to irregularly consume few caffeinated drinks (45 participants reported not to consume any caffeinated drinks currently at all).

included 20 words from the auditory material word list and 20 new words with the lures included among them. A sensitivity index d' was calculated using the formula $d' = Z(\text{hit rate}) - Z(\text{false alarm rate})$ in order to examine how well participants discriminated between old and new words. Hit rate was defined as $(\text{hits}/(\text{hits} + \text{misses}))$ (hits were the correctly identified old words; misses were the old words that were not recognized; Macmillan & Creelman, 1990). False alarm rate was defined as $(\text{false alarms}/(\text{false alarms} + \text{correct negative}))$ (false alarms were new words that were falsely identified as old words; correct negative were new words that were correctly identified as new words). For the Z transformation the formula $\text{NORM.S.INV}(\text{hit}) - \text{NORM.S.INV}(\text{false alarm})$ was used. Perfect scores were adjusted using the formula $1 - 1/(2n)$ for perfect hits and $1/(2n)$ for zero false alarms, with n being the number of total hits or false alarms (20 and 20, respectively) (Haatveit et al., 2010; Macmillan & Creelman, 1990). A higher d' score indicates that the signal was recognized better.

Retention of information was assessed for both memory tasks with visual and auditory material using a delayed free recall test for the two sets of learned words. This was conducted by telephone 24 h after the session. Participants were informed about the upcoming late recall task but were asked not to actively try to retain the words.

2.4 | Behavioral data analysis

2.4.1 | Analysis of single memory tasks

To analyze the stimulant-induced changes on memory, we first analyzed the outcome measures for each memory task separately in a General Linear Mixed Model (GLMM) (for a discussion of the advantages using the GLMM approach, see Barr et al., 2013; Baayen et al., 2008). The memory measures which served as dependent variables for the single GLMMs were (1) d' from the implicit memory task; (2–3) the amount of correctly remembered items in the memory task of visual material during early (early_visRecall) and late recall (late_visRecall); (4–5) the amount of correctly remembered items in the memory task of audio material during early (early_audRecall) and late recall (late_audRecall) and (6–7) the amount of falsely remembered critical lures (false memories) in the memory task of audio material during early (early_false_audRecall) and late recall (late_false_audRecall). The d' was modeled assuming a Gaussian error distribution. Except for d' , all other outcome measures (2–7) were count variables which is why we modeled those variables assuming a Poisson error distribution of the model residuals with the default log link function (Gardner et al., 1995).

The main independent variable of interest was the condition (binary: stimulant vs. placebo). Other covariates of no interest included into the model were (1) the group (a categorical variable indicating the specific stimulant group MOD, MPH or CAF) and (2) the medication order (a categorical variable indicating in which order the participants received either the placebo or the stimulant first). Subjects were modeled as random intercepts.

To assess the influence of the condition (stimulant vs. placebo) on all seven memory outcome measures, we performed likelihood ratio tests of the full model with the condition variable against the baseline model without the condition variable. The baseline model included the group variable, order (stimulant or placebo in first session) and the random subject effect. The full model was identical to the baseline model with the exception of the additional condition variable. The p -values for the individual models were obtained from likelihood ratio tests. Additionally, all obtained p -values from the likelihood ratio tests were corrected for multiple comparisons using the false discovery rate (Benjamini & Hochberg, 1995).

All statistical analyses were carried out in R (version 4.0.3, R Core Team, 2014). The GLMMs and linear models were estimated using the glmmTMB package (version 1.0.2.1; see Brooks et al., 2017). Visual inspection of residual plots did not reveal any obvious deviations from homoscedasticity or normality for all models. The behavioral data, the extracted within-connectivity values per cluster as well as the analysis code are publicly available on github <https://github.com/MaxiBecker/Neuroenhancement>.

2.4.2 | Measurement model for latent memory change factor

To relate stimulant-induced changes in memory to changes in resting state FC, we calculated a latent memory change factor from the difference scores (stimulant – placebo) of the individual memory measures. The latent memory change factor represents the shared covariance of the individual memory tasks and is therefore a more robust measure for memory change as a function of the stimulants. We included the difference values of the following four memory measures into the final model: d' , early_audRecall, late_audRecall, early_visRecall (see Section 2.4.1). To reach a good model fit, we excluded the difference values of the following variables: late_visRecall, early_false_audRecall and late_false_audRecall and fixed the error variance of the variable early_audRecall to one (see Figure 3).

The latent factor was estimated within a confirmatory factor analysis (CFA) in R using the lavaan package (version 0.6–7, Yves Rosseel, 2012). Model fit was evaluated via multiple fit indices: the χ^2 goodness-of-fit statistic, Bentler's comparative fit index (CFI), root mean square error of approximation (RMSEA) and standardized root-mean-square residual (SRMR). Accepted thresholds indicating good model fit are $\text{RMSEA} \leq .05$, $\text{SRMR} < 0.1$ and $\text{CFI} \geq .95$ (Hu & Bentler, 1998, 1999; Schermelleh-Engel et al., 2014).

2.5 | rs-fMRI parameters

Imaging was performed on a Siemens 3T Magnetom Trio Scanner (Siemens Healthcare, Erlangen, Germany) using an echo planar protocol with a 12-channel radiofrequency head coil. Functional resting

state images were obtained using a T2*-weighted echo planar imaging (EPI) sequence sensitive to blood oxygen level dependent (BOLD) contrast (TR = 2000 ms, TE = 30 ms, slice thickness = 3 mm, image matrix = 64×64 , FOV = 216 mm, flip angle = 80° , voxel size = 3 mm^3 , 36 axial slices). The participants were instructed to relax and keep their eyes closed for both resting state scans. Each scan contained a total of 150 volumes.

Structural images were collected using a three-dimensional T1-weighted magnetization prepared gradient-echo sequence (MPRAGE) [repetition time (TR) = 2500 ms; echo time (TE) = 4.77 ms; slice thickness = 1 mm, acquisition matrix = $256 \times 256 \times 192$, flip angle = 7° ; FOV = 256 mm, voxel size = 1 mm^3].

2.6 | rs-fMRI preprocessing and denoising

rs-fMRI state data was preprocessed, denoised, and subsequently analyzed using CONN—an open source connectivity toolbox (version 19c, Whitfield-Gabrieli & Nieto-Castanon, 2012).

2.6.1 | Preprocessing

The data was processed using CONN's default preprocessing pipeline which includes the following steps: Functional scans were realigned and slice-time corrected. Realignment includes coregistering all scans and resampling them to the first reference image using b-spline interpolation (see Andersson et al., 2001). Outlier scans were identified using framewise displacement and CONN's default parameters (framewise displacement above 0.9 mm or global BOLD signal changes above 5 SDs).

Functional and structural data was normalized into MNI space and segmented into gray matter, white matter as well as CSF tissue. Finally, the functional data was smoothed using an 8 mm full width half maximum Gaussian kernel to increase BOLD signal to noise-ratio.

2.6.2 | Denoising

Confounding effects to the estimated BOLD signal were first estimated and subsequently removed for each voxel in each subject and for each condition as well as each session using Ordinary Least Squares regression. For this purpose, an anatomical component-based noise correction procedure (CompCor) as implemented in CONN was used. Potential confounding effects included: six estimated subject-motion parameters including their first-order derivatives, noise components from areas of cerebral white matter and cerebrospinal fluid, constant, or linear session effects as well as the abovementioned identified outlier scans. Note, no global signal regression was applied to avoid the risk of artificially introducing anticorrelations (negative connectivity) into the FC estimates (Chai et al., 2012). Furthermore, the effects of denoising were further manually evaluated using CONN's quality control plots. Finally, the resulting time series were additionally band-pass filtered to .008–.09 Hz.

2.7 | rs-fMRI data analysis

2.7.1 | First-level analysis in CONN

We adopted a whole brain data-driven approach because we did not have a specific hypothesis where in the brain FC should increase as a function of all three stimulants. To investigate those stimulant-induced changes in FC on a network level, that is, the whole brain, we parcellated the brain into 268 Regions of Interest (ROIs) using the Shen atlas (Shen et al., 2013). The Shen atlas is parcellated based on resting state FC and therefore ideal for network analysis using FC (Finn et al., 2015; Rosenberg et al., 2016; Shen et al., 2013). For this reason, the Shen atlas was preferred to CONN's default FSL Harvard-Oxford atlas that contains only 132 ROIs derived from structural data (106 cortical and subcortical areas and 26 cerebellar areas from the AAL Atlas) (Finn et al., 2015; Rosenberg et al., 2016; Shen et al., 2013).

The time series for each ROI were acquired by mean averaging the BOLD time series of each voxel that belongs to the respective ROI. ROI-to-ROI connectivity matrices (268×268) were computed for each participant (48), for each condition (drug vs. placebo), each scan (2) and each session (2) separately. Each element in each connectivity matrix represents the strength of FC between a pair of ROIs and was defined as the Fisher-transformed bivariate correlation coefficient between the preprocessed and denoised BOLD time series of two ROIs.

2.7.2 | Second-level analysis in CONN

Rather than focusing on single connections between all possible pairs of ROIs, we adopted a hierarchical cluster analysis as implemented in CONN which categorizes ROIs into clusters based on ROI-to-ROI functional similarity and anatomical proximity metrics (Sørensen, 1948). This method is based on a complete-linkage metric meaning that the distance between two clusters is defined as the maximum distance between all pairs of ROIs within the two clusters. CONN's default distance measure that we used is defined as $D = 0.95 \times D_{\text{func}} + 0.05 \times D_{\text{anat}}$, where D_{func} is the squared Euclidean distance between the connectivity patterns for every pair of ROIs (averaged across all subjects), and D_{anat} represents the squared Euclidean distance between the anatomical centroid coordinates for every pair of ROI.

To infer stimulant-induced changes in FC on these groups/clusters of ROIs, a multivariate parametric General Linear Model was performed for all connections included in each of these groups/clusters of ROIs (in terms of within- and between-network connectivity sets, see Jafri et al., 2008). The resulting *F*-statistic for each pair of related ROI-groups/clusters was thresholded at an FDR-corrected cluster level $p < .05$ (Benjamini & Hochberg, 1995). To characterize the pattern of individual connections, a post hoc uncorrected $p < .05$ height threshold (connection-level) was applied.

To investigate stimulant induced changes on FC, we contrasted the stimulant condition (2 scans) with the placebo condition (2 scans)

averaged for each stimulant group (between subject contrast: $0.33 \times \text{MOD} + 0.33 \times \text{MPH} + 0.33 \times \text{CAF}$).

We additionally investigated whether there were whole-brain stimulant-induced changes in FC that relate to stimulant-induced changes in general memory. For this, we first extracted the subject-specific factor loadings of the latent memory change factor (see Section 2.4.2). Before entering the second level in CONN, the values were normalized, mean centered and orthogonalized to all three stimulant groups. Subsequently, the values were correlated with every group/cluster of related ROIs in the brain while controlling for differences between the groups (between subject contrast: $1 \times \text{Memory} + 0 \times \text{MOD} + 0 \times \text{MPH} + 0 \times \text{CAF}$). Finally, those correlations were contrasted between the stimulant condition (2 scans) and the placebo condition (2 scans).

For exploratory purposes, the resulting (two) clusters from the group analysis in CONN were further examined to investigate to what extent they also relate to change in latent memory performance. To this end, we extracted the within-FC of each cluster (mean average of all ROI-pairs that belong to the respective cluster). For each cluster, we subtracted the resulting within-FC values of the stimulant condition from the placebo condition. Subsequently, the difference values of those clusters were entered into a separate linear model together with three covariates of no interest to predict change in latent memory performance (i.e., the individual factor loadings from latent memory change factor). Due to high collinearity [$r = .53$, 95% CI (0.392–0.649)], the difference values of both clusters could not be entered as predictors into the same linear model. Therefore, we performed two separate linear models each predicting change in latent memory performance. The additional covariates per model were a binary variable indicating the respective scan and the order in which the subjects received either the stimulant or the placebo (binary variable: stimulant first vs. placebo first). The p -values for the individual models were obtained from likelihood ratio tests. That is to say, a baseline model including only the two covariates of no interest [scan and order (stimulant or placebo in first session)] was compared with the full model 1 including the two covariates and the difference value of the first cluster. The baseline model was also compared with the full model 2 including the two covariates and the difference value of the second cluster. To control for multiple comparison, all obtained p -values from the likelihood ratio tests were corrected using the false discovery rate (Benjamini & Hochberg, 1995).

3 | RESULTS

3.1 | Effects of stimulants on memory

Details on the model estimates from the individual GLMMs are depicted in Tables S1–S3.

3.1.1 | Memory task with visual material

On average, participants correctly recalled 35.21 ($SD = 18.24$) words in the stimulant and 31.69 ($SD = 16.88$) words in the placebo

condition. This difference in early memory recall was statistically significant [$Chi^2(1) = 8.622$, $p\text{-FDR} = .008$].

After 24 h, participants correctly recalled 21.53 ($SD = 17.61$) words in the stimulant and 16.77 ($SD = 13.33$) words in the placebo condition on average. This difference in late memory recall was also significant [$Chi^2(1) = 30.87$, $p\text{-FDR} = .000$] (see Figure 1).

3.1.2 | Memory task with audio material

On average, participants correctly recalled 40.60 ($SD = 9.64$) words in the stimulant and 37.21 ($SD = 10.21$) words in the placebo condition. This difference in early memory recall was statistically significant [$Chi^2(1) = 7.116$, $p\text{-FDR} = .013$].

After 24 h, participants correctly recalled 28.96 ($SD = 11.97$) words in the stimulant and 24.35 ($SD = 10.48$) words in the placebo condition on average. This difference in late memory recall was significant [$Chi^2(1) = 19.11$, $p\text{-FDR} = .000$] (see Figure 1).

3.1.3 | Memory task with audio material—false memory

Furthermore, on average participants falsely remembered 0.91 ($SD = 0.89$) lure words in the stimulant condition and 1.38 ($SD = 1.18$) lure words in the placebo condition. This difference in early false memory recall was statistically significant [$Chi^2(1) = 4.43$, $p\text{-FDR} = .041$] (see Figure 1).

After 24 h, participants falsely remembered 1.18 ($SD = 1.05$) lure words in the stimulant and 1.44 ($SD = 1.11$) lure words in the placebo condition. However, this difference in late false memory recall was not statistically significant [$Chi^2(1) = 2.17$, $p\text{-FDR} = .141$].

3.1.4 | Implicit memory task

On average, d' in the stimulant condition was 2.29 ($SD = 0.76$) whereas d' was 2.01 ($SD = .71$) in the placebo condition. This difference was statistically significant [$Chi^2(1) = 5.880$, $p\text{-FDR} = .015$] (see Figure 1).

3.1.5 | Measurement model for latent memory change factor

The model converged normally after 31 iterations. All variables loaded in the same direction (positive). Except for *early_visRecall*, all standardized factor loadings were within moderate to high range (difference values for d' : $\lambda = .479$; *early_audRecall*: $\lambda = .992$; *late_audRecall*: $\lambda = .567$; *early_visRecall*: $\lambda = .181$), suggesting that those variables contributed significantly to the latent memory change factor (see Figure 3, left panel). The exact test statistic did not reach significance ($\chi^2 = 0.588$, $p = .899$), indicating that the model was not significantly different from the data. Additionally, practical fit indices (CFI = 1.0,




FIGURE 1 Marginal means of memory performance in memory tasks with visual and audio material as a function of condition plotted separately for all three stimulant groups. CAF, caffeine; MOD, modafinil; MPH, methylphenidate; error bars represent between-subject 95% confidence intervals; D_{prime} , d' from an implicit memory task; early_audRecall, early recall for memory task with audio material; late_audRecall, late recall for memory task with audio material; early_visRecall, early recall for memory task with visual material; early_false_audRecall, early recall for lures (false memory) in memory task with audio material

RSME = $<.001$, and SRMR = $.027$) also suggested a good fit of the model to our data.

3.2 | Effects of stimulants on FC

The hierarchical cluster analysis revealed two significant clusters of ROIs that were all positively correlated. The first cluster [$F(2,44) = 11.31$, $p\text{-FDR} = .0237$] consisted of 26 connections from predominantly medial parietal regions being functionally connected to right prefrontal regions as a function of all three stimulants combined (see Figure 2, cluster 1). A post hoc analysis of the condition and group specific effect sizes revealed that those parietal-prefrontal regions were negatively correlated in the placebo condition, while this negative correlation was significantly reduced in the stimulant condition (see Figure 2, cluster 2). The second cluster [$F(2,44) = 12.25$, $p\text{-FDR} = .0237$] consisted of 10 connections from predominantly lateral parietal regions that were positively functionally connected to right prefrontal regions (see Figure 2). A post hoc analysis of the condition and group specific effect sizes revealed that FC generally increased as a function of all three stimulants (see Figure 2).

Table S5 lists all ROIs belonging to each cluster including their individual T -statistics. To demonstrate that the results from the hierarchical cluster analysis (as shown in Figure 2) were not just driven by

one stimulant, we additionally plotted both clusters separately for each group, each condition and each scan in Figure S4.

For exploratory purposes, we investigated whether stimulant-induced change in FC in both clusters predicts the amount of latent memory change. The first cluster [$\text{Chi}^2(1) = 7.33$, $p\text{-FDR} = .016$] negatively predicts latent memory change. The negative relationship is reasonable given that the ROI pairs in this cluster in the placebo condition are more negatively connected than in the stimulant condition (see Figure 2, lower left panel). In contrast, FC change in the second cluster [$\text{Chi}^2(1) = 0.07$, $p\text{-FDR} = 0.799$] does not predict latent memory change (see Figure 2, lower right panel). Details on the model estimates from the linear model are presented in Table S4. For the sake of completeness, the relationship between latent memory change and the respective cluster plotted separately for each group is presented in Figure S5.

3.3 | Link between stimulant-induced changes in whole brain rs-FC and memory

When investigating which brain areas correlate more strongly with stimulant-induced changes in latent memory performance in the stimulant compared with the placebo condition, one cluster survived multiple comparisons on a whole brain level [$F(2,43) = 11.75$, p -




FIGURE 2 Effects of caffeine, modafinil, and methylphenidate on functional connectivity. Upper panel: results from hierarchical cluster analysis. Two clusters significantly differed between the stimulant and placebo condition. First and second refer to the first and second resting state scan, respectively. The red lines represent significant correlations of the time series between the regions of interest (gray spheres). Lower panel: change in functional connectivity from the placebo to the stimulant condition correlated significantly negatively with change in latent memory performance from the placebo to the stimulant condition in cluster 1 but not in cluster 2. * indicates significance at a p -level $< .05$ (FDR-corrected)

$FDR = .037$]. This cluster consisted of 16 connections between ROIs in visual (lingual, calcarine and middle occipital cortex) and medial temporal lobe visual areas [hippocampus, parahippocampus (PHC), and superior temporal pole] (see Figure 3). A post hoc analysis revealed that ROIs from both areas were negatively correlated in the placebo condition, while this negative correlation was significantly reduced in the stimulant condition (see Figure 3). Table S5 lists all ROIs belonging to each cluster including their individual t -statistics.

4 | DISCUSSION

In this study, we investigated stimulant-induced effects of MPH, MOD, and CAF on memory and related FC. Despite their different modes of action, all three stimulants are considered cognitive neuroenhancers due to their positive effects on attention, cognitive control and specifically memory (Franke & Lieb, 2010; for a critical review on this matter, see Repantis et al., 2010). However, mnemonic and brain-functional effects of cognitive neuroenhancers (e.g., MPH, MOD, and CAF) are usually studied separately. While this expands our knowledge of stimulant-specific effects on behavior and brain

function, the question remains, whether there are shared cognitive enhancement effects over and above the individual stimulant. We speculated that one cognitive enhancement mechanism could be the stimulants' direct or indirect impact on extracellular levels of catecholamines in the brain. Specifically, dopamine has been shown to shape large-scale neural networks including prefrontal regions underlying cognitive functions. For this reason, we utilized behavioral and resting state fMRI data from a double-blind randomized placebocontrolled study (see Repantis et al., 2021) and investigated the combined effects of MPH, MOD and CAF on memory and FC at rest (Repantis et al., 2021). We hypothesized that memory performance should increase and large-scale neural networks including prefrontal regions should be modulated as a function of all three stimulants. Additionally, we assumed that stimulant-induced FC changes specific to memory should involve medial temporal lobe structures due to their consistent association with memory formation (for a meta-analysis, see Grady, 2020).

As predicted, we found enhanced mnemonic performance in different memory tasks in the stimulant compared with the placebo condition. That is to say, implicit memory measured as d' was enhanced. Correct recall for visual and audio material was also enhanced in the




FIGURE 3 Correlation of stimulant-induced changes in whole-brain rs-FC with changes in latent memory. Left panel: measurement model for latent memory change factor. d' , d prime from implicit memory task; e AR, early recall for memory task with audio material; I AR, late recall for memory task with audio material; e VR, early recall for memory task with visual material; * signifies statistical significance at the $p < .05$ level. Right panel: results from whole brain hierarchical cluster analysis. Negative correlation of occipital–medial temporal lobe cluster with the latent memory change factor is significantly reduced in the stimulant compared with the placebo condition. First and second refer to the first and second resting state scan, respectively. The red lines represent significant correlations of the time series between the regions of interest (gray spheres)

stimulant condition immediately after encoding as well as 24 h later. Furthermore, false memory for lures was reduced in the stimulant-condition, but only after immediate not during late recall. Further congruent with our hypotheses, we found a stimulant-induced effect of increased FC. Hierarchical cluster analysis revealed two main clusters of connections between predominantly prefrontal and parietal ROIs indicating FC change between the FPN and DMN at rest (see Figure 2 and Table S5). In one cluster, FC between predominantly lateral parietal and right prefrontal ROIs was increased in the stimulant compared with the placebo condition. In the other cluster, negative FC between predominantly medial parietal (Precuneus, Posterior Cingulum [PCC]) DMN and right prefrontal FPN ROIs was decreased in the stimulant compared with the placebo condition. A post hoc analysis revealed that the amount of negative connectivity reduction from the stimulant to the placebo condition predicted the amount of performance change in latent memory. On a whole brain level, however, one specific cluster including connections between medial temporal lobe areas (e.g., hippocampus and PHC) also belonging to the DMN and visual areas was significantly related to stimulant-induced latent memory change. That is to say, in this cluster negative FC was significantly reduced in the stimulant compared with the placebo condition.

4.1 | Stimulant-induced changes in coupling between FPN and DMN

The DMN is a set of cortical regions comprising hubs mostly located in the ventromedial prefrontal, inferior parietal lobule and PCC/Precuneus (Greicius et al., 2003). Those nodes exhibit coherent

fluctuations during resting state and have been associated with a range of cognitive processes including self-oriented mental activity and autobiographical as well as episodic memory retrieval (Buckner et al., 2008; Sestieri et al., 2011). Prior studies report that this network can be modulated by different stimulants such as MPH (Liddle et al., 2011; Silberstein et al., 2016) as well as MOD (Minzenberg et al., 2011; Schmaal et al., 2013) and CAF (Wong et al., 2012; Wu et al., 2014).

The FPN comprises regions such as the anterior insular cortex, anterior prefrontal, dorsolateral prefrontal and dorsomedial superior frontal cortex/anterior cingulate and the anterior inferior parietal lobule (Dang et al., 2012). It has been primarily associated with attention and cognitive control (Badre & D'Esposito, 2007; Cabeza et al., 2008; Dodds, Morein-Zamir, & Robbins, 2011; Scolarì, Seidl-Rathkopf, & Kastner, 2015). However, lidaka et al. have shown evidence that this network is also associated with successful retrieval in episodic memory (lidaka et al., 2006). FC within the FPN has been shown to be modulated by different stimulants such as MOD (Esposito et al., 2012; Visser et al., 2019); CAF (Haller et al., 2013; Koppelstaetter et al., 2008) and MPH (Mehta et al., 2000; Mueller et al., 2014). In fact, Schmidt and colleagues report a broad recruitment of frontoparietal regions when MOD and MPH are consumed compared with placebo (Schmidt et al., 2017).

At rest, the DMN typically exhibits anticorrelated activity with the FPN as we have observed in cluster 1 in the placebo condition (Chai et al., 2012; Fox et al., 2005; Uddin et al., 2009). Importantly however, the FPN has been found to flexibly couple with the DMN, depending on the task domain supporting goal-directed cognitive processes (Spreng et al., 2010). Dopaminergic signal transmission may

function as one underlying mechanism in this regard. Evidence suggests that dopamine enhances coupling between the FPN and DMN at rest (Dang et al., 2012). This finding is in line with our assumption that one neurobiological denominator of the cognitive enhancement effect could lie in the stimulants' ability to increase extracellular levels of catecholamines (e.g., dopamine, noradrenaline) in the brain. Catecholaminergic innervations range across the brain with high overlapping concentrations in hippocampus and prefrontal cortex (for review see Ranjbar-Slamloo & Fazlali, 2020). Hence, it may be the ability of specifically FPN's right prefrontal hubs to flexibly couple and decouple from the DMN depending on the task demands that relate to the cognitive enhancement effects associated with all three stimulants (CAF, MOD, and MPH).

Consequently, we found that the amount of negative connectivity reduction between predominantly right prefrontal DMN and medial posterior FPN regions (Precuneus, PCC) significantly predicted stimulant-induced latent memory change. While the right IFG is implicated in attention to facilitate goal-directed behavior (Dodds, Morein-Zamir, & Robbins, 2011), parietal regions of the DMN support episodic memory retrieval (Sestieri et al., 2011). Hence, decreasing the anticorrelation (negative connectivity) between both regions may be associated with increased attention during successful encoding of the stimulus material, leading to enhanced memory performance in the stimulant condition. However, the functional associations coupling between the nodes of the FPN and DMN seem to be specific because we only found a latent memory change related association between medial (not lateral) parietal DMN and right prefrontal FPN regions. This is congruent with studies, showing that the DMN is not a functionally homogenous unity but that different nodes within this network contribute distinctively to cognitive processes like cognitive control (Laird et al., 2009; Leech et al., 2011).

Further evidence, supporting the assumed relationship between catecholamines especially dopamine, FPN-DMN coupling and memory comes from patients with attention-deficit/hyperactivity-disorder (ADHD). ADHD has been associated with weakened prefrontal cortex function (Arnsten, 2006), altered DMN-FPN interactions (Sripada et al., 2014) and deficient attention-related encoding and retrieval processes (Ortega et al., 2020). Importantly, ADHD is successfully treated with MPH by increasing the extracellular concentration of catecholamines in prefrontal, striatal, and other brain areas (Mehta et al., 2000).

4.2 | Memory-related stimulant-induced FC changes between visual and medial temporal lobe regions

Episodic memory is critically dependent on the MTL and its functional connections to the cortex. For example, anatomical studies find many direct connections between the PHC and the DMN (Burwell, 2000; Witter et al., 2000). The MTL is also strongly functionally connected to early and late visual areas (Córdova, Tomparly, & Turk-Browne, 2016; Wang et al., 2016).

Cordova et al. (2016) show that memory retrieval can be facilitated by attention-related increased coupling between visual areas and MTL regions during encoding of visual stimuli. In line with those results, we find stimulant-induced modulations in FC between MTL and visual areas to be directly related to memory enhancement. Those stimulant-induced modulations may also be associated with attention, however, we did not observe increased positive but decreased negative connectivity between visual and MTL regions. While positive connectivity between brain regions at rest is assumed to reflect synchronized activity between those regions, the physiological mechanisms underlying negative connectivity are still debated (Goelman et al., 2014). Next to non-neuronal hemodynamic processes, one physiological source for negative connectivity could be neuronal inhibition (Bianciardi et al., 2011; Devor et al., 2007; Shmuel et al., 2006). Hence, the decreased negative connectivity between visual and MTL regions in the stimulant condition could reflect reduced inhibition between those brain regions facilitating encoding of the stimulus material in both memory tasks. In both tasks, participants had to memorize visually or aurally presented words. In fact, the Lingual Gyrus—one visual region identified within the hierarchical cluster—plays an important role in the recognition of words (Mechelli et al., 2000) and has also been found to support encoding of words via visual imagery (Leshikar et al., 2012). However, further research is necessary to investigate negative connectivity between MTL and visual regions in the context of memory.

4.3 | Limitations

It is worth mentioning three study limitations that may reduce the generalizability and interpretability of the results. First, only male participants were recruited to avoid differential menstrual cycle effects on the two study sessions 7 days apart. This selectivity also reduces study-irrelevant within-subject variability regarding the rs-fMRI data (Lisofsky et al., 2015). However, research including women is necessary to test whether the stimulant-induced behavioral and FC results generalize to the whole population. Similarly, we only included participants with no to low/irregular caffeine consumption into the study to avoid cognitive and physiological side effects due to caffeine withdrawal during the time period of the experiment (see James, 2014; James & Rogers, 2005). However, about 90% of the general U.S. population report to regularly consume caffeine (Meredith et al., 2013).

Second, the sample size per stimulant group ($n = 16$) was rather low, thus complicating any interpretation of differential effects of any of the three stimulant. We did not find evidence for systematic differences between the groups in the sense that one stimulant group was driving the obtained overall results on memory and FC. However, given that the combined cognitive enhancement effect of all three stimulant groups only explained about 4%–10% of the variance in memory, it cannot be excluded that there are systematic differences between the stimulant groups if the group size was increased (e.g., $n = 100$).

Third, we only used self-reports to screen the participants for confounding variables such as caffeine, prescription stimulant or illicit substance consumption or sleep quality. More objective measures such as sleep diaries or actigraphy to evaluate participants' sleeping patterns as well as a urine drug screen prior to testing would have been a better control for those confounding variables.

5 | CONCLUSION

We investigated cognitive enhancement effects of stimulants (CAF, MPH, and MOD) on memory and related large-scale neural networks. Our preliminary results indicate that reduced negative FC between the FPN and DMN as well as between visual and MTL regions reflect a cognitive enhancement effect on the neural system-level specific to stimulant-induced memory enhancement. Further research is required to test, whether those enhanced memory-related changes in FC are generalizable to other pharmacological neuroenhancers. Finally, further research is required to investigate the shared stimulant-induced effects on different neurotransmitter systems such as catecholamines (dopamine) that give rise to such large-scale neural-network changes.

AUTHOR CONTRIBUTIONS

Dimitris Repantis, Martin Dresler and S.K. conceived the study. Dimitris Repantis performed the research and acquired the data. Maxi Becker analyzed the data and wrote the article. Maxi Becker, Dimitris Repantis, Martin Dresler and Simone Kühn contributed to revisions of the article.

ACKNOWLEDGMENTS

This work was supported by a grant (no. 85648) of the Volkswagen Foundation, Germany. The Volkswagen Foundation had no role in the design, data collection, data analysis, data interpretation or writing of the article. S.K. has been funded by the German Science Foundation (DFG SFB 936/C7), the European Research Council (ERC-2016-StG-Self-Control-677804) and a Fellowship from the Jacobs Foundation (JRF 2016-2018). M.B. was supported by the Einstein Stiftung Berlin (EPP-2017-423). We acknowledge support by the Open Access Publication Fund of the Humboldt-University Berlin. We thank Lukas C. Adam for help in collecting the data. Open Access funding enabled and organized by Projekt DEAL.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

The behavioral data, the extracted within-connectivity values per cluster as well as the analysis code are publicly available on github <https://github.com/MaxiBecker/Neuroenhancement>.

ORCID

Maxi Becker  <https://orcid.org/0000-0002-9430-4077>

Martin Dresler  <https://orcid.org/0000-0001-7441-3818>

Simone Kühn  <https://orcid.org/0000-0001-6823-7969>

REFERENCES

- Andersson, J. L., Hutton, C., Ashburner, J., Turner, R., & Friston, K. (2001). Modeling geometric deformations in EPI time series. *Neuroimage*, 13(5), 903–919.
- Arnsten, A. F. T. (2006). Stimulants: Therapeutic actions in ADHD. *Neuropsychopharmacology*, 31, 2376–2383.
- Baayen, R. H., Davidson, D. J., & Bates, D. M. (2008). Mixed-effects modeling with crossed random effects for subjects and items. *Journal of Memory and Language*, 59(4), 390–412.
- Badre, D., & D'Esposito, M. (2007). Functional magnetic resonance imaging evidence for a hierarchical organization of the prefrontal cortex. *Journal of Cognitive Neuroscience*, 19(12), 2082–2099.
- Barr, D. J., Levy, R., Scheepers, C., & Tily, H. J. (2013). Random effects structure for confirmatory hypothesis testing: Keep it maximal. *Journal of Memory and Language*, 68(3), 255–278.
- Bastoji, H., & Jouvet, M. (1988). Successful treatment of idiopathic hypersomnia and narcolepsy with modafinil. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 12(5), 695–700.
- Battleday, R. M., & Brem, A. K. (2015). Modafinil for cognitive neuro-enhancement in healthy non-sleep-deprived subjects: A systematic review. *European Neuropsychopharmacology*, 25(11), 1865–1881.
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Statistical Methodological)*, 57(1), 289–300.
- Bentivoglio, M., & Morelli, M. (Eds.). (2005). *The organization and circuits of mesencephalic dopaminergic neurons and the distribution of dopamine receptors in the brain*. Elsevier.
- Berridge, C. W., Devilbiss, D. M., Andrzejewski, M. E., Arnsten, A. F., Kelley, A. E., Schmeichel, B., Hamilton, C., & Spencer, R. C. (2006). Methylphenidate preferentially increases catecholamine neurotransmission within the prefrontal cortex at low doses that enhance cognitive function. *Biological Psychiatry*, 60, 1111–1120.
- Bianciardi, M., Fukunaga, M., van Gelderen, P., de Zwart, J. A., & Duyn, J. H. (2011). Negative BOLD-fMRI signals in large cerebral veins. *Journal of Cerebral Blood Flow and Metabolism*, 31, 401–412.
- Birn, R. M., Converse, A. K., Rajala, A. Z., Alexander, A. L., Block, W. F., McMillan, A. B., ... Populin, L. C. (2019). Changes in endogenous dopamine induced by methylphenidate predict functional connectivity in nonhuman primates. *Journal of Neuroscience*, 39(8), 1436–1444.
- Bijsterbosch, J., Harrison, S., Duff, E., Alfaro-Almagro, F., Woolrich, M., & Smith, S. (2017). Investigations into within-and between-subject resting-state amplitude variations. *NeuroImage*, 159, 57–69.
- Boivin, D. B., Montplaisir, J., Petit, D., Lambert, C., & Lubin, S. (1993). Effects of modafinil on symptomatology of human narcolepsy. *Clinical Neuropharmacology*, 16(1), 46–53.
- Borota, D., Murray, E., Keceli, G., Chang, A., Watabe, J. M., Ly, M., Toscano, J. P., & Yassa, M. A. (2014). Post-study caffeine administration enhances memory consolidation in humans. *Nature Neuroscience*, 17(2), 201–203.
- Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R Journal*, 9(2), 378–400.
- Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network: anatomy, function, and relevance to disease. *Annals of the New York Academy of Sciences*, 1124(1), 1–38.
- Burwell, R. D. (2000). The parahippocampal region: Corticocortical connectivity. *Annals of the New York Academy of Sciences*, 911, 25–42.
- Buysse, D. J., Reynolds, C. F., III, Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989). The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Research*, 28(2), 193–213.
- Cabeza, R., Ciaramelli, E., Olson, I. R., & Moscovitch, M. (2008). The parietal cortex and episodic memory: An attentional account. *Nature Reviews Neuroscience*, 9(8), 613–625.

- Caviola, L., & Faber, N. S. (2015). Pills or push-ups? Effectiveness and public perception of pharmacological and non-pharmacological cognitive enhancement. *Frontiers in Psychology, 6*(23), 435.
- Cera, N., Tartaro, A., & Sensi, S. L. (2014). Modafinil alters intrinsic functional connectivity of the right posterior insula: A pharmacological resting state fMRI study. *PLoS One, 9*(9), e107145.
- Chai, X. J., Castañón, A. N., Öngür, D., & Whitfield-Gabrieli, S. (2012). Anticorrelations in resting state networks without global signal regression. *NeuroImage, 59*(2), 1420–1428.
- Compton, W. M., Han, B., Blanco, C., Johnson, K., & Jones, C. M. (2018). Prevalence and correlates of prescription stimulant use, misuse, use disorders, and motivations for misuse among adults in the United States. *American Journal of Psychiatry, 175*(8), 741–755.
- Córdova, N. I., Tompary, A., & Turk-Browne, N. B. (2016). Attentional modulation of background connectivity between ventral visual cortex and the medial temporal lobe. *Neurobiology of learning and memory, 134*, 115–122.
- Dang, L. C., O'Neil, J. P., & Jagust, W. J. (2012). Dopamine supports coupling of attention-related networks. *Journal of Neuroscience, 32*(28), 9582–9587.
- Devor, A., Tian, P., Nishimura, N., Teng, I. C., Hillman, E. M., Narayanan, S. N., Ulbert, I., Boas, D. A., Kleinfeld, D., & Dale, A. M. (2007). Suppressed neuronal activity and concurrent arteriolar vasoconstriction may explain negative blood oxygenation level-dependent signal. *Journal of Neuroscience, 27*(16), 4452–4459.
- Dodds, C. M., Morein-Zamir, S., & Robbins, T. W. (2011). Dissociating inhibition, attention, and response control in the frontoparietal network using functional magnetic resonance imaging. *Cerebral cortex, 21*(5), 1155–1165.
- Esposito, R., Cilli, F., Pieramico, V., Ferretti, A., Macchia, A., Saggino, A., Ciavardelli, D., Manna, A., Navarra, R., Cieri, F., Stuppia, L., Tartaro, A., & Sensi, S. (2012). Acute effects of modafinil on cognition and brain resting state network activity in young healthy subjects. *PLoS One, 8*(7), e69224.
- Ferré, S. (2008). An update on the mechanisms of the psychostimulant effects of caffeine. *Journal of Neurochemistry, 105*(4), 1067–1079.
- Finn, E. S., Shen, X., Scheinost, D., Rosenberg, M. D., Huang, J., Chun, M. M., Papademetris, X., & Constable, R. T. (2015). Functional connectome fingerprinting: Identifying individuals using patterns of brain connectivity. *Nature Neuroscience, 18*(11), 1664–1671.
- Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van, E. D. C., & Raichle, M. E. (2005). The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proceedings of the National Academy of Sciences of the United States of America, 102*, 9673–9678.
- Franke, A., & Lieb, K. (2010). Pharmakologisches neuroenhancement und "Hirndoping". *Bundesgesundheitsbl, 53*, 853–860. <https://doi.org/10.1007/s00103-010-1105-0>
- Franke, A. G., Gränsmark, P., Agricola, A., Schühle, K., Rommel, T., Sebastian, A., Balló, H. E., Gorbulev, S., Gerdes, C., Frank, B., Ruckes, C., Tüscher, O., & Lieb, K. (2017). Methylphenidate, modafinil, and caffeine for cognitive enhancement in chess: A double-blind, randomised controlled trial. *European Neuropsychopharmacology, 27*(3), 248–260.
- Fredholm, B. B., Battig, K., Holmen, J., Nehlig, A., & Zvartau, E. E. (1999). Actions of caffeine in the brain with special reference to factors that contribute to its widespread use. *Pharmacological Reviews, 51*, 83–133.
- Gardner, W., Mulvey, E. P., & Shaw, E. C. (1995). Regression analyses of counts and rates: Poisson, overdispersed Poisson, and negative binomial models. *Psychological Bulletin, 118*(3), 392–404.
- Goelman, G., Gordon, N., & Bonne, O. (2014). Maximizing negative correlations in resting-state functional connectivity MRI by time-lag. *PLoS One, 9*(11), e111554.
- Grady, C. L. (2020). Meta-analytic and functional connectivity evidence from functional magnetic resonance imaging for an anterior to posterior gradient of function along the hippocampal axis. *Hippocampus, 30*(5), 456–471.
- Greicius, M. D., Krasnow, B., Reiss, A. L., & Menon, V. (2003). Functional connectivity in the resting brain: A network analysis of the default mode hypothesis. *Proceedings of the National Academy of Sciences, 100*(1), 253–258.
- Griefahn, B., Künemund, C., Bröde, P., & Mehnert, P. (2001). Zur validität der deutschen übersetzung des morningness-eveningness-questionnaires von horne und östberg: the validity of a german version of the morningness-eveningness-questionnaire developed by Horne and Östberg. *Somnologie, 5*(2), 71–80.
- Haatveit, B. C., Sundet, K., Hugdahl, K., Ueland, T., Melle, I., & Andreassen, O. A. (2010). The validity of d prime as a working memory index: Results from the "Bergen n-back task". *Journal of Clinical and Experimental Neuropsychology, 32*(8), 871–880.
- Haller, S., Rodriguez, C., Moser, D., Toma, S., Hofmeister, J., Sinanaj, I., Van De Ville, D., Giannakopoulos, P., & Lovblad, K. O. (2013). Acute caffeine administration impact on working memory-related brain activation and functional connectivity in the elderly: A BOLD and perfusion MRI study. *Neuroscience, 250*, 364–371.
- Horne, J. A., & Östberg, O. (1976). A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *International Journal of Chronobiology, 4*, 97–110.
- Hu, L., & Bentler, P. M. (1998). Fit indices in covariance structure modeling: Sensitivity to underparameterized model misspecification. *Psychological Methods, 3*, 424–453.
- Hu, L. T., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal, 6*(1), 1–55.
- Iidaka, T., Matsumoto, A., Nogawa, J., Yamamoto, Y., & Sadato, N. (2006). Frontoparietal network involved in successful retrieval from episodic memory. Spatial and temporal analyses using fMRI and ERP. *Cerebral Cortex, 16*(9), 1349–1360.
- Ilieva, I. P., Hook, C. J., & Farah, M. J. (2015). Prescription stimulants' effects on healthy inhibitory control, working memory, and episodic memory: A meta-analysis. *Journal of Cognitive Neuroscience, 27*, 1069–1089.
- Irwin, C., Khalesi, S., Desbrow, B., & McCartney, D. (2020). Effects of acute caffeine consumption following sleep loss on cognitive, physical, occupational and driving performance: A systematic review and meta-analysis. *Neuroscience & Biobehavioral Reviews, 108*, 877–888.
- Jafri, M. J., Pearlson, G. D., Stevens, M., & Calhoun, V. D. (2008). A method for functional network connectivity among spatially independent resting-state components in schizophrenia. *NeuroImage, 39*(4), 1666–1681.
- James, J. E. (2014). Caffeine and cognitive performance: Persistent methodological challenges in caffeine research. *Pharmacology Biochemistry and Behavior, 124*, 117–122.
- James, J. E., & Keane, M. A. (2007). Caffeine, sleep and wakefulness: Implications of new understanding about withdrawal reversal. *Human Psychopharmacology: Clinical and Experimental, 22*(8), 549–558.
- James, J. E., & Rogers, P. J. (2005). Effects of caffeine on performance and mood: Withdrawal reversal is the most plausible explanation. *Psychopharmacology, 182*(1), 1–8.
- Koppelstaetter, F., Poeppel, T. D., Siedentopf, C. M., Ischebeck, A., Verius, M., Haala, I., Mottaghy, F. M., Rhomberg, P., Golaszewski, S., Gotwald, T., Lorenz, I. H., Kolbitsch, C., Felber, S., & Krause, B. J. (2008). Does caffeine modulate verbal working memory processes? An fMRI study. *NeuroImage, 39*(1), 492–499.
- Kuczenski, R., & Segal, D. S. (1997). Effects of methylphenidate on extracellular dopamine, serotonin, and norepinephrine: Comparison with amphetamine. *Journal of Neurochemistry, 68*(5), 2032–2037.
- Laird, A. R., Eickhoff, S. B., Li, K., Robin, D. A., Glahn, D. C., & Fox, P. T. (2009). Investigating the functional heterogeneity of the default mode network using coordinate-based meta-analytic modeling. *Journal of Neuroscience, 29*(46), 14496–14505.
- Leech, R., Kamourieh, S., Beckmann, C. F., & Sharp, D. J. (2011). Fractionating the default mode network: Distinct contributions of the ventral

- and dorsal posterior cingulate cortex to cognitive control. *Journal of Neuroscience*, 31(9), 3217–3224.
- Leshikar, E. D., Duarte, A., & Hertzog, C. (2012). Task-selective memory effects for successfully implemented encoding strategies. *PLoS One*, 7(5), 0038160. <https://doi.org/10.1371/journal.pone.0038160>
- Liddle, E. B., Hollis, C., Batty, M. J., Groom, M. J., Totman, J. J., Liotti, M., Scerif, G., & Liddle, P. F. (2011). Task-related default mode network modulation and inhibitory control in ADHD: Effects of motivation and methylphenidate. *Journal of Child Psychology and Psychiatry*, 52(7), 761–771.
- Linssen, A. M., Sambeth, A., Vuurman, E. F., & Riedel, W. J. (2014). Cognitive effects of methylphenidate in healthy volunteers: A review of single dose studies. *International Journal of Neuropsychopharmacology*, 17(6), 961–977.
- Lisofsky, N., Mårtensson, J., Eckert, A., Lindenberger, U., Gallinat, J., & Kühn, S. (2015). Hippocampal volume and functional connectivity changes during the female menstrual cycle. *NeuroImage*, 118, 154–162. <https://doi.org/10.1016/j.neuroimage.2015.06.012>
- Macmillan, N. A., & Creelman, C. D. (1990). Response bias: Characteristics of detection theory, threshold theory, and “nonparametric” indexes. *Psychological Bulletin*, 107(3), 401–413.
- Manalo, R. V., & Medina, P. (2018). Caffeine protects dopaminergic neurons from dopamine-induced neurodegeneration via synergistic adenosine-dopamine D2-like receptor interactions in transgenic *Caenorhabditis elegans*. *Frontiers in Neuroscience*, 12, 137.
- Mechelli, A., Humphreys, G. W., Mayall, K., Olson, A., & Price, C. J. (2000). Differential effects of word length and visual contrast in the fusiform and lingual gyri during reading. *Proceedings of the Biological Sciences*, 267(1455), 1909–1913.
- Mehta, M. A., Owen, A. M., Sahakian, B. J., Mavaddat, N., Pickard, J. D., & Robbins, T. W. (2000). Methylphenidate enhances working memory by modulating discrete frontal and parietal lobe regions in the human brain. *Journal of Neuroscience*, 20(6), RC65.
- Meredith, S. E., Juliano, L. M., Hughes, J. R., & Griffiths, R. R. (2013). Caffeine use disorder: A comprehensive review and research agenda. *Journal of Caffeine Research*, 3(3), 114–130.
- Minzenberg, M. J., & Carter, C. S. (2008). Modafinil: a review of neurochemical actions and effects on cognition. *Neuropsychopharmacology*, 33(7), 1477–1502.
- Mueller, S., Costa, A., Keeser, D., Pogarell, O., Berman, A., Coates, U., Reiser, M. F., Riedel, M., Möller, H. J., Ettinger, U., & Meindl, T. (2014). The effects of methylphenidate on whole brain intrinsic functional connectivity. *Human Brain Mapping*, 35(11), 5379–5388.
- Müller, U., Rowe, J. B., Rittman, T., Lewis, C., Robbins, T. W., & Sahakian, B. J. (2013). Effects of modafinil on non-verbal cognition, task enjoyment and creative thinking in healthy volunteers. *Neuropharmacology*, 64, 490–495.
- Nehlig, A. (2010). Is caffeine a cognitive enhancer? *Journal of Alzheimer's Disease*, 20, 85–94.
- Ortega, R., López, V., Carrasco, X., Escobar, M. J., García, A. M., Parra, M. A., & Aboitiz, F. (2020). Neurocognitive mechanisms underlying working memory encoding and retrieval in attention-deficit/hyperactivity disorder. *Scientific Reports*, 10(1), 1–13.
- Pauls, A. M., O'Daly, O. G., Rubia, K., Riedel, W. J., Williams, S. C., & Mehta, M. A. (2012). Methylphenidate effects on prefrontal functioning during attentional-capture and response inhibition. *Biological Psychiatry*, 72(2), 142–149.
- Pigeau, R., Naitoh, P., Buguet, A., McCann, C., Baranski, J., Taylor, M., ... Mack, I. (1995). Modafinil, d-amphetamine and placebo during 64 hours of sustained mental work. I. Effects on mood, fatigue, cognitive performance and body temperature. *Journal of Sleep Research*, 4(4), 212–228.
- Rack-Gomer, A. L., Liao, J., & Liu, T. T. (2009). Caffeine reduces resting-state BOLD functional connectivity in the motor cortex. *NeuroImage*, 46(1), 56–63.
- Randall, D. C., Viswanath, A., Bharania, P., Elsabagh, S. M., Hartley, D. E., Shneerson, J. M., & File, S. E. (2005). Does modafinil enhance cognitive performance in young volunteers who are not sleep-deprived? *Journal of Clinical Psychopharmacology*, 25(2), 175–179.
- Ranganath, C., Heller, A., Cohen, M. X., Brozinsky, C. J., & Rissman, J. (2005). Functional connectivity with the hippocampus during successful memory formation. *Hippocampus*, 15(8), 997–1005.
- Ranjbar-Slamloo, Y., & Fazlali, Z. (2020). Dopamine and noradrenaline in the brain; overlapping or dissociate functions? *Frontiers in Molecular Neuroscience*, 12, 334.
- Repantis, D., Bovy, L., Ohla, K., Kühn, S., & Dresler, M. (2021). Cognitive enhancement effects of stimulants: A randomized controlled trial testing methylphenidate, modafinil, and caffeine. *Psychopharmacology*, 238(2), 441–451.
- Repantis, D., Schlattmann, P., Laisney, O., & Heuser, I. (2010). Modafinil and methylphenidate for neuroenhancement in healthy individuals: A systematic review. *Pharmacological Research*, 62(3), 187–206.
- Riedel, W. J., Blokland, A. (2015). *Cognitive Enhancement*. In: Kantak KM, Wettstein JG (eds) *Cognitive Enhancement*. Springer International Publishing Switzerland, Handbook of Experimental Pharmacology, pp 215–236
- Roediger, H. L., & McDermott, K. B. (1995). Creating false memories: Remembering words not presented in lists. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 21(4), 803.
- Rosenberg, M. D., Finn, E. S., Scheinost, D., Papademetris, X., Shen, X., Constable, R. T., & Chun, M. M. (2016). A neuromarker of sustained attention from whole-brain functional connectivity. *Nature Neuroscience*, 19(1), 165–171.
- Rosseel, Y. (2012). Lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, 48(2), 1–36.
- Schermelleh-Engel, K., Kerwer, M., & Klein, A. G. (2014). Evaluation of model fit in nonlinear multilevel structural equation modeling. *Frontiers in Psychology*, 5, 181.
- Schmaal, L., Goudriaan, A. E., Joos, L., Krüse, A. M., Dom, G., van den Brink, W., & Veltman, D. J. (2013). Modafinil modulates resting-state functional network connectivity and cognitive control in alcohol-dependent patients. *Biological Psychiatry*, 73(8), 789–795.
- Scoriels, L., Jones, P. B., & Sahakian, B. J. (2013). Modafinil effects on cognition and emotion in schizophrenia and its neurochemical modulation in the brain. *Neuropharmacology*, 64, 168–184.
- Scolari, M., Seidl-Rathkopf, K. N., & Kastner, S. (2015). Functions of the human frontoparietal attention network: Evidence from neuroimaging. *Current opinion in behavioral sciences*, 1, 32–39.
- Schmidt, A., Müller, F., Dolder, P. C., Schmid, Y., Zanchi, D., Liechti, M. E., & Borgwardt, S. (2017). Comparative effects of methylphenidate, modafinil, and MDMA on response inhibition neural networks in healthy subjects. *International Journal of Neuropsychopharmacology*, 20(9), 712–720.
- Sestieri, C., Corbetta, M., Romani, G. L., & Shulman, G. L. (2011). Episodic memory retrieval, parietal cortex, and the default mode network: Functional and topographic analyses. *Journal of Neuroscience*, 31(12), 4407–4420.
- Shen, X., Tokoglu, F., Papademetris, X., & Constable, R. T. (2013). Groupwise whole-brain parcellation from resting-state fMRI data for network node identification. *NeuroImage*, 82, 403–415.
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., ... Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (MINI): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of clinical psychiatry*, 59(20), 22–33.
- Shmuel, A., Augath, M., Oeltermann, A., & Logothetis, N. K. (2006). Negative functional MRI response correlates with decreases in neuronal activity in monkey visual area V1. *Nature Neuroscience*, 9, 569–577.
- Silberstein, R. B., Pipingas, A., Farrow, M., Levy, F., & Stough, C. K. (2016). Dopaminergic modulation of default mode network brain functional

- connectivity in attention deficit hyperactivity disorder. *Brain and Behavior*, 6(12), e00582.
- Sørensen, T. (1948). A method of establishing groups of equal amplitude in plant sociology based on similarity of species and its application to analyses of the vegetation on Danish commons. *Biologiske Skrifter/Kongelige Danske Videnskabernes Selskab*, 5, 1–34.
- Spencer, R. C., Devilbiss, D. M., & Berridge, C. W. (2015). The cognition-enhancing effects of psychostimulants involve direct action in the prefrontal cortex. *Biological Psychiatry*, 77, 940–950.
- Spreng, R. N., Stevens, W. D., Chamberlain, J. P., Gilmore, A. W., & Schacter, D. L. (2010). Default network activity, coupled with the frontoparietal control network, supports goal-directed cognition. *NeuroImage*, 53(1), 303–317.
- Sripada, C. S., Kessler, D., & Angstadt, M. (2014). Lag in maturation of the brain's intrinsic functional architecture in attention-deficit/hyperactivity disorder. *Proceedings of the National Academy of Sciences of the United States of America*, 111, 14259–14264.
- Sripada, C. S., Kessler, D., Welsh, R., Angstadt, M., Liberzon, I., Phan, K. L., & Scott, C. (2013). Distributed effects of methylphenidate on the network structure of the resting brain: A connectomic pattern classification analysis. *NeuroImage*, 81, 213–221.
- Tal, O., Diwakar, M., Wong, C. W., Olafsson, V., Lee, R., Huang, M. X., & Liu, T. (2013). Caffeine-induced global reductions in resting-state BOLD connectivity reflect widespread decreases in MEG connectivity. *Frontiers in Human Neuroscience*, 7, 63.
- Talbot, M. (2009). Brain gain: The underground world of neuroenhancing drugs. *The New Yorker*, 32–43.
- Theunissen, E. L., de la Asuncion Elvira, J., van den Bergh, D., & Ramaekers, J. G. (2009). Comparing the stimulant effects of the H1-antagonist fexofenadine with 2 psychostimulants, modafinil and methylphenidate. *Journal of Clinical Psychopharmacology*, 29(5), 439–443.
- Uddin, L. Q., Clare Kelly, A. M., Biswal, B. B., Xavier Castellanos, F., & Milham, M. P. (2009). Functional connectivity of default mode network components: Correlation, anticorrelation, and causality. *Human Brain Mapping*, 30(2), 625–637.
- Ullrich, S., de Vries, Y. C., Kühn, S., Repantis, D., Dresler, M., & Ohla, K. (2015). Feeling smart: Effects of caffeine and glucose on cognition, mood and self judgment. *Physiology and Behavior*, 151, 629–637.
- Visser, M. M., Goodin, P., Parsons, M. W., Lillicrap, T., Spratt, N. J., Levi, C. R., & Bivard, A. (2019). Modafinil treatment modulates functional connectivity in stroke survivors with severe fatigue. *Scientific reports*, 9(1), 1–9.
- Volkow, N. D., Fowler, J. S., Logan, J., Alexoff, D., Zhu, W., Telang, F., Wang, G. J., Jayne, M., Hooker, J. M., Wong, C., Hubbard, B., Carter, P., Warner, D., King, P., Shea, C., Xu, Y., Muench, L., & Apelskog-Torres, K. (2009). Effects of modafinil on dopamine and dopamine transporters in the male human brain: Clinical implications. *JAMA*, 301(11), 1148–1154.
- Volkow, N. D., Wang, G. J., Fowler, J. S., Gatley, S. J., Logan, J., Ding, Y. S., Hitzemann, R., & Pappas, N. (1998). Dopamine transporter occupancies in the human brain induced by therapeutic doses of oral methylphenidate. *American Journal of Psychiatry*, 155(10), 1325–1331.
- Volkow, N. D., Wang, G. J., Fowler, J. S., Logan, J., Gerasimov, M., Maynard, L., Ding, Y., Gatley, S. J., Gifford, A., & Franceschi, D. (2001). Therapeutic doses of oral methylphenidate significantly increase extracellular dopamine in the human brain. *Journal of Neuroscience*, 21(2), RC121.
- Volkow, N. D., Wang, G. J., Logan, J., Alexoff, D., Fowler, J. S., Thanos, P. K., Wong, C., Casado, V., Ferre, S., & Tomasi, D. (2015). Caffeine increases striatal dopamine D 2/D 3 receptor availability in the human brain. *Translational Psychiatry*, 5(4), e549.
- Wagner, J. G. (1973). Intrasubject variation in elimination half-lives of drugs which are appreciably metabolized. *Journal of Pharmacokinetics and Biopharmaceutics*, 1(2), 165–173.
- Wais, P. E. (2008). fMRI signals associated with memory strength in the medial temporal lobes: A meta-analysis. *Neuropsychologia*, 46(14), 3185–3196.
- Walsh, J. K., Muehlbach, M. J., Humm, T. M., Dickins, Q. S., Sugerma, J. L., & Schweitzer, P. K. (1990). Effect of caffeine on physiological sleep tendency and ability to sustain wakefulness at night. *Psychopharmacology*, 101(2), 271–273.
- Wang, S. F., Ritchey, M., Libby, L. A., & Ranganath, C. (2016). Functional connectivity based parcellation of the human medial temporal lobe. *Neurobiology of Learning and Memory*, 134, 123–134.
- Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: A functional connectivity toolbox for correlated and anticorrelated brain networks. *Brain Connectivity*, 2(3), 125–141.
- Witter, M. P., Wouterlood, F. G., Naber, P. A., & Van Haeften, T. (2000). Anatomical organization of the parahippocampal-hippocampal network. *Annals of the New York Academy of Sciences*, 911(1), 1–24.
- Wong, C. W., Olafsson, V., Tal, O., & Liu, T. T. (2012). Anti-correlated networks, global signal regression, and the effects of caffeine in resting-state functional MRI. *NeuroImage*, 63(1), 356–364.
- Wood, S., Sage, J. R., Shuman, T., & Anagnostaras, S. G. (2014). Psychostimulants and cognition: A continuum of behavioral and cognitive activation. *Pharmacological Reviews*, 66, 193–221.
- Wu, W. C., Lien, S. H., Chang, J. H., & Yang, S. C. (2014). Caffeine alters resting-state functional connectivity measured by blood oxygenation level-dependent MRI. *NMR in Biomedicine*, 27(4), 444–452.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Becker, M., Repantis, D., Dresler, M., & Kühn, S. (2022). Cognitive enhancement: Effects of methylphenidate, modafinil, and caffeine on latent memory and resting state functional connectivity in healthy adults. *Human Brain Mapping*, 43(14), 4225–4238. <https://doi.org/10.1002/hbm.25949>