CORRECTION

Correction: Modeling of *In-Utero* and *Intra-Partum* Transmissions to Evaluate the Efficacy of Interventions for the Prevention of Perinatal HIV

Patumrat Sripan, Sophie Le Coeur, Billy Amzal, Lily Ingsrisawang, Patrinee Traisathit, Nicole Ngo-Giang-Huong, Kenneth McIntosh, Tim R. Cressey, Suraphan Sangsawang, Boonsong Rawangban, Prateep Kanjanavikai, Jean-Marc Tréluyer, Gonzague Jourdain, Marc Lallemant, Saïk Urien

Fig 2 is incorrect. The duration of treatment should be in weeks instead of days. The authors have provided a corrected version here.

🔓 OPEN ACCESS

Fig 2. Diagnostic plots for viral load time-course model. Top: 2a and 2b: Observed versus model predicted viral load values (expressed as $\log_{10}$ copies) of the population and individual predictions respectively. Solid black circles, measure values; grey symbols, simulation of below the limit of quantification data. Line, identity line. Bottom: Visual predictive check plots. (2c) Women receiving only zidovudine (ZDV); (2d) women receiving zidovudine plus lopinavir/ritonavir (ZDV+LPV/r).The lines denote the median, 5th and 95th percentiles for the observed data. The grey areas stand for the 95% confidence intervals of the median, 5th and 95th model prediction percentiles.

Reference

1. Sripan P, Le Coeur S, Amzal B, Ingsrisawang L, Traisathit P, Ngo-Giang-Huong N, et al. (2015) Modeling of *In-Utero* and *Intra-Partum* Transmissions to Evaluate the Efficacy of Interventions for the Prevention of Perinatal HIV. PLoS ONE 10(5): e0126647. doi:10.1371/journal.pone.0126647 PMID: 25992639