

Bioinformatic identification of key genes and analysis of prognostic values in clear cell renal cell carcinoma

TING LUO^{1*}, XIAOYI CHEN^{1*}, SHUFEI ZENG², BAOZHANG GUAN¹, BO HU¹, YU MENG¹,
FANNA LIU¹, TAKSUI WONG¹, YONGPIN LU², CHEN YUN², BERTHOLD HOCHER³ and LIANGHONG YIN¹

¹Department of Nephrology, The First Affiliated Hospital of Jinan University, Jinan University, Guangzhou, Guangdong 510632, P.R. China; ²Department of Nephrology, Charité-Universitätsmedizin Berlin, Campus Mitte, D-10117 Berlin; ³Institute of Nutritional Sciences, University of Potsdam, D-14558 Potsdam, Germany

Received December 14, 2017; Accepted May 22, 2018

DOI: 10.3892/ol.2018.8842

Abstract. The present study aimed to identify new key genes as potential biomarkers for the diagnosis, prognosis or targeted therapy of clear cell renal cell carcinoma (ccRCC). Three expression profiles (GSE36895, GSE46699 and GSE71963) were collected from Gene Expression Omnibus. GEO2R was used to identify differentially expressed genes (DEGs) in ccRCC tissues and normal samples. The Database for Annotation, Visualization and Integrated Discovery was utilized for functional and pathway enrichment analysis. STRING v10.5 and Molecular Complex Detection were used for protein-protein interaction (PPI) network construction and module analysis, respectively. Regulation network analyses were performed with the WebGestalt tool. UALCAN web-portal was used for expression validation and survival analysis of hub genes in ccRCC patients from The Cancer Genome Atlas (TCGA). A total of 65 up- and 164 downregulated genes were identified as DEGs. DEGs were enriched with functional terms and pathways compactly related to ccRCC pathogenesis. Seventeen hub genes and one significant module were filtered out and selected from the PPI network. The differential expression of hub genes was verified in TCGA patients. Kaplan-Meier plot showed that high mRNA expression of enolase 2 (*ENO2*) was associated with short overall survival in ccRCC patients

($P=0.023$). High mRNA expression of cyclin D1 (*CCND1*) ($P<0.001$), fms related tyrosine kinase 1 (*FLT1*) ($P=0.004$), plasminogen (*PLG*) ($P<0.001$) and von Willebrand factor (*VWF*) ($P=0.008$) appeared to serve as favorable factors in survival. These findings indicate that the DEGs may be key genes in ccRCC pathogenesis and five genes, including *ENO2*, *CCND1*, *PLT1*, *PLG* and *VWF*, may serve as potential prognostic biomarkers in ccRCC.

Introduction

Renal cell carcinoma (RCC) accounts for 2-3% of all human malignancies (1). It is estimated that more than 338,000 people are diagnosed with RCC each year, with a 22% increase projected by 2020; there are more than 140,000 RCC-related deaths per year (2). Clear cell renal cell carcinoma (ccRCC) is the most common (~75%), lethal subtype of RCC (3). Over the past decade, with improved surgical procedures and the application of specific targeted drugs, the survival of RCC patient has markedly improved (4). However, early accurate diagnosis of ccRCC is still a great challenge and chemotherapeutic or radiotherapeutic resistance remains (4).

A comprehensive understanding of ccRCC initiation, progression and metastasis contributes to early diagnosis and precise treatment. Previous studies have demonstrated that mutations of *VHL* are significant drivers of ccRCC by regulating various biological processes, and *VHL* alterations are considered as prognostic markers in ccRCC (5). Moreover, targeted therapies associated with the pVHL/HIF pathway have been tested in phase 3 trials (4). *VHL* alterations alone are insufficient to cause the cancer, as ccRCC is a systemic biological disease. Sequencing studies have identified some other specific molecular genetic alterations of ccRCC, such as mutations of *TCEB1* (6), *PBRM1* (7) and abnormal expression of miR-92 (8), miR-210 (9). Further insights into the molecular biology of ccRCC could help us find some novel molecular biomarkers and potential targets for early diagnosis and precise treatment.

Gene expression profiling arrays make it possible to identify numerous differentially expressed genes in tumor samples compared to non-tumor samples at the same time. In this study, we performed an integrated bioinformatics analysis

Correspondence to: Professor Lianghong Yin, Department of Nephrology, The First Affiliated Hospital of Jinan University, Jinan University, 613 West Huangpu Avenue, Guangzhou, Guangdong 510632, P.R. China
E-mail: jnufu09@126.com

Dr Berthold Hocher, Institute of Nutritional Sciences, University of Potsdam, 114-116 Arthur-Scheunert-Allee, D-14558 Potsdam, Germany
E-mail: hocher@uni-potsdam.de

*Contributed equally

Key words: clear cell renal cell carcinoma, bioinformatics, differentially expressed genes, biomarkers, Kaplan-Meier plot

of three gene expression profiles and identified several differentially expressed genes (DEGs) in ccRCC tissues compared with normal controls. We executed functional and pathway enrichment analysis, protein-protein interaction (PPI) network analysis of DEGs and employed the Kaplan-Meier method to analyze survival associated with hub genes. We intended to provide further insights into the complex molecular biology of ccRCC pathogenesis and to identify new key genes that may be candidates for diagnostic biomarkers, prognostic indicators or potential targets of precise therapy.

Materials and methods

Data collection. Three gene expression profiles (GSE36895, GSE46699 and GSE71963) were acquired from Gene Expression Omnibus (GEO) database, a free public genomics data repository for array- and sequence-based data.

The array data of GSE36895 included 29 ccRCC tumor samples and 23 matched adjacent normal kidney cortices (10). GSE46699 was comprised of 126 samples including 65 ccRCC tumors and 61 patient-matched adjacent-normal tissues (11). GSE71963 contained 32 ccRCC tumor samples and 16 normal kidney samples (12).

Data processing. GEO2R, a tool for online analysis of GEO series based on the R programming language (13), was used to screen DEGs between the normal kidneys and ccRCC samples. Adjusted P-value (adj. *P*) and \log_2 Fold Change (\log_2 FCI) were used to select significant DEGs. adj. $P < 0.05$ and \log_2 FCI > 2 were chosen as the cutoff criteria.

Functional and pathway enrichment analysis. Gene ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis of DEGs was carried out using The Database for Annotation, Visualization and Integrated Discovery (DAVID) online (14,15). $P < 0.05$ was selected as the cutoff value.

PPI network construction and significant module analysis. STRING v10.5 was utilized for functional interaction analysis to construct a PPI network (16). Confidence scores > 0.7 were considered significant. Genes with degrees > 10 were selected as hub genes. The PPI network was visualized by Cytoscape software, and module of PPI network was screened by the Molecular Complex Detection (MCODE) in Cytoscape. The parameters were set as follows: Degree cutoff: 2, node score cutoff: 0.2, k-core: 2, and max. depth: 100 (17). The functional and pathway enrichment analysis of the significant module was carried out by DAVID.

Regulation network analyses. The miRNAs and transcription factors (TFs) that potentially regulated the DEGs were predicted using Overrepresentation Enrichment Analysis (ORA) in WebGestalt software (18). Then miRNA-target network and TF-target network were also visualized using Cytoscape software.

TCGA verification and survival analysis of hub genes. UALCAN, a tool for in-depth analyses of The Cancer Genome Atlas (TCGA) data, was utilized to verify the differences in




Figure 1. Identification of DEGs in three mRNA expression profiles (GSE36895, GSE46699 and GSE71963). DEGs, differentially expressed genes.

expression levels of hub genes (19). The correlation of hub genes with overall survival (OS) of ccRCC patients was examined by recruiting UALCAN as well. Patient data were categorized into two groups based on transcripts per million (TPM) value. The data with TPM greater than upper quartile were assigned to a high expression group and the others with TPM below upper quartile belonged to low/medium expression group. Survival analysis was performed by Kaplan-Meier method, and the log-rank test was carried out. $P < 0.05$ was selected as the cutoff value.

Results

Identification of DEGs in ccRCC. A total of 591, 325 and 1118 genes were extracted from the GSE36895, GSE46699 and GSE71963 datasets, respectively. There were 229 genes consistently differentially expressed in all three datasets (Fig. 1), including 65 upregulated DEGs and 164 downregulated DEGs in ccRCC tissues compared with normal kidney tissues (Table I).

GO analysis of DEGs in ccRCC. After performing GO analysis of DEGs with DAVID online, the DEGs were classified into three groups: biological process group, molecular function group and cellular component group. We found that the upregulated genes were mainly enriched in biological processes related to hypoxia, blood vessel morphogenesis and angiogenesis. The downregulated genes were commonly involved in functional terms associated with cellular components, metabolism and homeostasis.

Pathway enrichment analysis of DEGs in ccRCC. KEGG pathway enrichment analysis of DEGs was also conducted with DAVID online. KEGG results of the up- and downregulated genes were displayed in Tables II and III, respectively. The upregulated genes were mostly enriched in HIF-1 signaling pathway, PPAR signaling pathway, focal adhesion, coagulation cascades and AMPK signaling pathway. The downregulated genes were mainly enriched in metabolic pathways, collecting duct acid secretion, aldosterone-regulated sodium reabsorption, carbon metabolism and biosynthesis of antibiotics.

Table I. DEGs in ccRCC tissues compared with normal controls.

DEGs	Gene name
Upregulated	TNFAIP6, PFKP, NDUFA4L2, CXCR4 , NPTX2, C1QC, FLT1 , LOX, PDK1, COL23A1, CDCA2, GAS2L3, KCNK3, NETO2, FABP7, RNASET2, ANGPTL4, GJC1, SCD, HILPDA, LOXL2, DGCR5, CA9, EGLN3, ENO2 , TMEM45A, PPP1R3C, CAV2, VWF , CCND1 , ST8SIA4, C3, DIRAS2, IGFBP3, FABP5, LAMA4, SAP30, CD36, CTHRC1, GAL3ST1, HK2, VEGFA , SCARB1, AHNAK2, CAV1 , TGFBI, INHBB, ZNF395, PLOD2, TMCC1, PLXDC1, BHLHE41, CYP2J2, SPAG4, LPCAT1, CP, C1QB, FAM26F, APOC1, ENPP3, SLC6A3, ACKR3, ANGPT2, NOL3, ESM1
Downregulated	PTGER3, ERBB4, RALYL, L1CAM, XPNPEP2, SLC4A1, MPPED2, EHF, HMGCS2, HPD, GGACT, SLC7A13, HRG, UGT3A1, GATA3, TMEM174, SLC13A1, PROM2, CALB1, SUSD2, KCNJ1, SLC12A3, CRYAA, HSD11B2, DEFB1, GPC5, CYP27B1, UCHL1, FABP1, TMEM30B, CYP4F2, NELL1, MTURN, FGF9, NPHS2, PSAT1, SLC4A9, TFCP2L1, ALDH4A1 , SLC12A1 , ERP27, ALDH8A1, SCIN, TSPAN8, KL, AZGP1, SLC22A6, EFHD1, LOC100505985, CRHBP, AQP2 , ASS1, TACSTD2, PVALB, FOXI1, ABAT, TMEM52B, IRX2, MIOX, PIGR, ATP6V1G3, SEMA6D, S100A2, SCD5, MAL, FGF1 , SORD, DMRT2, TFAP2B, GLDC , FBP1, RASD1, PLPPR1, CYP4F3, GSTM3, ESRRG, SLC47A2, KNG1 , SLC34A1, MUC15, PTPRO, DPEP1, MECOM, ACSF2, CYP17A1, MT1G, PLG , UPP2, MFSD4A, SLC22A8, HAO2, ALDH6A1, MT1F, TMEM213, CHL1, EGF , DCXR, UMOD, ATP6V0D2, ANK2, HOGA1, DIO1, ELF5, SCNN1A, HSPA2, SOSTDC1, TYRP1, ENPP6, PCP4, GPC3, HS6ST2, CLDN8, PCK1, SLC5A2, NOX4, BMPR1B, G6PC, WNK4, ADH6, HEPA, CAM2, SOST, SH3GL2, SCNN1B, ALB , ALDOB, DCN , SCNN1G, KCNJ10, SLC13A3, SUCNR1, AFM, RAB25, ACPP, HPGD, FXYD4, DNER, RHCG, CYP4A11, CTXN3, KCNJ15, GRB14, PTH1R, GGT6, SLC26A7, C7, TMEM178A, OGDHL, ATP6V1B1, DUSP9, SERPINA5, SFRP1, CLCNKB, SLC7A8, SLC7A8, PIPOX, MAL2, PDE1A, TMPRSS2, GPAT3, PRODH2, FAM151A, EPCAM, MRO, ATP6V0A4

A total of 65 upregulated DEGs and 164 downregulated DEGs were identified in ccRCC tissues, compared with normal kidney tissues. The hub genes were shown in boldface. DEGs, differentially expressed genes; ccRCC, clear cell renal cell carcinoma.

Table II. KEGG pathway enrichment analysis of 65 upregulated DEGs.

Pathway	Name	P-value	Genes
hsa04066	HIF-1 signaling pathway	1.14x10 ⁻⁵	PDK1, FLT1, VEGFA, EGLN3, ENO2, HK2, ANGPT2
hsa03320	PPAR signaling pathway	4.19x10 ⁻⁴	CD36, SCD, FABP7, FABP5, ANGPTL4
hsa04510	Focal adhesion	7.01x10 ⁻⁴	CAV2, VWF, LAMA4, CAV1, CCND1, FLT1, VEGFA
hsa04610	Complement and coagulation cascades	5.81x10 ⁻³	C1QB, VWF, C3, C1QC
hsa04152	AMPK signaling pathway	2.70x10 ⁻²	CCND1, CD36, SCD, PFKP
hsa05150	Staphylococcus aureus infection	3.35x10 ⁻²	C1QB, C3, C1QC
hsa04151	PI3K-Akt signaling pathway	3.53x10 ⁻²	VWF, LAMA4, CCND1, FLT1, VEGFA, ANGPT2
hsa05230	Central carbon metabolism in cancer	4.57x10 ⁻²	PDK1, PFKP, HK2
hsa00010	Glycolysis/Gluconeogenesis	4.96x10 ⁻²	ENO2, PFKP, HK2

The pathways were ranked by P-value. KEGG, Kyoto Encyclopedia of Genes and Genomes; DEGs, differentially expressed genes.

PPI network construction and significant module analysis. A total of 169 genes of the 229 DEGs in all three datasets were filtered into the PPI network complex, containing 169 nodes and 432 edges (Fig. 2A). There were 44 upregulated genes and 125 downregulated genes among the 169 DEGs. Seventeen nodes with a degree >10 were identified as hub genes, such as *ALB*, *VEGFA*, *EGF*, *AQP2*, *ENO2*, *PLG*, *FLT1*, etc. (bold in Table I). The characteristic properties of the hub nodes based on analysis of the PPI network were tabulated in Table IV. These properties included degree, betweenness, closeness, stress

and average shortest path length. After performing module analysis by MCODE, the most significant module was screened out from the PPI network of DEGs, composed of 15 nodes and 54 edges (Fig. 2B). Functional and pathway enrichment analysis of nodes in the module was displayed in Table V. Most of these nodes were enriched in the functional terms related to substance transport and the pathways associated with cancer.

TF-DEG regulatory network. The DEG-associated transcriptional regulatory network was shown in Fig. 3A. A total of

Table III. KEGG pathway enrichment analysis of 164 downregulated DEGs.

Pathway	Name	P-value	Genes
hsa01100	Metabolic pathways	2.40x10 ⁻⁵	TYRP1, SORD, ASS1, OGDHL, ALDOB, UPP2, ADH6, ATP6V1B1, GPAT3, PIPOX, GLDC, CYP27B1, ALDH4A1, ATP6V0D2, HPD, ALDH6A1, KL, HOGA1, FBP1, PCK1, CYP4A11, CYP17A1, GGT6, G6PC, HMGCS2, HAO2, ABAT, PRODH2, CYP4F3, CYP4F2, ATP6V1G3, PSAT1, ATP6V0A4, DCXR
hsa04966	Collecting duct acid secretion	2.40x10 ⁻⁵	CLCNKB, SLC4A1, ATP6V1G3, ATP6V1B1, ATP6V0A4, ATP6V0D2
hsa04960	Aldosterone-regulated sodium reabsorption	1.51x10 ⁻⁴	FXVD4, HSD11B2, SCNN1G, SCNN1B, SCNN1A, KCNJ1
hsa01200	Carbon metabolism	3.81x10 ⁻³	ALDH6A1, OGDHL, ALDOB, HAO2, FBP1, PSAT1, GLDC
hsa01130	Biosynthesis of antibiotics	7.03x10 ⁻³	HMGCS2, ASS1, OGDHL, ALDOB, HAO2, FBP1, PSAT1, PCK1, GLDC
hsa00010	Glycolysis/gluconeogenesis	1.19x10 ⁻²	G6PC, ALDOB, FBP1, ADH6, PCK1
hsa04742	Taste transduction	2.17x10 ⁻²	PDE1A, SCNN1G, SCNN1B, SCNN1A
hsa05110	Vibrio cholerae infection	3.32x10 ⁻²	ATP6V1G3, ATP6V1B1, ATP6V0A4, ATP6V0D2
hsa00630	Glyoxylate and dicarboxylate metabolism	4.96x10 ⁻²	HAO2, HOGA1, GLDC

The pathways were ranked by P-value. KEGG, Kyoto Encyclopedia of Genes and Genomes; DEGs, differentially expressed genes.

90 nodes with 135 edges were contained in this regulation network, including 61 downregulated genes, 19 upregulated genes and 10 TFs.

miRNA-DEG regulatory network. In total, 6 miRNAs were filtered out (miR-144, miR-96, miR-503, miR-150, miR-383 and miR-338) (Fig. 3B). A total of 31 nodes and 28 edges were included in this regulatory network.

TCGA validation and the Kaplan-Meier plot. TCGA data of ccRCC patients were used via the UALCAN data portal. The hub genes identified from the PPI network were differentially expressed between ccRCC tissues and normal tissues (Fig. 4). The expression trends were identical within the three GEO datasets. Kaplan-Meier curve for overall survival of TCGA patients with ccRCC was obtained according to the low and high expression of each gene. The results showed that patients in the high mRNA expression group for *ENO2* had significantly worse OS than those in the low/medium expression group (P=0.023) (Fig. 5A). While high mRNA expression level of *CCND1* was associated with longer OS for ccRCC patients (P=0.000), as well as *FLT1* (P=0.004), *PLG* (P=0.000), and *VWF* (P=0.008) (Fig. 5B-E).

Discussion

The prognosis remains uncertain in ccRCC patients. Identifying novel potential biomarkers for early diagnosis, prognostic evaluation or targeted therapy may improve patient outcomes. Here we performed an in-depth analysis of three

expression profiles (with 126 ccRCC tissues and 100 normal controls) using bioinformatics method and identified 65 up- and 164 downregulated genes. Then we constructed a PPI network of DEGs and extracted 17 hub genes and one significant module from the PPI network. GO and KEGG pathway analysis revealed that the DEGs were commonly involved in functional terms and pathways related to the progression and prognosis of ccRCC. For example, hypoxia and HIF-1 pathway alterations are critical for the initiation and metastasis of ccRCC (20). Hypoxia could induce a series of tumor-related aberrations within cellular metabolism, apoptosis, migration and angiogenesis through dysregulation of HIF target genes (20). Drugs targeting the HIF-1 pathway have proven to be effective in treating ccRCC patients (21). In addition, metabolic pathways play a critical role in ccRCC progression according to previous studies, as well as glycolysis/gluconeogenesis, AMPK signaling pathway, and PI3K-Akt signaling pathway (22).

Interestingly, the *Staphylococcus aureus* infection pathway was found to be significant in our study. Growing evidence has indicated that bacterial infection is highly associated with certain human malignancies (23). It has been reported that lipoteichoic acids from *S. aureus* induce proliferation of two human non-small-cell lung cancer cell lines, A549 and H226 (24). However, the role of *S. aureus* infection in ccRCC still remains to be detected.

Using a Kaplan-Meier plot for survival analysis, the mRNA expression levels of *ENO2*, *CCND1*, *PLT1*, *PLG* and *VWF* were found to be significantly correlated with OS in ccRCC.




Figure 2. DEGs protein-protein interaction (PPI) network complex and one significant module obtained from PPI network. (A) DEGs PPI network containing 169 nodes and 432 edges. (B) One significant module composed of 15 nodes and 54 edges. Red nodes and green nodes stand for upregulated genes and downregulated genes, respectively. Lines represent the interaction between nodes. DEG, differentially expressed genes.

Enolase 2 (ENO2) encodes an enolase isoenzyme which is considered as a sensitive and specific biomarker for small-cell lung cancer (25,26). According to our KEGG results, *ENO2* was involved in several pathways compactly related to ccRCC pathogenesis such as glycolysis/gluconeogenesis, HIF-1 signaling pathway and metabolic pathways. In addition, *ENO2* is found to be induced by HIF-2a although suppression of its mRNA expression alone does not significantly inhibit the growth of the ccRCC cell line 786-O (27). Combining our survival analysis, we infer that *ENO2* may be an indicator in the diagnosis and prognosis rather than a potential target for therapy.

Cyclin D1 (CCND1) encodes an essential protein in the cell cycle which shows dual functions in cell growth. It is well-established that *CCND1* regulates the cell cycle transition from G1 to S phase by binding to CK4 and CDK6 (28,29). Previous studies suggest that the overexpression of *CCND1* promotes cell growth in many malignancies (30-34). Other studies have shown an apoptotic induction effect of *CCND1*. Consistent expression of an exogenous *CCND1* significantly inhibits cell proliferation (35) and induces apoptosis in mammary epithelial cell lines (36). Upregulated *CCND1* induces apoptosis of fibroblasts (37) and has a positive

Table IV. Topology properties of 17 hub genes.

Genes name	Degree	Betweenness centrality	Closeness centrality	Clustering coefficient	Stress	Average shortest path length
ALB	50	0.42	0.50	0.10	30,746	2.00
VEGFA	35	0.14	0.42	0.15	11,030	2.40
EGF	26	0.14	0.45	0.25	11,990	2.23
AQP2	19	0.20	0.41	0.23	15,956	2.44
ENO2	17	0.08	0.39	0.13	6,610	2.60
PLG	16	0.01	0.38	0.45	1,644	2.62
CAV1	15	0.05	0.39	0.29	4,140	2.57
KNG1	15	0.04	0.38	0.45	3,414	2.62
CXCR4	15	0.02	0.38	0.45	3,020	2.62
FLT1	15	0.01	0.39	0.51	1,474	2.58
VWF	14	0.00	0.37	0.52	582	2.67
GLDC	13	0.06	0.34	0.15	5,708	2.96
DCN	12	0.09	0.37	0.26	6,442	2.69
CCND1	12	0.04	0.38	0.47	2,944	2.65
SLC12A1	12	0.03	0.38	0.42	3,942	2.62
ALDH4A1	12	0.03	0.31	0.21	3,776	3.20
FGF1	11	0.02	0.37	0.53	1,592	2.67

The genes were ranked by degree.

Table V. Functional and pathway enrichment analyses of nodes in the significant module.

Term	Description	Count	P-value
GO:0006811	Ion transport	12	6.36x10 ⁻¹⁰
GO:0034220	Ion transmembrane transport	10	1.07 x10 ⁻⁰⁸
GO:0007588	Excretion	5	1.97 x10 ⁻⁰⁸
GO:0016324	Apical plasma membrane	7	4.25 x10 ⁻⁰⁸
GO:0015672	Monovalent inorganic cation transport	8	4.97 x10 ⁻⁰⁸
GO:0050878	Regulation of body fluid levels	8	6.29 x10 ⁻⁰⁸
GO:0030001	Metal ion transport	9	7.29 x10 ⁻⁰⁸
GO:0016324	Apical plasma membrane	7	1.68 x10 ⁻⁰⁷
GO:0055085	Transmembrane transport	10	1.70 x10 ⁻⁰⁷
GO:0006812	Cation transport	9	2.94 x10 ⁻⁰⁷
KEGG:hsa04960	Aldosterone-regulated sodium reabsorption	4	1.94x10 ⁻⁰⁵
KEGG:hsa04510	Focal adhesion	5	1.40x10 ⁻⁰⁴
KEGG:hsa05219	Bladder cancer	3	1.50x10 ⁻⁰³
KEGG:hsa04742	Taste transduction	3	1.81x10 ⁻⁰³
KEGG:hsa05212	Pancreatic cancer	3	3.73x10 ⁻⁰³
KEGG:hsa04066	HIF-1 signaling pathway	3	8.32x10 ⁻⁰³
KEGG:hsa04151	PI3K-Akt signaling pathway	4	1.14x10 ⁻⁰²
KEGG:hsa05205	Proteoglycans in cancer	3	3.22x10 ⁻⁰²
KEGG:hsa04015	Rap1 signaling pathway	3	3.52x10 ⁻⁰²
KEGG:hsa04014	Ras signaling pathway	3	4.03x10 ⁻⁰²
KEGG:hsa04060	Cytokine-cytokine receptor interaction	3	4.16x10 ⁻⁰²

Two GO categories including GO FAT and GO Direct was used for GO analysis. The top 10 GO terms were selected by P-value. If the term was filtered out by GO DIRECT and GO FAT at the same time, the more significant one would be selected. The GO terms and pathways were ranked by P-value. GO, gene ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes.




Figure 3. Regulatory network complex. (A) TF-DEG regulatory network containing 61 downregulated genes, 19 upregulated genes and 10 TFs. (B) miRNA-DEG regulatory network containing 31 nodes and 28 edges. Red nodes, green nodes, blue nodes and yellow nodes stand for upregulated genes, downregulated genes, TFs and miRNAs respectively. TF, transcription factor; miR, miRNA; DEG, differentially expressed genes.

correlation with a high apoptotic index in squamous cell carcinomas (38). Our analysis and previous studies show that *CCND1* is upregulated in ccRCC patients (39). Furthermore, it has been reported that reducing *CCND1* expression leads to a suppression of tumor growth in ccRCC (27). *CCND1* is considered as an oncogene in ccRCC. Interestingly, our results showed that high expression of *CCND1* was associated

with favorable prognosis in ccRCC. Similarly, *CCND1* is elevated and has a favorable effect on disease-free survival in papillary superficial bladder cancer (40). Two independent studies have shown that colon cancer patients with higher *CCND1* expression have better outcomes (41,42). The molecular mechanism of *CCND1* in cancer awaits further investigation.




Figure 4. Boxplots showing the expression of the 17 hub genes in healthy controls (n=72) and ccRCC tissues (n=533) of TCGA samples. The t-test was performed on the relevant results (* $P < 0.05$ and *** $P < 0.001$). ccRCC, clear cell renal cell carcinoma; TCGA, The Cancer Genome Atlas.

The importance of *VEGF* in RCC progression is well established and several VEGFR inhibitors such as sunitinib and sorafenib have proven to be significantly beneficial for progression-free survival (PFS) and OS in phase 3 trials (43,44). Recent research has demonstrated that *FLT1* (also known as *VEGFR-1*) protein expression in the tumor epithelium of localized ccRCC patients has a negative effect on prognosis (45). Other studies have found that high mRNA expression level of *FLT1* is significantly related to favorable PFS in metastatic ccRCC patients treated with sunitinib (46). In this study, we found that higher mRNA expression levels of *FLT1* in ccRCC tissue were associated with longer OS. The implication of *FLT1* in ccRCC remains unclear. It should be noted that *FLT1* can be generated as a transmembrane form and a soluble form. Soluble *FLT1* (*sFlt1*) lacks transmembrane and intracellular domains in contrast to the primary form, a full-length transmembrane receptor (47). Additionally, *sFlt1* is thought to be a natural antagonist of *VEGF*. Recent studies have found that *sFlt1* has an antitumor effect on several cancer cells (48-50). Enhanced *sFlt1* expression in the serum of breast cancer patients inhibits circulating tumor cells entering the peripheral blood, which may contribute to favorable outcomes (51). Herein we hypothesize that not transmembrane *FLT1* but *sFLT1* may have an antitumor effect on ccRCC and

the value of *sFlt1* in patient serum or urine may be worthy of further evaluation.

More and more evidence has demonstrated that plasminogen-plasmin system components are involved in tumor growth, invasion and metastasis by regulating angiogenesis and cell migration (52). The high levels of *uPA*, *uPAR* or *PAI-1* expression have proven to be prognostic biomarkers of poor outcome in many cancers, such as ovary cancer, breast cancer and renal cancer (53). The mRNA expression level of *PLG* in ccRCC patients was found to be downregulated in our analysis and other studies (54). Our results revealed that the ccRCC patients with a higher *PLG* mRNA expression had longer OS. Similar results have been reported in advanced ovarian cancer recently, and *PLG* was identified to be a favorable prognostic biomarker in this disease (55).

Another favorable biomarker in our analysis is *Von Willebrand Factor (VWF)*, which shows dual functions in angiogenesis and cancer metastasis according to previous data (56). *VWF* exhibits a pro-apoptotic effect on 769P, a ccRCC-derived cell line (57). While others have found that serum *VWF* levels are notably higher in progressive RCC patients compared with stable RCC patients (58). More studies should be done to clarify the link between *VWF* and ccRCC.




Figure 5. Kaplan-Meier survival curve for (A) *ENO2*, (B) *CCND1*, (C) *FLT1*, (D) *PLG* and (E) *VWF* expression levels in TCGA patients with ccRCC. The log-rank test was carried out on the relevant results. ccRCC, clear cell renal cell carcinoma; TCGA, The Cancer Genome Atlas.

The main limitation of our study is that exploration is done at a bioinformatics level, *in silico*. Future studies, especially biological experiments *in vitro* and *in vivo* are needed to validate the function of these DEGs in ccRCC.

In conclusion, through an integrated bioinformatics analysis of three gene profiles, we identified 229 DEGs, which may contain key genes in ccRCC pathogenesis. Five of the 17 hub genes including *ENO2*, *CCND1*, *PLT1*, *PLG* and *VWF* were filtered out through our analysis and may be potential prognostic biomarkers in ccRCC.

Acknowledgements

Not applicable.

Funding

This study was supported by the Guangdong Obers Blood Purification Academician Work Station (grant no. 2013B090400004); the Guangzhou Entrepreneurial Leader Talent (grant no. LCY201215); and the Guangdong Provincial Center for Clinical Engineering of Blood Purification (grant no. 507204531040).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

BH, LY and TL designed the study; TL, XC, SZ, BG, BH, YL and CY performed data analysis; YM, FL and TW performed literature research and data acquisition and participated in the data analysis; TL and XC wrote the manuscript; BH and LY revised the manuscript; YL and CY edited the manuscript and approved the final version of the manuscript; LY obtained funding. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

References

- Li P, Znaor A, Holcatova I, Fabianova E, Mates D, Wozniak MB, Ferlay J and Scelo G: Regional geographic variations in kidney cancer incidence rates in European countries. *Eur Urol* 67: 1134-1141, 2015.
- Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D and Bray F: Cancer incidence and mortality worldwide: Sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 136: E359-E386, 2015.
- Srinivasan R, Ricketts CJ, Sourbier C and Linehan WM: New strategies in renal cell carcinoma: Targeting the genetic and metabolic basis of disease. *Clin Cancer Res* 21: 10-17, 2015.
- Capitanio U and Montorsi F: Renal cancer. *Lancet* 387: 894-906, 2016.
- Gossage L and Eisen T: Alterations in VHL as potential biomarkers in renal-cell carcinoma. *Nat Rev Clin Oncol* 7: 277-288, 2010.
- Cancer Genome Atlas Research Network: Comprehensive molecular characterization of clear cell renal cell carcinoma. *Nature* 499: 43-49, 2013.
- Ibragimova I, Maradeo ME, Dulaimi E and Cairns P: Aberrant promoter hypermethylation of PBRM1, BAP1, SETD2, KDM6A and other chromatin-modifying genes is absent or rare in clear cell RCC. *Epigenetics* 8: 486-493, 2013.
- Valera VA, Walter BA, Linehan WM and Merino MJ: Regulatory effects of microRNA-92 (miR-92) on VHL gene expression and the hypoxic activation of miR-210 in clear cell renal cell carcinoma. *J Cancer* 2: 515-526, 2011.
- McCormick RI, Blick C, Ragoussis J, Schoedel J, Mole DR, Young AC, Selby PJ, Banks RE and Harris AL: miR-210 is a target of hypoxia-inducible factors 1 and 2 in renal cancer, regulates ISCU and correlates with good prognosis. *Br J Cancer* 108: 1133-1142, 2013.
- Peña-Llopis S, Vega-Rubín-de-Celis S, Liao A, Leng N, Pavia-Jiménez A, Wang S, Yamasaki T, Zhebker L, Sivanand S, Spence P, *et al*: BAP1 loss defines a new class of renal cell carcinoma. *Nat Genet* 44: 751-759, 2012.
- Eckel-Passow JE, Serie DJ, Bot BM, Joseph RW, Cheville JC and Parker AS: ANKS1B is a smoking-related molecular alteration in clear cell renal cell carcinoma. *BMC Urol* 14: 14, 2014.
- Takahashi M, Tsukamoto Y, Kai T, Tokunaga A, Nakada C, Hijjiya N, Uchida T, Daa T, Nomura T, Sato F, *et al*: Downregulation of WDR20 due to loss of 14q is involved in the malignant transformation of clear cell renal cell carcinoma. *Cancer Sci* 107: 417-423, 2016.
- Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, Marshall KA, Phillippy KH, Sherman PM, Holko M, *et al*: NCBI GEO: Archive for functional genomics data sets-update. *Nucleic Acids Res* 41 (Database Issue): D991-D995, 2013.
- Huang da W, Sherman BT and Lempicki RA: Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4: 44-57, 2009.
- Huang da W, Sherman BT and Lempicki RA: Bioinformatics enrichment tools: Paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 37: 1-13, 2009.
- Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, Santos A, Doncheva NT, Roth A, Bork P, *et al*: The STRING database in 2017: Quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res* 45: D362-D368, 2017.
- Bader GD and Hogue CW: An automated method for finding molecular complexes in large protein interaction networks. *BMC Bioinformatics* 4: 2, 2003.
- Wang J, Duncan D, Shi Z and Zhang B: WEB-based GEne SeT AnaLysis Toolkit (WebGestalt): Update 2013. *Nucleic Acids Res* 41: W77-W83, 2013.
- Chandrashekar DS, Bashel B, Balasubramanya SAH, Creighton CJ, Ponce-Rodriguez I, Chakravarthi BVSK and Varambally S: UALCAN: A portal for facilitating tumor subgroup gene expression and survival analyses. *Neoplasia* 19: 649-658, 2017.
- Schödel J, Grampp S, Maher ER, Moch H, Ratcliffe PJ, Russo P and Mole DR: Hypoxia, hypoxia-inducible transcription factors, and renal cancer. *Eur Urol* 69: 646-657, 2016.
- Escudier B, Szczylik C, Porta C and Gore M: Treatment selection in metastatic renal cell carcinoma: Expert consensus. *Nat Rev Clin Oncol* 9: 327-337, 2012.
- Ciccarese C, Brunelli M, Montironi R, Fiorentino M, Iacovelli R, Heng D, Tortora G and Massari F: The prospect of precision therapy for renal cell carcinoma. *Cancer Treat Rev* 49: 37-44, 2016.
- Zhu C, Wang Y, Cai C and Cai Q: Bacterial infection and associated cancers. *Adv Exp Med Biol* 1018: 181-191, 2017.
- Hattar K, Reinert CP, Sibelius U, Gökyildirim MY, Subtil FSB, Wilhelm J, Eul B, Dahlem G, Grimminger F, Seeger W and Grandel U: Lipoteichoic acids from *Staphylococcus aureus* stimulate proliferation of human non-small-cell lung cancer cells in vitro. *Cancer Immunol Immunother* 66: 799-809, 2017.
- Fizazi K, Cojean I, Pignon JP, Rixe O, Gatineau M, Hadeif S, Arriagada R, Baldeyrou P, Comoy E and Le Chevalier T: Normal serum neuron specific enolase (NSE) value after the first cycle of chemotherapy: An early predictor of complete response and survival in patients with small cell lung carcinoma. *Cancer* 82: 1049-1055, 1998.
- Oremek GM, Sauer-Eppel H and Bruzdziak TH: Value of tumour and inflammatory markers in lung cancer. *Anticancer Res* 27: 1911-1915, 2007.
- Zhang T, Niu X, Liao L, Cho EA and Yang H: The contributions of HIF-target genes to tumor growth in RCC. *PLoS One* 8: e80544, 2013.
- MacLachlan TK, Sang N and Giordano A: Cyclins, cyclin-dependent kinases and cdk inhibitors: Implications in cell cycle control and cancer. *Crit Rev Eukaryot Gene Expr* 5: 127-156, 1995.
- Alao JP: The regulation of cyclin D1 degradation: Roles in cancer development and the potential for therapeutic invention. *Mol Cancer* 6: 24, 2007.
- Arber N, Hibshoosh H, Moss SF, Sutter T, Zhang Y, Begg M, Wang S, Weinstein IB and Holt PR: Increased expression of cyclin D1 is an early event in multistage colorectal carcinogenesis. *Gastroenterology* 110: 669-674, 1996.
- Gautschi O, Ratschiller D, Gugger M, Betticher DC and Heighway J: Cyclin D1 in non-small cell lung cancer: a key driver of malignant transformation. *Lung Cancer* 55: 1-14, 2007.
- Bosch F, Jares P, Campo E, Lopez-Guillermo A, Piris MA, Villamor N, Tassies D, Jaffe ES, Montserrat E, Rozman C, *et al*: PRAD-1/cyclin D1 gene overexpression in chronic lymphoproliferative disorders: A highly specific marker of mantle cell lymphoma. *Blood* 84: 2726-2732, 1994.
- Faust JB and Meeker TC: Amplification and expression of the bcl-1 gene in human solid tumor cell lines. *Cancer Res* 52: 2460-2463, 1992.
- Buckley MF, Sweeney KJ, Hamilton JA, Sini RL, Manning DL, Nicholson RI, deFazio A, Watts CK, Musgrove EA and Sutherland RL: Expression and amplification of cyclin genes in human breast cancer. *Oncogene* 8: 2127-2133, 1993.
- Han EK, Sgambato A, Jiang W, Zhang YJ, Santella RM, Doki Y, Cacace AM, Schieren I and Weinstein IB: Stable overexpression of cyclin D1 in a human mammary epithelial cell line prolongs the S-phase and inhibits growth. *Oncogene* 10: 953-961, 1995.

36. Han EK, Begemann M, Sgambato A, Soh JW, Doki Y, Xing WQ, Liu W and Weinstein IB: Increased expression of cyclin D1 in a murine mammary epithelial cell line induces p27kip1, inhibits growth, and enhances apoptosis. *Cell Growth Differ* 7: 699-710, 1996.
37. Sofer-Levi Y and Resnitzky D: Apoptosis induced by ectopic expression of cyclin D1 but not cyclin E. *Oncogene* 13: 2431-2437, 1996.
38. Kotelnikov VM, Coon JS IV, Mundle S, Kelanic S, LaFollette S, Taylor S IV, Hutchinson J, Panje W, Caldarelli DD and Preisler HD: Cyclin D1 expression in squamous cell carcinomas of the head and neck and in oral mucosa in relation to proliferation and apoptosis. *Clin Cancer Res* 3: 95-101, 1997.
39. Tang SW, Chang WH, Su YC, Chen YC, Lai YH, Wu PT, Hsu CI, Lin WC, Lai MK and Lin JY: MYC pathway is activated in clear cell renal cell carcinoma and essential for proliferation of clear cell renal cell carcinoma cells. *Cancer Lett* 273: 35-43, 2009.
40. Sgambato A, Migaldi M, Faraglia B, De Aloysio G, Ferrari P, Ardito R, De Gaetani C, Capelli G, Cittadini A and Trentini GP: Cyclin D1 expression in papillary superficial bladder cancer: Its association with other cell cycle-associated proteins, cell proliferation and clinical outcome. *Int J Cancer* 97: 671-678, 2002.
41. Holland TA, Elder J, McCloud JM, Hall C, Deakin M, Fryer AA, Elder JB and Hoban PR: Subcellular localisation of cyclin D1 protein in colorectal tumours is associated with p21(WAF1/CIP1) expression and correlates with patient survival. *Int J Cancer* 95: 302-306, 2001.
42. Ogino S, Noshiro K, Irahara N, Kure S, Shima K, Baba Y, Toyoda S, Chen L, Giovannucci EL, Meyerhardt JA and Fuchs CS: A cohort study of cyclin D1 expression and prognosis in 602 colon cancer cases. *Clin Cancer Res* 15: 4431-4438, 2009.
43. Motzer RJ, Hutson TE, Tomczak P, Michaelson MD, Bukowski RM, Rixe O, Oudard S, Negrier S, Szczylik C, Kim ST, *et al*: Sunitinib versus interferon alfa in metastatic renal-cell carcinoma. *N Engl J Med* 356: 115-124, 2007.
44. Escudier B, Eisen T, Stadler WM, Szczylik C, Oudard S, Staehler M, Negrier S, Chevreau C, Desai AA, Rolland F, *et al*: Sorafenib for treatment of renal cell carcinoma: Final efficacy and safety results of the phase III treatment approaches in renal cancer global evaluation trial. *J Clin Oncol* 27: 3312-3318, 2009.
45. Klatte T, Seligson DB, LaRochelle J, Shuch B, Said JW, Riggs SB, Zomorodian N, Kabinavar FF, Pantuck AJ and Belldegrun AS: Molecular signatures of localized clear cell renal cell carcinoma to predict disease-free survival after nephrectomy. *Cancer Epidemiol Biomarkers Prev* 18: 894-900, 2009.
46. Beuselinck B, Jean-Baptiste J, Schöffski P, Couchy G, Meiller C, Rolland F, Allory Y, Joniau S, Verkarre V, Elaidi R, *et al*: Validation of VEGFR1 rs9582036 as predictive biomarker in metastatic clear-cell renal cell carcinoma patients treated with sunitinib. *BJU Int* 118: 890-901, 2016.
47. Kendall RL and Thomas KA: Inhibition of vascular endothelial cell growth factor activity by an endogenously encoded soluble receptor. *Proc Natl Acad Sci USA* 90: 10705-10709, 1993.
48. Miyake T, Kumasawa K, Sato N, Takiuchi T, Nakamura H and Kimura T: Soluble VEGF receptor 1 (sFLT1) induces non-apoptotic death in ovarian and colorectal cancer cells. *Sci Rep* 6: 24853, 2016.
49. Takano S, Ishikawa E, Matsuda M, Sakamoto N, Akutsu H, Yamamoto T and Matsumura A: The anti-angiogenic role of soluble-form VEGF receptor in malignant gliomas. *Int J Oncol* 50: 515-524, 2017.
50. Niu J, Wang Y, Wang J, Bin L and Hu X: Delivery of sFIT-1 engineered MSCs in combination with a continuous low-dose doxorubicin treatment prevents growth of liver cancer. *Aging (Albany NY)* 8: 3520-3534, 2016.
51. Vilsmaier T, Rack B, Janni W, Jeschke U and Weissenbacher T; SUCCESS Study Group: Angiogenic cytokines and their influence on circulating tumour cells in sera of patients with the primary diagnosis of breast cancer before treatment. *BMC Cancer* 16: 547, 2016.
52. Kwaan HC and McMahon B: The role of plasminogen-plasmin system in cancer. *Cancer Treat Res* 148: 43-66, 2009.
53. McMahon BJ and Kwaan HC: Components of the plasminogen-plasmin system as biologic markers for cancer. *Adv Exp Med Biol* 867: 145-156, 2015.
54. Schrödter S, Braun M, Syring I, Klümper N, Deng M, Schmidt D, Perner S, Müller SC and Ellinger J: Identification of the dopamine transporter SLC6A3 as a biomarker for patients with renal cell carcinoma. *Mol Cancer* 15: 10, 2016.
55. Zhao S, Dorn J, Napieralski R, Walch A, Diersch S, Kotzsch M, Ahmed N, Hooper JD, Kiechle M, Schmitt M and Magdolen V: Plasmin(ogen) serves as a favorable biomarker for prediction of survival in advanced high-grade serous ovarian cancer. *Biol Chem* 398: 765-773, 2017.
56. Mojiri A, Stoletov K, Carrillo MA, Willetts L, Jain S, Godbout R, Jurasz P, Sergi CM, Eisenstat DD, Lewis JD and Jahroudi N: Functional assessment of von Willebrand factor expression by cancer cells of non-endothelial origin. *Oncotarget* 8: 13015-13029, 2017.
57. Mochizuki S, Soejima K, Shimoda M, Abe H, Sasaki A, Okano HJ, Okano H and Okada Y: Effect of ADAM28 on carcinoma cell metastasis by cleavage of von Willebrand factor. *J Natl Cancer Inst* 104: 906-922, 2012.
58. Braybrooke JP, O'Byrne KJ, Propper DJ, Blann A, Saunders M, Dobbs N, Han C, Woodhull J, Mitchell K, Crew J, *et al*: A phase II study of razoxane, an antiangiogenic topoisomerase II inhibitor, in renal cell cancer with assessment of potential surrogate markers of angiogenesis. *Clin Cancer Res* 6: 4697-4704, 2000.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.