

Research article

Open Access

Anthracyclines, proteasome activity and multi-drug-resistance

Mirela R Fekete¹, William H McBride² and Frank Pajonk^{*2}

Address: ¹Department of Neurology, Bürgerhospital, Tunzhofer Str. 14-16, 70191 Stuttgart, Germany and ²Department of Radiation Oncology, Roy E. Coats Labs., David Geffen School of Medicine at UCLA, 10833 Le Conte Avenue, Los Angeles, CA90095-1714, USA

Email: Mirela R Fekete - mirelafekete@gmx.de; William H McBride - wmcbride@mednet.ucla.edu; Frank Pajonk* - fpajonk@mednet.ucla.edu

* Corresponding author

Published: 13 September 2005

Received: 24 May 2005

BMC Cancer 2005, 5:114 doi:10.1186/1471-2407-5-114

Accepted: 13 September 2005

This article is available from: <http://www.biomedcentral.com/1471-2407/5/114>

© 2005 Fekete et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: P-glycoprotein is responsible for the ATP-dependent export of certain structurally unrelated compounds including many chemotherapeutic drugs. Amplification of P-glycoprotein activity can result in multi-drug resistance and is a common cause of chemotherapy treatment failure. Therefore, there is an ongoing search for inhibitors of P-glycoprotein. Observations that cyclosporin A, and certain other substances, inhibit both the proteasome and P-glycoprotein led us to investigate whether anthracyclines, well known substrates of P-gp, also inhibit the function of the proteasome.

Methods: Proteasome function was measured in cell lysates from ECV304 cells incubated with different doses of verapamil, doxorubicin, daunorubicin, idarubicin, epirubicin, topotecan, mitomycin C, and gemcitabine using a fluorogenic peptide assay. Proteasome function in living cells was monitored using ECV304 cells stably transfected with the gene for an ubiquitin/green fluorescent protein fusion protein. The ability of the proteasome inhibitor MG-132 to affect P-glycoprotein function was monitored by fluorescence due to accumulation of daunorubicin in P-glycoprotein overexpressing KB 8-5 cells.

Results: Verapamil, daunorubicin, doxorubicin, idarubicin, and epirubicin inhibited 26S chymotrypsin-like function in ECV304 extracts in a dose-dependent fashion. With the exception of daunorubicin, 20S proteasome function was also suppressed. The proteasome inhibitor MG-132 caused a dose-dependent accumulation of daunorubicin in KB 8-5 cells that overexpress P-glycoprotein, suggesting that it blocked P-glycoprotein function.

Conclusion: Our data indicate that anthracyclines inhibit the 26S proteasome as well as P-glycoprotein. Use of inhibitors of either pathway in cancer therapy should take this into consideration and perhaps use it to advantage, for example during chemosensitization by proteasome inhibitors.

Background

Multi-drug-resistance (MDR) is a common reason for chemotherapy treatment failure in breast cancer, leukemia, and non-Hodgkin lymphoma patients. MDR can often be attributed to over-expression of the *mdr1* gene

that codes for an ATP-dependent, transmembrane P-glycoprotein (P-gp) efflux pump pathway, which rapidly exports many structurally unrelated drugs from the cell, including anthracyclines [1,2].

Numerous pre-clinical and clinical studies using P-gp modulating compounds like verapamil, cyclosporin A, reserpine, staurosporine, propafenone, phenoxazine, chloroquine, phenothiazine and their derivatives have been undertaken to overcome MDR and several substances have been identified that are effective *in vitro* (reviewed in [3]). However, to revert MDR *in vivo*, most MDR-modulating drugs require serum concentrations that have unacceptable toxicity and therefore they are currently not used in standard chemotherapy regimens. The development of better, less toxic inhibitors might be aided by insights into the specificity of these inhibitors for other molecules and the spectrum of molecules bound by P-glycoprotein.

Two of the most commonly used MDR-modulating substances are verapamil and cyclosporin A (CsA), or their derivatives. Interestingly, CsA has recently been identified as an inhibitor of the 26S proteasome [4]. The 26S proteasome is a highly conserved multicatalytic protease responsible for ATP- and ubiquitin-dependent degradation of all short-lived and 70–90% of all long lived proteins including cyclin A, B and E, p21 and p27, p53, cJun, cFos, and IκB. As such, the 26S proteasome controls cell cycle, signal transduction pathways, apoptosis and major functions of the immune system. Indeed some of the immunosuppressive properties of CsA, such as decreases in the expression of MHC-I molecules on the surface of target cells [5] and apoptotic death of lymphocytes through inhibition of the transcription factor NF-κB [6], may be due to its inhibitory effect on proteasome function. Vinblastine, a known P-gp substrate has also been shown to inhibit proteasome activity [7]. And, remarkably, the HIV protease inhibitor ritonavir was identified as an inhibitor of P-gp [8] and the proteasome [9]. Since CsA and ritonavir have been shown to inhibit both proteasome and P-gp activities, we questioned whether there was cross specificity between P-gp and proteasome activities. Cross specificity might explain effects of P-gp inhibitors on multiple cellular parameters that seem extrinsic to a pumping function of P-gp. Insights into substrate cross specificity of P-gp could offer a basis for the development of more selective P-gp inhibitors. They could also indicate reasons for the toxicity of these inhibitors, and why they affect cellular functions other than those related to P-gp.

Using an *in vitro* model, we show that anthracyclines and verapamil both inhibit proteasome function. Additionally, we demonstrate that the proteasome inhibitor MG-132 inhibits P-gp function, thereby increasing the uptake of doxorubicin in the cytoplasm and the nucleus.

Methods

Cell culture

KB 8.5 human epitheloid carcinoma cells that overexpress P-gp were a generous gift from Dr. Peter Hafkemeyer

(University Clinic Freiburg, Germany). Every 21 days P-gp-positive KB 8.5 cells were selected by addition of colchicine (10 ng/ml, Sigma). 24 hours before drug treatment cells were plated into 6-well plates (Costar) at a density of 10^6 cells/well.

EVC 304 human bladder carcinoma cells and PC-3 prostate cancer cells were obtained from the German Microorganism and Tissue Culture Collection (German collection of microorganism and cell cultures, DSMZ, Braunschweig). Cells were grown in 75 cm² flasks (Falcon) at 37°C in a humidified atmosphere at 5 % CO₂ in DMEM medium (Sigma) supplemented with 10 % heat inactivated FCS (Sigma) and 1 % penicillin/streptomycin (Gibco BRL).

Drug treatment

Stock solutions of all cytotoxic drugs were obtained from the hospital pharmacy of the University Clinic Freiburg. MG-132 (Calbiochem) was dissolved at 10 mM in DMSO and stored as small aliquots (10–30 μl) at -20°C. In drug accumulation assays doxorubicin (10 μM), daunorubicin (2–16 μM) or MG-132 (0.5–50 μM, 0.5% DMSO) were added to cells at the indicated times. Control cells were subjected to DMSO treatment alone (0.5 %).

Proteasome function assays

20S and 26S proteasome function was measured as described previously [20]. Briefly, cells were washed with PBS, then with buffer I (50 mM Tris, pH 7.4, 2 mM DTT, 5 mM MgCl₂, 2 mM ATP), and pelleted by centrifugation. Glass beads and homogenization buffer (50 mM Tris, pH 7.4, 1 mM DTT, 5 mM MgCl₂, 2 mM ATP, 250 mM sucrose) were added and vortexed for 1 minute. Beads and cell debris were removed by centrifugation at 1,000 × g for 5 minutes and 10,000 × g for 20 minutes. Protein concentration was determined by the BCA protocol (Pierce). One hundred μg protein of each sample was diluted with buffer I to a final volume of 1000 μl and the fluorogenic proteasome substrate SucLLVY-7-amido-4-methylcoumarin (chymotrypsin-like, Sigma) was added in a final concentration of 80 μM in 1% DMSO. To access 20S function, buffer I was replaced by an ATP-free buffer containing SDS (20 mM HEPES, pH 7.8; 0.5 mM EDTA, 0.03% SDS) [10]. Cleavage activity was monitored continuously by detection of free 7-amido-4-methylcoumarin using a fluorescence plate reader (Gemini, Molecular Devices) at 380/460 nm and 37°C. As controls for drug studies, 7-amido-4-methylcoumarin (AMC, 2 μM) was incubated with drugs in buffer I without cell extracts and measurements of proteasome function were corrected when necessary.

Drug accumulation assay

Total cellular daunorubicin content and accumulation of doxorubicin in the cytoplasm and nucleus were determined as described elsewhere [11] with some minor modifications. Growth medium on cells was replaced by PBS for 40 minutes at 37°C. This was replaced by fresh PBS containing daunorubicin or doxorubicin and MG-132 or anthracyclines alone. In some experiments, cells were washed with PBS after daunorubicin treatment and incubated in PBS containing MG-132 for an additional 40 minutes at 37°C. After drug treatment, the cells were washed twice with PBS, re-suspended in either 4 ml lysis buffer (0.3 M sucrose, 0.05 mM EGTA pH 8.0, 60 mM KCl, 15 mM NaCl, 15 mM HEPES pH 7.5, 150 µM spermine, 50 µM spermidine) containing 20 µl triton X-100 for nuclear isolation or 400 µl of 50% ethanol in 1 M HCl (v/v) for whole cell lysis. For the latter, cells were vortexed and diluted with water to a final volume of 1.4 ml. The cells in lysis buffer were mixed and left on ice for 15 minutes before centrifuging. The nuclei (pellet) were then vortexed with 400 µl HCl/isopropanol. Fluorescence derived from daunorubicin or doxorubicin was measured in quadruplicates of 200 µl using a fluorescence plate reader (Gemini, Molecular Devices) at 480/575 nm.

Transfection

ECV304 cells were plated at a density of 250,000 cells/well into six-well plates twelve hours before transfection. Cells were transfected with 5 µg of a plasmid (pEGFP-N1, Clontech) coding for an ubiquitin (Ub)-R-GFP fusion protein under control of a CMV promoter [12] (a kind gift from Dr. M. Masucci, Karolinska Institute, Sweden) using the Superfect transfection kit (Qiagen) and following the manufacturer's instructions. Transfected cells were maintained in DMEM (10 % FSC, 1 % penicillin/streptomycin) supplemented with 500 µg/ml G418 (Sigma) and clones were obtained. Expression of Ub-R-GFP was analyzed by flow cytometry (FL1-H, FACSCalibur, Becton Dickinson) using CellQuest Software before and after treatment with the proteasome inhibitor MG-132 (50 µM) for 10 hours at 37°C. Clone #10 (ECV304/10), which showed low background and high MG-132-induced expression of Ub-R-GFP, was used for inhibition experiments.

Statistics

Experimental data are presented as mean ± standard error of the mean from at least three independent experiments. A p-value <0.05 in a two-sided student's t-test was considered as 'statistically significant'.

Results

Verapamil is an inhibitor of 20S and 26S proteasome function

In order to test the hypothesis that the P-gp inhibitor verapamil inhibits proteasome function, proteasome extracts




Figure 1

Verapamil is an inhibitor of 26S proteasome function. Incubation of crude extracts of ECV304 cells containing proteasomes with different doses of verapamil (50, 60, 80, 100, 200 µM) inhibited proteolysis of the chymotrypsin-like substrate SucLLVY-AMC in a dose-dependent manner, indicating inhibition of 26S proteasome function.

of ECV304 and PC-3 cells were incubated with different concentrations of the drug (0, 50, 100 and 200 µM) and immediately tested for their chymotrypsin-like activity against the fluorogenic substrate SucLLVY-7-amido-4-methylcoumarin. There was a dose-dependent inhibition of MG-132-sensitive 26S (Fig. 1A) and 20S (data not shown) proteasome function, consistent with a direct inhibitory effect of verapamil on the proteasome.

Anthracyclines inhibit 20S and 26S proteasome function in a dose-dependent manner

Since verapamil, vinblastine, and CsA have been found to inhibit 20S and 26S proteasome function and vinblastine and CsA serve as substrates of P-gp [3], we asked if anthracyclines in general have an inhibitory effect on this protease. When crude extracts of ECV304 cells were incubated with different doses (0 – 100 µM) of the anthracyclines doxorubicin, daunorubicin, idarubicin and epirubicin we observed dose-dependent inhibition of 26S proteasome function with IC₅₀ values of 65.5 µM for doxorubicin, 13.7 µM for daunorubicin, 38.6 µM for idarubicin and 29.2 µM for epirubicin (Table 1). Topotecan, mitomycin C, and gemcitabine had no measurable effect on 26S proteasome function (data not shown). 20S proteasome function was inhibited by doxorubicin (IC₅₀ 5.8 µM), idarubicin (IC₅₀ 92 µM), epirubicin (IC₅₀ 12.5 µM) but not by daunorubicin (Table 2).

Table 1: Chymotryptic 26S Proteasome Activity in Lysates from ECV 304 Cells

μM	Doxorubicin	Daunorubicin	Epirubicin	Idarubicin
0	1	1	1	1
6.25	$0.92 \pm 0.03^*$	$0.85 \pm 0.05^*$	0.79 ± 0.1 n.s.	0.97 ± 0.03 n.s.
12.5	$0.84 \pm 0.05^*$	$0.53 \pm 0.11^*$	0.57 ± 0.31 n.s.	0.96 ± 0.02 n.s.
25	$0.72 \pm 0.04^{**}$	$0.22 \pm 0.09^{**}$	$0.59 \pm 0.14^*$	0.79 ± 0.12 n.s.
50	$0.59 \pm 0.1^*$	$0.12 \pm 0.06^{**}$	$0.36 \pm 0.2^*$	$0.28 \pm 0.09^{**}$
100	$0.36 \pm 0.03^{***}$	$0.11 \pm 0.06^{**}$	$0.27 \pm 0.17^*$	$0.1 \pm 0.06^{**}$

n.s. not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (two-sided student's t-test)

Table 2: Chymotryptic 20S Proteasome Activity in Lysates from ECV 304 Cells

μM	Doxorubicin	Daunorubicin	Epirubicin	Idarubicin
0	1	1	1	1
6.25	$0.49 \pm 0.1^*$	0.62 ± 0.19 n.s.	0.71 ± 0.21 n.s.	$0.67 \pm 0.07^*$
12.5	$0.28 \pm 0.09^{**}$	0.64 ± 0.07 n.s.	0.46 ± 0.2 n.s.	$0.67 \pm 0.08^*$
25	$0.28 \pm 0.13^*$	0.81 ± 0.05 n.s.	$0.37 \pm 0.08^{**}$	$0.69 \pm 0.05^{**}$
50	$0.24 \pm 0.06^{**}$	0.95 ± 0.04 n.s.	$0.33 \pm 0.07^{**}$	$0.67 \pm 0.05^{**}$
100	$0.25 \pm 0.07^{**}$	1.05 ± 0.1 n.s.	$0.35 \pm 0.07^{**}$	$0.49 \pm 0.11^*$

n.s. not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (two-sided student's t-test)

In order to demonstrate if this inhibition could be observed in living cells, we incubated ECV304/10 cells, stably transfected with an expression plasmid for an Ub-GFP fusion protein with doxorubicin (100 μM) for 12 hours. When analyzed by fluorescence microscopy, the cells showed perinuclear accumulation of doxorubicin while GFP accumulated throughout the cytoplasm, indicating inhibition of proteasome function (Fig. 2).

MG-132 treatment reverts multi-drug-resistance in P-gp expressing KB 8-5 cells

The human epitheloid carcinoma cell line KB 8-5 is a well-characterized tumor cell line that over-expresses mdr-1 with associated MDR. Preliminary experiments showed that treatment of KB 8.5 cells with the reversible proteasome inhibitor MG-132 (3.125 to 50 μM) induced apoptosis within 24 hours. This is in accord with numerous studies reporting induction of apoptosis in cancer cells by proteasome inhibitors [13], and indicated that MG-132 enters KB 8-5 cells and that they are not abnormally resistant to its effects based on enhanced P-gp function. After 45 minutes of incubation with MG-132 (50 μM), no morphological signs of toxicity were observed. KB 8-5 cells treated with different doses of MG-132 and daunorubicin (10 μM) for 45 minutes showed increased, dose-dependent accumulation of daunorubicin in the cytoplasm (e.g.

a 4-fold increase at 50 μM MG-132, Fig. 3) indicating that MG-132 could block P-gp function. This was further supported by the observation that incubation of ECV304 cell with MG-132 (25 μM) caused an increased uptake of doxorubicin in the cytoplasm and in the nuclear fraction of the cells (Fig. 4).

Discussion

The observations that CsA [4] and vinblastine [7] have inhibitory effects on the cleavage activity of the 26S proteasome led us investigate the effects of anthracycline anticancer agents and verapamil on the activity of this protease. Verapamil caused a concentration-dependent inhibition of 20S and 26S function. Additionally, we found a concentration-dependent inhibition of 26S proteasome function for all four anthracyclines tested. Comparable results showing doxorubicin to be a non-competitive inhibitor of the proteasome have been reported previously [14]. With the exception of daunorubicin, anthracyclines also inhibited 20S chymotryptic function in a dose-dependent manner. It is known that doxorubicin is co-transported into the nucleus along with proteasomes [15,16] but our observation of a general direct inhibitory effect of anthracycline anticancer agents on the proteasome sheds a totally new light on the actions of these drugs.




Figure 2
Anthracyclines are inhibitors of proteasome function. Incubation of ECV304 cells stably transfected with an Ub-GFP fusion protein with daunorubicin (100 μ M, 16 h), caused accumulation of GFP throughout the cytoplasm (lower picture), indicating proteasome inhibition in living cells while untreated controls cells showed only little accumulation of GFP (A/B). Daunorubicin accumulated in the perinuclear region (C).

The inhibitory effects of the reversible inhibitor of the proteasome, MG-132, on P-glycoprotein function, supports the view that P-glycoprotein and the proteasome can both be targeted by this new class of chemotherapeutic drugs. This was further supported by the observation that verapamil, another established inhibitor of P-gp, inhibited the chymotryptic 20S and 26S function of the proteasome. The fact that both P-glycoprotein and proteasome activities can both be regulated by pro-inflammatory




Figure 3
MG-132 treatment of KB 8-5 causes intracellular accumulation of anthracyclines. Incubation of KB 8-5 cells, which overexpress P-gp, with increasing doses of MG-132 (0, 6.25, 12.5, 25, 50 μ M) caused a dose-dependent accumulation of daunorubicin, as measured by fluorescence, indicating inhibition of P-gp function by this proteasome inhibitor.




Figure 4
Accumulation of doxorubicin in the presence or absence of MG-132 (25 μ M) in the cytoplasm and the nuclear fraction of ECV304 cells.

cytokines and oxidative stress suggests [17-21] that studies on co-ordinate regulation of these activities might be illuminating.

These findings lead to interesting possibilities with respect to the possible use of proteasome inhibitors, which are just entering their first clinical trials [22,23], in combination therapy, as well as to the mechanism of action and toxicity of P-gp inhibitors. Using an *in vitro* system, we showed that the proteasome inhibitor MG-132 caused intracellular accumulation of anthracyclines, indicating inhibition of P-gp function. Proteasome inhibitors may interfere with drug-resistance at additional levels as P-gp and also topoisomerase II are degraded in a proteasome-

dependent manner and degradation is blocked by proteasome inhibitors [24,25]. However, given the long half-life of P-gp of 14–24 hours [26], the effects observed in our study after short-time incubation of the cells with MG-132 are probably not caused by an increased degradation of P-gp. The extent of the increase of anthracyclin accumulation in *mdr1*-overexpressing KB-8.5 treated with 25 μ M concentrations of MG-132 cells in our study was comparable to the effect of verapamil at 50 μ M [27]. Future studies have to clarify if similar effects can be obtained using clinically used proteasome inhibitors at concentrations typically reached in the serum of patients.

Tumor cells in general exhibit altered patterns of expression of proteasome subunits and their distribution between cytoplasm and nucleus often differs from normal cells [28–30]. This may explain why specific proteasome inhibitors like PS-341 are usually clinically well tolerated. Inhibition of proteasome function induces apoptosis of tumor cells [31–34] and sensitizes the surviving tumor cells to the actions of both chemotherapy [35] and radiation therapy [36,37]. Therefore, proteasome inhibitors might overcome P-gp-related MDR, with accompanying chemo- and radiosensitizing effects. Also, since tumor microvasculature expresses high levels of *mdr-1* [38,39], the possibility exists that the neovasculature is a target for these drugs *in vivo*. On the other hand, direct inhibition of proteasome function might be an additional major mechanism of action for anthracyclines. Such inhibition could contribute to their ability to enhance the efficacy of other chemotherapeutic drugs, independent of their ability to reverse MDR.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

MF carried out the molecular studies and helped to draft the manuscript. WMB participated in the design of the study and helped to draft the manuscript. FP designed the study and drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This investigation was supported in part by Grants of the German Research Foundation (DFG) Pa 723/1 Pa 723/3 (FP) and PHS Grant number CA-87887 awarded to WMc by the National Cancer Institute;

References

- Gottesman MM, Fojo T, Bates SE: **Multidrug resistance in cancer: role of ATP-dependent transporters.** *Nat Rev Cancer* 2002, **2(1)**:48-58.
- Lacayo NJ, Duran GE, Sikic BI: **Modulation of resistance to idarubicin by the cyclosporin PSC 833 (valsopodar) in multidrug-resistant cells.** *J Exp Ther Oncol* 2003, **3(3)**:127-135.
- Ambudkar SV, Dey S, Hrycyna CA, Ramachandra M, Pastan I, Gottesman MM: **Biochemical, cellular, and pharmacological aspects of the multidrug transporter.** *Annu Rev Pharmacol Toxicol* 1999, **39**:361-398.
- Meyer S, Kohler NG, Joly A: **Cyclosporine A is an uncompetitive inhibitor of proteasome activity and prevents NF-kappa B activation.** *FEBS Lett* 1997, **413(2)**:354-358.
- Harding CV, France J, Song R, Farah JM, Chatterjee S, Iqbal M, Siman R: **Novel dipeptide aldehydes are proteasome inhibitors and block the MHC-I antigen-processing pathway.** *J Immunol* 1995, **155(4)**:1767-1775.
- Marienfeld R, Neumann M, Chuvpilo S, Escher C, Kneitz B, Avots A, Schimpl A, Serfling E: **Cyclosporin A interferes with the inducible degradation of NF-kappa B inhibitors, but not with the processing of p105/NF-kappa B1 in T cells.** *Eur J Immunol* 1997, **27(7)**:1601-1609.
- Piccinini M, Tazartes O, Mezzatesta C, Ricotti E, Bedino S, Grosso F, Dianzani U, Tovo PA, Mostert M, Musso A, Rinaudo MT: **Proteasomes are a target of the anti-tumor drug vinblastine.** *Biochem J* 2001, **356(Pt 3)**:835-841.
- Drewe J, Gutmann H, Fricker G, Torok M, Beglinger C, Huwyler J: **HIV protease inhibitor ritonavir: a more potent inhibitor of P-glycoprotein than the cyclosporine analog SDZ PSC 833.** *Biochem Pharmacol* 1999, **57(10)**:1147-1152.
- Schmidtke G, Holzhtuter HG, Bogoy M, Kairies N, Groll M, de Giuli R, Emch S, Groettrup M: **How an inhibitor of the HIV-1 protease modulates proteasome activity.** *J Biol Chem* 1999, **274(50)**:35734-35740.
- Stein RL, Melandri F, Dick L: **Kinetic characterization of the chymotryptic activity of the 20S proteasome.** *Biochemistry* 1996, **35(13)**:3899-3908.
- Gieseler F, Nussler V, Brieden T, Kunze J, Valsamas S: **Intracellular pharmacokinetics of anthracyclines in human leukemia cells: correlation of DNA-binding with apoptotic cell death.** *Int J Clin Pharmacol Ther* 1998, **36(1)**:25-28.
- Dantuma NP, Lindsten K, Glas R, Jellne M, Masucci MG: **Short-lived green fluorescent proteins for quantifying ubiquitin/proteasome-dependent proteolysis in living cells.** *Nat Biotechnol* 2000, **18(5)**:538-543.
- Shinohara K, Tomioka M, Nakano H, Tone S, Ito H, Kawashima S: **Apoptosis induction resulting from proteasome inhibition.** *Biochem J* 1996, **317(Pt 2)**:385-388.
- Kiyomiya K, Satoh J, Horie H, Kurebe M, Nakagawa H, Matsuo S: **Correlation between nuclear action of anthracycline anticancer agents and their binding affinity to the proteasome.** *Int J Oncol* 2002, **21(5)**:1081-1085.
- Kiyomiya K, Matsuo S, Kurebe M: **Mechanism of specific nuclear transport of adriamycin: the mode of nuclear translocation of adriamycin-proteasome complex.** *Cancer Res* 2001, **61(6)**:2467-2471.
- Kiyomiya K, Matsuo S, Kurebe M: **Proteasome is a carrier to translocate doxorubicin from cytoplasm into nucleus.** *Life Sci* 1998, **62(20)**:1853-1860.
- Sukhai M, Piquette-Miller M: **Regulation of the multidrug resistance genes by stress signals.** *J Pharm Pharm Sci* 2000, **3(2)**:268-280.
- Stohwasser R, Kloetzel PM: **Cytokine induced changes in proteasome subunit composition are concentration dependent.** *Biol Chem* 1996, **377(9)**:571-577.
- Yang Y, Waters JB, Fruh K, Peterson PA: **Proteasomes are regulated by interferon gamma: implications for antigen processing.** *Proc Natl Acad Sci U S A* 1992, **89(11)**:4928-4932.
- Pajonk F, McBride WH: **Ionizing radiation affects 26s proteasome function and associated molecular responses, even at low doses.** *Radiother Oncol* 2001, **59(2)**:203-212.
- Grune T: **Oxidative stress, aging and the proteasomal system.** *Biogerontology* 2000, **1(1)**:31-40.
- Teicher BA, Ara G, Herbst R, Palombella VJ, Adams J: **The proteasome inhibitor PS-341 in cancer therapy.** *Clin Cancer Res* 1999, **5(9)**:2638-2645.
- Adams J, Palombella VJ, Elliott PJ: **Proteasome inhibition: a new strategy in cancer treatment.** *Invest New Drugs* 2000, **18(2)**:109-121.
- Zhang Z, Wu JY, Hait WN, Yang JM: **Regulation of the stability of P-glycoprotein by ubiquitination.** *Mol Pharmacol* 2004, **66(3)**:395-403.

25. Ogiso Y, Tomida A, Lei S, Omura S, Tsuruo T: **Proteasome inhibition circumvents solid tumor resistance to topoisomerase II-directed drugs.** *Cancer Res* 2000, **60(9)**:2429-2434.
26. Yoshimura A, Kuwazuru Y, Sumizawa T, Ikeda S, Ichikawa M, Usagawa T, Akiyama S: **Biosynthesis, processing and half-life of P-glycoprotein in a human multidrug-resistant KB cell.** *Biochim Biophys Acta* 1989, **992(3)**:307-314.
27. Ghauharali RI, Westerhoff HV, Dekker H, Lankelma J: **Saturable P-glycoprotein kinetics assayed by fluorescence studies of drug efflux from suspended human KB8-5 cells.** *Biochim Biophys Acta* 1996, **1278(2)**:213-222.
28. Amsterdam A, Pitzer F, Baumeister W: **Changes in intracellular localization of proteasomes in immortalized ovarian granulosa cells during mitosis associated with a role in cell cycle control.** *Proc Natl Acad Sci U S A* 1993, **90(1)**:99-103.
29. Kumatori A, Tanaka K, Inamura N, Sone S, Ogura T, Matsumoto T, Tachikawa T, Shin S, Ichihara A: **Abnormally high expression of proteasomes in human leukemic cells.** *Proc Natl Acad Sci U S A* 1990, **87(18)**:7071-7075.
30. Kanayama H, Tanaka K, Aki M, Kagawa S, Miyaji H, Satoh M, Okada F, Sato S, Shimbara N, Ichihara A: **Changes in expressions of proteasome and ubiquitin genes in human renal cancer cells.** *Cancer Res* 1991, **51(24)**:6677-6685.
31. Qiu JH, Asai A, Chi S, Saito N, Hamada H, Kirino T: **Proteasome inhibitors induce cytochrome c-caspase-3-like protease-mediated apoptosis in cultured cortical neurons.** *J Neurosci* 2000, **20(1)**:259-265.
32. Pasquini LA, Besio Moreno M, Adamo AM, Pasquini JM, Soto EF: **Lactacystin, a specific inhibitor of the proteasome, induces apoptosis and activates caspase-3 in cultured cerebellar granule cells.** *J Neurosci Res* 2000, **59(5)**:601-611.
33. Zhang XM, Lin H, Chen C, Chen BD: **Inhibition of ubiquitin-proteasome pathway activates a caspase-3-like protease and induces Bcl-2 cleavage in human M-07e leukaemic cells.** *Biochem J* 1999, **340(Pt 1)**:127-133.
34. Delic J, Masdehors P, Omura S, Cosset JM, Dumont J, Binet JL, Magdelenat H: **The proteasome inhibitor lactacystin induces apoptosis and sensitizes chemo- and radioresistant human chronic lymphocytic leukaemia lymphocytes to TNF-alpha-initiated apoptosis [see comments].** *Br J Cancer* 1998, **77(7)**:1103-1107.
35. Bold RJ, Virudachalam S, McConkey DJ: **Chemosensitization of pancreatic cancer by inhibition of the 26S proteasome.** *J Surg Res* 2001, **100(1)**:11-17.
36. Pajonk F, Pajonk K, McBride W: **Apoptosis and radiosensitization of Hodgkin's cells by proteasome inhibition.** *Int J Radiat Oncol Biol Phys* 2000, **47(4)**:1025-1032.
37. Russo SM, Tepper JE, Baldwin ASJ, Liu R, Adams J, Elliott P, Cusack JCJ: **Enhancement of radiosensitivity by proteasome inhibition: implications for a role of NF-kappaB.** *Int J Radiat Oncol Biol Phys* 2001, **50(1)**:183-193.
38. Iwahana M, Utoguchi N, Mayumi T, Goryo M, Okada K: **Drug resistance and P-glycoprotein expression in endothelial cells of newly formed capillaries induced by tumors.** *Anticancer Res* 1998, **18(4C)**:2977-2980.
39. Toth K, Vaughan MM, Peress NS, Slocum HK, Rustum YM: **MDRI P-glycoprotein is expressed by endothelial cells of newly formed capillaries in human gliomas but is not expressed in the neovasculature of other primary tumors.** *Am J Pathol* 1996, **149(3)**:853-858.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2407/5/114/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

