

Access this article online

Quick Response Code:



Website:

www.jehp.net

DOI:

10.4103/jehp.jehp\_1644\_20

# Introducing practical tools for fit to drive assessment of the elderly: A step toward improving the health of the elderly

Saiedeh Bahrampouri, Hamid Reza Khankeh, Seyed Ali Hosseini, Mohammadreza Mehmandar<sup>1</sup>, Abbas Ebadi<sup>2,3</sup>

## Abstract:

Today, as age increases, the demand for independent living has increased. Since driving is one of the safest and preferred ways for the elderly to travel, paying close attention to the accurate assessment of the elderly's driving ability can prevent traffic accidents in this age group. The purpose of this study was to identify and introduce practical tools for drive assessment fitness of the elderly. This systematic review was conducted according to Cochrane methodology and reported findings according to PRISMA. The following databases were searched from PubMed, ISI web of knowledge, Scopus, ProQuest, Medlib, SID, Magiran, Iran doc, and Iran Medex based on the population intervention comparison outcome method. The total records involving 12 main tools were assessed from 26 selected records in the final evaluation. The research findings indicated the selection of seven tools in the psycho-cognitive function domain such as TMT-B, Clock Drawing Test, MAZE, Montreal Cognitive Assessment, GDS-15, MMSE, and ACE-R, three tools in the sensory function domain such as Snellen, Confrontation Visual field, and Whispered Voice Test, and also two tools in motor function domain such as Rapid pace walk, and Manual test of the range of motion. The findings led to selecting practical, accurate, and fast tools for widespread use for the assessment of driving competencies of the elderly. Therefore, it is recommended that the selected tools be used in practical batteries to assess the driving skills of the elderly.

## Keywords:

Aged, competency, driver, driving, fit to drive, elder

## Introduction

According to the WHO, the population of over 60 in the world will almost double from 2015 to 2050 and will increase from 12% to 22%.<sup>[1]</sup> Prolonging life expectancy presents important opportunities for the elderly, family and even society as a whole,<sup>[2]</sup> which is expected to remain active as well as more. However, health is the determining factor in turning this stage of life into an opportunity or a threat.<sup>[3]</sup> Hence since 2015, the World Health Organization (WHO) has been using an approach called healthy

aging in the functional domains of the elderly.<sup>[2]</sup> Healthy aging is defined as the process of optimizing opportunities for mental, physical, and social health to enable older people to engage in social activities without discrimination and to enjoy a good, independent life.<sup>[4]</sup> As noted in WHO's report, there is no such thing as normal aging, and differences in the abilities and health needs of older people are not accidental and depend on their lifestyle throughout their lives.<sup>[4]</sup> An active lifestyle and maintaining social activities of the elderly is directly related to the process of healthy aging. In fact, with age, the type

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com

**How to cite this article:** Bahrampouri S, Khankeh HR, Hosseini SA, Mehmandar M, Ebadi A. Introducing practical tools for fit to drive assessment of the elderly: A step toward improving the health of the elderly. *J Edu Health Promot* 2021;10:463.

Health in Emergency and Disaster Research Center, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran, <sup>1</sup>Police University, NAJA Traffic Police, Tehran, Iran, <sup>2</sup>Behavioral Sciences Research Center, Life Style, Baqiyatallah University of Medical Sciences, Tehran, Iran, <sup>3</sup>Nursing Faculty, Baqiyatallah University of Medical Sciences, Tehran, Iran

## Address for correspondence:

Prof. Hamid Reza Khankeh, Health in Emergency and Disaster Research Center, University of Social Welfare and Rehabilitation, Kudakyar Alley, Daneshju Blvd., Evin, Tehran, Iran. E-mail: hamid.khankeh@ki.se

Received: 21-12-2020

Accepted: 08-03-2021

Published: 31-12-2021

of social activity of individuals changes.<sup>[5]</sup> In addition, globalization, technology development, urbanization, migration, and changing gender norms have also directly and indirectly affected the lives of the elderly.<sup>[1]</sup>

Driving is the most common, reliable and preferred mode of transportation for the elderly and provides them with freedom, safety, and access to other place and people.<sup>[6]</sup> This has led to a significant increase in the number of elderly drivers.<sup>[7]</sup> Driving is a continuous and simultaneous interaction between the driver and the environment, both of which can affect driving safety. Driving is done by performing coordinated tasks in the driver's sensory, cognitive, and musculoskeletal systems.<sup>[7]</sup> Aging reduces driving-related capabilities and coordination between the organs involved while driving, as well as driving skills. On the other hand, in the event of an accident, older drivers will suffer more damage.<sup>[8]</sup> It is also important to note that reducing age-related natural abilities, illness, medication, and other underlying factors can also affect the ability to drive in old age.<sup>[9-11]</sup> Therefore, the importance of assessment of driving competency of the elderly is not hidden from anyone. However, there is currently no single, agreed-upon instrument or tool in the world for measuring the driving skills of the elderly or predicting their driving outcomes.<sup>[12]</sup> Therefore, researchers in different parts of the world, according to self-goals and facilities, have combined different tools to be able to extract the most appropriate and accurate method for evaluating this capability.<sup>[13]</sup> Therefore, this study was conducted with the aim of introducing practical tools to assess the driving competencies in the elderly.

## Materials and Methods

A systematic review (SR) was conducted to provide an overview of the best available evidence. We did the search strategy based on the population intervention comparison outcome method and included all relevant articles without time limitation. The procedure was, then, followed by the PRISMA methods [Figure 1]. Nine databases including the PubMed, ISI web of knowledge, Scopus, ProQuest, Medlib, SID, Magiran, Iran doc, and IranMedex were searched. All databases were searched using the following keywords: Old People, Elder, Aged, Aged +80, Geriatric, Senior, Ageing, Aging, Old Age, Driver, Automobile Driving, Scale, measure, battery, screening tool, tool, Screening, Validity, Reliability, Questionnaire. A researcher (S. B.) conducted literature search. After removing duplications by Endnote X7 software, the titles, abstract, and full texts identified by the database searches were independently reviewed for eligibility by two researchers (S. B. and A. E.). In case of any dispute for screening and selection process, those were solved by discussion between the two researchers.




Figure 1: Research flow based on PRISMA methods

The review included all records linked with research objects, in English and Persian language, which (1) No need for specialized training; (2) Portability and easy portability; (3) No need for equipment or computer programs; (4) exist evidence of use in evaluating elderly drivers, and (6) psychometric criteria for the tool. Studies only published as an abstract were excluded due to the very high risk of bias in such designs.

The methodological quality of the included studies was critically assessed by the Downs and Black scale.<sup>[14]</sup> It is a checklist of 27 items to evaluate the risk of bias with each item scored "yes" (1 point), "no" (0 point) or "unable to determine" (0 point). The scale includes the aspects of reporting, external validity, internal validity, and power. The Downs and Black scale total, for the current review, was modified to 15 items, some items were not scored because of nonconformity in the items and included studies. Higher scores showed higher method quality.

Data were collected using data extraction form. If necessary, we contacted the authors for missing data, clarification, or both.

The study did not involve contact with humans, so the need for ethical approval was assess by university of Social Welfare and Rehabilitation Research Ethics Committee. This SR was not registered.

## Results

In total, data of 26 studies were included for qualitative analyses. Summary of results is indicated in Table 1. Relevant results such as design, sample size, population, and all tools that used in the study were extracted from

**Table 1: Summary of the reviewed studies**

Authors/year	Name of article or battery	Tools domain			
		Cognition/Mental	Sensory	Motor	Other
Engum <i>et al.</i> 1988 <sup>[53]</sup>	CBDI (Cognitive behavioral driver's inventory)	Visual reaction differential response visual reaction differential response reversed visual discrimination differential response II visual scanning	-	-	-
Wang <i>et al.</i> 2003/ AMA <sup>[54]</sup>	ADReS (Assessment of Driving-Related Skills)	TMT-B CDT	Snellen Confrontation test	ROM Muscle strength test RPW	-
Kantor <i>et al.</i> 2004 <sup>[55]</sup>	Model for predicting on the-road performance	TMT-B RT MMSE	-	Grip strength (WFL, not WFL)	-
Ball <i>et al.</i> 2006 <sup>[56]</sup>	Performance-Based Measures	Cued/delayed recall Scan test UFOV MFVP TMT-A TMT-B	-	RPW mobility questionnaire Foot-tap test ROM Arm reach	-
Langford <i>et al.</i> 2006 <sup>[57]</sup>	A re-assessment of older drivers as a road safety (GRIMPS and DriveABLE battery)	Test visual closure sub-test MFVP Cued recall, delayed recall scan test UFOV TMT-A TMT-B	Visual acuity (standard and low contrast)	RPW Arm reach Foot-tapping test Head neck rotation	-
Eby <i>et al.</i> 2007 <sup>[58]</sup>	Comprehensive battery of assessment instruments for older drivers	Ruler drop test MVPT CDTMMSE TMT-A TMT-B	Pelli-robson test Snellen "E" test Amsler Grid test Randot Stereoacuity test	RPW Arm reach test Clock reading test Jamar dynamometer test Nine-hole peg test	Medical history Road test
Molnar <i>et al.</i> 2007 <sup>[59]</sup>	Assessment Battery	MMSE Ottawa driving and dementia	-	Timed Toe Tap Test	Driving habits Medical
Stav <i>et al.</i> 2008 <sup>[60]</sup>	Predictive model	UFOV MMSE	contrast sensitivity slide-B	RPW	-
Wood <i>et al.</i> 2008 <sup>[61]</sup>	Multidomain tests	UFOV Color choice RT TMT-B	Dot motion threshold score	Knee extension strength score Sway path length	-
Keay <i>et al.</i> 2009 <sup>[62]</sup>	Factors that predict stopping or restricting driving in older drivers	GDS TMT-A Beery-buktenica developmental test of VMI	Pelli-Robson Cs chart	-	Driving history

Contd...

Table 1: Contd...

Authors/year	Name of article or battery	Tools domain			
		Cognition/Mental	Sensory	Motor	Other
Mathias <i>et al.</i> 2009 <sup>[63]</sup>	Cognitive predictors of unsafe driving	UFOV Ergovision Complex RT task Paper folding task Dot counting WMS visual reproduction Computerized Visual Attention Task	-	-	-
Barrash <i>et al.</i> 2010 <sup>[64]</sup>	Neuropsychological Tests	CFT TMT-A BD (Wechsler Adult Intelligence Scale-III BD test)	-	-	-
Dobbs <i>et al.</i> 2010 <sup>[65]</sup>	SIMARD-MD	Number conversion part a score Supermarket task score Repeat of word list score	-	-	-
AMA-2010 <sup>[66]</sup>	PODS (Plan For Older Drivers' Safety)	MoCA CDT Maze TMT-B	-	IADL ROM Get up and go Proprioception RPW	Driving history Medical history
Munro <i>et al.</i> 2010 <sup>[67]</sup>	Model Predicting Rate of Lane-Change Failure	Brief test of attention score Beery-Buktenicka test of visual-motor integration score	-	-	-
Dawson <i>et al.</i> 2010 <sup>[68]</sup>	Off-road neuropsychological battery	CFT-Copy CFT-Recall BD subtest of the Wechsler adult intelligence scale	Snellen chart	GP	-
Unsworth <i>et al.</i> 2012 <sup>[45]</sup>	OT-DORA battery	Road law and road craft test OT drive home Maze Test MMSE	Snellen chart Visual confrontation test Motor sequences screen - Selected test of Proprioception Short form McGill pain questionnaire and visual analogue scale and pain Diagram whispered voice test	Simulated accelerator-brake test Berg balance scale The motricity index right heel pivot test ROM - goniometry Tardieu scale of muscle tone Muscle strength scale	Medical drug
Anstey <i>et al.</i> 2012 <sup>[69]</sup>	Multifactorial Model of Driving Safety	Card rotation Paper folding Gestalt completion Snowy pictures Concealed words TMT-A TMT-B Number comparison task Digit symbol matching Letter cancellation tasks CRT	Bailey-lovie (logMAR) chart UFOV (Subtest 2)	-	Hazard perception test Hazard change detection task

Contd...

Table 1: Contd...

Authors/year	Name of article or battery	Tools domain			
		Cognition/Mental	Sensory	Motor	Other
Anderson <i>et al.</i> 2012 <sup>[70]</sup>	Neuropsychological Assessment of Driving Safety Risk in Older Adults With and Without Neurologic Disease	Digit-span backwards (adapted from WAISIII)			
		UFOV	Snellen chart	-	Road test
		GP	Pelli-Robson chart		
		CFT-copy			
		BD			
Dickerson <i>et al.</i> 2013 <sup>[71]</sup>	Assessment Tools Used by Driver Rehabilitation Specialists	BVRT-E			
		CFT-recall			
		Brake reaction	Road signs (Optec) b	ROM	Road test
		UFOV		Muscle tone	
		MMSE			
		TMT-A			
		TMT-B			
		MVPT			
		CDT			
		Letter-number cancellation			
Chaudhary <i>et al.</i> 2013 (NHTSA) <sup>[72]</sup>	Evaluating older drivers' skills	SBT			
		EFT	-	FRS	-
		RO-CFT			
		BDT			
		LCT			
		MNT			
		MVPT			
		Paper folding test			
		MDRS			
		TMT-A			
		TMT-B			
		BVRT			
		Cognitive flexibility test			
		SBT			
		Rey auditory verbal learning test			
		MMSE			
		CDR			
		CDT			
		TSRT			
		Nordic SDSA			
DST of the NAB					
Depression Scale					
CogStat					
Bowers <i>et al.</i> 2013 <sup>[73]</sup>	Clinical prediction of at-risk older drivers	MMSE	ETDRS chart	-	-
		UFOV	MARS chart		
Ferreira <i>et al.</i> 2013 <sup>[74]</sup>	Cognitive and psychomotor tests as predictors of on-road driving ability in older primary care patients	ACE-R	-	SDB	-
		SDSA			
		UFOV			
Ghasemi <i>et al.</i> 2015 <sup>[75]</sup>	Visual status	-	Snellen chart	-	-
			Confrontation test		
			D15 test		
			Pelli-Robson letter chart		

Contd...

Table 1: Contd...

Authors/year	Name of article or battery	Tools domain			
		Cognition/Mental	Sensory	Motor	Other
NHTSA-2016 <sup>[17]</sup>	Clinician's guide to assessing and counseling older drivers	CDT TMT-B MoCA maze test	Visual acuity Visual field	RPW ROM	-
Urlings <i>et al.</i> 2018 <sup>[12]</sup>	Predictive battery of tests for fitness to drive screenings	Knowledge of road signs	Snellen chart	Functional reach test	-

SMAS=short michigan alcoholism screening test, MVA=Motor vehicle administration, SIMARD-MD=Screen for the identification of cognitively impaired medically at-risk drivers, PODS=Plan for older drivers safety, MVPT=Motor-free visual perception test, TMT-A=Trail making test-A, TMT-B=Trail making test-B, MMSE=Mini mental state exam, VMI=Visualisation of missing information, MFVP=Motor free visual perception, MVPT=Motor free visual perceptual test, GDS=Geriatric depression score, HVL=Hopkins verbal-learning test, IVA=Integrated visual and auditory, CFT=Complex figure test, BD=Block design, MoCA=Montreal cognitive assessment, GP=Grooved pegboard, SBT=Short blessed test, EFT=Embedded figures test, RO-CFT=Rey-Osterrieth CFT, BDT=Block design test, LCT=Letter cancellation test, MNT=Maze navigation test, BVRT=Benton visual retention test, CDR=Clinical dementia rating, CDT=Clock drawing test, TSRT=Traffic sign recognition test, SDSA=Stroke driver screening assessment, DST=Driving scenes test, NAB=Neuropsychological assessment battery, ACE-R=Addenbrooke's cognitive examination revised, FRS=Functional Rating Scale, SDB=Senior drivers battery, VMI=Visual motor integration, ROM=Range of motion, RT=Reaction time, MDRS=Mattis Dementia Rating Scale, WFL=within functional limits., ETDRS=Early Treatment of Diabetic Retinopathy, MOMSSE=Mattis Organic Mental Status Syndrome Examination, ADL=activities of daily living, IADL=Instrumental activities of daily living, POMA=Performance Oriented Mobility Assessment, UFOV=useful field of view, WMS=Wechsler Memory Scale, OT-DORA=Occupational Therapy Driver Off-Road Assessment, CRT=choice reaction time, WAISIII=Wechsler Adult Intelligence Scale-Third Edition, BVRT-E=Benton Visual Retention Test

the selected records based on our research inclusion criteria. Selected studies led us to 12 main tools in three areas: sensory, psycho-cognitive, and motor. The list of tools of this study is shown in Table 2.

## Discussion and Conclusion

The findings of this study led to the introduction of 12 practical tools in the sensory, motor, and cognitive domains to assess the ability of the elderly to drive safely. We continue this section based on three main domains of study.

### Psycho-cognitive domain

The TMT-B has been widely used since 1944 for neuropsychological studies in clinical and research settings in Iran and the world. The validity has been evaluated and confirmed in various studies.<sup>[15,16]</sup> This test is used to assess driving competency and is a good predictor of driving accidents.<sup>[12,17,18]</sup> It is a pen and paper test. The participant is asked to connect a series of numbers and letters scattered on the page to each other alternately by a line. The time required to complete the test is recorded.<sup>[17]</sup>

The maze test is a pen and paper test and has different types. In this study, Dr. Snellger's maze test was selected. This test has been introduced in previous studies as a good predictor of driving accidents in the elderly living in the community.<sup>[19,20]</sup> In Iran, the mental maze test has been used to assess students' perceptual-motor skills.<sup>[21]</sup> To perform this test, the client must first complete a simple guide maze to familiarize themselves with the test. The main maze is then given to him. The test completion time is recorded in seconds.<sup>[17]</sup>

The MMSE was designed in 1975 by Folstein *et al.*<sup>[22]</sup> Since then, it has been used many times in various clinical and

research settings.<sup>[23]</sup> This test consists of 5 areas and 11 questions that examine the various dimensions of the subject's psychological and cognitive areas. Seyedian *et al.*<sup>[24]</sup> have reported its validity in Iran. The use of this tool to predict the driving status of the elderly has been recommended in previous studies.<sup>[25]</sup> This test takes about 5-10 min. The total score of this test is 30 points.

Clock Drawing Test is a very short and simple test for diagnosing cognitive disorders.<sup>[26]</sup> The validity and reliability of the test has been reviewed and confirmed in different countries.<sup>[27,28]</sup> In Iran, the validity and reliability of the test were reported by Sadeghipour Roodsari *et al.* (Reliability between evaluators 0.964 and kappa coefficient of 0.554 and concurrent validity of 0.782 with MMSE test).<sup>[29]</sup> This test is one of the most widely used tests to assess the driving status of the elderly, which has been used in many clinical and research settings.<sup>[17,30]</sup> However, so far 30 scoring methods have been reported for this test.<sup>[31]</sup> In order to perform this test, the subject is asked to draw a wall clock with all the numbers while the hands of the clock show 10 min past 11.<sup>[17]</sup>

The Montreal Cognitive Assessment Test (MOCA) is a cognitive screening questionnaire that measures cognitive skills over 10 min. Nasreddine *et al.* was designed it to diagnose mild cognitive impairment.<sup>[32]</sup> MOCA is one of the common tests used for cognitive assessment of the elderly in Iran and the world.<sup>[33,34]</sup> Its validity and reliability have been studied in the study of Chehrehnegar *et al.* in 2011.<sup>[35]</sup> It is also one of the most widely used tests to assess the condition of elderly drivers and predict their driving status.<sup>[17,36,37]</sup>

The Elderly Depression score (GDS-15), assesses depressive symptoms in the elderly. It is also one of the most widely used tests to study depression in Iran and the

**Table 2: List of selected tools**

Motor function domain	Sensory function domain	Psycho-cognitive function domain
RPW	Snellen	TMT-B
Manual test of range of motion	Confrontation visual field	Maze test
	Whispered voice test	CDT
		MoCA
		MMSE
		GDS-15
		ACE-R

RPW=Rapid pace walk, TMT-B=Trail making test-B, CDT=Clock drawing test, MoCA=Montreal cognitive assessment, MMSE=Mini-mental status exam, GDS=Geriatric depression score, ACE-R=Addenbrooke's Cognitive Examination-Revised

world.<sup>[38,39]</sup> Malakouti *et al.* reviewed the standardization of this questionnaire. In Iranian elderly and Cronbach's alpha coefficients of 0.9 and correlation coefficient of 0.58 were reported for the test-retest method.<sup>[39]</sup> The use of this test to evaluate the symptoms of depression in elderly drivers has also been reported.<sup>[40]</sup> In this test, the subject is asked to answer a questionnaire of 15 questions with two yes/no options. Counting the number of questions with a negative answer determines the test score. Addenbrooke's Cognitive Examination-Revised is a developed form of the MMSE which memory, language, and visuospatial subtests were extended, and fluency subtest was added. This summary version has the appropriate sensitivity and specificity for cognitive screening. psychometric properties of this test evaluated by Pouretamad *et al.* in Iran ( $r = 0.5, a = 0.84$ ).<sup>[41]</sup>

In this study, some psycho-cognitive tools were removed due to overlap with the measured components by selected tools. For example, the TMT-B test measures all components of the TMT-A test and in addition, assesses other components such as working memory, divided attention, and selective attention. Therefore, only the TMT-B test was retained as selected tools. In addition, the practicality and familiarity of the selected tools are other issues that are covered by the findings of this section.

### Sensory domain

Measuring visual acuity is the most basic and important method of measurement in visual examinations. The Snellen chart is considered as the most important clinical criterion indicating the quality of vision.<sup>[42]</sup> Dr. Hermann Snellen of the Netherlands in 1862 was developed first visual acuity chart. A special type of Snellen vision chart is the "rotational E" chart. All the symbols on the chart are capital letters E, located at 90° angles in different spatial directions. This test is the current standard in measuring visual acuity in Iran due to its availability, speed, and convenience.<sup>[43]</sup> The use of this test in assessing driving competence is common in Iran and the world and is currently used as a standard test for assessing visual acuity in examining assessment centers of police.<sup>[17,44]</sup>

The Confrontation Visual field is the simplest test to assess the field of view. This test has been used in

various studies to evaluate the field of vision of various age groups and work, especially drivers in Iran and the world.<sup>[17,45,46]</sup> According to the executive instructions, the criteria for medical qualification of applicants for various types of driving licenses in Iran, has been proposed as the only criterion for assessing the field of vision.<sup>[43]</sup> To perform this test, the examiner sits at a distance of 3 feet or 90 cm from the subject and at a level equal to him. Each eye is examined separately. The subject is asked to close his or her right eye. The examiner closes his left eye at the same time. Each (subject and examiner) must keep their gaze fixed on the other party's open eye. The examiner moves his hand in the subject's field of vision and points to the number 1, 2, or 5 with his fingers. It does this for every right, left, up, and down quadrilateral. Ask the subject to say the number shown.<sup>[47]</sup> The examiner shows any visual field defects by shading the area with a visual field defect.<sup>[17]</sup>

Since the introduction of Swan *et al.* in 1985, Whisper test has been used many times in clinical and research settings in different countries to assess the hearing status of drivers. Due to its ease of implementation and the lack of special tools and equipment, this test is one of the common tests for hearing assessment of drivers, especially elderly drivers.<sup>[48,49]</sup> In order to perform this test, the examiner stands at a distance of one hand (about 60 cm) from the right ear or the left ear from behind the patient. The nonexperimental ear canal is blocked by pressing a finger on the earlobe. The tester whispers the number-word and the combination of numbers and words in the exhaled state. The numbers are given in pairs to the test ear of the individual, and if there is more than 1 incorrect answer, the test result is considered positive (referral). In this case, 5 correct answers and more should be mentioned to be considered a negative screening result. The same process is repeated in the opposite ear.<sup>[50]</sup>

In this study, contrast sensitivity assessment, which is one of the important components for measuring driving ability, was omitted due to its specialization and the need for specialized equipment. However, examination of visual acuity and field can be associated with the diagnosis of primary vision problems and if necessary,

a specialized examination by an ophthalmologist can be used. Due to the fact that the selected tools in this study are used for screening, its ease of implementation and of course, the necessary accuracy should be considered. Since E-chart, Confrontation Visual field test and whispering tests are common and widely used tests in Iran and are also recommended to assess drivers' abilities, it seems that they can evaluate the components of this field well.

### Motor domain

Rapid pace walk evaluates the components of the motor domain in <3 min. The results of various studies indicate the predictive power of this test to assess driving competence, especially in the elderly.<sup>[17,51]</sup> Due to its easy implementation and without the need for specialized equipment, this test is one of the tools for assessing the motor system in a number of tools for assessing the competence of elderly drivers.<sup>[17,52]</sup> Reliability among the evaluators of this test was reported to be 0.87, 0.605.<sup>[48]</sup> In order to perform this test, the subject is asked to walk a distance of 10 feet, equivalent to 3 meters, in the fastest possible time and return to the starting point. If the client uses a cane or walker, he/she is asked to use his/her mobility aids during the test; and is recorded in the test results sheet. Test duration is recorded in seconds.

Range of Motion (ROM) examines the ROM of driving-related joints, including the neck, shoulders, elbows, fingers, and ankles. There are several methods for measuring the ROM of joints. However, in this study, due to the importance of speed and ease of testing, a qualitative method of measuring ROM was used. This method has also been used in the Clinical Guide for the Review and Advice of Elderly Drivers developed by the American Aging Association.<sup>[17]</sup> In order to perform the test, the subject is asked to: (1) Neck rotation: such as when turning the head back to move backward or to park. The subject is asked to turn his head backward. Now do the same for the other party. (2) Shoulder and elbow flexion: Imagine you are in control of a car. Do a full turn to the right and a full turn to the left. (3) Bending a finger: Fist both hands. (4) Plantar flexion of the ankle: Pretend you are depressing the accelerator pedal. Now repeat the same for the other leg. (5) Ankle dorsiflexion: Pull your toes toward your body. In other words, the movement of the back (dorsal) part of the foot towards the front of the leg. Range scoring is done simply by two options, "normal" and "abnormal" for each side. ROM is considered abnormal when the ROM is limited or despite pain or good ROM is accompanied by pain or pause.<sup>[17]</sup>

Despite the various tools for assessing motor and functional status, there is no agreed test or method for assessing elderly driving competence among researchers and specialists in the fields of rehabilitation and

transportation. The use of selected tests to assess the balance and strength of the lower limbs as well as ROM are common in Iran and the world and are most widely used tests to assess the motor function in elder drivers. Ease of execution, accuracy and no need for special equipment or tools are the features that exist in these two tools.

**Limitation and Strengths:** We attempted to use comprehensive search strategy and search several databases as well as grey literature; however, the possibility that evidence included in this review is subject to publication and related bias cannot be ruled out. In addition, the most important limitation of this study was the lack of access to the full text of some documents.

### Conclusion

Easy and accurate diagnosis of older drivers' disabilities with appropriate tools can be used to prevent traffic accidents and their consequences. Therefore, the use of practical tools in this field can help the health of elderly drivers, health system and the public. The findings of this study can be used to evaluate the driving abilities of the elderly in police clinical examination centers. It is also suggested that the present tools be used in order to conduct additional studies on the construction of a battery for elder driver assessment in Iran.

### Acknowledgment

We would like to thank Dr. Poursadeghiyan for his assistance.

### Financial support and sponsorship

This study was part of a PhD thesis supported by the university of Social Welfare and Rehabilitation.

### Conflicts of interest

There are no conflicts of interest.

### References

1. WHO. Ageing and Health: World Health Organization; 2018. Available from: <http://www.who.int/news-room/fact-sheets/detail/ageing-and-health>. [Last accessed on 2021 Apr 12].
2. Beard JR, Officer AM, Cassels AK. The World Report on Ageing and Health. USA: Oxford University Press; 2016.
3. WHO. Facts on Ageing and Health.: World Health Organization; 2017. Available from: <http://www.who.int/features/factfiles/ageing/en/>. [Last accessed on 2021 Apr 12]
4. WHO. Other Dimensions of the NCD Crisis: From Mental Health, Ageing, Dementia and Malnutrition to Deaths on the Roads, Violence and Disability. Ten Years in Public Health 2007-2017; 2017. Available from: <https://www.who.int/publications/10-year-review/ncd-other-dimensions/en/> [Last accessed on 2021 Apr 12]
5. Marcum CS. Age differences in daily social activities. Research on aging. 2013 Sep; 35 (5):612-40.
6. Makizako H, Shimada H, Hotta R, Doi T, Tsutsumimoto K,



- Nakakubo S, et al. Associations of near-miss traffic incidents with attention and executive function among older Japanese drivers. *Gerontology* 2018;64:495-502.
7. Koppel S, Berecki-Gisolf J. Car licensing trends of the baby boomer cohort (b. 1946-1965) Compared to earlier birth cohorts: Effects on the driving population in the state of Victoria, Australia. *Traffic Inj Prev* 2015;16:657-63.
  8. Dobbs BM. Aging baby boomers – A blessing or challenge for driver licensing authorities. *Traffic Inj Prev* 2008;9:379-86.
  9. Biglari H, Ebrahimi MH, Salehi M, Poursadeghiyan M, Ahmadzadeh I, Abbasi M. Relationship between occupational stress and cardiovascular diseases risk factors in drivers. *Int J Occup Med Environ Health* 2016;29:895-901.
  10. Abdoli N, Farnia V, Salemi S, Tatari F, Shakeri J, Basanj B, et al. Predictors of substance abuse among risky drivers: The role of personality characteristics and mental health. *J Educ Health Promot* 2018;7:163.
  11. Poursadeghiyan M, Mazloumi A, Saraji GN, Niknezhad A, Akbarzadeh A, Ebrahimi MH. Determination the levels of subjective and observer rating of drowsiness and their associations with facial dynamic changes. *Iran J Public Health* 2017;46:93-102.
  12. Urlings JHJ, Cuenen A, Brijs T, Lutin M, Jongen EMM. Aiding medical professionals in fitness-to-drive screenings for elderly drivers: Development of an office-based screening tool. *Int Psychogeriatr* 2018;30:1211-25.
  13. Gentzler MD, Smither JA. A literature review of major perceptual, cognitive, and/or physical test batteries for older drivers. *Work* 2012;41 Suppl 1:5381-3.
  14. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* 1998;52:377-84.
  15. Banazadeh N, Khalili N, Mazhari S, Anaridokht F. Validity and reliability of a persian version of the screen for cognitive impairment in psychiatry scale in patients with bipolar type one disorder. *Afzalipour J Clin Res* 2017;2:14-22.
  16. Tombaugh TN. Trail making test A and B: Normative data stratified by age and education. *Arch Clin Neuropsychol* 2004;19:203-14.
  17. National Highway Traffic Safety Administration. *Clinician's Guide to Assessing and Counseling Older Drivers*. 3<sup>rd</sup> edition. (Report No. DOT HS 812 228) Washington, DC: US Department of Transportation; 2016.
  18. Staplin L, Gish KW, Sifrit KJ. Using cognitive status to predict crash risk: Blazing new trails? *J Safety Res* 2014;48:19-25.
  19. Staplin L, Gish KW, Lococo KH, Joyce JJ, Sifrit KJ. The maze test: A significant predictor of older driver crash risk. *Accid Anal Prev* 2013;50:483-9.
  20. Ott BR, Davis JD, Papandonatos GD, Hewitt S, Festa EK, Heindel WC, et al. Assessment of driving-related skills prediction of unsafe driving in older adults in the office setting. *J Am Geriatr Soc* 2013;61:1164-9.
  21. Soleymani E, Sorbi MH. The comparison between mental maze function and bilateral nerve transmission learning (based on the mirror drawing test) on good and poor audiovisual memory of students. *Neuropsychology* 2018;3:9-24.
  22. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *Journal of psychiatric research*. 1975;12(3):189-98.
  23. Monroe T, Carter M. Using the Folstein Mini Mental State Exam (MMSE) to explore methodological issues in cognitive aging research. *Eur J Ageing* 2012;9:265-74.
  24. Seyedian M, Falah M, Norozian M, Nejat S, Delavar A, Ghasemzadeh HA. Preparing and determining the validity of the Persian version of the mini mental status test (Persian). *Sci J Med Syst Organ Islam Repub Iran* 2007;25:408-14.
  25. Crizzle AM, Classen S, Bédard M, Lanford D, Winter S. MMSE as a predictor of on-road driving performance in community dwelling older drivers. *Accid Anal Prev* 2012;49:287-92.
  26. Aprahamian I, Martinelli JE, Neri AL, Yassuda MS. The clock drawing test: A review of its accuracy in screening for dementia. *Dement Neuropsychol* 2009;3:74-81.
  27. Nair AK, Gavett BE, Damman M, Dekker W, Green RC, Mandel A, et al. Clock drawing test ratings by dementia specialists: Interrater reliability and diagnostic accuracy. *J Neuropsychiatry Clin Neurosci* 2010;22:85-92.
  28. Can SS, Gencay-Can A, Gunendi Z. Validity and reliability of the clock drawing test as a screening tool for cognitive impairment in patients with fibromyalgia. *Compr Psychiatry* 2012;53:81-6.
  29. Sadeghipour Roodsari M, Akbari Kamrani AA, Foroughan M, Mohammadi F, Karimloo M. Validity and reliability of the clock drawing test in older people (Persian). *Salmand Iran J Age* 2013;8:48-53.
  30. Ketelaars L, Pottel L, Lycke M, Goethals L, Ghekiere V, Santy L, et al. Use of the Freund clock drawing test within the Mini-Cog as a screening tool for cognitive impairment in elderly patients with or without cancer. *J Geriatric Oncol* 2013;4:174-82.
  31. Spenciore B, Alves H, Charchat-Fichman H. Scoring systems for the clock drawing test: A historical review. *Dement Neuropsychol* 2017;11:6-14.
  32. Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, et al. The Montreal Cognitive Assessment, MoCA: A brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* 2005;53:695-9.
  33. Yazdani S, Sharifi S, Foroughipour M, Kamyabgol A. The visuospatial episodic memory in Persian-speaking patients with multiple sclerosis. *J Mazandaran Univ Med Sci* 2019;29:82-95.
  34. Davis DH, Creavin ST, Yip JL, Noel-Storr AH, Brayne C, Cullum S. Montreal Cognitive Assessment for the diagnosis of Alzheimer's disease and other dementias. *Cochrane Database Syst Rev* 2015;(10):CD010775.
  35. Chehrehnegar N, Shams F, Zarshenas S, Kazemi F. Evaluating the reliability of the montreal cognitive assessment test and its agreement with mini mental state examination among healthy elderly (Persian). *J Res Rehabil Sci* 2012;7(5(SUPPLEMENT)); 674-80.
  36. Ma'u E, Cheung G. Ability of the maze navigation test, Montreal Cognitive Assessment, and trail making tests A & B to predict on-road driving performance in current drivers diagnosed with dementia. *N Z Med J* 2020;133:23-32.
  37. Kandasamy D, Williamson K, Carr DB, Abbott D, Betz ME. The utility of the Montreal Cognitive Assessment in predicting need for fitness to drive evaluations in older adults. *J Trans Health* 2019;13:19-25.
  38. Nyunt MS, Fones C, Niti M, Ng TP. Criterion-based validity and reliability of the Geriatric Depression Screening Scale (GDS-15) in a large validation sample of community-living Asian older adults. *Aging Ment Health* 2009;13:376-82.
  39. Malakouti K, Fathollahi P, Mirabzadeh A, Salavati M, Kahani S. Validation of geriatric depression scale (GDS-15) in Iran. *Res Med* 2006;30:361-9.
  40. Crizzle AM, Myers AM, Roy EA, Almeida QJ. Drivers with Parkinson's disease: Are the symptoms of PD associated with restricted driving practices? *J Neurol* 2013;260:2562-8.
  41. Pouretamad HR, Khatibi A, Ganjavi A, Shams J, Zarei M. Validation of Addenbrooke's cognitive examination (ACE) in a Persian-speaking population. *Dement Geriatr Cogn Disord* 2009;28:343-7.
  42. Alamdar M, Jafarzadehpur E, Mirzajani A, Yekta A. Prediction of the refractive error based on Snellen chart and OPD-scan III visual acuity. *J Paramed Sci Rehabil* 2018;7:33-43.
  43. The police force of the Islamic republic of Iran and ministry of health and medical education. [Executive instructions for physical

- and mental health (medical qualification) criteria for applicants for various types of driving licenses (Persian)]. Tehran: Health Regulations in Ministry of Health and Medical Education. 2009.
44. Behboudi H, Moghadam RS, Tiefeh N, Karkan MF. Vision disorders in drivers involved in traffic accidents. *J Ophthalmic Vis Res* 2017;12:451-2.
  45. Unsworth CA, Baker A, Taitz C, Chan SP, Pallant JF, Russell KJ, et al. Development of a standardised occupational therapy – Driver Off-Road Assessment Battery to assess older and/or functionally impaired drivers. *Aust Occup Ther J* 2012;59:23-36.
  46. Chan ML, Wong Y, Ng R, Koh GC. Medical conditions and driving fitness of older Singaporean taxi drivers. *Occup Med (Lond)* 2019;69:211-4.
  47. DVLA. Guidance: Visual Disorders: Assessing Fitness to Drive, Driver and Vehicle Licensing Agency. UK: Driver and Vehicle Licensing Agency; 2016. Available from: <https://www.gov.uk/guidance/visual-disorders-assessing-fitness-to-drive#minimum-standards-for-field-of-vision--all-drivers>. [Last accessed on 2020 May 30].
  48. Smith A, Marshall S, Porter M, Ha L, Bédard M, Gélinas I, et al. Stability of physical assessment of older drivers over 1 year. *Accid Anal Prev* 2013;61:261-6.
  49. Pozzi C, Lucchi E, Lanzoni A, Gentile S, Morghen S, Trabucchi M, et al. Why older people stop to drive? A cohort study of older patients admitted to a rehabilitation setting. *Aging Clin Exp Res* 2018;30:543-6.
  50. Farhadi M, Mahmoudian S, Phirozbakht M, Alaedini F, Rahimi F, Esmaelzadah M, et al. National Guide to Infant and Child Hearing loss Screening Program (Special instructions for experts and officials of comprehensive health service centers (urban/rural)) (Persian). Tehran: Ministry of Health and Medical Education; 2018.
  51. Posse C, McCarthy DP, Mann WC. A pilot study of interrater reliability of the Assessment of Driving-Related Skills: Older driver screening tool. *Top Geriatric Rehabil* 2006;22:113-20.
  52. Woolnough A, Salim D, Marshall SC, Weegar K, Porter MM, Rapoport MJ, et al. Determining the validity of the AMA guide: A historical cohort analysis of the assessment of driving related skills and crash rate among older drivers. *Accid Anal Prev* 2013;61:311-6.
  53. Engum ES, Cron L, Hulse CK, Pendergrass TM, Lambert W. Cognitive behavioral driver's inventory. *Cognitive rehabilitation*. 1988 Sep; 6 (5):34-50.
  54. Wang C, Kosinski CJ, Schwartzberg JG, Shanklin AV, Association AM. Physician's guide to assessing and counseling older drivers: American Medical Association; 2003.
  55. Kantor B, Mauger L, Richardson VE, Unroe KT. An analysis of an older driver evaluation program. *Journal of the American Geriatrics Society*. 2004;52 (8):1326-30.
  56. Ball KK, Roenker DL, Wadley VG, Edwards JD, Roth DL, McGwin Jr G, Raleigh R, Joyce JJ, Cissell GM, Dube T. Can high-risk older drivers be identified through performance-based measures in a Department of Motor Vehicles setting?. *Journal of the American Geriatrics Society*. 2006 Jan; 54 (1):77-84.
  57. Langford J, Koppel S, Charlton J, Fildes B, Newstead S. A re-assessment of older drivers as a road safety risk. *IATSS research*. 2006;30(1):27-37.
  58. Eby DW, Molnar LJ, Shope JT, Dellinger AM. Development and pilot testing of an assessment battery for older drivers. *Journal of safety research*. 2007;38(5):535-43.
  59. Molnar FJ, Marshall SC, Man-Son-Hing M, Wilson KG, Byszewski AM, Stiell I. Acceptability and concurrent validity of measures to predict older driver involvement in motor vehicle crashes: An emergency department pilot case-control study. *Accident Analysis & Prevention*. 2007;39 (5):1056-63.
  60. Stav WB, Justiss MD, McCarthy DP, Mann WC, Lanford DN. Predictability of clinical assessments for driving performance. *Journal of safety research*. 2008;39 (1):1-7.
  61. Wood JM, Anstey KJ, Kerr GK, Lacherez PF, Lord S. A multidomain approach for predicting older driver safety under in-traffic road conditions. *Journal of the American Geriatrics Society*. 2008;56 (6):986-93.
  62. Keay L, Munoz B, Turano KA, Hassan SE, Munro CA, Duncan DD, et al. Visual and cognitive deficits predict stopping or restricting driving: the Salisbury Eye Evaluation Driving Study (SEEDS). *Investigative ophthalmology & visual science*. 2009;50 (1):107-13.
  63. Mathias J, Lucas L. Cognitive predictors of unsafe driving in older drivers: a meta-analysis. *International psychogeriatrics*. 2009;21 (4):637.
  64. Barrash J, Stillman A, Anderson SW, Uc EY, Dawson JD, Rizzo M. Prediction of driving ability with neuropsychological tests: Demographic adjustments diminish accuracy. *Journal of the International Neuropsychological Society: JINS*. 2010;16 (4):679.
  65. Dobbs BM, Schopflocher D. The introduction of a new screening tool for the identification of cognitively impaired medically at-risk drivers: the SIMARD a modification of the DemTect. *Journal of primary care & community health*. 2010;1 (2):119-27.
  66. American Medical Association. Physician's guide to assessing and counseling older drivers. 2 ed. Chicago: American Medical Association; 2010.
  67. Munro CA, Jefferys J, Gower EW, Muñoz BE, Lyketsos CG, Keay L, et al. Predictors of lane change errors in older drivers. *Journal of the American Geriatrics Society*. 2010;58 (3):457-64.
  68. Dawson JD, Uc EY, Anderson SW, Johnson AM, Rizzo M. Neuropsychological predictors of driving errors in older adults. *Journal of the American Geriatrics Society*. 2010;58 (6):1090-6.
  69. Anstey KJ, Horswill MS, Wood JM, Hatherly C. The role of cognitive and visual abilities as predictors in the Multifactorial Model of Driving Safety. *Accident Analysis & Prevention*. 2012;45:766-74.
  70. Anderson SW, Aksan N, Dawson JD, Uc EY, Johnson AM, Rizzo M. Neuropsychological assessment of driving safety risk in older adults with and without neurologic disease. *Journal of clinical and experimental neuropsychology*. 2012;34 (9):895-905.
  71. Dickerson AE. Driving assessment tools used by driver rehabilitation specialists: Survey of use and implications for practice. *American Journal of Occupational Therapy*. 2013;67 (5):564-73.
  72. Chaudhary NK, Ledingham KA, Eby DW, Molnar LJ. Evaluating older drivers' skills. Preusser Research Group, Inc., 2013.
  73. Bowers AR, Anastasio RJ, Sheldon SS, O'Connor MG, Hollis AM, Howe PD, et al. Can we improve clinical prediction of at-risk older drivers? *Accident Analysis & Prevention*. 2013;59:537-47.
  74. Ferreira IS, Simões MR, Marôco J. The Addenbrooke's Cognitive Examination Revised as a potential screening test for elderly drivers. *Accident Analysis & Prevention*. 2012;49:278-86.
  75. Ghasemi M, Yazdi SHH, Heravian J, Jafarzadehpur E, Rezaee M. Comparison of visual status of Iranian military and commercial drivers. *Iranian Red Crescent Medical Journal*. 2015;17 (4).