



# BMJ Open Outcomes associated with older patients who present or develop delirium in the emergency department: protocol for a systematic review and meta-analysis

Sarah King <sup>1</sup>, Oddvar Uleberg <sup>1,2,3</sup>, Sindre A Pedersen,<sup>4</sup>  
Lars Petter Bjørnsen<sup>1,3</sup>

**To cite:** King S, Uleberg O, Pedersen SA, *et al.* Outcomes associated with older patients who present or develop delirium in the emergency department: protocol for a systematic review and meta-analysis. *BMJ Open* 2025;**15**:e095495. doi:10.1136/bmjopen-2024-095495

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-095495>).

Received 23 October 2024

Accepted 28 April 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

For numbered affiliations see end of article.

## Correspondence to

Dr Sarah King;  
[sarah.king@ntnu.no](mailto:sarah.king@ntnu.no)

## ABSTRACT

**Introduction** Delirium is commonly observed in older patients who are admitted to the emergency department (ED). Previous systematic reviews have identified poor outcomes associated with delirium in surgical, intensive care and other hospital settings, yet none have specifically considered the ED. This systematic review aims to examine associations between older patients who present or develop delirium in the ED and adverse outcomes within the hospital and after discharge.

**Methods and analysis** Searches will be conducted in MEDLINE, Embase, Web of Science, Cumulative Index to Nursing and Allied Health Literature, and the Cochrane Library. There will be no date or language restrictions. Key terms will include concepts related to delirium, the ED and older adults. Observational studies or non-intervention clinical studies will be included that compare outcomes in older patients (ie, ≥65 years) with and without delirium. Outcomes of interest will include length of hospital stay, non-home discharge (eg, nursing home/residential aged care facility), cognitive impairment, decreased physical function, mortality, readmission to hospital and quality of life measures. Two reviewers will independently screen the studies. Data extraction and quality assessment will be extracted by one reviewer and checked by a second reviewer, with any disagreements resolved by discussion or by a third reviewer. Where appropriate, data will be combined in a meta-analysis and a GRADE assessment will be made for each outcome. All methods will be guided by the Cochrane Handbook and the Centre for Reviews and Dissemination and reported following the Preferred Reporting Items for Systematic Review and Meta-Analysis statement as well as the recommendations set out by the Meta-analysis Of Observational Studies in Epidemiology group.

**Ethics and dissemination** As this systematic review will use published data, ethical approval is not required. The results will be disseminated through a peer-reviewed publication and conference presentations.

**PROSPERO registration number** CRD42024594975.

## INTRODUCTION

Delirium is a syndrome that is characterised by an acute decline in cognitive functioning and can present as hypoactive (eg, drowsy,

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ A rigorous systematic review methodology will be used to address the research question.
- ⇒ There will be no language or date restrictions to the search strategy.
- ⇒ The review includes observational studies, which are subject to bias and confounding.
- ⇒ Heterogeneity among the studies is expected and careful consideration of discrepancies will be needed to mitigate spurious conclusions.

lethargic, etc), hyperactive (eg, agitated, anxious, etc) or a fluctuating mix of these psychomotor subtypes. It has been estimated that almost one in four older acute medical hospital patients present or develop delirium.<sup>1</sup> Delirium is also commonly observed in the emergency department (ED),<sup>2</sup> with a recent systematic review reporting a prevalence rate of 15.2% in this setting (95% CI 12.5% to 18.0% (pooled crude rate based on 30 studies)).<sup>3</sup> Given that delirium is often unrecognised (eg, see El Hussein *et al*,<sup>4</sup> Lee *et al*,<sup>5</sup> Meged-Book *et al*,<sup>6</sup> and Barron and Holmes<sup>7</sup>), it is likely that some reported prevalence rates could represent underestimations.

Delirium is of concern because it has been associated with poorer outcomes in hospital settings, including mortality,<sup>8</sup> long-term cognitive decline,<sup>9 10</sup> and longer hospital stays and costs.<sup>11</sup> Previous systematic reviews (SRs) have been conducted that assess outcomes associated with delirium in surgical patients,<sup>12</sup> intensive care units<sup>13–15</sup> and hospitalised patients,<sup>16 17</sup> but none have specifically addressed the older ED patient, despite several new primary studies on this topic. A SR will be conducted by the Trondheim Emergency Department Research Group to examine associations between older patients who have a positive delirium diagnosis in the

ED and short- and long-term adverse outcomes. This SR specifically focuses on ED patients as they represent a distinct and vulnerable population. They often present with more acute medical or environmental stressors (eg, sepsis, polypharmacy) than those who develop delirium postoperatively or during a hospital stay. Moreover, the ED is a critical entry point for early delirium recognition—which could potentially improve prognosis and thus hospital length of stay. The ED thus presents specific challenges and opportunities for delirium research and interventions. By evaluating effect sizes of several outcomes derived from comparisons of older patients with and without delirium, it will be possible to assess the magnitude of the current problem in the ED context. This SR will be of relevance to clinicians and researchers who aim to prevent, detect and manage delirium in acute care patients.

## METHODS AND ANALYSIS

The primary research question of this SR is: What short- and long-term adverse outcomes are associated with older patients (ie,  $\geq 65$  years) who present or develop delirium in the ED compared with those who do not? To address this question, a SR will be undertaken according to guidance presented in the Cochrane Handbook<sup>18</sup> and the Centre for Reviews and Dissemination (CRD)<sup>19</sup> and reported following the Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA) statement<sup>20</sup> as well as the recommendations set out by the Meta-analysis Of Observational Studies in Epidemiology group.<sup>21</sup> This protocol was developed following the guideline of the Preferred Reporting Items for Systematic Review and Meta-analysis Protocols (PRISMA-P)<sup>22</sup> and is registered with PROSPERO (CRD42024594975).

## Inclusion and exclusion criteria

To be eligible for inclusion in the review, studies will have to meet all the following criteria in terms of population, exposure/intervention and comparator, outcomes, and study designs.

Studies that compare older patients (ie,  $\geq 65$  years) who are delirium positive (at ED arrival or during their visit) with those who are delirium negative will be eligible for inclusion. Delirium assessments must be made in the ED using a diagnostic tool or defined criteria (eg, Confusion Assessment Method, Brief Confusion Assessment Method, 3min Diagnostic Interview for Confusion Assessment Method, 4AT, Diagnostic and Statistical Manual of Mental Disorder Criteria criteria etc). Patients with delirium tremens, drug-induced delirium or hepatic encephalopathy will be excluded. Studies that specifically assess delirium after an ED visit will not be included. Studies examining older patients undergoing emergency surgery and older patients with COVID-19 will not be included as other SRs have been conducted in these population groups/settings.

Studies reporting one or more of the following short- (ie, within 30 days) or long-term (ie, up to 1 year) outcomes will be eligible for inclusion: length of hospital stay (or a measure of prolonged hospital stay), non-home discharge (eg, nursing home/residential aged care facility), cognitive impairment, decreased physical function, mortality (in hospital and out of hospital), readmission to hospital, and other quality of life (QoL) measures or an overall QoL assessment.

Prospective and retrospective observational studies or non-intervention clinical studies that compare outcomes between patients with and without delirium will be included. Controlled clinical trials or other types of intervention studies will be excluded, as well as case series, commentaries and letters to the editor.

Conference abstracts will only be included if there is no associated full publication and if adequate data are reported (eg, detailed information on the population, diagnostic criteria, and clear results with numerical data reported for those with and without delirium).

## Search strategy

The literature search will be conducted in MEDLINE, Embase, Web of Science, Cumulative Index to Nursing and Allied Health Literature, and the Cochrane Library. Key terms will include concepts related to delirium, the emergency department and older adults (online supplemental file 1). The search will not be limited by language or date. German, French and Scandinavian language papers will be assessed for inclusion and translated to English by members of the review team while Google Translate or DeepL will be used for all other languages. Grey literature will be searched in OAlster (OpenGrey, NYAM Grey Literature Report, and the British Library EThOS are not currently available for searching). Hand-searching will also be conducted to check references of relevant papers and reviews identified by the search. In addition, a citation search will be carried out to identify subsequent publications which have cited any of the included studies.

## Study selection and data extraction

The records retrieved from the searches will be de-duplicated in EndNote<sup>23</sup> and exported into Rayyan<sup>24</sup> for screening. During the first stage of study selection, titles and abstracts will be independently screened by two researchers against the inclusion/exclusion criteria. Full papers of all potentially relevant studies identified in the first stage will be screened by two reviewers working independently, with any discrepancies resolved by discussion between the reviewers or with the assistance of a third reviewer.

The number of studies identified by the search and excluded at various stages will be recorded and reported in a PRISMA study flow diagram.<sup>20</sup> After the second stage of screening, a table of excluded studies with detailed reasons for exclusion will be created.

Data to be extracted from each study will include information on the study (eg, study type, setting, country, objective, inclusion criteria, study methods including the delirium assessment tool used as well as who did the assessment and any information on their experience or training) and patient characteristics (eg, age, sex, main reason for ED visit, and for patients with delirium: subtype, severity), as well as the results and any potential confounding factors identified by the study authors. A data extraction form will be developed using Microsoft Excel<sup>25</sup> and piloted by two independent reviewers using a minimum of four studies. This process will also be used to check consistency in data extraction between the reviewers. The remaining studies will then be extracted by one reviewer and checked by a second reviewer. Any discrepancies will be resolved through discussion. If any relevant information is missing from the studies or if any data is unclear, the reviewers will attempt to contact the authors.

### Critical appraisal

The observational studies will be assessed using the Joanna Briggs Institute critical appraisal tools.<sup>26</sup> One reviewer will conduct the quality assessment, and a second reviewer will check the assessment with any discrepancies resolved through discussion or with the assistance of a third reviewer.

### Synthesis and analysis

Outcome comparisons between older patients with and without delirium will be presented.

Where possible and appropriate (ie, if the studies are clinically and statistically homogenous), data from the studies will be combined in a meta-analysis using Review-Manager (RevMan).<sup>27</sup> For continuous outcomes, means and SD will be collected and used to estimate study-specific and pooled mean differences with 95% CIs. For dichotomous outcomes, numerators and denominators will be collected, and Mantel-Haenszel risk ratios (RRs) or ORs and 95% CIs used to summarise effect sizes. If the scales/tools used to assess the outcomes differ between the studies, we will compute study-level standardised mean differences between comparison groups with 95% CIs. The results will be statistically pooled (where possible) using both fixed-effect and random-effects models and the results from these different models will be compared. Statistical heterogeneity will be assessed using the  $\chi^2$  test and the  $I^2$  statistic, and by examining the random effects between study variance ( $\text{Tau}^2$ ). If possible, sensitivity analyses will be performed to assess the robustness of results by excluding studies deemed to have a greater risk of bias. Comparisons between unadjusted and adjusted effect estimates will also be made. Funnel plots will be visually inspected to check for publication bias. For each outcome summarised, the GRADE system<sup>28</sup> will be used to provide an assessment of the quality of a body of evidence. One reviewer will rate the evidence using GRADEpro GDT<sup>29</sup>

and a second reviewer will check the assessment, with any discrepancies resolved through discussion.

Exploratory subgroup analyses will be performed where possible by date of publication, study type, delirium assessment method used in the study, assessor and training (if reported) and age of the study participants (if outcomes are categorised by age groups in the included studies). If there are any studies conducted in specific patient groups (eg, older patients with fractures), subgroup analysis may also be conducted by patient type.

A narrative synthesis of the studies will be performed if the studies are too diverse to perform a meta-analysis, or if only one study reported on an outcome and pooling cannot be undertaken.

### Patient and public involvement

None.

### ETHICS AND DISSEMINATION

As this systematic review will use published data, ethical approval is not required. The results will be disseminated through a peer-reviewed publication and conference presentations. If any minor amendments are made to this protocol during the review process, they will be reported in the final results of the paper.

### Author affiliations

<sup>1</sup>Trondheim Emergency Department Research Group (TEDRG), Department of Circulation and Medical Imaging, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

<sup>2</sup>Department of Research and Development, Norwegian Air Ambulance Foundation, Oslo, Norway

<sup>3</sup>Department of Emergency Medicine and Pre-Hospital Services, St. Olav's Hospital, Trondheim, Norway

<sup>4</sup>The Medicine and Health Library, Library Section for Research Support, Data and Analysis, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

X Oddvar Uleberg @uleodd

**Contributors** SK conceived the idea for the SR. All authors read and contributed to writing different aspects of the protocol. SK takes responsibility for the integrity of the work as a whole, from inception to published article. SK is responsible for the overall content as the guarantor.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, conduct, reporting or dissemination plans of this research.

**Patient consent for publication** Not applicable.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which

permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Sarah King <http://orcid.org/0000-0002-0637-7672>

Oddvar Uleberg <http://orcid.org/0000-0002-7913-5957>

#### REFERENCES

- Gibb K, Seeley A, Quinn T, *et al*. The consistent burden in published estimates of delirium occurrence in medical inpatients over four decades: a systematic review and meta-analysis study. *Age Ageing* 2020;49:352–60.
- Han JH, Wilson A, Ely EW. Delirium in the older emergency department patient: a quiet epidemic. *Emerg Med Clin North Am* 2010;28:611–31.
- Chen F, Liu L, Wang Y, *et al*. Delirium prevalence in geriatric emergency department patients: A systematic review and meta-analysis. *Am J Emerg Med* 2022;59:121–8.
- El Hussein MT, Hirst S, Stares R. Delirium in Emergency Departments: Is it Recognized? *J Emerg Nurs* 2021;47:809–17.
- Lee JS, Tong T, Chignell M, *et al*. Prevalence, management and outcomes of unrecognized delirium in a National Sample of 1,493 older emergency department patients: how many were sent home and what happened to them? *Age Ageing* 2022;51:afab214.
- Meged-Book T, Frenkel R, Nikonov A, *et al*. Delirium screening in the emergency department: evaluation and intervention. *Isr J Health Policy Res* 2024;13:16.
- Barron EA, Holmes J. Delirium within the emergency care setting, occurrence and detection: a systematic review. *Emerg Med J* 2013;30:263–8.
- Aung Thein MZ, Pereira JV, Nitchingham A, *et al*. A call to action for delirium research: Meta-analysis and regression of delirium associated mortality. *BMC Geriatr* 2020;20:325.
- Goldberg TE, Chen C, Wang Y, *et al*. Association of Delirium With Long-term Cognitive Decline: A Meta-analysis. *JAMA Neurol* 2020;77:1373–81.
- Wilcox ME, Girard TD, Hough CL. Delirium and long term cognition in critically ill patients. *BMJ* 2021;373:n1007.
- Dziegielewska C, Skead C, Canturk T, *et al*. Delirium and Associated Length of Stay and Costs in Critically Ill Patients. *Crit Care Res Pract* 2021;2021:6612187.
- Yan E, Veitch M, Saripella A, *et al*. Association between postoperative delirium and adverse outcomes in older surgical patients: A systematic review and meta-analysis. *J Clin Anesth* 2023;90:111221.
- Krewulak KD, Stelfox HT, Ely EW, *et al*. Risk factors and outcomes among delirium subtypes in adult ICUs: A systematic review. *J Crit Care* 2020;56:257–64.
- Salluh JIF, Wang H, Schneider EB, *et al*. Outcome of delirium in critically ill patients: systematic review and meta-analysis. *BMJ* 2015;350:h2538.
- Zhang Z, Pan L, Ni H. Impact of delirium on clinical outcome in critically ill patients: a meta-analysis. *Gen Hosp Psychiatry* 2013;35:105–11.
- Siddiqi N, House AO, Holmes JD. Occurrence and outcome of delirium in medical in-patients: a systematic literature review. *Age Ageing* 2006;35:350–64.
- Witlox J, Eurelings LSM, de Jonghe JFM, *et al*. Delirium in elderly patients and the risk of postdischarge mortality, institutionalization, and dementia: a meta-analysis. *JAMA* 2010;304:443–51.
- Higgins JPT, Chandler J, Cumpston M, *et al*, eds. *Cochrane handbook for systematic reviews of interventions version 6.5*. 2024.
- Glanville J, Sowden AJ, Kleijnen J. *Undertaking systematic reviews of research on effectiveness: CRD's guidance for those carrying out or commissioning reviews*. York: NHS Centre for Reviews and Dissemination, University of York, 2001.
- Page MJ, McKenzie JE, Bossuyt PM, *et al*. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
- Stroup DF, Berlin JA, Morton SC, *et al*. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000;283:2008–12.
- Moher D, Shamseer L, Clarke M, *et al*. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
- The EndNote Team. *EndNote*. 21st edn. Philadelphia, PA: Clarivate, 2013.
- Ouzzani M, Hammady H, Fedorowicz Z, *et al*. Rayyan-a web and mobile app for systematic reviews. *Syst Rev* 2016;5:210.
- Microsoft Excel. *Microsoft corporation*. 2019.
- Joanna Briggs Institute. *Checklist for systematic reviews and research syntheses*. 2017.
- The Cochrane Collaboration. *Review manager (RevMan)*. 2024.
- Balshem H, Helfand M, Schünemann HJ, *et al*. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011;64:401–6.
- GRADEpro GDT. Hamilton (ON): McMaster University (developed by Evidence Prime), 2023.