

RESEARCH ARTICLE

# Efficacy and Acceptability of Glycemic Control of Glucagon-Like Peptide-1 Receptor Agonists among Type 2 Diabetes: A Systematic Review and Network Meta-Analysis

Zhixia Li<sup>1</sup>, Yuan Zhang<sup>2</sup>, Xiaochi Quan<sup>1</sup>, Zhirong Yang<sup>1</sup>, Xiantao Zeng<sup>3</sup>, Linong Ji<sup>4</sup>, Feng Sun<sup>1</sup>✉\*, Siyan Zhan<sup>1</sup>✉\*

**1** Department of Epidemiology and Biostatistics, School of Public Health, Peking University Health Science Center, Beijing, China, **2** Department of Clinical Epidemiology and Biostatistics, McMaster University, 1280 Main Street West, Hamilton, ON, Canada, **3** Center for Evidence-based and Translational Medicine, Zhongnan Hospital, Wuhan University, Wuhan, China, **4** Department of Endocrinology and Metabolism, People's Hospital, Peking University, Beijing, China

✉ These authors contributed equally to this work.

\* [siyan-zhan@bjmu.edu.cn](mailto:siyan-zhan@bjmu.edu.cn) (SZ); [sunfeng@bjmu.edu.cn](mailto:sunfeng@bjmu.edu.cn) (FS)



## OPEN ACCESS

**Citation:** Li Z, Zhang Y, Quan X, Yang Z, Zeng X, Ji L, et al. (2016) Efficacy and Acceptability of Glycemic Control of Glucagon-Like Peptide-1 Receptor Agonists among Type 2 Diabetes: A Systematic Review and Network Meta-Analysis. PLoS ONE 11 (5): e0154206. doi:10.1371/journal.pone.0154206

**Editor:** Massimo Pietropaolo, Baylor College of Medicine, UNITED STATES

**Received:** July 9, 2015

**Accepted:** April 11, 2016

**Published:** May 9, 2016

**Copyright:** © 2016 Li et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This study is funded by National Natural Science Foundation of China (81302508) and Research Fund for the Doctoral Program of Higher Education (20120001110015).

**Competing Interests:** The authors have declared that no competing interests exist.

## Abstract

### Objective

To synthesize current evidence of the impact of Glucagon-like peptide-1 receptor agonists (GLP-1 RAs) on hypoglycemia, treatment discontinuation and glycemic level in patients with type 2 diabetes.

### Design

Systematic review and network meta-analysis.

### Data Sources

Literature search (Medline, Embase, the Cochrane library), website of clinical trial, bibliographies of published systematic reviews.

### Eligibility Criteria

Randomized controlled trials with available data comparing GLP-1 RAs with placebo or traditional anti-diabetic drugs in patients with type 2 diabetes.

### Data Synthesis

Traditional pairwise meta-analyses within DerSimonian-Laird random effects model and network meta-analysis within a Bayesian framework were performed to calculate odds ratios for the incidence of hypoglycemia, treatment discontinuation, HbA1c<7.0% and

HbA1c<6.5%. Ranking probabilities for all treatments were estimated to obtain a treatment hierarchy using the surface under the cumulative ranking curve (SUCRA) and mean ranks.

## Results

78 trials with 13 treatments were included. Overall, all GLP-1 RAs except for albiglutide increased the risk of hypoglycemia when compared to placebo. Reduction in the incidence of hypoglycemia was found for all GLP-1 RAs versus insulin (except for dulaglutide) and sulphonylureas. For the incidence of treatment discontinuation, increase was found for exenatide, liraglutide, lixisenatide and taspoglutide versus placebo, insulin and sitagliptin. For glycemic level, decrease was found for all GLP-1 RAs versus placebo. Dulaglutide, exenatide long-acting release (exe\_lar), liraglutide and taspoglutide had significant lowering effect when compared with sitagliptin (HbA1c<7.0%) and insulin (HbA1c<6.5%). Finally, according to SUCRAs, placebo, thiazolidinediones and albiglutide had the best decrease effect on hypoglycemia; sulphonylureas, sitagliptin and insulin decrease the incidence of treatment discontinuation most; exe\_lar and dulaglutide had the highest impact on glycemic level among 13 treatments.

## Conclusions

Among 13 treatments, GLP-1 RAs had a significant reduction with glycemic level but a slight increase effect on hypoglycemia and treatment discontinuation. While albiglutide had the best decrease effect on hypoglycemia and treatment discontinuation among all GLP-1 RAs. However, further evidence is necessary for more conclusive inferences on mechanisms underlying the rise in hypoglycemia.

## Introduction

An increasing number of patients with type 2 diabetes mellitus (T2DM) are being treated with glucagon-like peptide-1 receptor agonists (GLP-1 RAs), a new class of anti-diabetic agents based on incretin therapy[1, 2]. GLP-1 RAs are analogues of GLP-1, which could stimulate insulin secretion, improve insulin resistance and slow down gastrointestinal motility [3–5]. Exenatide (Byetta; Eli Lilly & Co.), liraglutide (Victoza; Novo Nordisk), the two earliest GLP-1 RAs, were approved by the United States Food and Drug Administration (FDA) in 2005 and 2010, respectively [6, 7]. Albiglutide (Tanzeum/Eperzan, GSK) and lixisenatide (Lyxumia, Sanofi) were approved by European Medical Agency (EMA) in 2013. Recently, Dulaglutide (Trulicity; Eli Lilly & Co.) was approved by FDA in 2014. Taspoglutide is currently in phase III clinical trials.

According to the International Diabetes Federation (IDF) in 2013, 387 million people are currently diagnosed with diabetes and there is a projected rise to 592 million people in the world living with diabetes by the year 2035[8]. It means that more and more people will need to be prescribed anti-diabetes medication to help achieve the recommended HbA1c target of <6.5% (National Institute for Health and Clinical Excellence (NICE), 2008) or HbA1c target of <7.0% (American Diabetes Association, (ADA))[9] to avoid the devastating complications of poor diabetes control. Patients with poorly controlled glycemic level would greatly increase the risk of hypoglycemia [10–12] and treatment discontinuation [13–15]. Therefore, an ideal anti-diabetic treatment would be one that can couple the achievement of glycemic control with a

low propensity for causing hypoglycemia and treatment discontinuation. Indeed, several clinical trials and meta-analyses [16–21] for GLP-1 RAs have demonstrated the lowering effect of glycemic levels as well as raised hypoglycemia and treatment discontinuation, although the mechanisms are not very clearly understood. However, since there are so much medicines to choose, which is better for clinical decision is still unknown. So there is a need to include all kinds of GLP-1 RAs simultaneously to assess the impact on hypoglycemia and treatment discontinuation between any two of them.

Therefore, we collected all randomized controlled trials (RCTs) of comparing GLP-1 RAs with placebo or traditional anti-diabetic drugs. A conventional pairwise meta-analysis was performed to summarize current evidence for the effect of GLP-1 RAs on hypoglycemia, treatment discontinuation and glycemic level in patients with T2DM. Additional network meta-analysis was conducted to assess the robustness of the pairwise meta-analysis, supplement missing evidence of head-to-head comparisons by combining both direct and indirect evidence and rank treatments in the evidence network.

## Method

### Systematic review registration

PROSPERO register, CRD42014015328

### Search strategy

In consultation with a medical librarian, a search strategy for MEDLINE, EMBASE and the Cochrane library (from inception to June 1st, 2014) was established. The following search strategy for Ovid-MEDLINE was adapted for other databases:

1. exp glucagon-like peptide-1 agonists/
2. (glucagon like peptide\* or GLP-1).tw.
3. (exenatide or liraglutide or albiglutide or taspoglutide or lixisenatide or LY2189265).tw.
4. randomized controlled trial.pt.
5. (randomized or randomised).tw.
6. (1 or 2 or 3) and (4 or 5)

In addition, completed but unpublished trials were identified from [www.clinicaltrials.gov](http://www.clinicaltrials.gov) website using the similar search strategy. The bibliographies of published systematic reviews were also searched. All relevant authors and principal manufacturers were contacted to supplement incomplete reports of the original papers or to provide new data for unpublished studies.

### Study selection

All the studies included are in English and they are eligible for inclusion only if they were RCTs involving GLP-1 RAs, active anti-diabetic drugs or placebo with complete data on hypoglycemia, treatment discontinuation or glycemic level. Trials are excluded if only they meet one of the following: (1) trials are not RCT (e.g., review, expert comment, editor opinion, new agent introduction, single case report, or case series); (2) if several studies included the same clinical trial, we only include the one which had the longest follow-up time and excluded the other early studies; (3) experimentation on animals or in vitro; (4) not conducted in T2DM; (5) pharmacokinetics research; (6) trials underway, unfinished, or suspended; (7) economical evaluation research; (8) other unrelated researches. These studies were approved by the local ethics

committees and written informed consent was obtained from all the patients. The eligibility of studies for inclusion criteria was assessed independently by four reviewers (ZXL, YZ, XCQ and ZRY) in duplicate.

## Data extraction and quality evaluation

Data were extracted using ADDIS software[22] with respect to trial information (author, publication year, sample size, trial duration, types of intervention and control), population characteristics (background therapy, diabetes duration, age, baseline level of HbA1c), reported outcomes (Number of hypoglycemia, treatment discontinuation, HbA1c<7.0% and HbA1c<6.5% events in each group) and information on methodology. Four investigators (ZXL, ZRY, XCQ and XTZ) extracted data independently, in duplicate. Any discrepancies were resolved by consensus between the two independent reviewers or by a senior investigator (FS).

Quality of studies was assessed according to JADAD scale[23], including adequate method for randomization, appropriate blinding procedures, and detailed report of withdrawals. The JADAD score was not used as a selection criterion, but only for descriptive purpose.

## Data analysis

**Methods for direct treatment comparisons.** Traditional pairwise meta-analyses was performed using DerSimonian-Laird random effects model[24]. Odds ratio (OR) for hypoglycemia, treatment discontinuation, HbA1c<7.0% and HbA1c<6.5% with 95% confidence interval (CI) were calculated as effect measures. For studies that did not report intention-to-treat, we analyzed outcomes as all-patients randomized. The  $I^2$ -statistic was calculated as a measure of the proportion of the overall variation that is attributable to between-study heterogeneity[25].

**Methods for indirect and mixed comparisons.** A random-effects network meta-analysis within a Bayesian framework[26] was performed to evaluate the relative effectiveness of each kind of GLP-1 RAs on hypoglycemia, HbA1c<7.0%, HbA1c<6.5% and the relative acceptability on treatment discontinuation. Bayesian network meta-analysis is a generalization of traditional meta-analysis that allows all evidence to be taken into account simultaneously (both direct and indirect). It can be applied whenever a connected network of evidence is available [26]. ORs for hypoglycemia, treatment discontinuation, HbA1c<7.0% and HbA1c<6.5% with 95% credible interval (CrI) were summarized. The posterior densities for all unknown parameters were estimated using MCMC (Markov chain Monte Carlo) for each model. Each chain used 40 000 iterations with a burn-in of 20 000.

Network meta-analyses enable estimation of the probability that each intervention is the best for each outcome. Probabilities for each treatment taking each possible rank were plotted in absolute rankograms or cumulative rankograms. Besides, the surface under the cumulative ranking curve (SUCRA)[27] were used to estimate the ranking probabilities for all treatments in order to obtain a treatment hierarchy. SUCRA is a percentage interpreted as the percentage of efficacy of a treatment on the outcome that would be ranked first without uncertainty, which is equal to 1 when the treatment is certain to be the best and 0 when it is certain to be the worst[27].

An absolute measure of fit  $\bar{D}_{res}$ , was considered to formally check the model's overall fit.  $\bar{D}_{res}$  is the posterior mean of the residual deviance (the deviance for the fitted model minus the deviance for the saturated model). Ideally, each data point should contribute about one to the posterior mean deviance so that it can be compared to the number of data points for the purpose of checking model fit[28].

Loop-specific approach was used to evaluate the presence of inconsistency locally in network meta-analysis models, that is, if the information of both sources of evidence is similar enough to be combined [29]. This method evaluates the consistency assumption in each closed loop of the network separately. Difference (inconsistency factor) with 95% CIs between direct and indirect estimations for a specific comparison was calculated to assess the presence of inconsistency in each loop. Inconsistency was defined as disagreement between direct and indirect evidence with a 95% CI excluding 0.

Analyses were conducted using STATA 11.0 (pairwise meta-analysis,  $I^2$  calculations and estimation of inconsistency), R 3.0.2 (SUCRA graphs) and WinBUGS 1.4.3 (network Meta-analysis and model fit).

## Results

### Study characteristics and evidence network

78 trials involving 13 treatments met the selection criteria. A total of 34685 patients contributed to the analysis. Flow chart of trials selection was shown in Fig 1.


**Study characteristics.** Table 1 summarized the characteristics of included 78 trials. The range of publication year was 2004–2014. Trial duration ranged from 26 to 234 weeks. The average age of included patients was 55.89 years [standard deviation (SD): 1.71], varied from 50.5 to 61.0 years. The median diabetes duration at baseline was 7.5 [interquartile range (IQR): 6.0–8.9] years. And the mean baseline glycemic level HbA1c was 8.2% (SD: 0.4%). Of the 78 trials included, albiglutide, dulaglutide, exenatide long-acting release (exe\_lar), exenatide, liraglutide, lixisenatide and taspoglutide were studied in 3, 3, 7, 25, 13, 9 and 8 trials, respectively. And 3 trials involved both exenatide and exe\_lar simultaneously. Besides, albiglutide and exenatide, albiglutide and liraglutide, dulaglutide and exenatide, exenatide and taspoglutide, exenatide and lixisenatide, liraglutide and exenatide, liraglutide and exe\_lar were both involved simultaneously in 1 trial, respectively.

Reporting quality of included studies varied. According to JADAD scale, the number of dropout and the methods used for randomization, allocation concealment and blinding were appropriately described in most cases (89.7%, 85.9%, 60.3% and 93.6%, respectively), although 34.6% (27/78) of trials were open label. Additionally, 88.5% (69/78) of trials used intention-to-treat analysis. (S1 Table). Overall, risk of bias is respectively low.

**Evidence network.** 13 treatments were analyzed, including 7 GLP-1 RAs (Albiglutide, Dulaglutide, Exe\_LAR, Exenatide, Liraglutide, Lixisenatide and Taspoglutide), 5 kinds of active anti-diabetic drugs (insulin, metformin (Met), sulphonylureas (SU), sitagliptin and thiazolidinediones (TZD)), and placebo. 85.90% (67/78) of trials were two-arm studies and the rest 14.10% (11/78) were multiple-arm studies (see Table 1 and Fig 2). Overall, 32932, 24919, 31588 and 23427 patients contributed to the analysis of hypoglycemia (Fig 2A, including 71 studies and 13 treatments), treatment discontinuation (Fig 2B, including 48 studies and 13 treatments), HbA1c < 7.0% (Fig 2C, including 67 studies and 13 treatments) and HbA1c < 6.5% (Fig 2D, including 48 studies and 13 treatments), respectively. Every group of GLP-1 RAs existed head-to-head (direct) comparison with placebo.

**Conventional meta-analysis of individual GLP-1 RAs.** Fig 3A–3D showed the effect of individual GLP-1 RAs on hypoglycemia, treatment discontinuation, HbA1c < 7.0% and HbA1c < 6.5% from direct pairwise meta-analysis.

Fig 3A displayed the effect on hypoglycemia. In comparison to placebo, dulaglutide, exenatide, liraglutide, lixisenatide and taspoglutide significantly increased the risk of hypoglycemia by 2.34 (95% CI: 1.39, 3.95), 2.26 (95%CI: 1.44, 3.56), 1.82 (95%CI: 1.31, 2.54), 1.70 (95%CI: 1.36, 2.13) and 3.13 (95%CI: 1.50, 6.53), respectively. No significant difference was found



**Fig 1. Flow chart of studies considered for inclusion, RCT = randomized controlled trial.** This Flow chart is based on PRISMA 2009 Flow Diagram [30].

doi:10.1371/journal.pone.0154206.g001

between albiglutide or exelgler versus placebo. Compared with insulin, exelgler (OR 0.36, 95% CI: 0.21, 0.62) and exenatide (OR 0.63, 95%CI: 0.51, 0.79) were associated with less hypoglycemia. No statistically significant difference was found between GLP-1 RAs and other active comparators in their effects on hypoglycemia.

Regarding treatment discontinuation, lixisenatide and tasoglutide significantly increased the incidence in comparison to placebo by 1.78 (95%CI: 1.30, 2.45) and 5.43 (95%CI: 2.96,

**Table 1. Characteristics of the 78 studies with 34685 patients included in the network Meta-analysis Duration > = 8w.**

NO	Study ID	Investigational treatments	Outcome reported	Size (pts)	Background therapy	Duration of trial(w)	Baseline information		
							Age (yrs)	Years of T2DM	HbA1c (%)
1	Ahren B, 2013[31]	Placebo,Lixisenatide	1, 2, 3, 4	680	Met	24	54.8	3.6	8.1
2	Ahren B, 2014[32]	Albiglutide,SU, placebo, Sitagliptin	1, 2, 3, 4	1012	Met	104	54.5	6	8.11
3	Apovian CM, 2010 [33]	Exenatide,Placebo	1, 2	194	Met/SU/SU+Met	24	54.8	5.5	7.6
4	Barnett AH, 2007 [34]	Exenatide,Insulin	1, 3, 4	276	Met/SU	16	54.9	7.4	9
5	Bergenstal R, 2009 [35]	Exenatide,Insulin	1, 3, 4	372	Met+SU	24	52.2	8.5	10.2
6	Bergenstal RM, 2010 [36]	Sitagliptin,TZD, Exe_LAR	1, 2, 3, 4	491	Met	26	52.5	6	8.6
7	Bergenstal RM, 2012[37]	Placebo,Sitagliptin, Taspoglutide	1, 2, 3	546	Met	24	56	5.9	8
8	Blevins T, 2011[38]	Exenatide,Exe_LAR	1, 2, 3, 4	252	Met+/-SU+/-TZD	24	55.5	7	8.45
9	Bolli G, 2013[39]	Placebo,Lixisenatide	1, 2, 3, 4	479	Met	24	56.1	6	8
10	Bunck MC, 2009 [40]	Exenatide,Insulin	1, 3	69	Met	52	58.3	4.9	7.5
11	Buse JB, 2004[41]	Exenatide,Placebo	1, 3	377	SU	30	55	6.3	8.6
12	Buse JB, 2009[42]	Liraglutide,Exenatide	1, 2, 3, 4	464	Met+/-SU	26	56.7	8.2	8.2
13	Buse JB, 2011[43]	Exenatide,Placebo	1, 2, 3, 4	259	Glar+/-Met/TZD	30	59	12	8.4
14	Buse JB, 2013[44]	Liraglutide,Exe_LAR	1, 2, 3	911	Met+/-SU+/-Piog	26	59	12	8.4
15	Charbonnel B, 2013[45]	Liraglutide,Sitagliptin	1, 2, 3, 4	650	Met	26	57.3	6	8.2
16	Davies M, 2013[46]	Exe_LAR,Insulin	1, 2, 3, 4	216	Met/Met+SU	26	58.5		8.4
17	Davies MJ, 2009 [47]	Exenatide,Insulin	1, 3, 4	234	Met+/- SU/TZD	26	56.5	8.7	8.6
18	Davis SN, 2007[48]	Exenatide,Insulin	1, 3	49	SU/Met	16	53	11	8.1
19	DeFronzo RA, 2005[49]	Exenatide,Placebo	1, 3	336	Met	30	53	5.8	8.2
20	DeFronzo RA, 2010[50]	Exenatide,TZD	1	90	Met	20	56	4.7	7.8
21	Derosa G, 2012 [51]	Exenatide,Placebo	1, 2	171	Met	48	57	7.7	8
22	Diamant M, 2012 [52]	Exe_LAR,Insulin	1, 2, 3, 4	466	Met/Met+SU	26	58	7.9	8.3
23	Drucker DJ, 2008 [53]	Exenatide,Exe_LAR	1, 2, 3	295	Met+/-SU+/-TZD	30	55	6.5	8.3
24	Fonseca VA, 2012 [54]	Placebo,Lixisenatide	1, 2, 3, 4	361	no	12	53.6	1.3	8
25	Gallwitz B, 2011 [55]	Exenatide,Insulin	1, 3, 4	354	Met/SU	26	57	5	7.9
26	Gallwitz B, 2012 [56]	SU,Exenatide	1, 2	1019	Met	234	56	5.6	7.45
27	Gao Y, 2009[57]	Exenatide,Placebo	1	466	Met/Met+SU	16	54.5	8	8.3
28	Garder A, 2011[58]	SU,Liraglutide	1, 3, 4	745	Met/SU/BG/Met+TZD	104	53	5.4	8.3
29	Heine RJ, 2005[59]	Exenatide,Insulin	1	549	Met+SU	26	58.9	9.6	8.2
30	Henry RR, 2013 [60]	Placebo,Taspoglutide	1, 2, 3, 4	320	Met+Piog	24	54.1	7.7	8.1

(Continued)

Table 1. (Continued)

NO	Study ID	Investigational treatments	Outcome reported	Size (pts)	Background therapy	Duration of trial(w)	Baseline information		
							Age (yrs)	Years of T2DM	HbA1c (%)
31	Hollander P, 2013 [61]	Placebo,Taspoglutide	1, 2, 3, 4	292	Met	24	53.5	5.08	7.55
32	Inagaki N, 2012[62]	Exe_LAR,Insulin	1, 3, 4	427	BG or BG + TZD	26	56.76	9.03	8.5
33	Iwamoto K, 2009 [63]	Placebo,Exe_LAR	3, 4	19	SU/BG/SU+BG/TZD	10	58	6	7.4
34	Ji LN, 2013[64]	Exenatide,Exe_LAR	2, 3, 4	678	Met, SU, TZD	26	55	8.2	8.7
35	Kadowaki T, 2009 [65]	Exenatide,Placebo	1, 3	114	SU/BG/SU+TZD/BG	12	60.3	11.8	8
36	Kadowaki T, 2011 [66]	Placebo,Exenatide	1, 3, 4	179	SU+/BG+/TZD	24	58.4	12	8.2
37	Kendall DM, 2005 [67]	Exenatide,Placebo	1, 3	743	Met/Met+SU	30	55.3	8.9	8.5
38	Kim D, 2007[68]	Placebo,Exe_LAR	1, 2, 3	29	Met	15	54	5	8.5
39	Li CJ, 2012[69]	Liraglutide,Insulin	1	84	Insulin	12	52	9	8.75
40	Liutkus J, 2010[70]	Exenatide,Placebo	1, 3, 4	165	TZD/TZD+Met	26	54.7	6.4	8.2
41	Marre M, 2009[71]	Liraglutide,Placebo, TZD	1, 2, 3, 4	808	Glimepiride	26	56.1	6.5	8.4
42	Mathieu C, 2014 [72]	Liraglutide,Insulin	2, 3	177	Insulin degludec (IDeg) OD + Met	52	61	12, 4	7.7
43	Moretto TJ, 2008 [73]	Placebo,Exenatide	1, 2, 3, 4	232	diet+exercise	24	54	1.7	7.8
44	Nauck M, 2014[74]	Dulaglutide, placebo, Sitagliptin	1, 2, 3, 4	921	diet+exercise/ monotherapy /Met +monotherapy	26/52	54	7	8.1
45	Nauck MA, 2007 [75]	Exenatide,Insulin	1, 2, 3	501	Met/SU	52	58.5	9.9	8.6
46	Nauck MA, 2009 [76]	Placebo,Taspoglutide	2, 3	197	Met	8	54	5.5	7.9
47	Nauck M, 2009[77]	SU,Liraglutide,Placebo	1, 2, 3, 4	845	Met	104	57	7.9	8.4
48	Nauk M, 2013[78]	Insulin,Taspoglutide	1, 2, 3, 4	1028	Met/SU	24	58	9.1	8.3
49	NCT00620282, 2011[79]	SU,Placebo,Liraglutide	1	49	Met	12	58.5	6.8	7.2
50	NCT00667732, 2013[80]	Exenatide,Placebo	1, 4	34	Met+/-Lantus Insulin				
51	NCT00701935, 2013[81]	Exenatide,Placebo	3	71		24	58		
52	Pinget M, 2013[82]	Placebo,Lixisenatide	1, 3, 4	484	Piog+/-Met	24	55.5	1.75	8.1
53	Pratley R, 2011[83]	Liraglutide,Sitagliptin	1, 3, 4	658	Met	52	55.3	6.2	8.4
54	Pratley RE, 2013 [84]	Taspoglutide,TZD	1, 3, 4	740	Met/SU/Met+SU	24	56.4	8.8	8.3
55	Pratley RE, 2014 [85]	Albiglutide,Liraglutide	1, 2, 3, 4	805		32	55.6	8.35	8.17
56	Ratner R, 2010[86]	Placebo, Taspoglutide	1, 2, 3, 4	129	Met	8	56.5	6.5	7.9
57	Ratner RE, 2010 [87]	Placebo,Lixisenatide	1, 3, 4	529	Met	13	56.5	7.1	7.5
58	Raz I, 2012[88]	Placebo,Taspoglutide	1, 2, 3, 4	354	no	24	54.8	2.4	7.6
59	Riddle MC,2013 [89]	Placebo,Lixisenatide	1, 2, 3, 4	446	Insulin Glar + Met	24	56	9.2	7.6

(Continued)



Table 1. (Continued)

NO	Study ID	Investigational treatments	Outcome reported	Size (pts)	Background therapy	Duration of trial(w)	Baseline information		
							Age (yrs)	Years of T2DM	HbA1c (%)
60	Riddle MC, 2013 [90]	Placebo,Lixisenatide	1, 2, 3, 4	495	Met+/-SU+/-TZD	24	57	12.5	8.4
61	Rosenstock J, 2013[91]	Exenatide, Taspoglutide	1, 2, 3	1149	TZD+/-Met	24	56	6.6	8.1
62	Rosenstock J, 2014[92]	Placebo,Lixisenatide	1, 2, 3	859	Met+/-SU	24	57.2	9.4	8.3
63	Rosenstock J, 2014[93]	Albiglutide, Insulin	1, 2, 3, 4	566	Insulin Glar	26	55.45	11	8.45
64	Rosenstock J, 2009[94]	Albiglutide,Exenatide, Placebo	1, 3	214	Met	16	54	4.9	8
65	Rosenstock J, 2013[95]	Exenatide,Lixisenatide	1, 2, 3, 4	634	Met	24	57.4	6.8	8
66	Russell-Jones D, 2009[96]	Placebo,Liraglutide, Insulin	1, 2, 3	576	Met+Glimepiride	26	57.6	9.5	8.3
67	Russell-Jones D, 2012[97]	Sitagliptin,Metformin, TZD,Exe_LAR	1, 3, 4	820	diet + exercise	26	53.8	2.7	8.5
68	Seino Y, 2008[98]	Liraglutide,Placebo	1, 2, 3, 4	211	oad	14	57	8	8.3
69	Seino Y, 2010[99]	SU,Liraglutide	3, 4	400	diet+exercise/ monotherapy	52	58.3	8.3	8.9
70	Seino Y, 2012[100]	Placebo,Lixisenatide	1, 2, 3, 4	311	Insulin/SU	24	58.4	13.9	8.53
71	Seino Y, 2014[101]	Albiglutide, placebo	2	211	diet+exercise/ monotherapy	16	57	7	8.55
72	Umpierrez G, 2014 [102]	Dulaglutide, Met	1, 2, 3, 4	807	diet + exercise	26/52	56	3	7.6
73	Umpierrez GE, 2011[103]	Placebo,LY	1, 2	262	oad	16	56.5	8.3	8.3
74	Wysham C, 2014 [104]	Dulaglutide, placebo, Exenatide	1, 2, 3, 4	976	Met+Piog	26	55.57	9	8.1
75	Yang W, 2011[105]	SU,Liraglutide	1, 2, 3	928	Met	16	53.3	7.5	8.6
76	Yuan GH, 2012 [106]	Exenatide,Metformin	1, 3, 4	59	no	26	50.5	<1 month	8.2
77	Zinman B, 2007 [107]	Exenatide,Placebo	1, 3	233	TZD+/-Met	16	56	8	7.9
78	Zinman B, 2009 [108]	Liraglutide,Placebo	1, 2, 3, 4	533	Met+Rosig	26	55	8.9	8.5


Note: LY: LY2189265, Dulaglutide. SU: sulphonylureas; TZD: thiazolidinedione; MET: metformin; Glar: glargine; Piog: pioglitazone; BG: biguanide; Rosig: rosiglitazone; oad: oral antidiabetic drug;—: unavailable information.

1: Hypoglycemia, 2: treatment discontinuation, 3:HbA1c<7.0%, 4: HbA1c<6.5%

doi:10.1371/journal.pone.0154206.t001

9.94), respectively. Significant increase was also found when exe\_lar versus insulin (OR 3.11 (95%CI: 1.43, 6.74)) and liraglutide versus SU (OR 3.48 (95%CI: 1.16, 10.48)). No statistically significant difference was found between other GLP-1 RAs versus placebo or active comparators in their effects on treatment discontinuation (Fig 3B).



Fig 3C displayed the effect on HbA1c <7%. In comparison to placebo, albiglutide, exe\_lar, exenatide, liraglutide, lixisenatide and taspoglutide significantly increased the incidence of HbA1c <7% by 3.31 (95% CI: 2.06, 5.32), 49.81 (95%CI: 5.54, 447.76), 3.62 (95%CI: 2.32, 5.65), 4.81 (95%CI: 3.46, 6.68), 3.13 (95%CI: 2.49, 3.92) and 7.25 (95%CI: 5.15, 10.20), respectively. No significant difference was found between dulaglutide versus placebo. Compared with



Lines connect the interventions that have been studied in head-to-head (direct) comparisons in the eligible RCTs. The width of the lines represents the cumulative number of RCTs for each pairwise comparison and the size of every node is proportional to the number of randomized participants (sample size). EXE\_LAR: exenatide long-acting release; SU: sulphonylureas; TZD: thiazolidinedione; MET: metformin.

**Fig 2. Evidence structure of eligible comparisons for network meta-analysis.**

doi:10.1371/journal.pone.0154206.g002



**Fig 3. Impact of individual GLP-1 receptor agonists on hypoglycemia, treatment discontinuation, HbA1c <7.0%, HbA1c <6.5% of direct pairwise meta-analysis.**

doi:10.1371/journal.pone.0154206.g003

insulin, exe\_lar (OR 1.92 (95%CI: 1.27, 2.92)) and liraglutide (OR 1.41 (95%CI: 1.03, 1.93)) were associated with higher incidence of HbA1c <7%. The incidence of HbA1c <7% was also increased when liraglutide versus sitagliptin (OR 1.91 (95%CI: 1.22, 3.00)). Besides, exenatide

decreased the incidence of HbA1c <7% in comparison with *exe\_lar* by 0.51 (95%CI: 0.33, 0.77). No statistically significant difference was found between other GLP-1 RAs versus placebo or active comparators in their effects on HbA1c <7%.

Regarding HbA1c <6.5%, exenatide, liraglutide, lixisenatide and taspoglutide were significantly increased the incidence in comparison with placebo by 3.14 (95% CI: 1.97, 5.01), 5.57 (95%CI: 2.75, 11.25), 3.36 (95%CI: 2.51, 4.50) and 6.89 (95%CI: 4.93, 9.62), respectively. No significant difference was found between albiglutide, dulaglutide or *exe\_lar* versus placebo. Besides, compared with insulin, *exe\_lar* (OR 3.55 (95%CI: 1.53, 8.23)) was associated with higher incidence of HbA1c<6.5%. Exenatide could also increase the incidence of HbA1c<6.5% when compared with *exe\_lar* (OR 0.40 (95%CI: 0.22, 0.73)). No statistically significant difference was found between GLP-1 RAs and other active comparators in their effects on HbA1c <6.5%. ORs with 95%CIs were listed in [Fig 3D](#).

## Network meta-analysis of individual GLP-1 RAs

Results of network meta-analysis among GLP-1 RAs, placebo and active comparators were displayed in [Fig 4](#). As shown in [Fig 4A](#), all GLP-1 RAs except for albiglutide increased the risk of hypoglycemia with range from 1.83 (95%CrI: 1.14, 2.95) to 2.71 (95%CrI: 1.92, 3.85) when compared with placebo. Compared with insulin, all GLP-1 RAs except for dulaglutide reduced the risk of hypoglycemia with range from 0.38 (95%CrI: 0.18, 0.78) to 0.63 (95%CrI: 0.43, 0.93). Similar findings were observed between all GLP-1 RAs and SU with ORs varied from 0.15 (95%CrI: 0.06, 0.35) to 0.25 (95%CrI: 0.13, 0.49). No statistically significant difference was found between GLP-1 RAs versus Met, sitagliptin or TZD.

Regarding treatment discontinuation, exenatide, liraglutide, lixisenatide and taspoglutide significantly increase the incidence when compared with placebo (range of ORs: 2.08 (95%CrI: 1.28, 3.57) to 5.00 (95%CrI: 2.86, 10.00)) or insulin (range of ORs: 2.28 (95%CrI: 1.02, 5.32) to 5.51 (95%CrI: 2.71, 12.08)). Similar associations were found when exenatide, *exe\_lar*, liraglutide, lixisenatide and taspoglutide were compared to SU (range of ORs: 2.33 (95%CrI: 1.04, 5.26) to 6.75 (95%CrI: 3.04, 15.60)). Exenatide, liraglutide, lixisenatide and taspoglutide were associated with higher incidence of treatment discontinuation with range from 2.33 (95%CrI: 1.01, 5.26) to 5.68 (95%CrI: 2.60, 12.59) when compared with sitagliptin. Increase was found between taspoglutide and TZD (OR 4.91 (95%CrI: 1.60, 16.02)) in their effects on the incidence of treatment discontinuation.

[Fig 4B](#) displayed the effect on HbA1c <7% and HbA1c <6.5%. For HbA1c <7%, all GLP-1 RAs significantly increased the incidence with range from 3.29 (95%CrI: 2.18, 4.97) to 8.39 (95%CrI: 3.67, 18.12) versus placebo. Dulaglutide, *exe\_lar*, liraglutide and taspoglutide were associated with higher incidence with the range from 1.96 (95%CrI: 1.15, 3.29) to 2.47 (95%CrI: 1.08, 5.35) when compared with sitagliptin. No statistically significant difference was found between GLP-1 RAs versus insulin, Met, SU or TZD.

Regarding HbA1c <6.5%, all GLP-1 RAs significantly increased the incidence with range from 3.45 (95%CrI: 2.50, 5.00) to 9.09 (95%CrI: 5.88, 16.67) in comparison to placebo. Dulaglutide, *exe\_lar*, liraglutide and taspoglutide were associated with higher incidence with the range from 2.11 (95%CrI: 1.23, 3.65) to 2.94 (95%CrI: 1.92, 4.55) when compared with insulin. Significant increase incidence of HbA1c <6.5% was found between *exe\_lar* and Met (OR 2.38 (95%CrI: 1.25, 4.76)). In comparison to SU, *exe\_lar* and liraglutide significantly increase the incidence of HbA1c <6.5% by 2.44 (95%CrI: 1.25, 4.76) and 1.82 (95%CrI: 1.16, 2.94), respectively. Besides, significantly increase incidence was found between dulaglutide, *exe\_lar*, liraglutide versus sitagliptin with range from 1.79 (95%CrI: 1.15, 2.78) to 2.33 (95%CrI: 1.43, 4.00). No statistically significant difference was found between GLP-1 RAs and TZD in their effects on HbA1c <6.5%.

<b>Albiglutide</b>	1.19 (0.42, 3.59)	2.07 (0.96, 4.34)	1.57 (0.66, 3.66)	0.82 (0.35, 1.76)	1.99 (0.98, 3.90)	1.87 (0.79, 4.31)	1.30 (0.23, 7.18)	0.67 (0.30, 1.48)	0.80 (0.35, 1.83)	0.92 (0.29, 2.90)	4.51 (1.95, 10.74)	0.90 (0.43, 1.77)
0.76 (0.29, 2.00)	<b>Dulaglutide</b> 1.73 (0.64, 4.16)	1.30 (0.45, 3.63)	0.69 (0.22, 1.91)	1.67 (0.61, 4.24)	1.57 (0.55, 4.14)	1.08 (0.26, 4.08)	0.56 (0.19, 1.52)	0.67 (0.26, 1.60)	0.77 (0.20, 2.76)	3.82 (1.28, 10.34)	0.75 (0.29, 1.80)	
0.60 (0.28, 1.24)	0.78 (0.39, 1.61)	<b>Exenatide</b> 0.76 (0.42, 1.41)	0.40 (0.20, 0.75)	0.96 (0.56, 1.69)	0.91 (0.47, 1.74)	0.62 (0.12, 3.34)	0.33 (0.16, 0.64)	0.39 (0.19, 0.80)	0.45 (0.15, 1.30)	2.20 (1.17, 4.30)	0.43 (0.26, 0.71)	
0.83 (0.34, 1.90)	1.07 (0.46, 2.52)	1.38 (0.78, 2.44)	<b>Exe_LAR</b> 0.53 (0.25, 1.04)	1.27 (0.65, 2.45)	1.19 (0.53, 2.70)	0.82 (0.15, 4.61)	0.43 (0.19, 0.96)	0.51 (0.23, 1.14)	0.59 (0.20, 1.68)	2.90 (1.30, 6.52)	0.57 (0.28, 1.12)	
<b>0.38</b> (0.18, 0.78)	0.49 (0.23, 1.07)	<b>0.63</b> (0.43, 0.93)	<b>0.46</b> (0.26, 0.80)	<b>Insulin</b> 2.42 (1.29, 4.78)	2.28 (1.02, 5.32)	1.57 (0.28, 9.11)	0.82 (0.37, 1.87)	0.98 (0.44, 2.28)	1.12 (0.36, 3.52)	5.51 (2.71, 12.08)	1.09 (0.56, 2.16)	
0.89 (0.43, 1.80)	1.16 (0.54, 2.51)	1.49 (0.92, 2.36)	1.08 (0.58, 1.97)	2.36 (1.41, 3.93)	<b>Liraglutide</b> 0.94 (0.46, 1.90)	0.65 (0.12, 3.45)	0.34 (0.18, 0.62)	0.40 (0.21, 0.80)	0.46 (0.17, 1.26)	2.28 (1.13, 4.70)	0.45 (0.26, 0.75)	
0.88 (0.36, 2.13)	1.15 (0.50, 2.66)	1.47 (0.83, 2.61)	1.07 (0.49, 2.30)	2.33 (1.23, 4.53)	0.99 (0.51, 1.92)	<b>Lixisenatide</b> 0.70 (0.13, 3.78)	0.36 (0.16, 0.81)	0.43 (0.19, 0.99)	0.49 (0.16, 1.54)	2.43 (1.15, 5.39)	0.48 (0.28, 0.78)	
0.91 (0.30, 2.83)	1.20 (0.49, 3.02)	1.52 (0.63, 3.90)	1.11 (0.44, 2.99)	2.42 (0.96, 6.48)	1.02 (0.40, 2.77)	1.04 (0.38, 2.97)	<b>Met</b> 0.52 (0.10, 2.90)	0.62 (0.12, 3.17)	0.72 (0.10, 4.74)	3.52 (0.64, 19.69)	0.69 (0.13, 3.58)	
<b>0.15</b> (0.06, 0.35)	<b>0.20</b> (0.08, 0.49)	<b>0.25</b> (0.13, 0.49)	<b>0.18</b> (0.08, 0.41)	<b>0.40</b> (0.19, 0.82)	<b>0.17</b> (0.09, 0.31)	<b>0.17</b> (0.08, 0.39)	<b>0.16</b> (0.06, 0.48)	<b>SU</b> 1.20 (0.55, 2.63)	1.37 (0.44, 4.33)	<b>6.75</b> (3.04, 15.60)	1.33 (0.68, 2.57)	
0.97 (0.42, 2.24)	1.28 (0.60, 2.82)	1.63 (0.88, 3.02)	1.19 (0.59, 2.43)	2.62 (1.34, 4.98)	1.10 (0.61, 1.97)	1.11 (0.52, 2.44)	1.07 (0.41, 2.76)	<b>6.50</b> (3.04, 13.92)	1.15 (0.38, 3.44)	<b>5.68</b> (2.60, 12.59)	1.11 (0.68, 2.18)	
1.15 (0.44, 2.98)	1.49 (0.61, 3.77)	1.92 (0.95, 3.95)	1.39 (0.64, 3.14)	3.05 (1.46, 6.56)	1.28 (0.62, 2.73)	1.30 (0.55, 3.05)	1.25 (0.45, 3.45)	<b>7.67</b> (3.14, 18.83)	1.17 (0.54, 2.56)	<b>4.91</b> (1.60, 16.02)	0.97 (0.33, 2.79)	
0.86 (0.36, 2.07)	1.13 (0.49, 2.64)	1.45 (0.81, 2.58)	1.06 (0.49, 2.18)	2.32 (1.23, 4.26)	0.97 (0.50, 1.86)	0.98 (0.47, 2.05)	0.94 (0.34, 2.55)	<b>5.78</b> (2.53, 12.99)	0.89 (0.43, 1.76)	0.75 (0.35, 1.59)	<b>2.20</b> (1.10, 0.35)	
1.62 (0.77, 3.37)	<b>2.11</b> (1.05, 4.34)	<b>2.71</b> (1.92, 3.85)	<b>1.97</b> (1.06, 3.70)	<b>4.31</b> (2.71, 7.02)	<b>1.83</b> (1.14, 2.95)	<b>1.84</b> (1.15, 3.00)	1.77 (0.69, 4.38)	<b>10.78</b> (5.50, 21.69)	1.66 (0.88, 3.08)	1.41 (0.69, 2.89)	<b>1.87</b> (1.07, 3.39)	<b>Placebo</b>

**Figure 4A.** Results of network meta-analysis for hypoglycemia (lower left quarter) and treatment discontinuation (upper right quarter)

<b>Albiglutide</b>	1.64 (0.77, 3.47)	0.82 (0.44, 1.54)	<b>2.11</b> (1.10, 4.23)	0.72 (0.39, 1.37)	1.59 (0.91, 2.79)	0.79 (0.40, 1.57)	0.88 (0.39, 1.96)	0.87 (0.45, 1.65)	0.90 (0.49, 1.63)	1.36 (0.68, 2.71)	1.52 (0.75, 3.09)	<b>0.23</b> (0.12, 0.40)
0.58 (0.22, 1.55)	<b>Dulaglutide</b> 0.50 (0.28, 0.90)	1.29 (0.70, 2.50)	<b>0.44</b> (0.23, 0.86)	0.98 (0.53, 1.84)	<b>0.48</b> (0.25, 0.96)	0.54 (0.29, 1.02)	0.53 (0.26, 1.12)	<b>0.55</b> (0.31, 0.99)	0.83 (0.43, 1.65)	0.93 (0.47, 1.91)	<b>0.14</b> (0.08, 0.25)	
1.32 (0.69, 2.57)	2.27 (0.98, 4.91)	<b>Exenatide</b> 2.59 (1.71, 4.04)	0.88 (0.59, 1.32)	<b>1.96</b> (1.24, 3.09)	0.96 (0.61, 1.53)	1.08 (0.56, 2.05)	1.07 (0.59, 1.92)	1.10 (0.67, 1.81)	1.67 (0.97, 2.87)	<b>1.87</b> (1.11, 3.18)	<b>0.28</b> (0.19, 0.39)	
0.60 (0.29, 1.22)	1.04 (0.43, 2.35)	0.46 (0.28, 0.71)	<b>Exe_LAR</b> 0.34 (0.22, 0.52)	0.76 (0.44, 1.26)	<b>0.37</b> (0.21, 0.65)	<b>0.42</b> (0.21, 0.80)	<b>0.41</b> (0.25, 0.70)	0.64 (0.37, 1.07)	0.72 (0.40, 1.26)	0.11 (0.06, 0.17)		
0.95 (0.49, 1.84)	1.64 (0.67, 3.77)	0.72 (0.48, 1.06)	1.58 (0.97, 2.59)	<b>Insulin</b> 2.21 (1.30, 3.70)	1.09 (0.63, 1.91)	1.22 (0.61, 2.44)	1.21 (0.63, 2.28)	1.24 (0.72, 2.14)	<b>1.88</b> (1.05, 3.38)	<b>2.11</b> (1.23, 3.65)	<b>0.31</b> (0.19, 0.48)	
0.73 (0.39, 1.39)	1.26 (0.53, 2.84)	<b>0.56</b> (0.35, 0.88)	1.22 (0.71, 2.11)	0.77 (0.48, 1.29)	<b>Liraglutide</b> 0.49 (0.29, 0.83)	0.55 (0.27, 1.10)	<b>0.55</b> (0.34, 0.86)	<b>0.56</b> (0.36, 0.87)	0.85 (0.50, 1.46)	0.95 (0.55, 1.68)	0.14 (0.09, 0.21)	
1.47 (0.70, 3.24)	<b>2.54</b> (1.03, 6.01)	1.12 (0.68, 1.85)	<b>2.44</b> (1.33, 4.65)	1.54 (0.87, 2.83)	<b>2.00</b> (1.14, 3.60)	1.11 (0.53, 2.32)	1.10 (0.57, 2.13)	1.14 (0.64, 2.02)	1.73 (0.94, 3.22)	<b>1.94</b> (1.10, 3.46)	<b>0.29</b> (0.20, 0.40)	
1.15 (0.43, 3.19)	1.98 (0.80, 4.81)	0.87 (0.37, 2.06)	1.91 (0.81, 4.73)	1.20 (0.50, 3.04)	1.57 (0.65, 3.91)	0.78 (0.31, 2.05)	<b>Met</b> 1.02 (0.45, 2.19)	1.02 (0.53, 1.99)	1.02 (0.77, 3.14)	1.74 (0.82, 3.71)	<b>0.26</b> (0.13, 0.50)	
1.18 (0.55, 2.54)	2.03 (0.80, 5.16)	0.89 (0.47, 1.72)	0.89 (0.97, 4.04)	1.24 (0.64, 2.48)	1.61 (0.95, 2.77)	0.80 (0.39, 1.66)	1.02 (0.37, 2.78)	<b>SU</b> 1.03 (0.58, 1.85)	1.03 (0.81, 3.04)	1.74 (0.90, 3.53)	<b>0.26</b> (0.14, 0.44)	
1.44 (0.70, 2.99)	<b>2.47</b> (1.08, 5.35)	1.09 (0.63, 1.86)	<b>2.39</b> (1.35, 4.29)	1.51 (0.85, 2.75)	<b>1.96</b> (1.15, 3.29)	0.98 (0.50, 1.86)	1.24 (0.52, 2.94)	1.22 (0.61, 2.41)	<b>Sitagliptin</b> 1.51 (0.88, 2.58)	1.69 (0.94, 3.13)	<b>0.25</b> (0.15, 0.40)	
0.75 (0.33, 1.75)	1.30 (0.51, 3.27)	0.58 (0.30, 1.11)	1.26 (0.65, 2.48)	0.80 (0.40, 1.61)	1.03 (0.54, 2.02)	0.52 (0.24, 1.10)	0.66 (0.25, 1.66)	0.64 (0.28, 1.47)	0.53 (0.27, 1.04)	<b>TZD</b> 1.11 (0.64, 2.00)	0.17 (0.10, 0.27)	
0.68 (0.33, 1.44)	1.17 (0.47, 2.74)	<b>0.52</b> (0.31, 0.84)	1.13 (0.62, 2.10)	0.72 (0.41, 1.26)	0.93 (0.53, 1.62)	<b>0.46</b> (0.25, 0.85)	0.59 (0.23, 1.50)	0.58 (0.28, 1.20)	0.47 (0.26, 0.86)	0.90 (0.44, 1.78)	<b>1.11</b> (0.64, 2.00)	<b>1.15</b> (0.09, 0.23)
<b>4.84</b> (2.54, 9.44)	<b>8.39</b> (3.67, 18.12)	<b>3.66</b> (2.66, 5.15)	<b>8.04</b> (4.99, 13.42)	<b>5.09</b> (3.33, 8.03)	<b>6.58</b> (4.37, 10.17)	<b>3.29</b> (2.18, 4.97)	<b>4.20</b> (1.77, 10.04)	<b>4.11</b> (2.22, 7.63)	<b>3.35</b> (2.05, 5.76)	<b>6.40</b> (3.36, 12.19)	<b>7.14</b> (4.58, 11.32)	<b>Placebo</b>

**Figure 4B.** Results of network meta-analysis for HbA1c <7.0% (lower left quarter) HbA1c and <6.5% (upper right quarter)

Note: Results represent odds ratios and 95% credible intervals for the incidence of HbA1c <7.0% (lower left quarter) and HbA1c <6.5% (upper right quarter). Treatments were reported in alphabetical order and the estimation was calculated as the row-defining treatment compared with the column-defining treatment. Exe\_LAR: exenatide long-acting release; MET: metformin; SU: sulphonylureas; TZD: thiazolidinedione.

**Fig 4. Results of network meta-analysis.** (A) Results of network meta-analysis for hypoglycemia (lower left quarter) and treatment discontinuation (upper right quarter). (B) Results of network meta-analysis for HbA1c <7.0% (lower left quarter) HbA1c and <6.5% (upper right quarter).

doi:10.1371/journal.pone.0154206.g004

### Ranking of 13 treatments on hypoglycemia, treatment discontinuation, HbA1c <7.0% and HbA1c <6.5%

Table 2 showed the mean values of SUCRA (S1 Fig) for providing the hierarchy of 13 treatments on hypoglycemia, treatment discontinuation, HbA1c <7.0% and HbA1c <6.5% based on the absolute rank probabilities. According to SUCRAs, placebo, TZD and albiglutide had the best decrease effect on hypoglycemia, with probability of 4.08%, 24.95% and 35.19% respectively. SU, sitagliptin and insulin had the lowest probability of treatment discontinuation with rates of 13.44%, 23.01% and 23.63%, respectively. With respect to HbA1c <7.0% and HbA1c <6.5%, exe\_lar and dulaglutide lowering glycaemic level most among 13 treatments.

**Table 2. Ranking: probability from SUCRA of efficacy and acceptability of glycemic control of different GLP-1s.**

CODE	Treatments	Hypoglycemia		Treatment discontinuation		HbA1c <7.0%		HbA1c <6.5%	
		SUCRA	Rank	SUCRA	Rank	SUCRA	Rank	SUCRA	Rank
1	Albiglutide	<b>0.3519</b>	<b>3</b>	0.3644	6	0.4989	7	0.4805	6
2	Dulaglutide	0.5555	10	0.4941	7	<b>0.8460</b>	<b>2</b>	<b>0.8186</b>	<b>2</b>
3	Exe_LAR	0.5027	9	0.6290	9	<b>0.8718</b>	<b>1</b>	<b>0.9531</b>	<b>1</b>
4	Exenatide	0.7519	11	0.7903	12	0.2802	10	0.3122	10
5	Insulin	0.9090	12	<b>0.2363</b>	<b>3</b>	0.5381	6	0.2145	12
6	Liraglutide	0.4444	6	0.7657	11	0.7421	4	<b>0.8149</b>	<b>3</b>
7	Lixisenatide	0.4461	7	0.7311	10	0.2239	12	0.2901	11
8	Met	0.4224	5	0.5147	8	0.4067	8	0.3779	8
9	Sitagliptin	0.3668	4	<b>0.2301</b>	<b>2</b>	0.2362	11	0.3905	7
10	SU	0.9994	13	<b>0.1344</b>	<b>1</b>	0.3690	9	0.3652	9
11	Taspoglutide	0.4597	8	0.9901	13	<b>0.7859</b>	<b>3</b>	0.7792	4
12	TZD	<b>0.2495</b>	<b>2</b>	0.3226	5	0.7011	5	0.7033	5
13	Placebo	<b>0.0408</b>	<b>1</b>	0.2972	4	0.0001	13	0.0000	13

Note: Rank: probability of being the best treatment, of being the second best, the third best and so on, among the 13 comparisons. SUCRA: surface under the cumulative ranking curve. As for hypoglycemia and treatment discontinuation first one means the best safety. As for HbA1c<7.0% and HbA1c<6.5% first one means has the best efficacy.

doi:10.1371/journal.pone.0154206.t002

### Model fit and inconsistency check

Statistical inconsistency between direct and indirect comparisons was generally low for four outcomes. Most loops (networks of three or four comparisons that arise when collating studies involving different selections of competing treatments) were consistent, since their 95% CIs included 0 according to the forest plots, which meant the direct estimation of the summary effect did not differentiate from the indirect estimation. The summary estimations of network meta-analysis are relatively robust.

The model fit was evaluated using the posterior mean of the residual deviance  $\bar{D}_{res}$ . The values of the  $\bar{D}_{res}$  for hypoglycemia, treatment discontinuation, HbA1c<7.0% and HbA1c<6.5% were 121.98, 83.34, 131.68 and 86.80 respectively, which were close to corresponding 152, 104, 145 and 104 of the number of data points for four outcomes, meaning that model's overall fit is relatively satisfactory.

### Discussion

Aside from adequate glycemic control, increasing attention is being paid to the hypoglycemia and treatment discontinuation effect of GLP-1 RAs recently [14, 15]. Our network meta-analysis suggested that all GLP-1 RAs significantly increase the risk of hypoglycemia compared with placebo (except for albiglutide), and reduce the risk of hypoglycemia compared with insulin (except for dulaglutide) and SU. In terms of the increasing incidence of treatment discontinuation, exenatide, liraglutide, lixisenatide and taspoglutide had significant effect when compared with either placebo, insulin, SU or sitagliptin, and exe\_lar only increased the incidence of treatment discontinuation significantly when compared with SU. This was accompanied by taspoglutide in comparison to TZD. Besides, all GLP-1 RAs decreased glycemic level compared with placebo, and dulaglutide, exe\_lar, liraglutide and taspoglutide had significant lowering effect

when compared with sitagliptin (HbA1c<7.0%) and insulin(HbA1c<6.5%). Regarding to HbA1c <6.5%, there was also a significant lowering effect for exenatide and liraglutide in comparison to SU and sitagliptin, dulaglutide in comparison to sitagliptin, exenatide in comparison to Met.

### Effect on hypoglycemia

Hypoglycemia is a common complication of intensive diabetes therapy, which could cause fall, seizure, coma, and even death[109]. The UK Prospective Diabetes Study (UKPDS) reported that maintenance of tight glycemic control in T2DM with insulin treated led to a significant increase in the incidence of hypoglycemia[110]. Our network meta-analysis showed that the significant increasing in the incidence of hypoglycemia was associated with all GLP-1 RAs except for albiglutide, which was consistent with Riddle's study [90]. Riddle's results showed that the incidence of symptomatic hypoglycemia was 28% for lixisenatide and 22% for placebo, and 1.2% subjects had severe hypoglycemia with lixisenatide vs. 0.0% with placebo. While, the beneficial hypoglycemia lowering effect of all GLP-1 RAs was observed when compared with insulin (except for dulaglutide) and SU, which was consistent with previous reviews[10, 111] and clinical trials[112]. Besides, studies also reported that the incidence of hypoglycemia was similar across GLP-1 RA treatment groups, and most of patients with hypoglycemia had the history of treating with concomitant SU therapy [111, 113].

To date, the mechanism of hypoglycemia for T2DM has not been clearly identified. It may involve complex regulation, but it has been shown that  $\beta$ -cell failure precede defects of  $\alpha$ -cell response to lowering glucagon levels in T2DM, indicating that the counter-regulatory effect of glucagon to hypoglycemia is impaired in T2DM[114, 115]. Fukuda's[116] study reported that the degree of  $\alpha$ -cell dysfunction is related with the lack of  $\beta$ -cell function in diabetes. Commonly, insulin represses glucagon secretion as a pulsatile manner, but this coordination is disrupted in patients with T2DM and it could potentially contribute to glucagon dysregulation [117]. So finally, the defect of an increment in glucagon secretion during hypoglycemia is the result of  $\beta$ -cell failure in advanced T2DM[118].

### Treatment discontinuation increasing effect

Our network meta-analysis showed that exenatide, liraglutide, lixisenatide and tasoglutide had significant increasing effect on the incidence of treatment discontinuation when compared with either placebo, insulin, SU or sitagliptin. Exenatide only increased the incidence of treatment discontinuation when compared with SU. This was accompanied by tasoglutide in comparison to TZD. Similar results were indicated in relevant clinical trials[113].

Several reasons may be account for this. Firstly, All GLP-1 RAs are injected subcutaneously, and cannot be administered orally. The incidence of treatment discontinuation among patients who had injection site adverse events was high[119]. Secondly, the adverse events of GLP-1 RAs like nausea, diarrhea, and vomiting, also account for the incidence of treatment discontinuation[120]. Especially for the most commonly occurred nausea, which usually lasts a long time, is a tough experience for T2DM to bear.

### Glycemic level lowering effect

The beneficial glycemic level lowering effect of all GLP-1 RAs in our analysis was consistent with previous studies [10, 121]. Scheen's [122] study reported that the HbA1c lowering potential for GLP-1 RAs is approximately at 1%–1.5% on average. A review of 8 head-to-head phase III clinical programs showed that the primary efficacy endpoint in all of the GLP-1 RAs was change in HbA1c from baseline with a noninferiority margin of 0.4%[111]. Similar results were

indicated in relevant clinical trials. The significant glycemic level lowering effect of EXQW was observed in series of DURATION trials, with mean reductions of -0.9% to -1.63% [18, 44, 123–126]. Liraglutide was found to lower HbA1c by -0.9 to -1.1% [127].

Besides, our study also found that dulaglutide, *exe\_lar*, liraglutide and taspoglutide had significant lowering effect when compared with sitagliptin (HbA1c < 7.0%) and insulin (HbA1c < 6.5%), which was consistent with Nauck's results, which showed that dulaglutide 0.75 mg reducing HbA1c by  $0.87\% \pm 0.06\%$  versus sitagliptin reducing HbA1c by  $0.39\% \pm 0.06\%$  ( $P, 0.001$ ) [128]. Regarding to HbA1c < 6.5%, our study also demonstrated that there was a significant lowering effect for *exe\_lar* and liraglutide in comparison to SU and sitagliptin, dulaglutide in comparison to sitagliptin, *exe\_lar* in comparison to Met.

## Strengths

A major strength of our study is the inclusion of a substantially greater number of trials of GLP-1 RAs than earlier meta-analysis [16, 20, 21], thus it is the largest completed evaluation of GLP-1 RAs' effect on hypoglycemia, treatment discontinuation and glycemic level to date. Furthermore, the network meta-analysis based on Bayesian model makes indirect comparison among multiple treatments available, especially when there are few trials for direct comparison between different anti-diabetic drugs, such as comparisons between dulaglutide and insulin in our study. Network meta-analysis has been proved to be the most appropriate method for multiple treatments comparison to date [26, 129]. In addition, the network technique enables the estimation of the probability that one intervention is the best for one outcome. Thus it can provide an explicit ranking when many treatments are competing for one outcome. Our study provided the ranks of GLP-1 RAs and traditional anti-diabetic drugs on hypoglycemia, treatment discontinuation and glycemic level for the first time.

## Limitations

Several limitations are worthy to be mentioned. First, only trials only in English were included, and our literature search was from inception to June 1st, 2014, and didn't include literatures published after June 1st, 2014, which may lead to potential publication bias and selection bias. Secondly, none of the trials included was specially designed to evaluate the effect of GLP-1 RAs on hypoglycemia, treatment discontinuation and glycemic level. Thirdly, the different duration of years of T2DM in 78 trials may cause heterogeneous, which may influence the different response to therapy and increase the possibility of hypoglycemia. Thus the results of our study should be considered as hypothesis generation, and any conclusions should be drawn with caution.

## Conclusion

In conclusion, our network meta-analysis presents the associations among GLP-1 RAs, traditional anti-diabetic drugs and placebo on hypoglycemia, treatment discontinuation and glycemic level. GLP-1 RAs had the lowering effect on glycemic level, increasing effect on hypoglycemia and treatment discontinuation. While, GLP-1 RAs were associated with lower incidence of hypoglycemia when compared with active comparators. However, further evidence is necessary for more conclusive inferences on mechanisms underlying the increasing in hypoglycemia.

## Supporting Information

**S1 Fig. Plots for ranking probability of different dosing of GLP-1s on impact of SBP, DBP, heart rate and hypertension.**

(PDF)



**S1 File. PRISMA 2009 Checklist.**

(PDF)

**S1 Table. Quality of included trials by adjusted Jadad scale.**

(PDF)

**Acknowledgments**

Funding from the National Natural Science Foundation of China (81302508) and Research Fund for the Doctoral Program of Higher Education (20120001110015) is gratefully acknowledged by all authors. We are grateful to Peking University, McMaster University, Center for Evidence-based and Translational Medicine of Zhongnan Hospital and their staff whose hard work made this study possible. Special thanks to all of the original study authors.

**Author Contributions**

Conceived and designed the experiments: FS SYZ. Performed the experiments: ZXL YZ XCQ ZRY XTZ. Analyzed the data: ZXL FS. Contributed reagents/materials/analysis tools: ZXL YZ XCQ ZRY XTZ LNJ FS SYZ. Wrote the paper: ZXL FS. Contributed to interpreting the results, draft reviewing, and finalizing the paper: LNJ SYZ.

**References**

1. Hansen KB, Knop FK, Holst JJ, Vilsboll T. Treatment of type 2 diabetes with glucagon-like peptide-1 receptor agonists. *Int J Clin Pract*. 2009; 63:1154–60. doi: [10.1111/j.1742-1241.2009.02086.x](https://doi.org/10.1111/j.1742-1241.2009.02086.x) PMID: [19624785](https://pubmed.ncbi.nlm.nih.gov/19624785/)
2. Goykhman S, Drincic A, Desmangles JC, Rendell M. Insulin Glargine: a review 8 years after its introduction. *Expert opinion on pharmacotherapy*. 2009; 10(4):705–18. Epub 2009/03/17. doi: [10.1517/14656560902775677](https://doi.org/10.1517/14656560902775677) PMID: [19284367](https://pubmed.ncbi.nlm.nih.gov/19284367/).
3. Nauck MA, Heimesaat MM, Behle K, Holst JJ, Nauck MS, Ritzel R, et al. Effects of glucagon-like peptide 1 on counterregulatory hormone responses, cognitive functions, and insulin secretion during hyperinsulinemic, stepped hypoglycemic clamp experiments in healthy volunteers. *The Journal of clinical endocrinology and metabolism*. 2002; 87(3):1239–46. Epub 2002/03/13. doi: [10.1210/jcem.87.3.8355](https://doi.org/10.1210/jcem.87.3.8355) PMID: [11889194](https://pubmed.ncbi.nlm.nih.gov/11889194/).
4. Vilsboll T, Krarup T, Madsbad S, Holst JJ. Both GLP-1 and GIP are insulinotropic at basal and postprandial glucose levels and contribute nearly equally to the incretin effect of a meal in healthy subjects. *Regulatory peptides*. 2003; 114(2–3):115–21. Epub 2003/07/02. PMID: [12832099](https://pubmed.ncbi.nlm.nih.gov/12832099/).
5. Zander M, Madsbad S, Madsen JL, Holst JJ. Effect of 6-week course of glucagon-like peptide 1 on glycaemic control, insulin sensitivity, and beta-cell function in type 2 diabetes: a parallel-group study. *Lancet*. 2002; 359(9309):824–30. Epub 2002/03/19. doi: [10.1016/s0140-6736\(02\)07952-7](https://doi.org/10.1016/s0140-6736(02)07952-7) PMID: [11897280](https://pubmed.ncbi.nlm.nih.gov/11897280/).
6. Wajsborg E, Tavarua A. Exenatide: clinical aspects of the first incretin-mimetic for the treatment of type 2 diabetes mellitus. *Expert opinion on pharmacotherapy*. 2009; 10(1):135–42. Epub 2009/02/25. doi: [10.1517/14656560802611832](https://doi.org/10.1517/14656560802611832) PMID: [19236187](https://pubmed.ncbi.nlm.nih.gov/19236187/).
7. Neumiller JJ, Campbell RK. Liraglutide: a once-daily incretin mimetic for the treatment of type 2 diabetes mellitus. *The Annals of pharmacotherapy*. 2009; 43(9):1433–44. Epub 2009/07/30. doi: [10.1345/aph.1M134](https://doi.org/10.1345/aph.1M134) PMID: [19638470](https://pubmed.ncbi.nlm.nih.gov/19638470/).
8. Diabetes facts and figures. Brussels: International Diabetes Federation. 2014 Available from: <http://www.idf.org/worlddiabetesday/toolkit/gp/facts-fig>. Accessed June 12, 2014.
9. Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the expert committee on the diagnosis and classification of diabetes mellitus. *Diabetes care*. 2003; 26 Suppl 1:S5–20. Epub 2002/12/28. PMID: [12502614](https://pubmed.ncbi.nlm.nih.gov/12502614/).
10. Inzucchi SE, Bergenstal RM, Buse JB, Diamant M, Ferrannini E, Nauck M, et al. Management of hyperglycemia in type 2 diabetes, 2015: a patient-centered approach: update to a position statement of the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes care*. 2015; 38(1):140–9. Epub 2014/12/30. doi: [10.2337/dc14-2441](https://doi.org/10.2337/dc14-2441) PMID: [25538310](https://pubmed.ncbi.nlm.nih.gov/25538310/).

11. Kachroo S, Kawabata H, Colilla S, Shi L, Zhao Y, Mukherjee J, et al. Association between hypoglycemia and fall-related events in type 2 diabetes mellitus: analysis of a U.S. commercial database. *Journal of managed care & specialty pharmacy*. 2015; 21(3):243–53. Epub 2015/03/03. PMID: [25726033](#).
12. Danaei G, Finucane MM, Lu Y, Singh GM, Cowan MJ, Paciorek CJ, et al. National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *Lancet*. 2011; 378(9785):31–40. Epub 2011/06/28. doi: [10.1016/s0140-6736\(11\)60679-x](#) PMID: [21705069](#).
13. Curkendall SM, Thomas N, Bell KF, Juneau PL, Weiss AJ. Predictors of medication adherence in patients with type 2 diabetes mellitus. *Current medical research and opinion*. 2013; 29(10):1275–86. Epub 2013/07/03. doi: [10.1185/03007995.2013.821056](#) PMID: [23815104](#).
14. Zarowitz BJ, Conner C. The intersection of safety and adherence: new incretin-based therapies in patients with type 2 diabetes mellitus. *Pharmacotherapy*. 2009; 29(12 Pt 2):55s–67s. Epub 2009/12/02. doi: [10.1592/phco.29.pt2.55S](#) PMID: [19947817](#).
15. Khunti K, Davies M. Glycaemic goals in patients with type 2 diabetes: current status, challenges and recent advances. *Diabetes, obesity & metabolism*. 2010; 12(6):474–84. Epub 2010/06/04. doi: [10.1111/j.1463-1326.2009.01186.x](#) PMID: [20518803](#).
16. Eng C, Kramer CK, Zinman B, Retnakaran R. Glucagon-like peptide-1 receptor agonist and basal insulin combination treatment for the management of type 2 diabetes: a systematic review and meta-analysis. *Lancet*. 2014; 384(9961):2228–34. Epub 2014/09/16. doi: [10.1016/s0140-6736\(14\)61335-0](#) PMID: [25220191](#).
17. Dungan KM, Povedano ST, Forst T, Gonzalez JG, Atisso C, Sealls W, et al. Once-weekly dulaglutide versus once-daily liraglutide in metformin-treated patients with type 2 diabetes (AWARD-6): a randomised, open-label, phase 3, non-inferiority trial. *Lancet*. 2014; 384(9951):1349–57. Epub 2014/07/16. doi: [10.1016/s0140-6736\(14\)60976-4](#) PMID: [25018121](#).
18. Diamant M, Van Gaal L, Guerci B, Stranks S, Han J, Malloy J, et al. Exenatide once weekly versus insulin glargine for type 2 diabetes (DURATION-3): 3-year results of an open-label randomised trial. *The lancet Diabetes & endocrinology*. 2014; 2(6):464–73. Epub 2014/04/16. doi: [10.1016/s2213-8587\(14\)70029-4](#) PMID: [24731672](#).
19. Pratley RE, Nauck MA, Barnett AH, Feinglos MN, Ovalle F, Harman-Boehm I, et al. Once-weekly albiglutide versus once-daily liraglutide in patients with type 2 diabetes inadequately controlled on oral drugs (HARMONY 7): a randomised, open-label, multicentre, non-inferiority phase 3 study. *The lancet Diabetes & endocrinology*. 2014; 2(4):289–97. Epub 2014/04/08. doi: [10.1016/s2213-8587\(13\)70214-6](#) PMID: [24703047](#).
20. Monami M, Marchionni N, Mannucci E. Glucagon-like peptide-1 receptor agonists in type 2 diabetes: a meta-analysis of randomized clinical trials. *European journal of endocrinology / European Federation of Endocrine Societies*. 2009; 160(6):909–17. Epub 2009/03/26. doi: [10.1530/eje-09-0101](#) PMID: [19318378](#).
21. Sun F, Yu K, Wu S, Zhang Y, Yang Z, Shi L, et al. Cardiovascular safety and glycemic control of glucagon-like peptide-1 receptor agonists for type 2 diabetes mellitus: a pairwise and network meta-analysis. *Diabetes research and clinical practice*. 2012; 98(3):386–95. Epub 2012/10/02. doi: [10.1016/j.diabres.2012.09.004](#) PMID: [23020934](#).
22. van Valkenhoef G, Lu G, de Brock B, Hillege H, Ades AE, Welton NJ. Automating network meta-analysis. *Res Synth Method*. 3:285–99.
23. Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gavaghan DJ, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Controlled clinical trials*. 1996; 17(1):1–12. Epub 1996/02/01. PMID: [8721797](#).
24. Dersimonian R, Laird N. Meta-analysis in clinical trials. *Controlled clinical trials*. 1986; 7:177–88. PMID: [3802833](#)
25. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; 21:1539–58. PMID: [12111919](#)
26. Lu G, Ades AE. Combination of direct and indirect evidence in mixed treatment comparisons. *Stat Med*. 2004; 23(20):3105–24. Epub 2004/09/28. doi: [10.1002/sim.1875](#) PMID: [15449338](#).
27. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of clinical epidemiology*. 2011; 64(2):163–71. Epub 2010/08/07. doi: [10.1016/j.jclinepi.2010.03.016](#) PMID: [20688472](#).
28. Spiegelhalter DJ, Best NG, Carlin BP, Van Der Linde A. Bayesian measures of model complexity and fit. *J Royal Stat Soci: Series B (Statistical Methodology)*. 2002; 64:583–639.
29. Song F, Altman DG, Glenny AM, Deeks JJ. Validity of indirect comparison for estimating efficacy of competing interventions: empirical evidence from published meta-analyses. *BMJ (Clinical research*

- ed). 2003; 326(7387):472. Epub 2003/03/01. doi: [10.1136/bmj.326.7387.472](https://doi.org/10.1136/bmj.326.7387.472) PMID: [12609941](https://pubmed.ncbi.nlm.nih.gov/12609941/); PubMed Central PMCID: [PMC150178](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC150178/).
30. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *International journal of surgery (London, England)*. 2010; 8(5):336–41. Epub 2010/02/23. doi: [10.1016/j.ijsu.2010.02.007](https://doi.org/10.1016/j.ijsu.2010.02.007) PMID: [20171303](https://pubmed.ncbi.nlm.nih.gov/20171303/).
  31. Ahren B, Leguizamo Dimas A, Miossec P, Saubadu S, Aronson R. Efficacy and Safety of Lixisenatide Once-Daily Morning or Evening Injections in Type 2 Diabetes Inadequately Controlled on Metformin (GetGoal-M). *Diabetes care*. 2013; 36(9):2543–50. doi: [10.2337/dc12-2006](https://doi.org/10.2337/dc12-2006) PMID: [23536584](https://pubmed.ncbi.nlm.nih.gov/23536584/); PubMed Central PMCID: [PMC3747937](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC3747937/).
  32. Ahren B, Johnson SL, Stewart M, Cirkel DT, Yang F, Perry C, et al. HARMONY 3: 104-Week Randomized, Double-Blind, Placebo- and Active-Controlled Trial Assessing the Efficacy and Safety of Albiglutide Compared With Placebo, Sitagliptin, and Glimepiride in Patients With Type 2 Diabetes Taking Metformin. *Diabetes care*. 2014. doi: [10.2337/dc14-0024](https://doi.org/10.2337/dc14-0024) PMID: [24898304](https://pubmed.ncbi.nlm.nih.gov/24898304/).
  33. Apovian CM, Bergenstal RM, Cuddihy RM, Qu Y, Lenox S, Lewis MS, et al. Effects of Exenatide Combined with Lifestyle Modification in Patients with Type 2 Diabetes. *American Journal of Medicine*. 2010; 123(5):468.e9–.e17.
  34. Barnett AH, Burger J, Johns D, Brodows R, Kendall DM, Roberts A, et al. Tolerability and efficacy of exenatide and titrated insulin glargine in adult patients with type 2 diabetes previously uncontrolled with metformin or a sulfonylurea: A multinational, randomized, open-label, two-period, crossover non-inferiority trial. *Clinical therapeutics*. 2007; 29(11):2333–48. PMID: [18158075](https://pubmed.ncbi.nlm.nih.gov/18158075/)
  35. Bergenstal R, Lewin A, Bailey T, Chang D, Gylvin T, Roberts V, et al. Efficacy and safety of biphasic insulin aspart 70/30 versus exenatide in subjects with type 2 diabetes failing to achieve glycemic control with metformin and a sulfonylurea. *Current Medical Research & Opinion*. 2009; 25(1):65–75. PMID: [19210140](https://pubmed.ncbi.nlm.nih.gov/19210140/).
  36. Bergenstal RM, Wysham C, Macconell L, Malloy J, Walsh B, Yan P, et al. Efficacy and safety of exenatide once weekly versus sitagliptin or pioglitazone as an adjunct to metformin for treatment of type 2 diabetes (DURATION-2): a randomised trial. *Lancet*. 2010; 376(9739):431–9. PMID: [20580422](https://pubmed.ncbi.nlm.nih.gov/20580422/). doi: [10.1016/S0140-6736\(10\)60590-9](https://doi.org/10.1016/S0140-6736(10)60590-9)
  37. Bergenstal RM, Forti A, Chiasson JL, Woloschak M, Boldrin M, Balena R. Efficacy and safety of tasoglutide versus sitagliptin for type 2 diabetes mellitus (T-Emerge 4 Trial). *Diabetes Therapy*. 2012; 3(1):1–19.
  38. Blevins T, Pullman J, Malloy J, Yan P, Taylor K, Schulteis C, et al. DURATION-5: exenatide once weekly resulted in greater improvements in glycemic control compared with exenatide twice daily in patients with type 2 diabetes. *Journal of Clinical Endocrinology & Metabolism*. 2011; 96(5):1301–10. Epub 2011/02/11. doi: [10.1210/jc.2010-2081](https://doi.org/10.1210/jc.2010-2081) PMID: [21307137](https://pubmed.ncbi.nlm.nih.gov/21307137/).
  39. Bolli GB, Munteanu M, Dotsenko S, Niemoeller E, Boka G, Wu Y, et al. Efficacy and safety of lixisenatide once daily vs. placebo in people with Type 2 diabetes insufficiently controlled on metformin (Get-Goal-F1). *Diabetic medicine: a journal of the British Diabetic Association*. 2013. doi: [10.1111/dme.12328](https://doi.org/10.1111/dme.12328) PMID: [24117597](https://pubmed.ncbi.nlm.nih.gov/24117597/).
  40. Bunck MC, Diamant M, Corner A, Eliasson B, Malloy JL, Shaginian RM, et al. One-year treatment with exenatide improves beta-cell function, compared with insulin glargine, in metformin-treated type 2 diabetic patients: a randomized, controlled trial. *Diabetes care*. 2009; 32(5):762–8. PMID: [19196887](https://pubmed.ncbi.nlm.nih.gov/19196887/); PubMed Central PMCID: [PMCSource: NLM. PMC2671094](https://pubmed.ncbi.nlm.nih.gov/PMC/PMCSource:.nlm.nih.gov/PMC2671094/). doi: [10.2337/dc08-1797](https://doi.org/10.2337/dc08-1797)
  41. Buse JB, Henry RR, Han J, Kim DD, Fineman MS, Baron AD, et al. Effects of exenatide (exendin-4) on glycemic control over 30 weeks in sulfonylurea-treated patients with type 2 diabetes. *Diabetes care*. 2004; 27(11):2628–35. PMID: [15504997](https://pubmed.ncbi.nlm.nih.gov/15504997/).
  42. Buse JB, Rosenstock J, Sesti G, Schmidt WE, Montanya E, Brett JH, et al. Liraglutide once a day versus exenatide twice a day for type 2 diabetes: a 26-week randomised, parallel-group, multinational, open-label trial (LEAD-6). *Lancet*. 2009; 374(9683):39–47. PMID: [19515413](https://pubmed.ncbi.nlm.nih.gov/19515413/). doi: [10.1016/S0140-6736\(09\)60659-0](https://doi.org/10.1016/S0140-6736(09)60659-0)
  43. Buse JB, Bergenstal RM, Glass LC, Heilmann CR, Lewis MS, Kwan AYM, et al. Use of twice-daily exenatide in Basal insulin-treated patients with type 2 diabetes: a randomized, controlled trial. *Ann Intern Med*. 2011; 154(2):103–12. PMID: [21138825](https://pubmed.ncbi.nlm.nih.gov/21138825/). doi: [10.7326/0003-4819-154-2-201101180-00300](https://doi.org/10.7326/0003-4819-154-2-201101180-00300)
  44. Buse JB, Nauck M, Forst T, Sheu WH, Shenouda SK, Heilmann CR, et al. Exenatide once weekly versus liraglutide once daily in patients with type 2 diabetes (DURATION-6): a randomised, open-label study. *Lancet*. 2013; 381(9861):117–24. Epub 2012/11/13. doi: [10.1016/s0140-6736\(12\)61267-7](https://doi.org/10.1016/s0140-6736(12)61267-7) PMID: [23141817](https://pubmed.ncbi.nlm.nih.gov/23141817/).
  45. Charbonnel B, Steinberg H, Eymard E, Xu L, Thakkar P, Prabhu V, et al. Efficacy and safety over 26 weeks of an oral treatment strategy including sitagliptin compared with an injectable treatment

- strategy with liraglutide in patients with type 2 diabetes mellitus inadequately controlled on metformin: a randomised clinical trial. *Diabetologia*. 2013; 56(7):1503–11. doi: [10.1007/s00125-013-2905-1](https://doi.org/10.1007/s00125-013-2905-1) PMID: [23604551](https://pubmed.ncbi.nlm.nih.gov/23604551/).
46. Davies M, Heller S, Sreenan S, Sapin H, Adetunji O, Tahbaz A, et al. Once-weekly exenatide versus once- or twice-daily insulin detemir: randomized, open-label, clinical trial of efficacy and safety in patients with type 2 diabetes treated with metformin alone or in combination with sulfonylureas. *Diabetes care*. 2013; 36(5):1368–76. doi: [10.2337/dc12-1333](https://doi.org/10.2337/dc12-1333) PMID: [23275363](https://pubmed.ncbi.nlm.nih.gov/23275363/); PubMed Central PMCID: PMC3631870.
  47. Davies MJ, Donnelly R, Barnett AH, Jones S, Nicolay C, Kilcoyne A. Exenatide compared with long-acting insulin to achieve glycaemic control with minimal weight gain in patients with type 2 diabetes: Results of the helping evaluate exenatide in patients with diabetes compared with long-acting insulin (HEELA) study. *Diabetes, Obesity and Metabolism*. 2009; 11(12):1153–62. doi: [10.1111/j.1463-1326.2009.01154.x](https://doi.org/10.1111/j.1463-1326.2009.01154.x) PMID: [19930005](https://pubmed.ncbi.nlm.nih.gov/19930005/)
  48. Davis SN, Johns D, Maggs D, Xu H, Northrup JH, Brodows RG. Exploring the substitution of exenatide for insulin in patients with type 2 diabetes treated with insulin in combination with oral antidiabetes agents. *Diabetes care*. 2007; 30(11):2767–72. PMID: [17595353](https://pubmed.ncbi.nlm.nih.gov/17595353/)
  49. DeFronzo RA, Ratner RE, Han J, Kim DD, Fineman MS, Baron AD. Effects of exenatide (exendin-4) on glycemic control and weight over 30 weeks in metformin-treated patients with type 2 diabetes. *Diabetes care*. 2005; 28(5):1092–100. PMID: [15855572](https://pubmed.ncbi.nlm.nih.gov/15855572/).
  50. DeFronzo RA, Triplitt C, Qu Y, Lewis MS, Maggs D, Glass LC. Effects of exenatide plus rosiglitazone on (beta)-cell function and insulin sensitivity in subjects with type 2 diabetes on metformin. *Diabetes care*. 2010; 33(5):951–7. doi: [10.2337/dc09-1521](https://doi.org/10.2337/dc09-1521) PMID: [20107105](https://pubmed.ncbi.nlm.nih.gov/20107105/)
  51. Derosa G, Franzetti IG, Querci F, Carbone A, Ciccarelli L, Piccinni MN, et al. Exenatide plus metformin compared with metformin alone on beta-cell function in patients with Type 2 diabetes. *Diabetic medicine: a journal of the British Diabetic Association*. 2012; 29(12):1515–23. doi: [10.1111/j.1464-5491.2012.03699.x](https://doi.org/10.1111/j.1464-5491.2012.03699.x) PMID: [22540883](https://pubmed.ncbi.nlm.nih.gov/22540883/).
  52. Diamant M, Van Gaal L, Stranks S, Guerci B, MacConell L, Haber H, et al. Safety and efficacy of once-weekly exenatide compared with insulin glargine titrated to target in patients with type 2 diabetes over 84 weeks. *Diabetes care*. 2012; 35(4):683–9. doi: [10.2337/dc11-1233](https://doi.org/10.2337/dc11-1233) PMID: [22357185](https://pubmed.ncbi.nlm.nih.gov/22357185/); PubMed Central PMCID: PMC3308312.
  53. Drucker DJ, Buse JB, Taylor K, Kendall DM, Trautmann M, Zhuang D, et al. Exenatide once weekly versus twice daily for the treatment of type 2 diabetes: a randomised, open-label, non-inferiority study. *Lancet*. 2008; 372(9645):1240–50. PMID: [18782641](https://pubmed.ncbi.nlm.nih.gov/18782641/). doi: [10.1016/S0140-6736\(08\)61206-4](https://doi.org/10.1016/S0140-6736(08)61206-4)
  54. Fonseca VA, Alvarado-Ruiz R, Raccach D, Boka G, Miossec P, Gerich JE. Efficacy and Safety of the Once-Daily GLP-1 Receptor Agonist Lixisenatide in Monotherapy: A randomized, double-blind, placebo-controlled trial in patients with type 2 diabetes (GetGoal-Mono). *Diabetes care*. 2012; 35(6):1225–31. Epub 2012/03/21. doi: [10.2337/dc11-1935](https://doi.org/10.2337/dc11-1935) PMID: [22432104](https://pubmed.ncbi.nlm.nih.gov/22432104/); PubMed Central PMCID: PMC3357248.
  55. Gallwitz B, Bohmer M, Segiet T, Molle A, Milek K, Becker B, et al. Exenatide twice daily versus premixed insulin aspart 70/30 in metformin-treated patients with type 2 diabetes: a randomized 26-week study on glycemic control and hypoglycemia. *Diabetes care*. 2011; 34(3):604–6. PMID: [21285388](https://pubmed.ncbi.nlm.nih.gov/21285388/); PubMed Central PMCID: PMCSource: NLM. PMC3041190 [Available on 03/01/12]. doi: [10.2337/dc10-1900](https://doi.org/10.2337/dc10-1900)
  56. Gallwitz B, Guzman J, Dotta F, Guerci B, Sim O R, Basson BR, et al. Exenatide twice daily versus glimepiride for prevention of glycaemic deterioration in patients with type 2 diabetes with metformin failure (EUREXA): an open-label, randomised controlled trial. *The Lancet*. 2012; 6736(12):1–9. doi: [10.1016/S0140-6736\(12\)60479-6](https://doi.org/10.1016/S0140-6736(12)60479-6) PMID: [22683137](https://pubmed.ncbi.nlm.nih.gov/22683137/).
  57. Gao Y, Yoon KH, Chuang L-M, Mohan V, Ning G, Shah S, et al. Efficacy and safety of exenatide in patients of Asian descent with type 2 diabetes inadequately controlled with metformin or metformin and a sulphonylurea. *Diabetes Research & Clinical Practice*. 2009; 83(1):69–76. PMID: [19019476](https://pubmed.ncbi.nlm.nih.gov/19019476/).
  58. Garber A, Henry RR, Ratner R, Hale P, Chang CT, Bode B, et al. Liraglutide, a once-daily human glucagon-like peptide 1 analogue, provides sustained improvements in glycaemic control and weight for 2 years as monotherapy compared with glimepiride in patients with type 2 diabetes. *Diabetes, obesity & metabolism*. 2011; 13(4):348–56. <http://dx.doi.org/10.1111/j.1463-1326.2010.01356.x>. PMID: [21205128](https://pubmed.ncbi.nlm.nih.gov/21205128/); PubMed Central PMCID: PMCSource: NLM. PMC3084519.
  59. Heine RJ, Van Gaal LF, Johns D, Mihm MJ, Widel MH, Brodows RG, et al. Exenatide versus insulin glargine in patients with suboptimally controlled type 2 diabetes: a randomized trial. *Ann Intern Med*. 2005; 143(8):559–69. PMID: [16230722](https://pubmed.ncbi.nlm.nih.gov/16230722/).
  60. Henry RR, Rosenstock J, Logan DK, Alessi TR, Luskey K, Baron MA. Randomized Trial of Continuous Subcutaneous Delivery of Exenatide by ITCA 650 Versus Twice-Daily Exenatide Injections in

- Metformin-Treated Type 2 Diabetes. *Diabetes care*. 2013; 36(9):2559–65. doi: [10.2337/dc12-2410](https://doi.org/10.2337/dc12-2410) PMID: [23645886](https://pubmed.ncbi.nlm.nih.gov/23645886/); PubMed Central PMCID: PMC3747935.
61. Hollander P, Lasko B, Barnett AH, Bengus M, Kanitra L, Pi-Sunyer FX, et al. Effects of tasoglutide on glycemic control and body weight in obese patients with type 2 diabetes (T-merge 7 study). *Obesity* (Silver Spring, Md). 2013; 21(2):238–47. Epub 2013/02/14. doi: [10.1002/oby.20042](https://doi.org/10.1002/oby.20042) PMID: [23404788](https://pubmed.ncbi.nlm.nih.gov/23404788/).
  62. Inagaki N, Atsumi Y, Oura T, Saito H, Imaoka T. Efficacy and safety profile of exenatide once weekly compared with insulin once daily in Japanese patients with type 2 diabetes treated with oral antidiabetes drug(s): results from a 26-week, randomized, open-label, parallel-group, multicenter, noninferiority study. *Clinical therapeutics*. 2012; 34(9):1892–908 e1. doi: [10.1016/j.clinthera.2012.07.007](https://doi.org/10.1016/j.clinthera.2012.07.007) PMID: [22884767](https://pubmed.ncbi.nlm.nih.gov/22884767/).
  63. Iwamoto K, Nasu R, Yamamura A, Kothare PA, Mace K, Wolka AM, et al. Safety, tolerability, pharmacokinetics, and pharmacodynamics of exenatide once weekly in Japanese patients with type 2 diabetes. *Endocrine Journal*. 2009; 56(8):951–62. PMID: [19706990](https://pubmed.ncbi.nlm.nih.gov/19706990/)
  64. Ji L, Onishi Y, Ahn CW, Agarwal P, Chou CW, Haber H, et al. Efficacy and safety of exenatide once-weekly vs exenatide twice-daily in Asian patients with type 2 diabetes mellitus. *Journal of Diabetes Investigation*. 2013; 4(1):53–61. doi: [10.1111/j.2040-1124.2012.00238.x](https://doi.org/10.1111/j.2040-1124.2012.00238.x) PMID: [24843631](https://pubmed.ncbi.nlm.nih.gov/24843631/)
  65. Kadowaki T, Namba M, Yamamura A, Sowa H, Wolka AM, Brodows RG. Exenatide exhibits dose-dependent effects on glycemic control over 12 weeks in Japanese patients with suboptimally controlled type 2 diabetes. *Endocrine Journal*. 2009; 56(3):415–24. PMID: [19194050](https://pubmed.ncbi.nlm.nih.gov/19194050/)
  66. Kadowaki T, Namba M, Imaoka T, Yamamura A, Goto W, Boardman MK, et al. Improved glycemic control and reduced bodyweight with exenatide: A double-blind, randomized, phase 3 study in Japanese patients with suboptimally controlled type 2 diabetes over 24 weeks. *Journal of Diabetes Investigation*. 2011; 2(3):210–7. doi: [10.1111/j.2040-1124.2010.00084.x](https://doi.org/10.1111/j.2040-1124.2010.00084.x) PMID: [24843486](https://pubmed.ncbi.nlm.nih.gov/24843486/)
  67. Kendall DM, Riddle MC, Rosenstock J, Zhuang D, Kim DD, Fineman MS, et al. Effects of exenatide (exendin-4) on glycemic control over 30 weeks in patients with type 2 diabetes treated with metformin and a sulfonylurea. *Diabetes care*. 2005; 28(5):1083–91. PMID: [15855571](https://pubmed.ncbi.nlm.nih.gov/15855571/).
  68. Kim D, MacConell L, Zhuang D, Kothare PA, Trautmann M, Fineman M, et al. Effects of once-weekly dosing of a long-acting release formulation of exenatide on glucose control and body weight in subjects with type 2 diabetes. *Diabetes care*. 2007; 30(6):1487–93. PMID: [17353504](https://pubmed.ncbi.nlm.nih.gov/17353504/)
  69. Li CJ, Li J, Zhang QM, Lv L, Chen R, Lv CF, et al. Efficacy and safety comparison between liraglutide as add-on therapy to insulin and insulin dose-increase in Chinese subjects with poorly controlled type 2 diabetes and abdominal obesity. *Cardiovascular diabetology*. 2012;11.
  70. Liutkus J, Rosas Guzman J, Norwood P, Pop L, Northrup J, Cao D, et al. A placebo-controlled trial of exenatide twice-daily added to thiazolidinediones alone or in combination with metformin. *Diabetes, Obesity and Metabolism*. 2010; 12(12):1058–65. doi: [10.1111/j.1463-1326.2010.01251.x](https://doi.org/10.1111/j.1463-1326.2010.01251.x) PMID: [20977576](https://pubmed.ncbi.nlm.nih.gov/20977576/)
  71. Marre M, Shaw J, Br, x00E, ndle M, Bebakar WMW, et al. Liraglutide, a once-daily human GLP-1 analogue, added to a sulphonylurea over 26 weeks produces greater improvements in glycaemic and weight control compared with adding rosiglitazone or placebo in subjects with Type 2 diabetes (LEAD-1 SU). *Diabetic Medicine*. 2009; 26(3):268–78. PMID: [19317822](https://pubmed.ncbi.nlm.nih.gov/19317822/). doi: [10.1111/j.1464-5491.2009.02666.x](https://doi.org/10.1111/j.1464-5491.2009.02666.x)
  72. Mathieu C, Rodbard HW, Cariou B, Handelsman Y, Philis-Tsimikas A, Francisco AM, et al. A comparison of adding liraglutide versus a single daily dose of insulin aspart to insulin degludec in subjects with type 2 diabetes (BEGIN: VICTOZA ADD-ON). *Diabetes, obesity & metabolism*. 2014. doi: [10.1111/dom.12262](https://doi.org/10.1111/dom.12262) PMID: [24443830](https://pubmed.ncbi.nlm.nih.gov/24443830/).
  73. Moretto TJ, Milton D, x00E, i R, Ridge TD, Macconell LA, et al. Efficacy and tolerability of exenatide monotherapy over 24 weeks in antidiabetic drug-naïve patients with type 2 diabetes: a randomized, double-blind, placebo-controlled, parallel-group study. *Clinical therapeutics*. 2008; 30(8):1448–60. PMID: [18803987](https://pubmed.ncbi.nlm.nih.gov/18803987/). doi: [10.1016/j.clinthera.2008.08.006](https://doi.org/10.1016/j.clinthera.2008.08.006)
  74. Nauck M, Weinstock RS, Umpierrez GE, Guerci B, Skrivaneck Z, Milicevic Z. Efficacy and Safety of Dulaglutide Versus Sitagliptin After 52 Weeks in Type 2 Diabetes in a Randomized Controlled Trial (AWARD-5). *Diabetes care*. 2014. doi: [10.2337/dc13-2761](https://doi.org/10.2337/dc13-2761) PMID: [24742660](https://pubmed.ncbi.nlm.nih.gov/24742660/).
  75. Nauck MA, Duran S, Kim D, Johns D, Northrup J, Festa A, et al. A comparison of twice-daily exenatide and biphasic insulin aspart in patients with type 2 diabetes who were suboptimally controlled with sulfonylurea and metformin: a non-inferiority study. *Diabetologia*. 2007; 50(2):259–67. PMID: [17160407](https://pubmed.ncbi.nlm.nih.gov/17160407/).
  76. Nauck MA, Ratner RE, Kapitza C, Berria R, Boldrin M, Balena R. Treatment with the human once-weekly glucagon-like peptide-1 analog tasoglutide in combination with metformin improves glycemic control and lowers body weight in patients with type 2 diabetes inadequately controlled with metformin alone: A double-blind placebo-controlled study. *Diabetes care*. 2009; 32(7):1237–43. doi: [10.2337/dc08-1961](https://doi.org/10.2337/dc08-1961) PMID: [19366970](https://pubmed.ncbi.nlm.nih.gov/19366970/)

77. Nauck M, Frid A, Hermansen K, Shah NS, Tankova T, Mitha IH, et al. Efficacy and safety comparison of liraglutide, glimepiride, and placebo, all in combination with metformin, in type 2 diabetes. *Diabetes care*. 2009; 32(1):84–90. CN-00754008. doi: [10.2337/dc08-1355](https://doi.org/10.2337/dc08-1355) PMID: [18931095](https://pubmed.ncbi.nlm.nih.gov/18931095/)
78. Nauck M, Horton E, Andjelkovic M, Ampudia-Blasco FJ, Parusel CT, Boldrin M, et al. Taspoglutide, a once-weekly glucagon-like peptide 1 analogue, vs. insulin glargine titrated to target in patients with Type 2 diabetes: an open-label randomized trial. *Diabetic medicine: a journal of the British Diabetic Association*. 2013; 30(1):109–13. Epub 2012/09/04. doi: [10.1111/dme.12003](https://doi.org/10.1111/dme.12003) PMID: [22937895](https://pubmed.ncbi.nlm.nih.gov/22937895/).
79. The Effect of Liraglutide on Endothelial Function in Subjects With Type 2 Diabetes Mellitus. 2011. <http://clinicaltrials.gov/ct2/show/NCT00620282?term=NCT00620282&rank=1>.
80. How Glargine Insulin, Oral Diabetes Medications and Exenatide May Improve Blood Sugar Control and Weight Gain in Type 2 Diabetics. 2013. <http://ClinicalTrials.gov/show/NCT00667732>.
81. Effect of Exenatide on Abdominal Fat Distribution in Patients With Type 2 Diabetes Pretreated With Metformin. 2013. <http://ClinicalTrials.gov/show/NCT00701935>.
82. Pinget M, Goldenberg R, Niemoeller E, Muehlen-Bartmer I, Guo H, Aronson R. Efficacy And Safety Of Lixisenatide Once Daily Versus Placebo In Type 2 Diabetes Insufficiently Controlled On Pioglitazone (Getgoal-P). *Diabetes, obesity & metabolism*. 2013; 15(11):1000–7. doi: [10.1111/dom.12121](https://doi.org/10.1111/dom.12121) PMID: [23627775](https://pubmed.ncbi.nlm.nih.gov/23627775/).
83. Pratley R, Nauck M, Bailey T, Montanya E, Cuddihy R, Filetti S, et al. One year of liraglutide treatment offers sustained and more effective glycaemic control and weight reduction compared with sitagliptin, both in combination with metformin, in patients with type 2 diabetes: A randomised, parallel-group, open-label trial. *International Journal of Clinical Practice*. 2011; 65(4):397–407. doi: [10.1111/j.1742-1241.2011.02656.x](https://doi.org/10.1111/j.1742-1241.2011.02656.x) PMID: [21355967](https://pubmed.ncbi.nlm.nih.gov/21355967/)
84. Pratley RE, Urosevic D, Boldrin M, Balena R. Efficacy and tolerability of taspoglutide versus pioglitazone in subjects with type 2 diabetes uncontrolled with sulphonylurea or sulphonylurea-metformin therapy: A randomized, double-blind study (T-emerge 6). *Diabetes, Obesity and Metabolism*. 2013; 15(3):234–40. doi: [10.1111/dom.12009](https://doi.org/10.1111/dom.12009) PMID: [22958426](https://pubmed.ncbi.nlm.nih.gov/22958426/)
85. Pratley RE, Nauck MA, Barnett AH, Feinglos MN, Ovalle F, Harman-Boehm I, et al. Once-weekly albiglutide versus once-daily liraglutide in patients with type 2 diabetes inadequately controlled on oral drugs (HARMONY 7): a randomised, open-label, multicentre, non-inferiority phase 3 study. *The Lancet Diabetes & Endocrinology*. 2014; 2(4):289–97.
86. Ratner R, Nauck M, Kapitza C, Asnagli V, Boldrin M, Balena R. Safety and tolerability of high doses of taspoglutide, a once-weekly human GLP-1 analogue, in diabetic patients treated with metformin: A randomized double-blind placebo-controlled study. *Diabetic Medicine*. 2010; 27(5):556–62. doi: [10.1111/j.1464-5491.2010.02990.x](https://doi.org/10.1111/j.1464-5491.2010.02990.x) PMID: [20536952](https://pubmed.ncbi.nlm.nih.gov/20536952/)
87. Ratner RE, Rosenstock J, Boka G. Dose-dependent effects of the once-daily GLP-1 receptor agonist lixisenatide in patients with Type 2 diabetes inadequately controlled with metformin: A randomized, double-blind, placebo-controlled trial. *Diabetic Medicine*. 2010; 27(9):1024–32. doi: [10.1111/j.1464-5491.2010.03020.x](https://doi.org/10.1111/j.1464-5491.2010.03020.x) PMID: [20722676](https://pubmed.ncbi.nlm.nih.gov/20722676/)
88. Raz I, Fonseca V, Kipnes M, Durrwell L, Hoekstra J, Boldrin M, et al. Efficacy and safety of taspoglutide monotherapy in drug-naive type 2 diabetic patients after 24 weeks of treatment: results of a randomized, double-blind, placebo-controlled phase 3 study (T-emerge 1). *Diabetes care*. 2012; 35(3):485–7. Epub 2012/02/04. doi: [10.2337/dc11-1942](https://doi.org/10.2337/dc11-1942) PMID: [22301126](https://pubmed.ncbi.nlm.nih.gov/22301126/).
89. Riddle MC, Forst T, Aronson R, Sauque-Reyna L, Souhami E, Silvestre L, et al. Adding Once-Daily Lixisenatide for Type 2 Diabetes Inadequately Controlled With Newly Initiated and Continuously Titrated Basal Insulin Glargine: A 24-Week, Randomized, Placebo-Controlled Study (GetGoal-Duo 1). *Diabetes care*. 2013; 36(9):2497–503. doi: [10.2337/dc12-2462](https://doi.org/10.2337/dc12-2462) PMID: [23564915](https://pubmed.ncbi.nlm.nih.gov/23564915/); PubMed Central PMCID: [PMC3747901](https://pubmed.ncbi.nlm.nih.gov/PMC3747901/).
90. Riddle MC, Aronson R, Home P, Marre M, Niemoeller E, Miossec P, et al. Adding once-daily lixisenatide for type 2 diabetes inadequately controlled by established basal insulin: a 24-week, randomized, placebo-controlled comparison (GetGoal-L). *Diabetes care*. 2013; 36(9):2489–96. Epub 2013/05/01. doi: [10.2337/dc12-2454](https://doi.org/10.2337/dc12-2454) PMID: [23628617](https://pubmed.ncbi.nlm.nih.gov/23628617/); PubMed Central PMCID: [PMC3747925](https://pubmed.ncbi.nlm.nih.gov/PMC3747925/).
91. Rosenstock J, Balas B, Charbonnel B, Bolli GB, Boldrin M, Ratner R, et al. The Fate of Taspoglutide, a Weekly GLP-1 Receptor Agonist, Versus Twice-Daily Exenatide for Type 2 Diabetes: The T-Emerge 2 Trial. *Diabetes care*. 2013; 36(3):498–504. doi: [10.2337/dc12-0709](https://doi.org/10.2337/dc12-0709) PMID: [23139373](https://pubmed.ncbi.nlm.nih.gov/23139373/); PubMed Central PMCID: [PMC3579343](https://pubmed.ncbi.nlm.nih.gov/PMC3579343/).
92. Rosenstock J, Hanefeld M, Shamanna P, Min KW, Boka G, Miossec P, et al. Beneficial effects of once-daily lixisenatide on overall and postprandial glycemic levels without significant excess of hypoglycemia in type 2 diabetes inadequately controlled on a sulfonylurea with or without metformin (Get-Goal-S). *Journal of diabetes and its complications*. 2014; 28(3):386–92. doi: [10.1016/j.jdiacomp.2014.01.012](https://doi.org/10.1016/j.jdiacomp.2014.01.012) PMID: [24650952](https://pubmed.ncbi.nlm.nih.gov/24650952/).

93. Rosenstock J, Fonseca VA, Gross JL, Ratner RE, Ahren B, Chow FC, et al. Advancing Basal Insulin Replacement in Type 2 Diabetes Inadequately Controlled With Insulin Glargine Plus Oral Agents: A Comparison of Adding Albiglutide, a Weekly GLP-1 Receptor Agonist, Versus Thrice-Daily Prandial Insulin Lispro. *Diabetes care*. 2014. doi: [10.2337/dc14-0001](https://doi.org/10.2337/dc14-0001) PMID: [24898300](https://pubmed.ncbi.nlm.nih.gov/24898300/).
94. Rosenstock J, Reusch J, Bush M, Yang F, Stewart M, Albiglutide Study G. Potential of albiglutide, a long-acting GLP-1 receptor agonist, in type 2 diabetes: a randomized controlled trial exploring weekly, biweekly, and monthly dosing. *Diabetes care*. 2009; 32(10):1880–6. PMID: [19592625](https://pubmed.ncbi.nlm.nih.gov/19592625/); PubMed Central PMCID: PMCSource: NLM. PMC2752910 [Available on 10/01/10]. doi: [10.2337/dc09-0366](https://doi.org/10.2337/dc09-0366)
95. Rosenstock J, Raccach D, Koranyi L, Maffei L, Boka G, Miossec P, et al. Efficacy and Safety of Lixisenatide Once Daily Versus Exenatide Twice Daily in Type 2 Diabetes Inadequately Controlled on Metformin: A 24-Week, Randomized, Open-Label, Active-Controlled Study (GetGoal-X). *Diabetes care*. 2013. doi: [10.2337/dc12-2709](https://doi.org/10.2337/dc12-2709) PMID: [23698396](https://pubmed.ncbi.nlm.nih.gov/23698396/).
96. Russell-Jones D, Vaag A, Schmitz O, Sethi BK, Lalic N, Antic S, et al. Liraglutide vs insulin glargine and placebo in combination with metformin and sulfonylurea therapy in type 2 diabetes mellitus (LEAD-5 met+SU): A randomised controlled trial. *Diabetologia*. 2009; 52(10):2046–55. doi: [10.1007/s00125-009-1472-y](https://doi.org/10.1007/s00125-009-1472-y) PMID: [19688338](https://pubmed.ncbi.nlm.nih.gov/19688338/)
97. Russell-Jones D, Cuddihy RM, Hanefeld M, Kumar A, Gonzalez JG, Chan M, et al. Efficacy and safety of exenatide once weekly versus metformin, pioglitazone, and sitagliptin used as monotherapy in drug-naïve patients with type 2 diabetes (DURATION-4): a 26-week double-blind study. *Diabetes care*. 2012; 35(2):252–8. Epub 2012/01/03. doi: [10.2337/dc11-1107](https://doi.org/10.2337/dc11-1107) PMID: [22210563](https://pubmed.ncbi.nlm.nih.gov/22210563/); PubMed Central PMCID: PMC3263915.
98. Seino Y, Rasmussen MF, Zdravkovic M, Kaku K. Dose-dependent improvement in glycemia with once-daily liraglutide without hypoglycemia or weight gain: A double-blind, randomized, controlled trial in Japanese patients with type 2 diabetes. *Diabetes research and clinical practice*. 2008; 81(2):161–8. doi: [10.1016/j.diabres.2008.03.018](https://doi.org/10.1016/j.diabres.2008.03.018) PMID: [18495285](https://pubmed.ncbi.nlm.nih.gov/18495285/)
99. Seino Y, Rasmussen MF, Nishida T, Kaku K. Efficacy and safety of the once-daily human GLP-1 analogue, liraglutide, vs glibenclamide monotherapy in Japanese patients with type 2 diabetes. *Current medical research and opinion*. 2010; 26(5):1013–22. doi: [10.1185/03007991003672551](https://doi.org/10.1185/03007991003672551) PMID: [20199137](https://pubmed.ncbi.nlm.nih.gov/20199137/)
100. Seino Y, Min KW, Niemoeller E, Takami A, Investigators EG-LAS. Randomized, double-blind, placebo-controlled trial of the once-daily GLP-1 receptor agonist lixisenatide in Asian patients with type 2 diabetes insufficiently controlled on basal insulin with or without a sulfonylurea (GetGoal-L-Asia). *Diabetes, obesity & metabolism*. 2012; 14(10):910–7. doi: [10.1111/j.1463-1326.2012.01618.x](https://doi.org/10.1111/j.1463-1326.2012.01618.x) PMID: [22564709](https://pubmed.ncbi.nlm.nih.gov/22564709/); PubMed Central PMCID: PMC3466411.
101. Seino Y, Inagaki N, Miyahara H, Okuda I, Bush M, Ye J, et al. A randomized dose-finding study demonstrating the efficacy and tolerability of albiglutide in Japanese patients with type 2 diabetes mellitus. *Current medical research and opinion*. 2014; 30(6):1095–106. doi: [10.1185/03007995.2014.896327](https://doi.org/10.1185/03007995.2014.896327) PMID: [24552155](https://pubmed.ncbi.nlm.nih.gov/24552155/).
102. Umpierrez G, Povedano ST, Manghi FP, Shurzinske L, Pechter V. Efficacy and Safety of Dulaglutide Monotherapy Versus Metformin in Type 2 Diabetes in a Randomized Controlled Trial (AWARD-3). *Diabetes care*. 2014. doi: [10.2337/dc13-2759](https://doi.org/10.2337/dc13-2759) PMID: [24842985](https://pubmed.ncbi.nlm.nih.gov/24842985/).
103. Umpierrez GE, Blevins T, Rosenstock J, Cheng C, Anderson JH, Bastyr EJ 3rd. The effects of LY2189265, a long-acting glucagon-like peptide-1 analogue, in a randomized, placebo-controlled, double-blind study of overweight/obese patients with type 2 diabetes: the EGO study. *Diabetes, obesity & metabolism*. 2011; 13(5):418–25. Epub 2011/01/22. doi: [10.1111/j.1463-1326.2011.01366.x](https://doi.org/10.1111/j.1463-1326.2011.01366.x) PMID: [21251180](https://pubmed.ncbi.nlm.nih.gov/21251180/).
104. Wysham C, Blevins T, Arakaki R, Colon G, Garcia P, Atisso C, et al. Efficacy and Safety of Dulaglutide Added on to Pioglitazone and Metformin Versus Exenatide in Type 2 Diabetes in a Randomized Controlled Trial (AWARD-1). *Diabetes care*. 2014. doi: [10.2337/dc13-2760](https://doi.org/10.2337/dc13-2760) PMID: [24879836](https://pubmed.ncbi.nlm.nih.gov/24879836/).
105. Yang W, Chen L, Ji Q, Liu X, Ma J, Tandon N, et al. Liraglutide provides similar glycaemic control as glimepiride (both in combination with metformin) and reduces body weight and systolic blood pressure in Asian population with type 2 diabetes from China, South Korea and India: A 16-week, randomized, double-blind, active control trial. *Diabetes, Obesity and Metabolism*. 2011; 13(1):81–8. doi: [10.1111/j.1463-1326.2010.01323.x](https://doi.org/10.1111/j.1463-1326.2010.01323.x) PMID: [21114607](https://pubmed.ncbi.nlm.nih.gov/21114607/)
106. Yuan GH, Song WL, Huang YY, Guo XH, Gao Y. Efficacy and tolerability of exenatide monotherapy in obese patients with newly diagnosed type 2 diabetes: A randomized, 26 weeks metformin-controlled, parallel-group study. *Chinese Medical Journal*. 2012; 125(15):2677–81. PMID: [22931974](https://pubmed.ncbi.nlm.nih.gov/22931974/)
107. Zinman B, Hoogwerf BJ, Duran Garcia S, Milton DR, Giaconia JM, Kim DD, et al. The effect of adding exenatide to a thiazolidinedione in suboptimally controlled type 2 diabetes: a randomized trial. *Annals of internal medicine*. 2007; 146(7):477–85. Epub 2007/04/04. PMID: [17404349](https://pubmed.ncbi.nlm.nih.gov/17404349/).

108. Zinman B, Gerich J, Buse JB, Lewin A, Schwartz S, Raskin P, et al. Efficacy and safety of the human glucagon-like peptide-1 analog liraglutide in combination with metformin and thiazolidinedione in patients with type 2 diabetes (LEAD-4 Met+TZD). [Erratum appears in *Diabetes Care*. 2010 Mar;33(3):692]. *Diabetes care*. 2009; 32(7):1224–30. PMID: [19289857](#); PubMed Central PMCID: PMCSource: NLM. PMC2699702. doi: [10.2337/dc08-2124](#)
109. Cryer PE. Hypoglycemia: pathophysiology, diagnosis and treatment. New York: Oxford University Press. 1997.
110. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) Group. *Lancet*. 1998; 352(9131):837–53. Epub 1998/09/22. PMID: [9742976](#).
111. Trujillo JM, Nuffer W, Ellis SL. GLP-1 receptor agonists: a review of head-to-head clinical studies. *Therapeutic advances in endocrinology and metabolism*. 2015; 6(1):19–28. Epub 2015/02/14. doi: [10.1177/2042018814559725](#) PMID: [25678953](#); PubMed Central PMCID: PMCPmc4321870.
112. Inagaki N, Atsumi Y, Oura T, Saito H, Imaoka T. Efficacy and safety profile of exenatide once weekly compared with insulin once daily in Japanese patients with type 2 diabetes treated with oral antidiabetic drug(s): results from a 26-week, randomized, open-label, parallel-group, multicenter, noninferiority study. *Clinical therapeutics*. 2012; 34(9):1892–908.e1. Epub 2012/08/14. doi: [10.1016/j.clinthera.2012.07.007](#) PMID: [22884767](#).
113. Montilla S, Marchesini G, Sammarco A, Trotta MP, Siviero PD, Tomino C, et al. Drug utilization, safety, and effectiveness of exenatide, sitagliptin, and vildagliptin for type 2 diabetes in the real world: data from the Italian AIFA Anti-diabetics Monitoring Registry. *Nutrition, metabolism, and cardiovascular diseases: NMCD*. 2014; 24(12):1346–53. Epub 2014/10/11. doi: [10.1016/j.numecd.2014.07.014](#) PMID: [25300980](#).
114. Dunning BE, Gerich JE. The role of alpha-cell dysregulation in fasting and postprandial hyperglycemia in type 2 diabetes and therapeutic implications. *Endocrine reviews*. 2007; 28(3):253–83. Epub 2007/04/06. doi: [10.1210/er.2006-0026](#) PMID: [17409288](#).
115. Rizza RA. Pathogenesis of fasting and postprandial hyperglycemia in type 2 diabetes: implications for therapy. *Diabetes*. 2010; 59(11):2697–707. Epub 2010/08/14. doi: [10.2337/db10-1032](#) PMID: [20705776](#); PubMed Central PMCID: PMCPmc2963523.
116. Fukuda M, Tanaka A, Tahara Y, Ikegami H, Yamamoto Y, Kumahara Y, et al. Correlation between minimal secretory capacity of pancreatic beta-cells and stability of diabetic control. *Diabetes*. 1988; 37(1):81–8. Epub 1988/01/01. PMID: [3275557](#).
117. Menge BA, Gruber L, Jorgensen SM, Deacon CF, Schmidt WE, Veldhuis JD, et al. Loss of inverse relationship between pulsatile insulin and glucagon secretion in patients with type 2 diabetes. *Diabetes*. 2011; 60(8):2160–8. Epub 2011/06/17. doi: [10.2337/db11-0251](#) PMID: [21677283](#); PubMed Central PMCID: PMCPmc3142077.
118. Segel SA, Paramore DS, Cryer PE. Hypoglycemia-associated autonomic failure in advanced type 2 diabetes. *Diabetes*. 2002; 51(3):724–33. Epub 2002/03/02. PMID: [11872673](#).
119. Home PD, Shamanna P, Stewart M, Yang F, Miller M, Perry C, et al. Efficacy and tolerability of albiglutide versus placebo or pioglitazone over 1 year in people with type 2 diabetes currently taking metformin and glimepiride: HARMONY 5. *Diabetes, obesity & metabolism*. 2015; 17(2):179–87. Epub 2014/11/20. doi: [10.1111/dom.12414](#) PMID: [25406730](#).
120. Gallwitz B. Benefit-risk assessment of exenatide in the therapy of type 2 diabetes mellitus. *Drug safety*. 2010; 33(2):87–100. Epub 2010/01/20. doi: [10.2165/11319130-000000000-00000](#) PMID: [20082536](#).
121. Lindamood CA, Taylor JR. Emerging New Therapies for the Treatment of Type 2 Diabetes Mellitus: Glucagon-like Peptide-1 Receptor Agonists. *Clinical therapeutics*. 2015; 37(3):483–93. Epub 2015/02/11. doi: [10.1016/j.clinthera.2015.01.003](#) PMID: [25659912](#).
122. Scheen AJ. GLP-1 receptor agonists or DPP-4 inhibitors: how to guide the clinician? *Annales d'endocrinologie*. 2013; 74(5–6):515–22. Epub 2013/04/11. doi: [10.1016/j.ando.2012.06.002](#) PMID: [23570814](#).
123. Buse JB, Drucker DJ, Taylor KL, Kim T, Walsh B, Hu H, et al. DURATION-1: exenatide once weekly produces sustained glycemic control and weight loss over 52 weeks. *Diabetes care*. 2010; 33(6):1255–61. Epub 2010/03/11. doi: [10.2337/dc09-1914](#) PMID: [20215461](#); PubMed Central PMCID: PMCPmc2875434.
124. Bergenstal RM, Wysham C, Macconell L, Malloy J, Walsh B, Yan P, et al. Efficacy and safety of exenatide once weekly versus sitagliptin or pioglitazone as an adjunct to metformin for treatment of type 2 diabetes (DURATION-2): a randomised trial. *Lancet*. 2010; 376(9739):431–9. Epub 2010/06/29. doi: [10.1016/s0140-6736\(10\)60590-9](#) PMID: [20580422](#).



125. Russell-Jones D, Cuddihy RM, Hanefeld M, Kumar A, Gonzalez JG, Chan M, et al. Efficacy and safety of exenatide once weekly versus metformin, pioglitazone, and sitagliptin used as monotherapy in drug-naive patients with type 2 diabetes (DURATION-4): a 26-week double-blind study. *Diabetes care*. 2012; 35(2):252–8. Epub 2012/01/03. doi: [10.2337/dc11-1107](https://doi.org/10.2337/dc11-1107) PMID: [22210563](https://pubmed.ncbi.nlm.nih.gov/22210563/); PubMed Central PMCID: [PMCPmc3263915](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC3263915/).
126. Blevins T, Pullman J, Malloy J, Yan P, Taylor K, Schulteis C, et al. DURATION-5: exenatide once weekly resulted in greater improvements in glycemic control compared with exenatide twice daily in patients with type 2 diabetes. *The Journal of clinical endocrinology and metabolism*. 2011; 96(5):1301–10. Epub 2011/02/11. doi: [10.1210/jc.2010-2081](https://doi.org/10.1210/jc.2010-2081) PMID: [21307137](https://pubmed.ncbi.nlm.nih.gov/21307137/).
127. Garber A, Henry RR, Ratner R, Hale P, Chang CT, Bode B. Liraglutide, a once-daily human glucagon-like peptide 1 analogue, provides sustained improvements in glycaemic control and weight for 2 years as monotherapy compared with glimepiride in patients with type 2 diabetes. *Diabetes, obesity & metabolism*. 2011; 13(4):348–56. Epub 2011/01/06. doi: [10.1111/j.1463-1326.2010.01356.x](https://doi.org/10.1111/j.1463-1326.2010.01356.x) PMID: [21205128](https://pubmed.ncbi.nlm.nih.gov/21205128/); PubMed Central PMCID: [PMCPmc3084519](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC3084519/).
128. Nauck M, Weinstock RS, Umpierrez GE, Guerci B, Skrivanek Z, Milicevic Z. Efficacy and safety of dulaglutide versus sitagliptin after 52 weeks in type 2 diabetes in a randomized controlled trial (AWARD-5). *Diabetes care*. 2014; 37(8):2149–58. Epub 2014/04/20. doi: [10.2337/dc13-2761](https://doi.org/10.2337/dc13-2761) PMID: [24742660](https://pubmed.ncbi.nlm.nih.gov/24742660/); PubMed Central PMCID: [PMCPmc4113177](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC4113177/).
129. Caldwell DM, Ades AE, Higgins JP. Simultaneous comparison of multiple treatments: combining direct and indirect evidence. *BMJ (Clinical research ed)*. 2005; 331(7521):897–900. Epub 2005/10/15. doi: [10.1136/bmj.331.7521.897](https://doi.org/10.1136/bmj.331.7521.897) PMID: [16223826](https://pubmed.ncbi.nlm.nih.gov/16223826/); PubMed Central PMCID: [PMCPmc1255806](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC1255806/).