



# Signal Integration in Quorum Sensing Enables Cross-Species Induction of Virulence in *Pectobacterium wasabiae*

Rita S. Valente, Pol Nadal-Jimenez,\* André F. P. Carvalho, Filipe J. D. Vieira, Karina B. Xavier

Instituto Gulbenkian de Ciência, Oeiras, Portugal

**ABSTRACT** Bacterial communities can sense their neighbors, regulating group behaviors in response to cell density and environmental changes. The diversity of signaling networks in a single species has been postulated to allow custom responses to different stimuli; however, little is known about how multiple signals are integrated and the implications of this integration in different ecological contexts. In the plant pathogen *Pectobacterium wasabiae* (formerly *Erwinia carotovora*), two signaling networks—the N-acyl homoserine lactone (AHL) quorum-sensing system and the Gac/Rsm signal transduction pathway—control the expression of secreted plant cell wall-degrading enzymes, its major virulence determinants. We show that the AHL system controls the Gac/Rsm system by affecting the expression of the regulatory RNA RsmB. This regulation is mediated by ExpR2, the quorum-sensing receptor that responds to the *P. wasabiae* cognate AHL but also to AHLs produced by other bacterial species. As a consequence, this level of regulation allows *P. wasabiae* to bypass the Gac-dependent regulation of RsmB in the presence of exogenous AHLs or AHL-producing bacteria. We provide *in vivo* evidence that this pivotal role of RsmB in signal transduction is important for the ability of *P. wasabiae* to induce virulence in response to other AHL-producing bacteria in multispecies plant lesions. Our results suggest that the signaling architecture in *P. wasabiae* was coopted to prime the bacteria to eavesdrop on other bacteria and quickly join the efforts of other species, which are already exploiting host resources.

**IMPORTANCE** Quorum-sensing mechanisms enable bacteria to communicate through small signal molecules and coordinate group behaviors. Often, bacteria have various quorum-sensing receptors and integrate information with other signal transduction pathways, presumably allowing them to respond to different ecological contexts. The plant pathogen *Pectobacterium wasabiae* has two N-acyl homoserine lactone receptors with apparently the same regulatory functions. Our work revealed that the receptor with the broadest signal specificity is also responsible for establishing the link between the main signaling pathways regulating virulence in *P. wasabiae*. This link is essential to provide *P. wasabiae* with the ability to induce virulence earlier in response to higher densities of other bacterial species. We further present *in vivo* evidence that this novel regulatory link enables *P. wasabiae* to join related bacteria in the effort to degrade host tissue in multispecies plant lesions. Our work provides support for the hypothesis that interspecies interactions are among the major factors influencing the network architectures observed in bacterial quorum-sensing pathways.

Bacteria use different signaling systems to communicate with one another, synchronizing gene expression and coordinating their functions to engage in group behaviors (1, 2). However, the mechanisms by which cells integrate and interpret multiple signals received by their neighbors and the surrounding environment, as well

Received 9 March 2017 Accepted 28 April 2017 Published 23 May 2017

**Citation** Valente RS, Nadal-Jimenez P, Carvalho AFP, Vieira FJD, Xavier KB. 2017. Signal integration in quorum sensing enables cross-species induction of virulence in *Pectobacterium wasabiae*. mBio 8:e00398-17. <https://doi.org/10.1128/mBio.00398-17>.


**Editor** Bonnie Bassler, Princeton University

**Copyright** © 2017 Valente et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Karina B. Xavier, [kxavier@igc.gulbenkian.pt](mailto:kxavier@igc.gulbenkian.pt).

\* Present address: Pol Nadal-Jimenez, Institute of Integrative Biology, University of Liverpool, Liverpool, United Kingdom

R.S.V. and P.N.J. contributed equally to this work.



**FIG 1** Major signal transduction pathways regulating PCWDEs in *Pectobacterium* spp. In *Pectobacterium* spp. at low cell density, when the AHL signals are low, *ExpR1* and *ExpR2* induce the transcription of *rsmA*, repressing expression of PCWDEs and the AHL synthase *ExpI*. As cell density increases, AHLs accumulate and inactivate *ExpR1* and *ExpR2*. As a result, *rsmA* transcription is no longer induced. The GacS/GacA system is also active at high cell density and promotes the transcription of *rsmB*, a noncoding RNA that blocks RsmA activity by sequestration. Inhibition of RsmA will, in turn, result in increased production of PCWDEs and induction of virulence. We show here that regulation of RsmB is not exclusively controlled by the GacS/GacA system but is also regulated by AHLs. In the absence of AHLs, *ExpR1* and *ExpR2* repress the expression of *rsmB*. Repression of *rsmB* mediated by *ExpR1* requires RsmA, while *ExpR2* represses RsmB in an RsmA-dependent and -independent manner. Therefore, at high cell density, *rsmB* expression increases as a result of the activation mediated by the GacS/GacA system and relief of the repression mediated by *ExpR1* and *ExpR2*. Gray and black lines/boxes indicate inactive and active pathways, respectively. Arrows indicate activation, while intersecting lines indicate repression. Solid lines indicate transcriptional regulation, while dashed lines indicate posttranscriptional and posttranslational mechanisms. Not all of the regulatory interactions displayed here have been proven to occur through direct interactions.

as the consequences of such integration, remain largely unknown. The *Pectobacterium* genus includes an important group of plant pathogens whose virulence is characterized by the production of a large array of pectolytic plant cell wall-degrading enzymes (PCWDEs). Two main signaling pathways are responsible for the coordinated production of these virulence factors: the quorum-sensing N-acyl homoserine lactone (AHL) system and the Gac/Rsm system (Fig. 1). These two signal transduction pathways are also used by multiple gammaproteobacteria to regulate virulence (3–8).

In *Pectobacterium wasabiae*, the AHL system includes *ExpI*, a LuxI-type AHL synthase, and two AHL receptors, *ExpR1* and *ExpR2*, which are LuxR homologues. These two receptors control the transcription of *rsmA*, a global repressor of virulence (9). Unlike in most AHL receptors, the DNA binding activity of *ExpR1* and *ExpR2* is repressed by ligand binding, and thus, AHLs prevent the transcription of *rsmA*, derepressing the expression of virulence mainly through the control of PCWDE production. These two quorum-sensing receptors have different ligand specificities: *ExpR1* recognizes the main AHL produced by *P. wasabiae* (3-oxo-C8-HSL, 3-oxo-octanoyl homoserine lactone) and to a lesser extent 3-oxo-C10-HSL (3-oxo-decanoyl homoserine lactone), *ExpR2* binds to a broader range of AHLs, such as C6-HSL (hexanoyl homoserine lactone), 3-oxo-C6-HSL, C7-HSL (heptanoyl homoserine lactone), 3-oxo-C8-HSL, and, to a lesser extent, C8-HSL (octanoyl homoserine lactone) and 3-oxo-C10-HSL (decanoyl homoserine lactone) (9–12). Some of these AHLs are produced by other pectolytic bacteria,

while *P. wasabiae* produces mainly 3-oxo-C8-HSL and trace amounts of C8-HSL and oxo-C6-HSL (9). Therefore, it has been proposed that ExpR1 responds mainly to the cognate signal (3-oxo-C8-HSL) while ExpR2 responds to the cognate signal but also to AHLs produced by other species (11).

The Gac/Rsm system comprises the two-component GacS/GacA system (also named ExpS/ExpA in *Pectobacterium* species) and the noncoding regulatory RNA RsmB. Upon activation, GacS phosphorylates the transcriptional regulator GacA, which, in turn, activates the transcription of *rsmB*. RsmB binds to RsmA, neutralizing its mRNA binding activity (13, 14) and derepressing the translation of all RsmA targets, including that of PCWDEs (11, 15, 16).

Despite many efforts to identify the signal(s) responsible for the activation of the Gac/Rsm system, the precise chemical structure of this signal(s) remains elusive (17). Accumulation of intermediates of the Krebs cycle and production of short-chain fatty acids at low pH have been shown to stimulate the system. However, the relevant stimuli leading to the accumulation of these metabolites and activation of the system at neutral pH have not been identified (18–20). Recently, extracellular potassium and calcium have been described as important environmental signals that trigger activation of the Gac/Rsm system in *P. wasabiae* and *Pseudomonas aeruginosa*, respectively (21, 22). Nonetheless, other signals, presumably self-produced molecules, that accumulate in the extracellular growth medium are still required for full activation (23).


In *Pectobacterium* spp., the *rsmA* promoter is the only known direct target of ExpR1 and ExpR2 (10, 11, 24). At low cell density, the absence of AHLs results in ExpR1 and ExpR2 binding to the *rsmA* promoter region, activating its transcription and repressing virulence, while at high cell density, high AHL concentrations result in inhibition of the binding of ExpR1 and ExpR2 to the *rsmA* promoter, preventing *rsmA* transcription. Repression of RsmA expression by AHLs together with its sequestration by RsmB, results in inhibition of RsmA-mediated regulation and consequent virulence expression (10, 11, 24). Thus, in *Pectobacterium* spp., both pathways converge at the level of RsmA: while the AHL system controls *rsmA* transcription (11, 25), Gac/Rsm posttranscriptionally regulates RsmA activity via the binding of RsmB to RsmA (14), (Fig. 1). Additionally, there is evidence of feedback inhibition mediated by RsmA where, at low cell density, when RsmA levels are high, the mRNA of *expl* is targeted by RsmA, decreasing *Expl* production (15).

Here we investigated the mechanisms of signal integration in *P. wasabiae*, a bacterium known to rely on the integration of multiple signals to regulate virulence expression and assess population density (3). We report that the AHL network also controls the Gac/Rsm system by modulating the transcription of *rsmB*. We show that this novel level of regulation is essential for the ability of *P. wasabiae* to respond to the presence of other AHL-producing bacteria in multispecies plant lesions. By studying the architecture of the major signaling networks regulating virulence in *P. wasabiae*, we provide *in vivo* evidence that integration of multiple signaling pathways can be important in the regulation of bacterial group behaviors in multispecies communities.

## RESULTS

### Two signaling pathways are required to control virulence in *P. wasabiae*.

Previous genetic studies targeting genes affecting virulence and PCWDE production in *Pectobacterium* spp. identified transposon insertions in *expl*, *gacS*, and *gacA* (13, 26, 27). However, while a *gacA::Tn10* mutant was completely avirulent, a *gacS::Tn10* mutant was partially attenuated. We determined the virulence phenotype in *gacA* and *gacS* full-deletion mutants of *P. wasabiae* SCC3193 (28). Polygalacturonase and pectate lyase are two of the main PCWDEs in *Pectobacterium* spp. Analysis of *pehA* (polygalacturonase gene) promoter expression (Fig. 2A), enzymatic activity of the pectate lyase (Pel) (Fig. 2B), and virulence in a potato maceration infection model in *P. wasabiae* revealed that virulence was fully abolished in both the *gacA* and *gacS* deletion mutants (Fig. 2C). Supporting the possibility that the transposon insertion in the *gacS::Tn10* mutant described previously is not a full loss-of-function mutant (27). We have also constructed




**FIG 2** Both *expl* and the Gac/Rsm system are required for virulence in *P. wasabiae*. Virulence expression in WT *P. wasabiae* SCC3193 and *expl*, *gacS*, *gacA*, and *rsmB* mutants was measured with a  $P_{pehA}::gfp$  reporter fusion (A), the Pel enzymatic assay (B), and a potato infection assay at 48 h (C). Growth curves of the strains used are shown in panel D. \*\*\*,  $P < 0.001$  by Mann-Whitney test ( $n = 8$ ). Holm-Bonferroni correction was used for multiple comparisons. a.u., arbitrary units. Error bars represent the standard deviation of the mean. For panels A, B, and D,  $n = 3$ . Abs, absorbance.

a mutant with full deletion of *rsmB*, the major target of the GacS/GacA two-component system, and evaluated its virulence. The phenotype of the *rsmB* deletion mutant was the same as that of the *gacA* and *gacS* mutants (Fig. 2). None of these mutants had growth defects, as shown by the growth curves in Fig. 2D. Additionally, in the *expl* mutant that had been described as being fully avirulent (26), expression of *pehA*, Pel activity, and virulence were as low as in all of the Gac/Rsm deletion mutants (Fig. 2). These results demonstrated that both signaling pathways are required to activate virulence and are dependent on one another. Therefore, to understand the molecular basis of this observation, we decided to investigate the existence of regulatory links between the two pathways. As *Expl* is a target of *RsmA* (15), we hypothesized that the GacS/GacA system could control AHL levels. We measured AHL levels in the different Gac/Rsm pathway mutants, confirming lower production in the *gacS*, *gacA*, and *rsmB* mutants (see Fig. S1 in the supplemental material). These results provided an explanation of how the AHL quorum-sensing network requires the Gac/Rsm pathway to activate virulence. Next, we investigated whether the AHL network could also regulate the Gac/Rsm pathway, since such a link between the two pathways could explain the avirulence phenotype observed in an *expl* mutant.


#### AHLs control *RsmB*, the major target of the GacS/GacA two-component system.

A clue to a possible link between the AHL and the Gac/Rsm pathway came from a genetic screening performed in our laboratory that revealed *expl* as a potential regulator of *RsmB* (21). To explore this possibility, we analyzed *rsmB* transcription in wild-type (WT) and AHL quorum-sensing pathway mutants in the background of the




**FIG 3** Expression of *rsmB* in *P. wasabiae* is regulated by AHLs.  $P_{rsmB}::gfp$  expression was measured in WT *P. wasabiae* SCC3193 and *expl*, *explR1 explR2*, *expl explR1 explR2*, and *gacA* mutants. Complementation with AHLs was performed with 2  $\mu$ M 3-oxo-C8-HSL. a.u., arbitrary units. Error bars represent the standard deviation of the mean.  $n = 3$ .

*P. wasabiae* SCC3193 strain, both in the absence and presence of exogenous 3-oxo-C8-HSL (the major AHL produced by *P. wasabiae*). As shown in Fig. 3, WT *P. wasabiae* regulates *rsmB* transcription in a cell density-dependent manner. At early stages of growth, the transcription of *rsmB*, measured by *gfp* expression in WT cells carrying a  $P_{rsmB}::gfp$  promoter fusion, is turned off and is activated only after 6 h of growth as cells reach high density (optical density at 600 nm [OD<sub>600</sub>] of >1) (circles, Fig. 2D and 3). In contrast, in the *expl* mutant, expression was much lower than in the WT at all stages of growth and only slightly higher than in the *gacA* mutant, which is thought to be the main activator of *rsmB* (closed triangles and diamonds, respectively, Fig. 2D and 3). The transcription of *rsmB* in the *expl* mutant could be induced by exogenous addition of 3-oxo-C8-HSL to the growth medium (open triangles, Fig. 3). This AHL-dependent regulation relies on the known AHL receptors (ExpR1 and ExpR2), as shown by the high levels of *rsmB* transcription and the lack of response to 3-oxo-C8-HSL observed in the *expl explR1 explR2* triple mutant (Fig. 3). We next determined if this AHL-dependent regulation of *rsmB* could influence virulence factor production. In the *expl* mutant, Pel activity is low, even in cultures grown to a high cell density (Fig. 4). But as expected, addition of AHLs can restore induction of the production of these enzymes to WT levels, as observed by the levels of Pel activity obtained in the *expl* mutant cultured in the presence of exogenously supplied 3-oxo-C8-HSL (Fig. 4). Accordingly, Pel activity was



**FIG 4** Regulation of Pel activity by AHLs requires RsmB. Pel enzymatic assay was measured in WT *P. wasabiae* SCC3193 and *explR1 explR2*, *expl*, and *expl rsmB* mutants. Complementation with AHLs was performed with 2  $\mu$ M 3-oxo-C8-HSL. *rsmB* expression was induced with 1  $\mu$ M IPTG. Cells were collected when cultures reached an OD<sub>600</sub> of approximately 3.5. Error bars represent the standard deviation of the mean.  $n = 7$ . Abs, absorbance.



**FIG 5** Expression of *rsmB* is controlled by Expr2 in an RsmA-independent manner. Comparison of the effects of AHLs on  $P_{rsmB::gfp}$  expression in the presence (A) or absence (B) of RsmA. Complementation with AHLs was performed with  $2 \mu\text{M}$  3-oxo-C8-HSL. Cells were collected when cultures reached an  $\text{OD}_{600}$  of approximately 3.5. a.u., arbitrary units. Error bars represent the standard deviation of the mean.  $n = 7$ .

also derepressed in an *expr1 expr2* mutant. In contrast, in the absence of *rsmB*, addition of AHLs could not restore Pel activity, as demonstrated by the addition of 3-oxo-C8-HSL to the *expl rsmB* double mutant. Additionally, overexpression of *rsmB* under the control of a *lac* promoter in an *expl* mutant, in the absence of AHLs, was sufficient to fully rescue Pel activity (Fig. 4). These results demonstrate that AHLs regulate *rsmB* and that this regulation is essential for the AHL-dependent control of virulence.


**Expr2 is responsible for the RsmA-independent control of RsmB.** We next determined the different contributions of Expr1 and Expr2 to the AHL-dependent regulation of *rsmB* transcription. As shown in Fig. 5A, both Expr1 and Expr2 regulate the AHL-dependent expression of *rsmB*. Nonetheless, it is obvious that repression mediated by Expr2 is stronger than that of Expr1 because the presence of the Expr2 receptor (observed in an *expl expr1* double mutant) is sufficient to fully repress *rsmB* expression, while Expr1 is only responsible for a slight reduction in the *rsmB* levels (as observed in the *expl expr2* double mutant). These results differ from those obtained with the *rsmA::gfp* promoter fusion, which show that, in accordance with previous work (24), the transcription of *rsmA* is regulated equally by Expr1 and Expr2 (see Fig. S3). Because in other bacteria, like *Pseudomonas fluorescens* and *Escherichia coli*, RsmA/CsrA proteins were proposed to influence the expression of their cognate regulatory RNAs

through feedback mechanisms (29–31), we investigated whether the AHL-dependent regulation of *rsmB* could take place via RsmA. To test this possibility, we constructed an *rsmA* deletion in the *expl* background. As in other bacteria, the *rsmA* mutant exhibits a severe growth defect (see Fig. S2), making it difficult to compare expression with other mutants (17). To overcome this problem, we selected for a spontaneous mutant in the *expl rsmA* full-deletion background that grows similarly to the WT. This mutant, which we named *rsmA<sup>sup</sup> expl*, is an *rsmA expl* double-deletion mutant with a second-site mutation that partially rescues the growth defect of the *rsmA expl* mutant (see Fig. S2). Our results showed that expression of  $P_{rsmB}::gfp$  in the *rsmA<sup>sup</sup> expl* mutant was higher than in the *expl* mutant, indicating that RsmA might repress *rsmB* transcription. However, since RsmA is a posttranscriptional regulator, this regulation is likely to occur indirectly via a transcriptional activator of *rsmB*. Importantly, in the *rsmA<sup>sup</sup> expl* mutant, expression of  $P_{rsmB}::gfp$  still responded to AHLs (Fig. 5B), providing evidence that the AHL-dependent regulation of *rsmB* transcription still occurs in the absence of RsmA. Next, we determined if the RsmA-independent control of *rsmB* transcription is dependent on ExpR1 and/or ExpR2. We constructed two additional mutants, the *rsmA<sup>sup</sup> expl expr1* and *rsmA<sup>sup</sup> expl expr2* mutants, and analyzed their  $P_{rsmB}::gfp$  expression levels in the presence or absence of AHLs. Our results demonstrated that in the mutant where *expr2* was still present (the *rsmA<sup>sup</sup> expl expr1* mutant), *rsmB* expression still responded to the addition of 3-oxo-C8-HSLs, while in the mutant expressing only ExpR1, addition of 3-oxo-C8-HSL had no effect on *rsmB* levels (Fig. 5). These results, together with the results presented in Fig. S3, show that while in the presence of RsmA, both ExpR1 and ExpR2 can mediate the AHL-dependent regulation of *rsmB*, in the absence of RsmA, ExpR1 no longer regulates *rsmB* expression in response to AHLs, while ExpR2 maintains this regulation. Therefore, we concluded that ExpR2, but not ExpR1, is the regulator responsible for the AHL-dependent regulation of RsmB in the absence of RsmA. We identified one potential ExpR binding site (*lux* box) upstream of *rsmB* (see Fig. S5), which indicates that ExpR2 can regulate *rsmB* transcription directly, but further work is necessary to prove this interaction.

As discussed below, this level of regulation mediated by ExpR2 and not ExpR1 provides the first evidence that these regulators actually regulate different targets. Former studies reported that ExpR1 and ExpR2 have the same regulatory response to different signal inputs to repress virulence by directly activating the transcription of *rsmA* (11) (see Fig. S3), and there was no previous evidence that these regulators had other targets besides RsmA (11, 32). However, our results show that ExpR2, and not ExpR1, regulates *rsmB* transcription in an RsmA-independent manner.

**AHLs are sufficient to fully restore virulence in a *gacA* mutant.** Our results showed that AHLs regulate RsmB. This finding, together with the fact that the levels of AHLs are low in *gacS* and *gacA* mutants (see Fig. S1), prompted us to determine if exogenously supplied AHLs could restore *rsmB* transcription and virulence in GacS/GacA system mutants. Addition of 3-oxo-C8-HSL to a *gacA* or *gacS* mutant restored *rsmB* expression to levels similar to those of the WT strain (Fig. 6A). Subsequently, we corroborated that PCWDE activity could also be rescued to WT levels in a *gacA* mutant by the addition of 3-oxo-C8-HSLs (Fig. 6B). However, and in agreement with our results indicating that RsmB plays a major role in this regulation, addition of 3-oxo-C8-HSL to the *rsmB* mutant could not restore Pel activity (Fig. 6B). These results showed that while these two signaling networks of *P. wasabiae* tightly control virulence expression, exogenous addition of AHLs could bypass the requirement for activation of the GacS/GacA system.


**Cross-species activation of RsmB by AHL-producing bacteria triggers early virulence expression in *P. wasabiae*.** The induction of *rsmB* transcription and virulence in a *gacA* mutant by exogenously supplied AHLs, together with the finding that ExpR2, the receptor capable of sensing foreign AHLs, participates in this regulation, suggested that this level of regulation could be important for cross-species activation in multispecies communities. To test this possibility, we analyzed the expression of *rsmB*



**FIG 6** AHL complementation restores *rsmB* expression and Pel activity to WT levels in the absence of the GacS/GacA two-component system. (A) Effects of AHLs on  $P_{rsmB}::gfp$  expression in *P. wasabiae* SCC3193 *gacA* and *gacS* mutants compared to that in the WT. (B) Effects of AHLs on Pel activity in *P. wasabiae* SCC3193 *gacA* and *rsmB* mutants compared to that in the WT. Complementation with AHLs was performed with 2  $\mu$ M 3-oxo-C8-HSL. Cells were collected when cultures reached an OD<sub>600</sub> of approximately 3.5. a.u., arbitrary units. Error bars represent the standard deviation of the mean.  $n = 7$ . Abs, absorbance.

and *pehA* with *gfp* promoter fusions in *P. wasabiae* when cocultured with other peptolytic bacteria producing similar AHLs. We compared the density-dependent regulation of *rsmB* and *pehA* expression in a *P. wasabiae* SCC3193 reporter strain in pure culture with that in cocultures with *Pectobacterium carotovorum* Ecc71 and Ecc15, which produce 3-oxo-C6-HSL as the major AHL (12). As shown in Fig. 7, when *P. wasabiae* carrying the  $P_{rsmB}::gfp$  or  $P_{pehA}::gfp$  plasmid was diluted to a low density in the presence of a 10-fold higher density of *P. carotovorum* Ecc71 or Ecc15, neither the *rsmB* nor the *pehA* gene in *P. wasabiae* was repressed and both continued to be expressed. This response was in contrast to what happened when *P. wasabiae* was growing alone or in the presence of AHL-deficient strain Ecc71 or Ecc15. In cocultures of WT *P. wasabiae* with an *expl* mutant of *P. carotovorum* Ecc71 or Ecc15, a typical density response was observed as the expression of both *rsmB* and *pehA* was turned off at low cell density upon dilution into fresh medium. These results indicate that when grown in coculture with Ecc71 or Ecc15, *P. wasabiae* no longer interprets only its own cell density but instead, by sensing the overall concentration of AHLs present in the environment, induces virulence according to the total density of the related and unrelated AHL-producing bacteria. We next confirmed that this cross-species activation by AHL-






**FIG 7** Induction of *rsmB* and *pehA* in multispecies cultures. Shown is the cell density-dependent activation of  $P_{rsmB}$  (A) and  $P_{pehA}$  (B) in WT *P. wasabiae* (*Pw*) SCC3193 in the presence of a 10-fold excess of AHL-producing *P. carotovorum* (*Pc*) strain Ecc71 or Ecc15 and the corresponding *expl* (AHL<sup>-</sup>) mutant of *P. carotovorum*. The final densities of *P. wasabiae* are lower in the multispecies cultures than in the monocultures because the yield of *P. wasabiae* is lower in the presence of the other bacteria. This is a representative experiment from three independent experiments. a.u., arbitrary units.

producing bacteria that enabled the activation of *rsmB* and *pehA* transcription in *P. wasabiae* at low cell densities was independent of GacA. As demonstrated above (Fig. 2A and 3), when *P. wasabiae* was grown in pure culture, the expression of both *pehA* and *rsmB* was much lower in the *gacA* mutant than in the WT. However, as shown in Fig. 8, *gacA* was no longer required for induction of *rsmB* or *pehA* when *P. wasabiae* SCC3193 was grown in coculture with either of the two *P. carotovorum* strains tested because induction of *rsmB* and *pehA* was observed when the *P. wasabiae gacA* mutant, carrying the reporter plasmid, was mixed with a 10-fold higher density of *P. carotovorum* Ecc71 or Ecc15 (Fig. 8). These inductions were dependent on the AHLs produced by the two *P. carotovorum* species, as induction of the two *P. wasabiae* reporters was not observed in the cocultures with *expl* mutants of *P. carotovorum* Ecc71 and Ecc15 (Fig. 8). Importantly, *rsmB* was essential for the AHL-dependent response in multispecies cultures, as no activation was observed in the *rsmB* mutant, even in the presence of AHL-producing pectobacteria (Fig. 8B). Furthermore, consistent with the results showing that AHL-dependent regulation of RsmB requires ExpR2 but not ExpR1 (Fig. 5B), induction of *rsmB* by the AHLs produced by *P. carotovorum* Ecc71 was still observed when the *P. wasabiae expr1* mutant was cocultured with *P. carotovorum*, but not when the *P. wasabiae expr2* mutant was used (see Fig. S4).

These results demonstrated that although the GacS/GacA two-component system was essential for activation of full expression and production of PCWDEs in *P. wasabiae* monocultures, these quorum-sensing-regulated traits could be activated in *P. wasabiae* via RsmB in multispecies cultures, even in the absence of the GacS/GacA system if the other species produced related AHLs. Therefore, these results show that cross-species activation in *P. wasabiae* by other AHL-producing bacteria can override the need for the GacS/GacA system and induce virulence even before the cell density of *P. wasabiae* has reached the quorum threshold.

***P. wasabiae* virulence is higher in multispecies lesions with AHL-producing bacteria.** We finally asked whether the ability of *P. wasabiae* to respond to foreign AHLs would have an effect on the enzymatic maceration of plant tissue by *P. wasabiae* in potatoes infected with a mixture of *P. wasabiae* and *P. carotovorum* Ecc15 (at a 1:10 ratio). To assess tissue maceration caused by induction of the expression of PCWDEs produced only by *P. wasabiae*, avirulent Ecc15 *rsmB* mutant or *rsmB expl* double mutant bacteria (Fig. 9A) were used in the mixture. It was clear that production of PCWDEs by *P. wasabiae* was higher in tubers infected with mixtures of *P. wasabiae* and *P. carotovorum* Ecc15 capable of producing AHLs (WT *P. wasabiae* plus *P. carotovorum rsmB*) than in the potatoes with mixtures of *P. wasabiae* and the Ecc15 strain that does not produce AHLs (WT *P. wasabiae* plus *P. carotovorum rsmB expl*) (Fig. 9A). Moreover,




**FIG 8** Cross-species induction of *rsmB* and *pehA* in the absence of the Gac system. Activation of  $P_{rsmB}$  (A) and  $P_{pehA}$  (B) in *P. wasabiae* SCC3193 *gacA* (gray bars) and *rsmB* (striped bars) mutants in the presence of a 10-fold excess of AHL-producing bacteria and the corresponding *expl* mutants. WT *P. wasabiae* (black bars) and *gacA* and *rsmB* mutants grown in monocultures are shown as references for comparison. *Pw* stands for *P. wasabiae*, and *Pc* stands for *P. carotovorum*. Cells were collected when cultures of the *P. wasabiae* reporter strain reached approximately  $5 \times 10^7$  CFU/ml. a.u., arbitrary units. Error bars represent the standard deviation of the mean.  $n = 3$ .

potatoes infected with *P. wasabiae* monocultures (WT *P. wasabiae*) showed lower levels of maceration than potatoes infected with the same number and density of *P. wasabiae* cells in the mixture with AHL-producing strain Ecc15 (WT *P. wasabiae* plus *P. carotovorum rsmB*) (Fig. 9A). We further confirmed that, in line with our *in vitro* data (Fig. 8B), the AHL-mediated activation of *P. wasabiae* virulence observed in the multispecies infections with the Ecc15 strain that produces AHLs is independent of GacA, as the virulence of a *P. wasabiae gacA* mutant could still be induced by the AHLs produced by Ecc15 (Fig. 9B). Importantly, both *rsmB* and *expl2* were essential for the AHL-dependent response in multispecies infections, as no additional induction of virulence was observed in the *P. wasabiae rsmB* or *expl2* mutant either in the absence or in the presence of AHL-producing Ecc15 bacteria (Fig. 9B).

## DISCUSSION

In this study, we analyzed the integration of the two major signaling networks controlling virulence in *P. wasabiae*. Our results showed that the AHL synthase *Expl*, as well as all of the components of the Gac/Rsm system, is essential for activation of virulence in this bacterium. These findings provided evidence of a previously unreported link between these two signaling networks. Indeed, our studies revealed that the Gac/Rsm pathway affects the levels of AHLs, but most importantly, we discovered that AHLs control the levels of the regulatory RNA *RsmB*. This additional level of



**FIG 9** The virulence in *P. wasabiae* is higher in AHL-producing multispecies potato lesions and requires *rsmB*. Comparison of tissue maceration in potato tubers at 24 h caused by WT *P. wasabiae* (Pw) SCC3193 (A) and the *gacA*, *rsmB*, and *expR2* mutants (B) in mono-infections or in coinfections with a 10-fold excess of an AHL-producing (Ecc15 *rsmB*) or nonproducing (Ecc15 *rsmB expl*) *P. carotovorum* (Pc) strain. The maceration phenotypes of Ecc15 *rsmB* and *rsmB expl* mutants are also shown in panel A. \*\*,  $P \leq 0.01$ ; \*\*\*,  $P < 0.001$ ; ns, not significant (Mann-Whitney test;  $n \geq 5$ ). Holm-Bonferroni correction was used for multiple comparisons.

regulation provides an explanation for the avirulent phenotype of *expl*. In the absence of AHLs, *ExpR2* represses *rsmB* and activation of the GacS/GacA system is not sufficient to relieve this inhibition. These results also showed that *ExpR1* and *ExpR2* regulate different targets. Earlier studies identified *RsmA* as the only known target of *ExpR1* and *ExpR2*. We show that while *ExpR1* and *ExpR2* regulate *RsmA* expression to similar levels, only *ExpR2* is required for AHL-dependent regulation of *rsmB* transcription. Given the positions of *expR1* and *expR2* in the genome, where *expR1* is linked to *expl* and *expR2* is located elsewhere, and the fact that while *ExpR1* is specific to 3-oxo-C8-HSL, *ExpR2* also responds to AHLs produced by other species, it is thought that *ExpR1* is the original partner of *Expl* and that a later acquisition of *ExpR2* probably enabled the gain of a novel function (11). On the basis of our results, we speculate that the *RsmA*-independent control of *ExpR2* on *rsmB* expression could be the novel function, or at least one of the novel functions, that provided a potential benefit of maintaining *ExpR2*.

The discovery of the regulation of *rsmB* expression through AHLs led us to show that exogenously supplied AHLs could override the requirement of the GacS/GacA system inducing *rsmB* expression. This regulation of *rsmB* in *P. wasabiae* is crucial for the response to AHLs in this bacterium because virulence can be restored to WT levels by the addition of AHLs in all of the mutants tested, except in the *rsmB* mutant. Overall, these results demonstrate that *RsmB*, previously thought to be regulated mainly by the GacS/GacA two-component system, is a key component in the integration of the sensory information contained in the AHL and GacS/GacA signaling pathways in *P. wasabiae*.

The finding that *ExpR2*, the receptor that enables *P. wasabiae* to respond to AHLs

from other species, has an additional function in the AHL-dependent regulation of RsmB that differentiates it from ExpR1 led us to hypothesize that having RsmB under the control of the AHL system could be relevant in multispecies communities. The further evidence that addition of AHLs was sufficient to restore the induction of Pel activity in a *gacA* mutant prompted us to test whether cross-species activation by AHL-producing bacteria could override the need for the GacS/GacA system in *P. wasabiae*. We examined *rsmB* and *pehA* expression in various *P. wasabiae* reporter strains mixed with a higher cell density of two *P. carotovorum* strains (Ecc71 and Ecc15). Unlike *P. wasabiae*, which produces mainly 3-oxo-C8-HSL as the main AHL, *P. carotovorum* produces 3-oxo-C6-HSL as the major AHL (12). We observed that although *P. wasabiae* *gacA* and *rsmB* mutants had low levels of PCWDEs in monocultures, in multispecies cultures with WT *P. carotovorum*, *pehA* expression was induced in the *P. wasabiae* *gacA* mutant by the AHLs produced by *P. carotovorum* but not in the *P. wasabiae* *rsmB* mutant. Therefore, RsmB, but not GacA, was essential for the AHL-dependent cross-species activation of virulence. These findings indicate that by overriding the need to activate the GacS/GacA system in multispecies cultures, WT *P. wasabiae* is primed, independently of its own population density, to induce virulence expression even at very low cell densities in response to the AHLs produced by other species. On the contrary, under conditions where no other AHL-producing bacteria are present, the GacS/GacA two-component system, which also regulates the rate of AHL production, might be important to provide the necessary buffer to delay activation of the production of virulence factors (such as PCWDEs), ensuring that the population engages in the production of these common goods only when the quorum has been reached and thus when it is productive to do so.

We speculate that integration of these two signal transduction networks in *P. wasabiae* could have evolved to provide a fast response to changing ecological situations. Different pathogen species are often isolated from diseased tubers within the same affected crops (33) and the same potato tubers (34). Therefore, it is plausible that multispecies lesions can occur in nature. Under such a hypothetical scenario, it is conceivable that, in soil, *P. wasabiae* will encounter plants already colonized by large numbers of related AHL-producing bacteria. In such niches, *P. wasabiae*, and probably other pectobacteria, might profit by responding to AHLs produced by others and by being primed to join the community in the effort to degrade the plant cell wall barrier. It is not obvious whether cooperation or competition is more likely to take place among the different bacteria within lesions, but sensing the presence of other species might be beneficial in both scenarios. These arguments are supported by our observation that *P. wasabiae* can induce the expression of virulence by sensing AHLs produced by other bacteria present at higher concentrations in multispecies potato lesions and by the fact that potato maceration caused by induction of virulence in *P. wasabiae* was higher in the multispecies lesions than in the lesions infected with the same cell densities of *P. wasabiae* in monocultures. It has been suggested that transmission of soft rot disease is driven by insect vectors harboring important plant pathogens (35, 36), with these insects being easily attracted to decaying plant material. Therefore, bigger lesions should increase the chances of attracting insects and other organisms that can serve as transmission vectors. Moreover, transmission could be facilitated by direct contamination through spreading of rotting material to adjoining tubers. Therefore, we reason that higher tissue maceration is potentially better for bacterial transmission and thus bacterial fitness.

Interestingly, an extra layer of regulation ensures that the signaling pathways regulating virulence in *P. wasabiae* respond to quorum-sensing signals only in the presence of damaged plant tissue. This level of virulence regulation is provided by the regulator KdgR, which represses the transcription of PCWDEs in the absence of plant signals (30, 37). The presence of plant cell wall breakdown products (supplied in our *in vitro* experiments by the addition of polygalacturonic acid [PGA] to the culture medium) is required to relieve KdgR repression and activate virulence. In *P. wasabiae*, this level of regulation might be essential also to ensure that the population will not

respond to all AHL-producing bacteria. Because of the KdgR control, by responding to AHLs only in the presence of plant signals, presumably, *P. wasabiae* will only engage with microorganisms that are also participating in the degradation of the plant tissue. The multiple levels of regulation to control virulence described here show how signal transduction networks are tightly regulated to ensure that virulence is only activated in the right context and illustrate how ecological interactions can have such an important role in the regulation of virulence.

*Vibrio cholerae* and *Vibrio harveyi* are other examples of bacteria where different cell-to-cell signals are integrated into a single-output pathway (38, 39). In a recent report, Jung et al. demonstrated that in *V. cholerae*, the absence of any of the four signal receptors can be compensated for by the other three, maintaining *in vivo* colonization levels similar to those of the WT (38). Thus, in contrast to what happens in *P. wasabiae* (Fig. 1), in *V. cholerae*, one signaling system is enough to induce WT colonization levels in the infection assays used (38). Moreover, another difference stems from the fact that WT *V. cholerae* appears highly resistant to signal perturbation, as an exogenous supply of the cholera autoinducer could not induce a response as long as the other systems were present. These findings were interpreted as a strategy adopted by *V. cholerae* to be less susceptible to environmental interference, in particular, to signals from other bacteria (38). We concluded that in pectobacteria, the architecture of the cell-to-cell communication networks functions contrary to that of *V. cholerae*, as it allows input from species in the community to alter gene expression.

The present study and studies of *Vibrio* spp. exemplify how bacteria use different cell-to-cell signaling architectures to provide alternative outputs, presumably to optimize survival in their different ecological niches (40–42). We argue that our work supports the conclusion that the ecological factors are likely to be among the major factors dictating the constraints on the architecture integrating multiple quorum-sensing signal networks. Bacteria may have evolved to respond to the presence of other species, as this might be the case in *P. wasabiae*, which responds to other pectolytic bacteria, or to have robust systems that are optimized to avoid interference from neighbors, as has been proposed for *V. cholerae*. Therefore, our expectations are that analyses of the signaling networks such as the one described here will help us to understand how different species react to their neighbors and respond under different ecological conditions.

## MATERIALS AND METHODS

**Bacterial strains, plasmids, and culture conditions.** The strains, plasmids, and primers used in this study are listed in Tables S1 and S2. All of the *P. wasabiae* strains used are derived from SCC3193 (28). *P. wasabiae* and *P. carotovorum* strains were grown at 30°C with aeration in Luria-Bertani (LB) broth supplemented with 0.4% PGA (Sigma P3850). PGA is required to induce the expression of virulence. *E. coli* DH5 $\alpha$  was used for routine cloning procedures. This strain was grown at 37°C with aeration in LB broth. Antibiotics were used at the following concentrations (mg liter<sup>-1</sup>): ampicillin, 100; kanamycin, 50; spectinomycin (Spec), 50; chloramphenicol, 25; streptomycin, 100.

**Genetic and molecular techniques.** To construct the  $P_{rsmB}::gfp$  reporter (see Fig. S5), the fragment containing the  $P_{rsmB}::gfp$  fusion was amplified from pRSV206 (21) with P0528 and P0665 and cloned into pOM1 (43). To construct the  $P_{rsmA}::gfp$  reporter, the promoter region of *rsmA* was amplified with primers P0766 and P0656 and cloned into pUC18 (44). The *gfp* fragment obtained by PCR from pCMW1 (45) was cloned downstream of pUC18- $P_{rsmA}$ . Subsequently, the  $P_{rsmA}::gfp$  fragment was amplified with P0766 and P0665 and cloned into pOM1. To construct the  $P_{pehA}::gfp$  reporter, the promoter region of *pehA* from SCC3193 was amplified with P1028 and P0658 and cloned into pUC18. Subsequently, the *gfp* fragment obtained by PCR from pCMW1 was cloned downstream of pUC18- $P_{pehA}$ . Next, the  $P_{pehA}::gfp$  fragment was amplified with P1028 and P0665 and cloned into pOM1. *P. wasabiae* SCC3193 deletion mutants were constructed by chromosomal gene replacement with an antibiotic marker with the  $\lambda$ -Red recombinase system (46, 47) as previously described (21). For *rsmB::kan*, *expR1::kan*, and *expR2::kan* deletions, plasmid pKD4 (47) was used as the template to amplify the *kan* gene. The *rsmA* suppressor (*rsmA*<sup>sup</sup> *expl*; see Table S3) was obtained by selection of overgrown spontaneous mutants in LB agar plates and used as recipients for electroporation of the *expR1::kan* or *expR2::kan* fragment. Similarly to *P. wasabiae* SCC3193, *P. carotovorum* Ecc15 deletion mutants were also constructed by chromosomal gene replacement with an antibiotic marker. The plasmid used to express the arabinose-inducible  $\lambda$ -Red recombinase system (pLIPS) in Ecc15 was constructed by introducing the system into the pOM1 vector. The  $\lambda$ -Red recombinase system (composed of the *araC*, *gam*, *bet*, *exo*, and *af60A* genes, a total of 3,421 bp) was amplified by PCR from vector pKD46 with primers P1108 and P1109 and cloned into pOM1.

PCRs for cloning purposes were performed with the proofreading Herculase II polymerase (Agilent) for SCC3193 or with Bio-X-ACT (BioLine) for Ecc15. Other PCRs were performed with Dream Taq polymerase (Fermentas). Digestions were done with Fast Digest Enzymes (Fermentas), and ligations were performed with T4 DNA ligase (New England Biolabs). All cloning steps were performed with *E. coli* DH5 $\alpha$ . All mutants and constructs were confirmed by sequencing at the Instituto Gulbenkian de Ciéncia sequencing facility.

**Pectate lyase activity assay.** To analyze extracellular pectate lyase (Pel) activity, overnight cultures were diluted to a starting OD<sub>600</sub> of 0.05 in LB broth supplemented with 0.4% PGA. At the time points indicated, aliquots were collected to evaluate growth and to analyze Pel activity in the supernatants by the previously described thiobarbituric acid colorimetric method (21, 48).

***P. wasabiae* virulence assay.** Virulence was analyzed by assessing the maceration of potato tubers with the protocol adapted from reference 49. Potatoes were washed and surface sterilized by being soaked for 10 min in 10% bleach, followed by 10 min in 70% ethanol. Overnight cultures in LB broth were washed twice and diluted to an OD<sub>600</sub> of 0.05 in phosphate-buffered saline (PBS). For multispecies infections, overnight cultures of *P. wasabiae* and *P. carotovorum* were washed twice and diluted in PBS to OD<sub>600</sub>s of 0.05 and 0.5, respectively, separately or in mixtures. Thirty-microliter aliquots were then used to inoculate the previously punctured potatoes. Potato tubers were incubated at 28°C at a relative humidity above 90% for 24 or 48 h, as specified. After incubation, potatoes were sliced and macerated tissue was collected and weighed.

**Promoter expression analyses.** *P. wasabiae* strains containing the different promoter-*gfp* plasmids were grown overnight in LB broth plus PGA and Spec and inoculated into fresh medium at a starting OD<sub>600</sub> of 0.05. At the times indicated, aliquots were collected to monitor growth and assess gene expression. As previously described (21), for the analyses of gene expression, aliquots of the cultures were diluted 1:100 in PBS and expression was measured by flow cytometry (LSRFortessa; BD) and analyzed with Flowing Software v 2.5.1. A minimum of 5,000 green fluorescent protein (GFP)-positive single cells were acquired per sample. Expression of the promoter-*gfp* fusions is reported as the median GFP expression of GFP-positive single cells in arbitrary units.

**Cross-species expression experiments.** The interspecies signaling effects on the activation of *rsmB* expression were tested similarly to those of single species. Briefly, WT and *gacA* and *rsmB* mutant *P. wasabiae* strains harboring either plasmid *P*<sub>*rsmB*</sub>::*gfp* or *P*<sub>*pehA*</sub>::*gfp* and WT and *expl* mutant *P. carotovorum* Ecc15 and Ecc71 strains were grown overnight in LB broth plus 0.4% PGA. *P. wasabiae* strains (reporters) and *P. carotovorum* strains were diluted in fresh medium to OD<sub>600</sub>s of 0.05 and 0.5, respectively (1:10). The diluted cultures were maintained separately for 2 h to allow the reporter strain to reduce the AHL concentration and mimic the low cell density state. After 2 h, the *P. wasabiae* cells were centrifuged and resuspended in the *P. carotovorum* high cell density cultures. Aliquots of the multispecies cultures were diluted 1:100 in PBS, and expression was measured by flow cytometry as explained above. The two populations were clearly distinguishable by the presence or absence of GFP fluorescence. *P. wasabiae* growth was quantified by determining the number of CFU per milliliter of LB agar supplemented with Spec.

**Statistical analysis.** Data were analyzed with GraphPad Prism 6 software and R program v 3.0.2. The Mann-Whitney test was performed to determine statistical significance, and *P* values were adjusted with the Holm-Bonferroni correction for multiple comparisons. An adjusted *P* value of <0.05 was used as the cutoff for statistical significance.

## SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/mBio.00398-17>.

**TEXT S1**, PDF file, 0.4 MB.

**FIG S1**, PDF file, 0.3 MB.

**FIG S2**, PDF file, 0.2 MB.

**FIG S3**, PDF file, 0.3 MB.

**FIG S4**, PDF file, 0.2 MB.

**FIG S5**, PDF file, 0.8 MB.

**TABLE S1**, PDF file, 0.6 MB.

**TABLE S2**, PDF file, 0.4 MB.

**TABLE S3**, PDF file, 0.5 MB.

## ACKNOWLEDGMENTS

We thank Joana Amaro for technical assistance and Roberto Balbontín, Brian Hammer, and Joao B. Xavier for suggestions and helpful comments on the manuscript. We are very grateful to Andres Mãe (Institute of Molecular and Cell Biology) for sharing protocols and strains and Tapio Palva (University of Helsinki) for providing strains used in this study.

This work was supported by the Fundação Para a Ciéncia e Tecnologia (FCT) (PTDC/BIA-BCM/101585/2008) and the Howard Hughes Medical Institute (international

early career scientist grant HHMI 55007436). R.V. was supported by award SFRH/BD/33570/2008; F.V. was supported by SRFH/BD/113986/2015 within the SCOPE of the Ph.D. program Molecular Biosciences PD/00133/2012, both funded by FCT; and P.N. was supported by Marie Curie Intra-European grant PIEF-GA-2011-301365.

## REFERENCES

- Rutherford ST, Bassler BL. 2012. Bacterial quorum sensing: its role in virulence and possibilities for its control. *Cold Spring Harb Perspect Med* 2:a012427. <https://doi.org/10.1101/cshperspect.a012427>.
- Papenfort K, Bassler BL. 2016. Quorum sensing signal-response systems in Gram-negative bacteria. *Nat Rev Microbiol* 14:576–588. <https://doi.org/10.1038/nrmicro.2016.89>.
- Bhatt S, Edwards AN, Nguyen HTT, Merlin D, Romeo T, Kalman D. 2009. The RNA binding protein CsrA is a pleiotropic regulator of the locus of enterocyte effacement pathogenicity island of enteropathogenic *Escherichia coli*. *Infect Immun* 77:3552–3568. <https://doi.org/10.1128/IAI.00418-09>.
- Altier C, Suyemoto M, Lawhon SD. 2000. Regulation of *Salmonella enterica* serovar Typhimurium invasion genes by *csrA*. *Infect Immun* 68:6790–6797. <https://doi.org/10.1128/IAI.68.12.6790-6797.2000>.
- Johnston C, Pegues DA, Hueck CJ, Lee A, Miller SI. 1996. Transcriptional activation of *Salmonella typhimurium* invasion genes by a member of the phosphorylated response-regulator superfamily. *Mol Microbiol* 22:715–727. <https://doi.org/10.1046/j.1365-2958.1996.d01-1719.x>.
- Reimann C, Beyeler M, Latifi A, Winteler H, Foglino M, Lazdunski A, Haas D. 1997. The global activator GacA of *Pseudomonas aeruginosa* PAO positively controls the production of the autoinducer N-butyryl homoserine lactone and the formation of the virulence factors pyocyanin, cyanide, and lipase. *Mol Microbiol* 24:309–319. <https://doi.org/10.1046/j.1365-2958.1997.3291701.x>.
- Pessi G, Williams F, Hindle Z, Heurlier K, Holden MTG, Cámara M, Haas D, Williams P. 2001. The global posttranscriptional regulator RsmA modulates production of virulence determinants and N-acylhomoserine lactones in *Pseudomonas aeruginosa*. *J Bacteriol* 183:6676–6683. <https://doi.org/10.1128/JB.183.22.6676-6683.2001>.
- Vakulskas CA, Potts AH, Babitzke P, Ahmer BMM, Romeo T. 2015. Regulation of bacterial virulence by Csr (Rsm) systems. *Microbiol Mol Biol Rev* 79:193–224. <https://doi.org/10.1128/MMBR.00052-14>.
- Brader G, Sjöblom S, Hyytiäinen H, Sims-Huopaniemi K, Palva ET. 2005. Altering substrate chain length specificity of an acylhomoserine lactone synthase in bacterial communication. *J Biol Chem* 280:10403–10409. <https://doi.org/10.1074/jbc.M408603200>.
- Cui Y, Chatterjee A, Hasegawa H, Chatterjee AK. 2006. *Erwinia carotovora* subspecies produce duplicate variants of ExpR, LuxR homologs that activate *rsmA* transcription but differ in their interactions with N-acylhomoserine lactone signals. *J Bacteriol* 188:4715–4726. <https://doi.org/10.1128/JB.00351-06>.
- Sjöblom S, Brader G, Koch G, Palva ET. 2006. Cooperation of two distinct ExpR regulators controls quorum sensing specificity and virulence in the plant pathogen *Erwinia carotovora*. *Mol Microbiol* 60:1474–1489. <https://doi.org/10.1111/j.1365-2958.2006.05210.x>.
- Chatterjee A, Cui Y, Hasegawa H, Leigh N, Dixit V, Chatterjee AK. 2005. Comparative analysis of two classes of quorum-sensing signaling systems that control production of extracellular proteins and secondary metabolites in *Erwinia carotovora* subspecies. *J Bacteriol* 187:8026–8038. <https://doi.org/10.1128/JB.187.23.8026-8038.2005>.
- Murata H, Chatterjee A, Liu Y, Chatterjee AK. 1994. Regulation of the production of extracellular pectinase, cellulase, and protease in the soft rot bacterium *Erwinia carotovora* subsp. *carotovora*: evidence that *aepH* of *E. carotovora* subsp. *carotovora* 71 activates gene expression in *E. carotovora* subsp. *carotovora*, *E. carotovora* subsp. *atroseptica*, and *Escherichia coli*. *Appl Environ Microbiol* 60:3150–3159.
- Hyytiäinen H, Montesano M, Palva ET. 2001. Global regulators ExpA (GacA) and KdgR modulate extracellular enzyme gene expression through the RsmA-rsmB system in *Erwinia carotovora* subsp. *carotovora*. *Mol Plant Microbe Interact* 14:931–938. <https://doi.org/10.1094/MPMI.2001.14.8.931>.
- Cui Y, Chatterjee A, Liu Y, Dumenyo CK, Chatterjee AK. 1995. Identification of a global repressor gene, *rsmA*, of *Erwinia carotovora* subsp. *carotovora* that controls extracellular enzymes, N-(3-oxohexanoyl)-L-homoserine lactone, and pathogenicity in soft-rotting *Erwinia* spp. *J Bacteriol* 177:5108–5115. <https://doi.org/10.1128/jb.177.17.5108-5115.1995>.
- Cui Y, Chatterjee A, Chatterjee AK. 2001. Effects of the two-component system comprising GacA and GacS of *Erwinia carotovora* subsp. *carotovora* on the production of global regulatory rsmB RNA, extracellular enzymes, and harpinEcc. *Mol Plant Microbe Interact* 14:516–526. <https://doi.org/10.1094/MPMI.2001.14.4.516>.
- Lapouge K, Schubert M, Allain FH-t, Haas D. 2008. Gac/Rsm signal transduction pathway of gamma-proteobacteria: from RNA recognition to regulation of social behaviour. *Mol Microbiol* 67:241–253. <https://doi.org/10.1111/j.1365-2958.2007.06042.x>.
- Takeuchi K, Kiefer P, Reimann C, Keel C, Dubuis C, Rolli J, Vorholt JA, Haas D. 2009. Small RNA-dependent expression of secondary metabolism is controlled by Krebs cycle function in *Pseudomonas fluorescens*. *J Biol Chem* 284:34976–34985. <https://doi.org/10.1074/jbc.M109.052571>.
- Chavez RG, Alvarez AF, Romeo T, Georgellis D. 2010. The physiological stimulus for the BarA sensor kinase. *J Bacteriol* 192:2009–2012. <https://doi.org/10.1128/JB.01685-09>.
- Lawhon SD, Maurer R, Suyemoto M, Altier C. 2002. Intestinal short-chain fatty acids alter *Salmonella typhimurium* invasion gene expression and virulence through BarA/SirA. *Mol Microbiol* 46:1451–1464. <https://doi.org/10.1046/j.1365-2958.2002.03268.x>.
- Valente RS, Xavier KB. 2015. The Trk potassium transporter is required for RsmB-mediated activation of virulence in the phytopathogen *Pectobacterium wasabiae*. *J Bacteriol* 198:248–255. <https://doi.org/10.1128/JB.00569-15>.
- Broder UN, Jaeger T, Jenal U. 2016. LadS is a calcium-responsive kinase that induces acute-to-chronic virulence switch in *Pseudomonas aeruginosa*. *Nat Microbiol* 2:16184. <https://doi.org/10.1038/nmicrobiol.2016.184>.
- Dubuis C, Haas D. 2007. Cross-species GacA-controlled induction of antibiosis in pseudomonads. *Appl Environ Microbiol* 73:650–654. <https://doi.org/10.1128/AEM.01681-06>.
- Cui Y, Chatterjee A, Hasegawa H, Dixit V, Leigh N, Chatterjee AK. 2005. ExpR, a LuxR homolog of *Erwinia carotovora* subsp. *carotovora*, activates transcription of *rsmA*, which specifies a global regulatory RNA-binding protein. *J Bacteriol* 187:4792–4803. <https://doi.org/10.1128/JB.187.14.4792-4803.2005>.
- Köiv V, Mäe A. 2001. Quorum sensing controls the synthesis of virulence factors by modulating *rsmA* gene expression in *Erwinia carotovora* subsp. *carotovora*. *Mol Genet Genomics* 265:287–292. <https://doi.org/10.1007/s004380000413>.
- Pirhonen M. 1991. Identification of pathogenicity determinants of *Erwinia carotovora* subsp. *carotovora* by transposon mutagenesis. *Mol Plant Microbe Interact* 4:276. <https://doi.org/10.1094/MPMI.4.4.276>.
- Eriksson AR, Andersson RA, Pirhonen M, Palva ET. 1998. Two-component regulators involved in the global control of virulence in *Erwinia carotovora* subsp. *carotovora*. *Mol Plant Microbe Interact* 11:743–752. <https://doi.org/10.1094/MPMI.1998.11.8.743>.
- Pirhonen M, Heino P, Helander I, Harju P, Palva ET. 1988. Bacteriophage T4 resistant mutants of the plant pathogen *Erwinia carotovora*. *Microb Pathog* 4:359–367. [https://doi.org/10.1016/0882-4010\(88\)90063-0](https://doi.org/10.1016/0882-4010(88)90063-0).
- Reimann C, Valverde C, Kay E, Haas D. 2005. Posttranscriptional repression of GacS/GacA-controlled genes by the RNA-binding protein RsmE acting together with RsmA in the biocontrol strain *Pseudomonas fluorescens* CHA0. *J Bacteriol* 187:276–285. <https://doi.org/10.1128/JB.187.1.276-285.2005>.
- Bejerano-Sagie M, Xavier KB. 2007. The role of small RNAs in quorum sensing. *Curr Opin Microbiol* 10:189–198. <https://doi.org/10.1016/j.mib.2007.03.009>.
- Yakhnin H, Yakhnin AV, Baker CS, Sineva E, Berezin I, Romeo T, Babitzke P. 2011. Complex regulation of the global regulatory gene *csrA*: CsrA-mediated translational repression, transcription from five promoters by  $E\sigma^{70}$  and  $E\sigma(S)$ , and indirect transcriptional activation by CsrA and  $E\sigma(S)$ , and indirect transcriptional activation by CsrA. *Mol Microbiol* 81:689–704. <https://doi.org/10.1111/j.1365-2958.2011.07723.x>.
- Barnard AML, Salmond GPC. 2007. Quorum sensing in *Erwinia* species.

- Anal Bioanal Chem 387:415–423. <https://doi.org/10.1007/s00216-006-0701-1>.
33. Dees MW, Lebecka R, Perminow JIS, Czajkowski R, Grupa A, Motyka A, Zoledowska S, Śliwka J, Lojkowska E, Brurberg MB. 2017. Characterization of *Dickeya* and *Pectobacterium* strains obtained from diseased potato plants in different climatic conditions of Norway and Poland. *Eur J Plant Pathol* 148:1–13. <https://doi.org/10.1007/s10658-016-1140-2>.
  34. Ranjan RK, Singh D, Baranwal VK. 2016. Simultaneous detection of brown rot- and soft rot-causing bacterial pathogens from potato tubers through multiplex PCR. *Curr Microbiol* 73:652–659. <https://doi.org/10.1007/s00284-016-1110-0>.
  35. Kloepper JW, Brewer JW, Harrison MD. 1981. Insect transmission of *Erwinia carotovora* var. *carotovora* and *Erwinia carotovora* var. *atroseptica* to potato plants in the field. *Am Potato J* 58:165–175. <https://doi.org/10.1007/BF02854416>.
  36. Molina JJ, Harrison MD, Brewer JW. 1974. Transmission of *Erwinia carotovora* var. *atroseptica* by *Drosophila melanogaster* Meig. I. Acquisition and transmission of the bacterium. *Am Potato J* 51:245–250. <https://doi.org/10.1007/BF02851435>.
  37. Newton JA, Fray RG. 2004. Integration of environmental and host-derived signals with quorum sensing during plant-microbe interactions. *Cell Microbiol* 6:213–224. <https://doi.org/10.1111/j.1462-5822.2004.00362.x>.
  38. Jung SA, Chapman CA, Ng WL. 2015. Quadruple quorum-sensing inputs control *Vibrio cholerae* virulence and maintain system robustness. *PLoS Pathog* 11:e1004837. <https://doi.org/10.1371/journal.ppat.1004837>.
  39. Mehta P, Goyal S, Long T, Bassler BL, Wingreen NS. 2009. Information processing and signal integration in bacterial quorum sensing. *Mol Syst Biol* 5:325. <https://doi.org/10.1038/msb.2009.79>.
  40. Cornforth DM, Papat R, McNally L, Gurney J, Scott-Phillips TC, Ivans A, Diggle SP, Brown SP. 2014. Combinatorial quorum sensing allows bacteria to resolve their social and physical environment. *Proc Natl Acad Sci U S A* 111:4280–4284. <https://doi.org/10.1073/pnas.1319175111>.
  41. Drees B, Reiger M, Jung K, Bischofs IB. 2014. A modular view of the diversity of cell-density-encoding schemes in bacterial quorum-sensing systems. *Biophys J* 107:266–277. <https://doi.org/10.1016/j.bpj.2014.05.031>.
  42. Even-Tov E, Bendori SO, Valastyan J, Ke X, Pollak S, Bareia T, Ben-Zion I, Bassler BL, Eldar A. 2016. Social evolution selects for redundancy in bacterial quorum sensing. *PLOS Biol* 14:e1002386. <https://doi.org/10.1371/journal.pbio.1002386>.
  43. Espéli O, Moulin L, Boccard F. 2001. Transcription attenuation associated with bacterial repetitive extragenic BIME elements. *J Mol Biol* 314:375–386. <https://doi.org/10.1006/jmbi.2001.5150>.
  44. Yanisch-Perron C, Vieira J, Messing J. 1985. Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* 33:103–119. [https://doi.org/10.1016/0378-1119\(85\)90120-9](https://doi.org/10.1016/0378-1119(85)90120-9).
  45. Waters CM, Bassler BL. 2006. The *Vibrio harveyi* quorum-sensing system uses shared regulatory components to discriminate between multiple autoinducers. *Genes Dev* 20:2754–2767. <https://doi.org/10.1101/gad.1466506>.
  46. Murphy KC. 1998. Use of bacteriophage lambda recombination functions to promote gene replacement in *Escherichia coli*. *J Bacteriol* 180:2063–2071.
  47. Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci U S A* 97:6640–6645. <https://doi.org/10.1073/pnas.120163297>.
  48. Sherwood RT. 1966. Pectin lyase and polygalacturonase production by *Rhizoctonia solani* and other fungi. *Phytopathology* 279–286.
  49. McMillan GP, Hedley D, Fyffe L, Pérombelon MCM. 1993. Potato resistance to soft-rot erwinias is related to cell wall pectin esterification. *Physiol Mol Plant Pathol* 42:279–289. <https://doi.org/10.1006/pmpp.1993.1026>.