



# Corrigendum: Lipidomic Analysis of TRPC1 Ca<sup>2+</sup>-Permeable Channel-Knock Out Mouse Demonstrates a Vital Role in Placental Tissue Sphingolipid and Triacylglycerol Homeostasis Under Maternal High-Fat Diet

Michael R. Bukowski<sup>1\*</sup>, Brij B. Singh<sup>2</sup>, James N. Roemmich<sup>1</sup> and Kate J. Claycombe-Larson<sup>1\*</sup>

<sup>1</sup> USDA-ARS Grand Forks Human Nutrition Research Center, Grand Forks, ND, United States, <sup>2</sup> School of Dentistry, UT Health Science Center San Antonio, San Antonio, TX, United States

## OPEN ACCESS

### Approved by:

Frontiers Editorial Office,  
Frontiers Media SA, Switzerland

### \*Correspondence:

Michael R. Bukowski  
michael.bukowski@usda.gov  
Kate J. Claycombe-Larson  
kate.larson@usda.gov

### Specialty section:

This article was submitted to  
Pediatric Endocrinology,  
a section of the journal  
Frontiers in Endocrinology

Received: 20 May 2022

Accepted: 23 May 2022

Published: 28 June 2022

### Citation:

Bukowski MR, Singh BB,  
Roemmich JN and Claycombe-  
Larson KJ (2022) Corrigendum:  
Lipidomic Analysis of TRPC1 Ca<sup>2+</sup>-  
Permeable Channel-Knock Out  
Mouse Demonstrates a Vital Role in  
Placental Tissue Sphingolipid and  
Triacylglycerol Homeostasis Under  
Maternal High-Fat Diet.  
Front. Endocrinol. 13:949419.  
doi: 10.3389/fendo.2022.949419

**Keywords:** infusion lipidomics, placental lipidome, TRPC1, sphingolipid metabolism, triacylglycerol

## A Corrigendum on

### Lipidomic Analysis of TRPC1 Ca<sup>2+</sup>-Permeable Channel-Knock Out Mouse Demonstrates a Vital Role in Placental Tissue Sphingolipid and Triacylglycerol Homeostasis Under Maternal High-Fat Diet

By Bukowski MR, Singh BB, Roemmich JN and Claycombe-Larson KJ (2022) *Front. Endocrinol.* 13:854269. doi: 10.3389/fendo.2022.854269

The original article contained an error. In the first sentence of the section on *Animal Protocol Design and Approval*, the control mouse strain was incorrectly identified as C57BL/6; the actual mouse strain used was B6129SF2/J. The sentence “Two-month-old female C57BL/6 mice (Envigo, Indianapolis, IN) were fed diets containing either 16% (normal-fat, NF) or 45% fat (high-fat, HF) for 12 weeks (**Table S1**).” should have read “Two-month-old female B6129SF2/J mice (Envigo, Indianapolis, IN) were fed diets containing either 16% (normal-fat, NF) or 45% fat (high-fat, HF) for 12 weeks (**Table S1**).”

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

**Publisher’s Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Bukowski, Singh, Roemmich and Claycombe-Larson. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.