

Nicotinic acetylcholine receptors and cancer (Review)

NINGNING DANG, XIANGUANG MENG and HAIYAN SONG

Department of Dermatology, Jinan Central Hospital Affiliated to Shandong University, Jinan, Shandong 250013, P.R. China

Received November 11, 2015; Accepted March 1, 2016

DOI: 10.3892/br.2016.625

Abstract. Nicotine, the primary addictive constituent of cigarettes, is believed to contribute to cancer promotion and progression through the activation of nicotinic acetylcholine receptors (nAChRs), which are membrane ligand-gated cation channels. nAChRs activation can be triggered by the neurotransmitter Ach, or certain other biological compounds, such as nicotine. In recent years, genome-wide association studies have indicated that allelic variation in the $\alpha 5$ - $\alpha 3$ - $\beta 4$ nAChR cluster on chromosome 15q24-15q25.1 is associated with lung cancer risk. The role of nAChRs in other types of cancer has also been reported. The present review highlights the role of nAChRs in types of human cancer.

Contents

1. Introduction
2. Lung cancer
3. Other cancers
4. Conclusion

1. Introduction

Acetylcholine receptors (AChRs), similar to a number of other ligand-activated neurotransmitter receptors in neuronal system, consist of 2 subtypes: Metabotropic muscarinic receptors and ionotropic nicotinic receptors (1). These 2 subtypes can be activated by the endogenous neurotransmitter ACh (1). The first subtype, the metabotropic muscarinic receptors, is G protein-coupled seven-transmembrane proteins, which directly alter the cellular homeostasis of phospholipase C, inositol-3-phosphate, cyclic adenosine monophosphate and free calcium. Activation of metabotropic AChRs is relatively slow (from msec to sec). The other AChR subtype

is the fast ionotropic cationic nicotinic AChRs (nAChRs). These ion channels are sensitive to activation by acetylcholine and nicotine, and their activation is extremely fast (msec to sub-msec) (1). nAChRs consist of 5 subunits, comprising 10 α subunits (from $\alpha 1$ to $\alpha 10$), 4 β subunits (from $\beta 1$ to $\beta 4$), 1 δ , and 1 ϵ or γ subunit. The receptors can form homo- or hetero-pentamers, and enclose a central ion channel. The nAChR subunit composition also has a role in regulating the function of nAChRs (1-3). All the nAChRs show permeability to various cations, such as Na^+ , Ca^{2+} and K^+ , with $\alpha 7$ nAChR showing the highest permeability for Ca^{2+} . As well as channel activity, a number of other intracellular events and various downstream signaling pathways can also be regulated by nAChRs, such as Ca^{2+} /calmodulin, mitogen-activated protein kinase (MAPK), protein kinase C and vascular endothelial growth factor (VEGF) (4-6). Different nAChR subtypes also have roles in control distinct physiological processes, such as maintenance, metabolic tone and control of carcinogenesis.

nAChR genes are expressed not only in neuronal systems, but also in numerous non-neuronal tissues cells such as skin, pancreas and lung, suggesting that nAChRs may have roles in other biological processes in addition to synaptic transmission (7). In fact, nAChRs have been found not only to perform their classic function at neuromuscular junctions, but also to function in the regulation of cellular processes such as cell proliferation (7) and cell death (1,3,8). Decreased survival rates in cancer are believed to be affected by the nAChR pathway activation, influencing tumor cell proliferation, apoptosis (3,8), proinvasive and angiogenic phenotypes (9) and epithelial-mesenchymal transition (EMT). This review summarizes the current evidence of the role of nAChRs in types of human cancer.

2. Lung cancer

Tobacco smoking is known to be the major cause of lung cancer worldwide (10). Numerous large genome-wide association studies have found an association between single-nucleotide polymorphism (SNP) variation at 15q24-15q25.1 and susceptibility to lung cancer (11-18). There are three nicotinic acetylcholine receptor (*CHRNA*) genes at this locus. Studies from the International Agency for Research on Cancer (Lyon, France), the MD Anderson Cancer (Houston, TX, USA), and deCODE Genetics (Reykjavik, Iceland) have shown that there is a nearly 14% increase in lung cancer susceptibility associated with nAChR cluster variations, regardless of

Correspondence to: Dr Ningning Dang, Department of Dermatology, Jinan Central Hospital Affiliated to Shandong University, 105 Jiefang Road, Jinan, Shandong 250013, P.R. China
E-mail: ningningdangxj@sina.com

Key words: nicotinic acetylcholine receptor, cancer, tobacco

smoking status (12). In addition, a non-synonymous variant, rs16969968, in *CHRNA5*, which results in an amino acid substitution (D398N) in the second intracellular loop of the protein, can increase lung cancer risk >30% (12). In the *CHRNA3/CHRNA4* intergenic region, the SNP rs8023462 can interfere with *CHRNA3/CHRNA4* gene expression by interacting with GATA transcription factors (19). Overexpression of *CHRNA3* by ectopic expression was shown to induce apoptosis (20). Knockdown of *CHRNA3* by short hairpin RNA (shRNA) in lung cancer cells abolished the response of the cells to apoptosis-inducing agents, suggesting the importance of this gene in human cancer (20). *CHRNA3* knockdown may result in overexpression of *CHRNA7* and *CHRNA5* (20), while abnormal *CHRNA4* methylation was found to be insufficient to induce significant silencing of the gene, and could not downregulate gene expression. One study in 28 squamous cell carcinomas (SCCs) of the lung showed increased levels of *CHRNA5* and *CHRNA3* transcripts, together with higher ACh levels, which is associated with increased levels of choline acetyltransferase transcripts and decreased levels of cholinesterase transcripts. Incubation of the H520 SCC cell line with nicotine increased ACh secretion, expression of nAChRs ($\alpha 7$ and $\beta 4$), and activity of those expressed nAChRs. Darifenacin, an M3 mAChR-selective antagonist, was able to abolish the proliferative effects of nicotine and ACh on lung cancer growth by downregulating the activation of MAPK (21). Human non-small cell lung cancer (NSCLC) tissues and cancer cells were shown to express $\alpha 7$ -nAChR (22). Nicotine can cause Ca^{2+} influx into lung cancer cells by binding to $\alpha 7$ -nAChR and triggering membrane depolarization, which activates voltage-gated Ca^{2+} channels and subsequently activates the MAPK pathway, which may result in increased expression of the B-cell lymphoma-2 protein and downregulation of apoptosis (23). Subsequently, the MAPK pathway activates the transcription factor nuclear factor- κB (NF- κB) and promotes cancer cell proliferation. Expression of $\alpha 7$ -nAChR is also associated with activation of the Rb-Raf-1/phospho-extracellular signal-regulated kinase (ERK)/phospho-p90RSK pathway. In certain cases, $\alpha 7$ -nAChR antagonists were able to inhibit proliferation of human NSCLC cells via mitochondria-associated apoptosis (22). Expression of the $\beta 2$ and $\beta 4$ nAChR subunits was also been found in all adenocarcinoma (AC) and SCC tissue samples; in ACs, all $\alpha 5$, $\alpha 7$, $\beta 2$ and $\beta 4$ nAChR subunits were detected in glandular structures, while in SCCs, the $\alpha 5$, $\beta 2$ and $\beta 4$ subunits were mainly expressed in tumor cells at invasive fronts (24). $\alpha 7$ nAChR inhibited cell proliferation in well-differentiated NSCLC cells, but promoted cell proliferation by mediating the pro-proliferative activity of nicotine in poorly differentiated NSCLC cells (24). The $\alpha 7$ nAChR and heteromeric $\alpha 5^* \beta 2^* \beta 4^*$ nAChRs are involved in tumor progression by stimulating invasion and regulating proliferation (24). Li *et al.* (25) reported that nicotine can promote tumor growth and cause resistance to erlotinib in the PC9 xenograft model by activating the $\alpha 1$ nAChR and epidermal growth factor (EGFR) pathways. In A549 cells, use of shRNA or the selective antagonist methoctramine to block M2 mAChR (M2R) signaling resulted in inhibition of tumor cell proliferation *in vitro* and *in vivo*. Blocking of M2R signaling decreased MAPK and AKT phosphorylation.

Further studies indicated that M2R signaling also promotes EMT *in vitro* and *in vivo* (26). Chernyavsky *et al.* (27) reported that cell membrane (cm)-nAChRs could promote growth of lung cancer cells, and could also promote growth factor signaling [$\alpha 7$ cm-nAChR synergizes with EGF, $\alpha 3$ synergizes with VEGF, $\alpha 4$ synergizes with insulin-like growth factor I (IGF-I) and VEGF, and $\alpha 9$ synergizes with EGF, IGF-I and VEGF]; and simultaneous activation of cm- and mitochondrial-nAChRs produces a combination of growth-promoting and anti-apoptotic signals. Activation of M3R by ACh was also found to promote cell proliferation, invasion and migration of lung cancer cells by activating the EGFR/phosphoinositide-3-kinase (PI3K)/AKT pathway and interleukin-8 (28). Recently, it was reported that cyclooxygenase-2-derived prostaglandin E2 increased $\alpha 7$ nAChR expression in NSCLC cells, and this effect was regulated by Jun kinase, PI3K, and protein kinase A signals (29). This finding shows a novel link between prostanoids and cholinergic signaling (29).

3. Other cancers

Tobacco smoking is also a risk factor for pancreatic cancer. Studies have shown that nAChR activation can enhance pancreatic tumor-promoting events, and that multiple pathways are involved (30). Nicotine treatment causes a significant reduction in expression of GATA6 and Mist1, both of which are regulators of acinar cell differentiation. Nicotine treatment can not only induce de-differentiation of acinar cells, but can also increase pancreatic progenitor cell activity, by increasing expression of the stem cell genes *Sox9* and *ALDH* and enhancing cancer stem cell populations. All the effects of nicotine occur through activation of $\alpha 7$ nAChR, followed by activation of the AKT and MAPK signaling pathways (30). As in the nervous system, nAChRs have roles in regulating the synthesis and release of catecholamines in normal pancreatic ductal cells and in pancreatic cancer cells. Nicotine was shown to induce catecholamine secretion, which was dependent on the $\alpha 3$, $\alpha 5$ and $\alpha 7$ nAChR subunits (31-33). Additional studies also found that in pancreatic cancer cells, nicotine activated ERK1/2, a well-known proliferation and survival-related signaling pathway (34-36).

Nicotine also inhibits γ -aminobutyric acid (GABA) synthesis and secretion in normal pancreatic cells and cancer cells in a time-dependent manner. GABA is able to inhibit pancreatic tumor growth *in vivo* and *in vitro*, which can be facilitated through the $\alpha 4$ nAChR subunit (31,33,37). In addition to its proliferative effects, nicotine promotes migration and invasion of pancreatic cancer cells by upregulating the mucin MUC4 in the CD18/HPAF pancreatic cancer cell line (38). The nicotine-induced increase in MUC4 expression is dose-dependent, acting through $\alpha 7$ nAChR activation and subsequently activation of Janus kinase 2, signal transducer and activator of transcription 3 (STAT3), and ERK1/2 (38). Nicotine treatment also enhances the association of the MUC4 promoter with the transcription factors E2F1 and STAT1 (39).

Through induction of osteopontin (OPN) synthesis and secretion, nicotine can also promote cell proliferation and metastasis in pancreatic cancer cell lines. Studies on pancreatic cancer cell lines showed that treatment with nicotine

resulted in increased proliferation and concomitant OPN gene promoter activation, mRNA expression and protein secretion (40). This effect occurred via nAChR receptor activation, with subsequent phosphorylation of ERK1/2 (40). It has also been found that nicotine-mediated migration, invasion and metastasis of pancreatic cancer cells is dependent on OPN, through inducing matrix metalloproteinase-9 and VEGF (41).

In gastric cancer, nicotine and nicotine-derived nitrosamine ketone (NNK) were found to enhance AGS cell proliferation significantly through $\alpha 7$ nAChR and β -adrenergic receptors. Nicotine promotes cell proliferation, which is dependent on Erk1/2 activation, while NNK-induced cell growth is dependent on p38 MAPK. In addition, nicotine and NNK mediates cyclooxygenase-2 induction to modulate cell cycle regulatory proteins, and contributes to cancer development. A selective cyclooxygenase-2 inhibitor, SC-236, was shown to block nicotine/NNK-induced cell proliferation (42). An *in vitro* study on the AGS gastric cancer cell line reported that when $\alpha 7$ nAChR was knocked down by siRNA, the cells were more resistant to 5-fluorouracil treatment (43).

ACh receptors are also expressed in human mesothelioma and normal mesothelial cells. Nicotine stimulation prompts cell growth through activation of nicotinic cholinergic receptors. Activation of $\alpha 7$ nAChR can cause Ca^{2+} influx into cells, which subsequently activates voltage-gated Ca^{2+} channels, which in turn activate the MAPK pathway. Subsequently, MAPK promotes cancer cell proliferation through activating transcription factor NF- κ B (4).

In human mammary epithelial cells, non-malignant MCF10A and malignant MCF7 breast cells were shown to express the $\alpha 3$, $\alpha 5$, $\alpha 7$, $\alpha 9$, $\alpha 10$, $\beta 1$, $\beta 2$, γ , δ and ϵ nAChR subunits and the M1, M3, M4 and M5 muscarinic receptor subtypes. The malignancy was associated with expression of the $\alpha 1$, $\alpha 4$ and $\beta 4$ nAChR subunits and the M2 subtype. Malignant transformation of basal epithelial cells was also associated with overexpression of $\alpha 7$ - and $\alpha 9$ -made nAChRs. NNK upregulated ERK1/2 phosphorylation, stimulated expression of the gene encoding the tumor promoter hepatocyte growth factor, downregulated expression of the tumor suppressor gene *CDKN2A* and induced tumorigenic transformation of MCF10A cells (44).

4. Conclusion

Taken together, the obtained results provide new insight into the molecular mechanisms of nAChR-mediated oncogenic effects on types of human cancer. It is clear that nAChRs has significant roles in the pathogenesis of cancers. Further analyses of nAChRs may contribute to clinical applications in the future. More *in vivo* studies are required to confirm their oncogenic effects.

References

- Albuquerque EX, Pereira EF, Alkondon M and Rogers SW: Mammalian nicotinic acetylcholine receptors: From structure to function. *Physiol Rev* 89: 73-120, 2009.
- Niu XM and Lu S: Acetylcholine receptor pathway in lung cancer: New twists to an old story. *World J Clin Oncol* 5: 667-676, 2014.
- Schuller HM: Is cancer triggered by altered signalling of nicotinic acetylcholine receptors? *Nat Rev Cancer* 9: 195-205, 2009.
- Trombino S, Cesario A, Margaritora S, Granone P, Motta G, Falugi C and Russo P: Alpha7-nicotinic acetylcholine receptors affect growth regulation of human mesothelioma cells: Role of mitogen-activated protein kinase pathway. *Cancer Res* 64: 135-145, 2004.
- Arredondo J, Chernyavsky AI, Jolkovsky DL, Pinkerton KE and Grando SA: Receptor-mediated tobacco toxicity: Cooperation of the Ras/Raf-1/MEK1/ERK and JAK-2/STAT-3 pathways downstream of alpha7 nicotinic receptor in oral keratinocytes. *FASEB J* 20: 2093-2101, 2006.
- Nakayama H, Numakawa T and Ikeuchi T: Nicotine-induced phosphorylation of Akt through epidermal growth factor receptor and Src in PC12h cells. *J Neurochem* 83: 1372-1379, 2002.
- Bierut LJ: Nicotine dependence and genetic variation in the nicotinic receptors. *Drug Alcohol Depend* 104 (Suppl 1): S64-S69, 2009.
- Catassi A, Servent D, Paleari L, Cesario A and Russo P: Multiple roles of nicotine on cell proliferation and inhibition of apoptosis: Implications on lung carcinogenesis. *Mutat Res* 659: 221-231, 2008.
- Ng MK, Wu J, Chang E, Wang BY, Katzenberg-Clark R, Ishii-Watabe A and Cooke JP: A central role for nicotinic cholinergic regulation of growth factor-induced endothelial cell migration. *Arterioscler Thromb Vasc Biol* 27: 106-112, 2007.
- Beasley MB, Brambilla E and Travis WD: The 2004 World Health Organization classification of lung tumors. *Semin Roentgenol* 40: 90-97, 2005.
- Amos CI, Wu X, Broderick P, Gorlov IP, Gu J, Eisen T, Dong Q, Zhang Q, Gu X, Vijayakrishnan J, *et al*: Genome-wide association scan of tag SNPs identifies a susceptibility locus for lung cancer at 15q25.1. *Nat Genet* 40: 616-622, 2008.
- Hung RJ, McKay JD, Gaborieau V, Boffetta P, Hashibe M, Zaridze D, Mukeria A, Szeszenia-Dabrowska N, Lissowska J, Rudnai P, *et al*: A susceptibility locus for lung cancer maps to nicotinic acetylcholine receptor subunit genes on 15q25. *Nature* 452: 633-637, 2008.
- Thorgeirsson TE, Geller F, Sulem P, Rafnar T, Wiste A, Magnusson KP, Manolescu A, Thorleifsson G, Stefansson H, Ingason A, *et al*: A variant associated with nicotine dependence, lung cancer and peripheral arterial disease. *Nature* 452: 638-642, 2008.
- McKay JD, Hung RJ, Gaborieau V, Boffetta P, Chabrier A, Byrnes G, Zaridze D, Mukeria A, Szeszenia-Dabrowska N, Lissowska J, *et al*; EPIC Study: Lung cancer susceptibility locus at 5p15.33. *Nat Genet* 40: 1404-1406, 2008.
- Wang Y, Broderick P, Webb E, Wu X, Vijayakrishnan J, Matakidou A, Qureshi M, Dong Q, Gu X, Chen WV, *et al*: Common 5p15.33 and 6p21.33 variants influence lung cancer risk. *Nat Genet* 40: 1407-1409, 2008.
- Broderick P, Wang Y, Vijayakrishnan J, Matakidou A, Spitz MR, Eisen T, Amos CI and Houlston RS: Deciphering the impact of common genetic variation on lung cancer risk: A genome-wide association study. *Cancer Res* 69: 6633-6641, 2009.
- Liu P, Vikis HG, Wang D, Lu Y, Wang Y, Schwartz AG, Pinney SM, Yang P, de Andrade M, Petersen GM, *et al*: Familial aggregation of common sequence variants on 15q24-25.1 in lung cancer. *J Natl Cancer Inst* 100: 1326-1330, 2008.
- Landi MT, Chatterjee N, Yu K, Goldin LR, Goldstein AM, Rotunno M, Mirabello L, Jacobs K, Wheeler W, Yeager M, *et al*: A genome-wide association study of lung cancer identifies a region of chromosome 5p15 associated with risk for adenocarcinoma. *Am J Hum Genet* 85: 679-691, 2009.
- Flora AV, Zambrano CA, Gallego X, Miyamoto JH, Johnson KA, Cowan KA, Stitzel JA and Ehringer MA: Functional characterization of SNPs in CHRNA3/B4 intergenic region associated with drug behaviors. *Brain Res* 1529: 1-15, 2013.
- Paliwal A, Vaissière T, Kraus A, Cuenin C, Cros MP, Zaridze D, Mukeria A, Boffetta P, Hainaut P, Brennan P, *et al*: Aberrant DNA methylation links cancer susceptibility locus 15q25.1 to apoptotic regulation and lung cancer. *Cancer Res* 70: 2779-2788, 2010.
- Song P, Sekhon HS, Fu XW, Maier M, Jia Y, Duan J, Proskosil BJ, Gravett C, Lindstrom J, Mark GP, *et al*: Activated cholinergic signaling provides a target in squamous cell lung carcinoma. *Cancer Res* 68: 4693-4700, 2008.
- Paleari L, Catassi A, Ciarlo M, Cavalieri Z, Bruzzo C, Servent D, Cesario A, Chessa L, Cilli M, Piccardi F, *et al*: Role of alpha7-nicotinic acetylcholine receptor in human non-small cell lung cancer proliferation. *Cell Prolif* 41: 936-959, 2008.

23. Heusch WL and Maneckjee R: Signalling pathways involved in nicotine regulation of apoptosis of human lung cancer cells. *Carcinogenesis* 19: 551-556, 1998.
24. Medjber K, Freidja ML, Grelet S, Lorenzato M, Maouche K, Nawrocki-Raby B, Birembaut P, Polette M and Tournier JM: Role of nicotinic acetylcholine receptors in cell proliferation and tumour invasion in broncho-pulmonary carcinomas. *Lung Cancer* 87: 258-264, 2015.
25. Li H, Wang S, Takayama K, Harada T, Okamoto I, Iwama E, Fujii A, Ota K, Hidaka N, Kawano Y, *et al*: Nicotine induces resistance to erlotinib via cross-talk between $\alpha 1$ nAChR and EGFR in the non-small cell lung cancer xenograft model. *Lung Cancer* 88: 1-8, 2015.
26. Zhao Q, Gu X, Zhang C, Lu Q, Chen H and Xu L: Blocking M2 muscarinic receptor signaling inhibits tumor growth and reverses epithelial-mesenchymal transition (EMT) in non-small cell lung cancer (NSCLC). *Cancer Biol Ther* 16: 634-643, 2015.
27. Chernyavsky AI, Shchepotin IB, Galitovkiy V and Grando SA: Mechanisms of tumor-promoting activities of nicotine in lung cancer: Synergistic effects of cell membrane and mitochondrial nicotinic acetylcholine receptors. *BMC Cancer* 15: 152, 2015.
28. Xu R, Shang C, Zhao J, Han Y, Liu J, Chen K and Shi W: Activation of M3 muscarinic receptor by acetylcholine promotes non-small cell lung cancer cell proliferation and invasion via EGFR/PI3K/AKT pathway. *Tumour Biol* 36: 4091-4100, 2015.
29. Zhong X, Fan Y, Ritzenthaler JD, Zhang W, Wang K, Zhou Q and Roman J: Novel link between prostaglandin E2 (PGE2) and cholinergic signaling in lung cancer: The role of c-Jun in PGE2-induced $\alpha 7$ nicotinic acetylcholine receptor expression and tumor cell proliferation. *Thorac Cancer* 6: 488-500, 2015.
30. Schaal C, Padmanabhan J and Chellappan S: The role of nAChR and calcium signaling in pancreatic cancer initiation and progression. *Cancers (Basel)* 7: 1447-1471, 2015.
31. Al-Wadei MH, Al-Wadei HA and Schuller HM: Effects of chronic nicotine on the autocrine regulation of pancreatic cancer cells and pancreatic duct epithelial cells by stimulatory and inhibitory neurotransmitters. *Carcinogenesis* 33: 1745-1753, 2012.
32. Al-Wadei MH, Al-Wadei HA and Schuller HM: Pancreatic cancer cells and normal pancreatic duct epithelial cells express an autocrine catecholamine loop that is activated by nicotinic acetylcholine receptors $\alpha 3$, $\alpha 5$, and $\alpha 7$. *Mol Cancer Res* 10: 239-249, 2012.
33. Al-Wadei HA, Plummer HK III and Schuller HM: Nicotine stimulates pancreatic cancer xenografts by systemic increase in stress neurotransmitters and suppression of the inhibitory neurotransmitter gamma-aminobutyric acid. *Carcinogenesis* 30: 506-511, 2009.
34. Hermann PC, Sancho P, Cañamero M, Martinelli P, Madriles F, Michl P, Gress T, de Pascual R, Gandia L and Guerra C: Nicotine promotes initiation and progression of KRAS-induced pancreatic cancer via Gata6-dependent dedifferentiation of acinar cells in mice. *Gastroenterology* 147: 1119-1133.e4, 2014.
35. Bose C, Zhang H, Udupa KB and Chowdhury P: Activation of p-ERK1/2 by nicotine in pancreatic tumor cell line AR42J: Effects on proliferation and secretion. *Am J Physiol Gastrointest Liver Physiol* 289: G926-G934, 2005.
36. Chowdhury P and Walker A: A cell-based approach to study changes in the pancreas following nicotine exposure in an animal model of injury. *Langenbecks Arch Surg* 393: 547-555, 2008.
37. Schuller HM, Al-Wadei HA and Majidi M: GABA B receptor is a novel drug target for pancreatic cancer. *Cancer* 112: 767-778, 2008.
38. Momi N, Ponnusamy MP, Kaur S, Rachagani S, Kunigal SS, Chellappan S, Ouellette MM and Batra SK: Nicotine/cigarette smoke promotes metastasis of pancreatic cancer through $\alpha 7$ nAChR-mediated MUC4 upregulation. *Oncogene* 32: 1384-1395, 2013.
39. Kunigal S, Ponnusamy MP, Momi N, Batra SK and Chellappan SP: Nicotine, IFN- γ and retinoic acid mediated induction of MUC4 in pancreatic cancer requires E2F1 and STAT-1 transcription factors and utilize different signaling cascades. *Mol Cancer* 11: 24, 2012.
40. Chipitsyna G, Gong Q, Anandanadesan R, Alnajjar A, Batra SK, Wittel UA, Cullen DM, Akhter MP, Denhardt DT, Yeo CJ, *et al*: Induction of osteopontin expression by nicotine and cigarette smoke in the pancreas and pancreatic ductal adenocarcinoma cells. *Int J Cancer* 125: 276-285, 2009.
41. Lazar M, Sullivan J, Chipitsyna G, Gong Q, Ng CY, Salem AF, Aziz T, Witkiewicz A, Denhardt DT, Yeo CJ, *et al*: Involvement of osteopontin in the matrix-degrading and proangiogenic changes mediated by nicotine in pancreatic cancer cells. *J Gastrointest Surg* 14: 1566-1577, 2010.
42. Shin VY, Jin HC, Ng EK, Yu J, Leung WK, Cho CH and Sung JJ: Nicotine and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone induce cyclooxygenase-2 activity in human gastric cancer cells: Involvement of nicotinic acetylcholine receptor (nAChR) and beta-adrenergic receptor signaling pathways. *Toxicol Appl Pharmacol* 233: 254-261, 2008.
43. Chen WY, Huang CY, Cheng WL, Hung CS, Huang MT, Tai CJ, Liu YN, Chen CL and Chang YJ: Alpha 7-nicotinic acetylcholine receptor mediates the sensitivity of gastric cancer cells to 5-fluorouracil. *Tumour Biol* 36: 9537-9544, 2015.
44. Kalantari-Dehaghi M, Parnell EA, Armand T, Bernard HU and Grando SA: The nicotinic acetylcholine receptor-mediated reciprocal effects of the tobacco nitrosamine NNK and SLURP-1 on human mammary epithelial cells. *Int Immunopharmacol* 29: 99-104, 2015.