




Draft Genome Sequence of *Corynebacterium pseudotuberculosis* Strain PA07 Biovar *ovis*, Isolated from a Sheep Udder in Amazonia

Fabrício Almeida Araújo, Joana Montezano Marques,
Vitória Almeida Gonçalves de Moura, Maria Paula Cruz Schneider,
Soraya Silva Andrade, Alyne Cristina Sodré Lima,  Luis Carlos Guimarães,
Adriana Ribeiro Carneiro Folador, Artur Silva, Rommel T. J. Ramos

University of Pará, Center of Genomics and System Biology, Laboratory of Genomic and Bioinformatics, Belém, Pará, Brazil

ABSTRACT In this work, we present the draft genome sequence of *Corynebacterium pseudotuberculosis* strain PA07 biovar *ovis*, isolated from a caseous secretion from a sheep udder in Pará, Brazil. The genome contains 2,320,235 bp, 52.2% G+C content, 2,191 coding sequences (CDSs), five pseudogenes, 48 tRNAs, and three rRNAs.

Corynebacterium pseudotuberculosis is a Gram-positive, facultative intracellular, pleomorphic, nonsporulating, noncapsulated, nonmotile bacterium that, together with *Mycobacterium*, *Nocardia*, and *Rhodococcus*, forms the CMNR group which shares specific characteristics, such as (i) high G+C content and (ii) specific cell wall organization, characterized by the presence of peptidoglycan, arabinogalactan, and mycolic acids (1, 2).

C. pseudotuberculosis is the etiological agent of caseous lymphadenitis (CLA), ulcerative lymphangitis, mastitis (clinical and subclinical), and ulcerative dermatitis. CLA is characterized by the formation of abscesses in the superficial lymph nodes and subcutaneous tissues (3, 4). CLA is a widespread disease that has been reported all over the world (5). This disease is associated with direct economic losses due to the progressive reduction in weight gain, depreciated wool and skin, reduced milk production, and eventually death caused by toxemia of the infected animals. Moreover, this bacterium presents significant zoonotic potential (5).

C. pseudotuberculosis presents some variable tools for persistence and escape from the immune system of the host (3). Thus, the molecular study of the genome of this pathogen and the search for new target genes is an important step for vaccine development (6, 7).

Herein, we present the draft genome of *C. pseudotuberculosis* strain PA07 biovar *ovis*, isolated from a Santa Inês sheep udder in Pará, Brazil. To the best of our knowledge, we report the first recovery of *C. pseudotuberculosis* from an abscess located in sheep udder in the state of Pará. This strain is part of the collection of Laboratory of Genomics and System Biology located at the Federal University of Pará, Belém, Pará, Brazil.

The genome sequencing was performed using an Ion Torrent personal genome machine (PGM) (Thermo, Fisher) with a fragment library. The *de novo* assembly strategy was performed using SPAdes genome assembler software version 3.9.0 (8). The assembly generated seven contigs with 2,320,235 bp. The contigs were automatically annotated using Rapid Annotations using Subsystems Technology (RAST) (9). The genome has 2,191 coding sequences (CDSs), five pseudogenes, 48 tRNAs, three rRNAs, and a G+C content of 52.2%.

Received 16 January 2017 Accepted 19 January 2017 Published 23 March 2017

Citation Araújo FA, Marques JM, de Moura VAG, Schneider MPC, Andrade SS, Lima ACS, Guimarães LC, Folador ARC, Silva A, Ramos RTJ. 2017. Draft genome sequence of *Corynebacterium pseudotuberculosis* strain PA07 biovar *ovis*, isolated from a sheep udder in Amazonia. Genome Announc 5:e00040-17. <https://doi.org/10.1128/genomeA.00040-17>.

Copyright © 2017 Araújo et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Rommel T. J. Ramos, rommelthiago@gmail.com.

Accession number(s). The *C. pseudotuberculosis* whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession no. [MQVG00000000](#). The version described in this paper is MQVG01000000.

ACKNOWLEDGMENTS

This study was supported by the National Council for Scientific and Technological Development (CNPq), Coordination for the Improvement of Higher Education Personnel (CAPES), Foundation for Supporting Research in the State of Pará (FAPESPA), and by the Genome and Proteome Network of the State of Pará (RPGP).

REFERENCES

1. Rebouças MF, Loureiro D, Bastos BL, Moura-Costa LF, Hanna SA, Azevedo V, Meyer R, Portela RW. 2013. Development of an indirect ELISA to detect *Corynebacterium pseudotuberculosis* specific antibodies in sheep employing T1 strain culture supernatant as antigen. *Pesq Vet Bras* 33:1296–1302. <https://doi.org/10.1590/S0100-736X2013001100002>.
2. Dorella FA, Pacheco LGC, Oliveira SC, Miyoshi A, Azevedo V. 2006. *Corynebacterium pseudotuberculosis*: microbiology, biochemical properties, pathogenesis and molecular studies of virulence. *Vet Res* 37:201–218. <https://doi.org/10.1051/vetres:2005056>.
3. Baird GJ, Fontaine MC. 2007. *Corynebacterium pseudotuberculosis* and its role in ovine caseous lymphadenitis. *J Comp Pathol* 137:179–210. <https://doi.org/10.1016/j.jcpa.2007.07.002>.
4. Yeruham I, Elad D, Friedman S, Perl S. 2003. *Corynebacterium pseudotuberculosis* infection in Israeli dairy cattle. *Epidemiol Infect* 131:947–955. <https://doi.org/10.1017/S095026880300894X>.
5. Soares SC, Silva A, Trost E, Blom J, Ramos R, Carneiro A, Ali A, Santos AR, Pinto AC, Diniz C, Barbosa EG, Dorella FA, Aburjaile F, Rocha FS, Nascimento KK, Guimarães LC, Almeida S, Hassan SS, Bakhtiar SM, Pereira UP, Abreu VAC, Schneider MPC, Miyoshi A, Tauch A, Azevedo V. 2013. The pan-genome of the animal pathogen *Corynebacterium pseudotuberculosis* reveals differences in genome plasticity between the biovar *ovis* and *equi* strains. *PLoS One* 8:e53818. <https://doi.org/10.1371/journal.pone.0053818>.
6. D'Afonseca V, Moraes PM, Dorella FA, Pacheco LG, Meyer R, Portela RW, Miyoshi A, Azevedo V. 2008. A description of genes of *Corynebacterium pseudotuberculosis* useful in diagnostics and vaccine applications. *Genet Mol Res* 7:252–260. <https://doi.org/10.4238/vol7-1gmr438>.
7. Dorella FA, Gala-Garcia A, Pinto AC, Sarrouh B, Antunes CA, Ribeiro D, Aburjaile FF, Fiaux KK, Guimarães LC, Seyffert N, El-Aouar RA, Silva R, Hassan SS, Castro TLP, Marques WS, Ramos R, Carneiro A, de Sá P, Miyoshi A, Azevedo V, Silva A. 2013. Progression of “OMICS” methodologies for understanding the pathogenicity of *Corynebacterium pseudotuberculosis*: the Brazilian experience. *Comput Struct Biotechnol J* 6:1–7. <https://doi.org/10.5936/csbj.201303013>.
8. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
9. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.