

Risk factors for intraocular pressure rise following phacoemulsification

Muge Coban-Karatas, Selcuk Sizmaz, Rana Altan-Yaycioglu, Handan Canan, Yonca Aydın Akova

Purpose: This study was designed to analyze the risk factors resulting in high intraocular pressure (IOP), which was accepted as IOP higher than 22 mmHg, following uncomplicated phacoemulsification. **Materials and Methods:** The records of 812 eyes of 584 patients who underwent uncomplicated phacoemulsification were evaluated. There were 330 men and 254 women ranging between the age of 26 and 89 years (65.4 ± 9.8 years). The preoperative, postoperative first day (day 1), first week (day 7), and first month (day 30) IOP values were analyzed. Data on history of diabetes, glaucoma, pseudoexfoliation (PXF), incision site, capsular staining with trypan blue, and surgeon were recorded. A multinomial regression analysis was performed to analyse the relationship of the factors with postoperative high IOP. **Results:** The mean IOP was 15.6 ± 4.3 mmHg preoperatively. Postoperatively that were changed to 19.7 ± 9.0 mmHg at day 1, 12.7 ± 4.5 mmHg at day 7, and 12.8 ± 3.7 mmHg at day 30. The factors such as surgeon, presence of PXF, diabetes, surgical incision site, and trypan blue were not related to the postoperative high IOP ($P > 0.05$, in all). The only factor that related to high IOP at all visits was glaucoma ($P < 0.005$). **Conclusion:** According to our results, preoperative diagnosis of glaucoma seems to be the only factor to affect the postoperative IOP higher than 22 mmHg.

Key words: Intraocular pressure rise, phacoemulsification, risk factors

Intraocular pressure (IOP) may rise even after uncomplicated cataract surgery which might need intervention.^[1,2] The response in IOP to phacoemulsification is biphasic, with a transient immediate rise followed by a modest long-term decrease.^[3-5] Postoperative IOP usually peaks 5–7 h after surgery and returns to normal levels in 1–3 days.^[3,6,7] Although transient, the elevated IOP can cause ocular pain, may increase the risk of sight threatening complications such as retinal vascular occlusion, progressive field loss in advanced glaucoma, and anterior ischemic optic neuropathy in susceptible patients.^[8,9] Several risk factors such as glaucoma, viscoelastic agent, and surgical procedure for postoperative IOP rise have been identified.^[3,10] Our aim in this study was to analyze the risk factors that might result in IOP rise following uneventful phacoemulsification.

Materials and Methods

We retrospectively evaluated the records of 812 eyes of 584 consecutive patients, who underwent uncomplicated clear corneal incision phacoemulsification surgery in our clinic between May 2004 and July 2008. There were 330 men and 254 women ranging between the age of 26 and 89 years (65.4 ± 9.8 years).

All the surgical procedures were performed by two surgeons (RAY and SS). All the patients had peribulbar anesthesia

Presented in ASCRS Symposium on Cataract, IOL and Refractive Surgery April 3–8, 2009.

Department of Ophthalmology, Baskent University, Ankara, Turkey

Correspondence to: Prof. Muge Coban-Karatas, Department of Ophthalmology, Baskent University, Faculty of Medicine Teaching and Research Center, Dadaloğlu Mah. 39 sk. No: 6 Yüreğir, Adana, Turkey. E-mail: bkaratas99@hotmail.com

Manuscript received: 08.01.11; **Revision accepted:** 03.11.11

and uncomplicated phacoemulsification surgery *via* 2.85 mm corneal incision placed either temporal, or oblique. Two side port incisions were made to allow access of a second instrument. Dispersive viscoelastic (Viscoat, Alcon, TX, USA) was used to fill the anterior chamber. Continuous curvilinear capsulorhexis was performed routinely. In mature cataracts before performing capsulorhexis, trypan blue was used to dye the capsule. During phacoemulsification, a stop and chop or phaco-chop technique was used. Following, the corneal incision was enlarged to 4.0 mm. The capsule was filled with a cohesive viscoelastic (Provisc, Alcon, TX, USA), and a foldable posterior chamber intraocular lens was implanted in the capsular bag. Viscoelastic was thoroughly removed via bimanual irrigation/aspiration (I/A), and a rock and roll technique was used for the removal of viscoelastic behind the IOL.

The preoperative as well as postoperative first day, first week, and first month visit IOP values were measured by non-contact tonometer (Full Auto Tonometer TX-F, Canon Inc, USA) and recorded. IOP was grouped as normal (≤ 22 mmHg) and high (≥ 22 mmHg). All the glaucoma patients were primary open angle glaucoma. Patients that suffer from other types of glaucoma were excluded from the study. Patients who were receiving antiglaucomatous medication also continued to receive the same medication before and after surgery. None of the patients was given prophylactic medication before the surgery. Antiglaucoma drugs that incite more inflammation like prostaglandin analogs were stopped 2 weeks before surgery and an antiglaucoma drop that does not incite inflammation was prescribed. Antiglaucoma drugs that incite more inflammation like prostaglandin analogs were stopped 2 weeks before surgery. An antiglaucoma drug that does not incite inflammation was prescribed and continued for 1 month after surgery. In addition, acetazolamide was not given after the surgery.

All data were entered to SPSS software (Statistical Package for the Social Sciences, version 10.0, SPSS Inc, Chicago, IL, USA)

Access this article online

Website:

www.ijo.in

DOI:

10.4103/0301-4738.99997

Quick Response Code:



and the changes in IOP were compared with paired Student's *t*-test. Data on prior history of glaucoma, pseudoexfoliation (PXF), incision site, anterior capsular staining, and surgeon were recorded. The relation of these factors with postoperative high IOP at each visit (days 1, 7, and 30) were evaluated using multinomial regression analysis. The level of significance was set at <0.05.

This study was approved by Baskent University Institutional Review Board and Ethics Committee (project no: KA10/31).

Results

The mean preoperative IOP was 15.6 ± 4.3 mmHg (ranged from 7 to 36 mmHg). At day 1, the IOP increased significantly to 19.7 ± 9.0 mmHg (ranged between 6 and 58 mmHg; $P < 0.001$, 95% CI -4.68 to -3.52). The mean IOP was 12.7 ± 4.5 mmHg (ranged between 6 and 37 mmHg) at day 7, and 12.8 ± 3.7 mmHg (ranged between 6 and 34) at day 30. Compared to the preoperative values, days 7 and 30 IOP values were significantly low ($P < 0.001$, 95% CI 2.54–3.17 for day 7, and 2.47–3.03 for day 30).

At day 1, the high IOP was detected in 249 (30.7%) eyes with a mean of 30.7 ± 7.5 mmHg. At day 7, the number of patients with high IOP decreased to 26 eyes (8.8%). Similarly, at day 30 only 16 (1.2%) eyes had the high IOP.

The incidences of high IOP with possible associated factors are shown on Table 1. Diabetes was present in 100 eyes (12.3%), PXF was observed in 77 eyes (9.5%), and glaucoma in 60 eyes (7.4%). In 477 eyes (58.7%), the operation was performed through a temporal incision, and in 335 eyes (41.3%) a superior oblique incision was performed. Trypan blue was used in 43 eyes (5.3%).

With the multinomial regression analysis, none of the factors

was related to the high IOP postoperatively ($P < 0.05$), except glaucoma. Glaucoma was the only factor related to high IOP values during the postoperative period ($P = 0.004$ at day 1, 0.001 at day 7, and < 0.001 at day 30). Of the 60 glaucomatous eyes, all were receiving the glaucoma medication. The IOP was high in 30 eyes (51.7%) in the first day, in 7 eyes (12.1%) in the first week, and in 6 (10.3%) patients in the first month.

Discussion

Transient IOP rise may be observed in the early postoperative period after uneventful cataract surgery.^[11-13] In this study, we retrospectively evaluated the risk factors that might affect the IOP rise following uncomplicated phacoemulsification surgery. The IOP measurements were recorded at the postoperative days 1, 7, and 30. We accepted 22 mmHg as the cut-off point for high IOP. Thirty percent of all eyes had high IOP at day 1. Among all the factors such as surgeon, diabetes, PXF, glaucoma, incision site, and trypan blue use, the only factor related to postoperative high IOP values was glaucoma ($P \leq 0.001$).

Our results indicate that short-term postoperative IOP was higher in eyes with primary open angle glaucoma than in nonglaucomatous eyes ($P \leq 0.001$). In another study, glaucoma was reported as a risk factor for pressure rise after phacoemulsification and the incidence of substantial elevation in IOP was similar between the eyes with primary open angle glaucoma and those with pseudoexfoliative glaucoma.^[14] Tong and Miller^[15] retrospectively investigated the preoperative and postoperative IOP measurements of 385 consecutive eyes having uneventful phacoemulsification. Patients with preoperative diagnosis of glaucoma had significantly higher IOP at postoperative first week. Yasutani *et al.*^[14] also mentioned that a substantial increase in IOP occurred in approximately 13% of the eyes with open angle glaucoma 1 day after phacoemulsification surgery.

Table 1: The distribution of IOP values as normal (≤ 22 mmHg) or high (> 22 mmHg) at each visit (day 1, day 7, and day 30) according to the investigated factors such as surgeon, pseudoexfoliation (PXF), glaucoma, PXF and glaucoma, diabetes (DM), incision [so: Superior oblique, t: Temporal], trypan dye [1: Present; 2: Absent]

		IOP (mmHg)								
		Day 1			Day 7			Day 30		
		≤ 22	> 22	<i>P</i>	≤ 22	> 22	<i>P</i>	≤ 22	> 22	<i>P</i>
Surgeon	RAY	342	155	0.374	478	19	0.187	490	7	0.363
	SS	221	94		308	7		309	6	
PXF	1	514	221	0.660	711	24	0.633	723	12	0.372
	2	49	28		75	2		76	1	
Glaucoma	1	533	219	0.004*	733	19	0.001*	745	7	<0.001*
	2	30	30		53	7		54	6	
PXF and Glaucoma	1	552	239	0.491	766	25	0.435	779	12	0.702
	2	11	10		20	1		20	1	
DM	1	493	219	0.783	688	24	0.321	700	12	0.477
	2	70	30		98	2		99	1	
Incision	so	242	93	0.088	324	11	0.685	330	5	0.487
	t	321	156		462	15		469	8	
Trypan	1	540	229	0.053	746	23	0.257	760	9	0.098
	2	23	20		40	3		41	2	

The relation of associated factors with high IOP was evaluated using the multinomial regression analysis. *P*: Probability value, *: Probability less than 0.05

In our study, PXF with normal IOP was not significantly correlated with high IOP even in the early postoperative period after uneventful phacoemulsification. Also in two studies, it has been reported that there was no significant difference in the IOP after phacoemulsification in eyes with and without PXF.^[16,17] On the other hand, a long-term lasting reduction in mean IOP occurred in PXF eyes.^[18-21] Damji *et al.*^[21] demonstrated that in the 2 year follow-up, the patients with PXF have a greater IOP lowering effect following phacoemulsification than those without, and the authors concluded that this effect was correlated with the volume of irrigating fluid utilized at the time of surgery. In addition to these results, Cimetta and Cimetta^[18] operated a group of 39 open angle, nonglaucomatous eyes with cataract and PXF syndrome and a control group of open angle, nonglaucomatous eyes with cataract, using a standard phaco technique along with bimanual anterior capsule PXF material aspiration. Phacoemulsification with anterior capsule PXF material aspiration significantly reduced the mean diurnal IOP in the PXF group lasting one year postoperatively. In another retrospective comparative study, 1122 eyes with PXF, 240 with glaucoma and 882 without glaucoma underwent uneventful phacoemulsification. A long-term reduction in mean IOP occurred in PXF eyes with or without glaucoma.^[19] However, IOP rise in the early postoperative period was noted after phacoemulsification in nonglaucomatous eyes with PXF and IOP control was advised.^[22-24] This early IOP rise may be due to severe inflammation after cataract surgery in eyes with PXF due to pathological iris vessels with an increased permeability for protein. In addition, there were eyes with an elevated IOP without inflammation in the PXF group in which the mechanism of IOP rise could not be identified.^[23] However, in this study, with or without glaucoma, we did not observe any relation of PXF to high IOP.

Anterior capsular staining with trypan blue in our study was not associated with high IOP at any postoperative visit ($P > 0.05$). In a preliminary study, 25 eyes of 25 patients with a unilateral mature or hypermature cataract, trypan blue dye was used to stain the anterior capsule. Adverse reactions related to the dye such as raised IOP were not observed in the immediate postoperative period or at the end of the mean follow-up of 3 months.^[25] In a comparative study, phacoemulsification of 82 patients who had white mature cataract in one eye and senile cataract in the other were operated. Trypan blue dye was used in the white mature cataract and not used in the fellow eye. Postoperative IOP was not significantly different in the two groups.^[26] In a previous study,^[15] wound construction, anesthesia type, the eye operated on, patient age, and sex did not significantly influence the postoperative pressure change. Similarly in this study, there was no correlation between history of diabetes, surgeon, and incision site ($P > 0.05$). Surgeon's experience was reported as an important factor for the postoperative IOP rise. The mean pressure rise in eyes operated by experienced surgeons was about half the pressure rise in eyes operated by beginners, as beginners often perform intraocular surgery in a more traumatizing manner than experienced surgeons.^[6] In our study, there was no relation between high IOP and surgeon. This was most probably related to the similar experience of two surgeons in present series.

As for the incision site, a study that compares temporal sclerocorneal or clear corneal incision, postoperative IOP was significantly higher in the sclerocorneal tunnel group than in

the clear corneal incision group.^[27] In our group, all surgeries were performed via clear corneal incision, so we are unable to compare with different types of incision.

In conclusion, our study demonstrated that following uneventful phacoemulsification, the diagnosis of glaucoma was the only risk factor for IOP higher than 22 mmHg. On the other hand, according to our results, PXF was not a risk factor for high IOP. We believe that patients who have glaucoma are at risk, and should be monitored closely for high IOP following phacoemulsification surgery.

References

- Cohen VM, Demetria H, Jordan K, Lamb RJ, Vivian AJ. First day post-operative review following uncomplicated phacoemulsification. *Eye (Lond)* 1998;12:634-6.
- Dinakaran S, Desai SP, Raj PS. Is the first postoperative day review necessary following uncomplicated phacoemulsification surgery? *Eye (Lond)* 2000;14:364-6.
- Fang EN, Kass MA. Increased intraocular pressure after cataract surgery. *Semin Ophthalmol* 1994;9:235-42.
- Tennen DG, Masket S. Short- and long-term effect of clear corneal incision on intraocular pressure. *J Cataract Refract Surg* 1996;22:568-70.
- Obstbaum SA. Cataract surgery and its effect on intraocular pressure [editorial]. *J Cataract Refract Surg* 1999;25:877.
- Bömer TG, Lagreze WD, Funk J. Intraocular pressure rise after phacoemulsification with posterior chamber lens implantation: Effect of prophylactic medication, wound closure, and surgeon's experience. *Br J Ophthalmol* 1995;79:809-13.
- Jürgens I, Mattheu A, Castilla M. Ocular hypertension after cataract surgery: A comparison of three surgical techniques and two viscoelastics. *Ophthalmic Surg Lasers* 1997;28:30-6.
- Hayreh SS. Anterior ischemic optic neuropathy. IV. Occurrence after cataract extraction. *Arch Ophthalmol* 1980;98:1410-6.
- Savage JA, Thomas JV, Belcher CD 3rd, Simmons RJ. Extracapsular cataract extraction and posterior chamber intraocular lens implantation in glaucomatous eyes. *Ophthalmology* 1985;92:1506-16.
- Morgan RK, Skuta GL. Viscoelastic-related glaucomas. *Semin Ophthalmol* 1994;9:229-34.
- Gross JG, Meyer DR, Robin AL, Filar AA, Kelley JS. Increased intraocular pressure in the immediate postoperative period after extracapsular cataract extraction. *Am J Ophthalmol* 1988;105:466-9.
- Mc Guigan LJ, Gottsch J, Stark WJ, Maumenee AE, Quigley HA. Extracapsular cataract extraction and posterior chamber lens implantation in eyes with preexisting glaucoma. *Arch Ophthalmol* 1986;104:1301-8.
- Barron BA, Busin M, Page C, Bergsma DR, Kaufman HE. Comparison of the effects of Viscoat and Healon on postoperative intraocular pressure. *Am J Ophthalmol* 1985;100:377-84.
- Yasutani H, Hayashi K, Hayashi H, Hayashi F. Intraocular pressure rise after phacoemulsification surgery in glaucoma patients. *J Cataract Refract Surg* 2004;30:1219-24.
- Tong JT, Miller KM. Intraocular pressure change after sutureless phacoemulsification and foldable posterior chamber lens implantation. *J Cataract Refract Surg* 1998;24:256-62.
- Dosso AA, Bonvin ER, Leuenberger PM. Exfoliation syndrome and phacoemulsification. *J Cataract Refract Surg* 1997;23:122-5.
- Fine IH, Hoffman RS. Phacoemulsification in the presence of pseudoexfoliation: Challenges and options. *J Cataract Refract Surg* 1997;23:160-5.
- Cimetta DJ, Cimetta AC. Intraocular pressure changes after clear

- corneal phacoemulsification in nonglaucomatous pseudoexfoliation syndrome. *Eur J Ophthalmol* 2008;18:77-81
19. Shingleton BJ, Laul A, Nagao K, Wolff B, O'Donoghue M, Eagan E, *et al.* Effect of phacoemulsification on intraocular pressure in eyes with pseudoexfoliation: Single surgeon series. *J Cataract Refract Surg* 2008;34:1834-41.
 20. Altan-Yaycioglu R, Canan H, Pelit A, Akova YA. Intraocular pressure in pseudoexfoliative eyes following phacoemulsification. *J Cataract Refract Surg* 2009;25:952-4.
 21. Damji KF, Konstas AG, Liebmann JM, Hodge WG, Ziakas NG, Giannikakis S, *et al.* Intraocular pressure following phacoemulsification in patients with and without exfoliation syndrome: A 2 year prospective study. *Br J Ophthalmol* 2006;90:1014-8.
 22. Pohjalainen T, Vesti E, Ulusitalo RJ, Laatikainen L. Intraocular pressure after phacoemulsification and intraocular lens implantation in nonglaucomatous eyes with and without exfoliation. *J Cataract Refract Surg* 2001;27:426-31.
 23. Akinci A, Batman C, Zilelioglu O. Phacoemulsification in pseudoexfoliation syndrome. *Ophthalmologica* 2008;222:112-6.
 24. Levkovitch-Verbin H, Hahot-Wilner Z, Burla N, Melamed S, Goldenfeld M, Bar-Sela SM. Intraocular pressure elevation within the first 24 hours after cataract surgery in patients with glaucoma or exfoliation syndrome. *Ophthalmology* 2008;115:104-8.
 25. Kothari K, Jain SS, Shah NJ. Anterior capsular staining with trypan blue for capsulorhexis in mature and hypermature cataracts. A preliminary study. *Indian J Ophthalmol* 2001;49:177-80.
 26. Ermiş SS, Oztürk F, Inan UU. Comparing the efficacy and safety of phacoemulsification in white mature and other types of senile cataracts. *Br J Ophthalmol* 2003;87:1356-9.
 27. Schwenn O, Dick HB, Krummenauer F, Krist R, Pfeiffer N. Intraocular pressure after small incision cataract surgery: Temporal sclerocorneal versus clear corneal incision. *J Cataract Refract Surg* 2001;27:421-5.

Cite this article as: Coban-Karatas M, Sizmaz S, Altan-Yaycioglu R, Canan H, Akova YA. Risk factors for intraocular pressure rise following phacoemulsification. *Indian J Ophthalmol* 2013;61:115-8.

Source of Support: Nil. **Conflict of Interest:** Nil.

New features on the journal's website

Optimized content for mobile and hand-held devices

HTML pages have been optimized of mobile and other hand-held devices (such as iPad, Kindle, iPod) for faster browsing speed.

Click on **[Mobile Full text]** from Table of Contents page.

This is simple HTML version for faster download on mobiles (if viewed on desktop, it will be automatically redirected to full HTML version)

E-Pub for hand-held devices

EPUB is an open e-book standard recommended by The International Digital Publishing Forum which is designed for reflowable content i.e. the text display can be optimized for a particular display device.


Click on **[EPub]** from Table of Contents page.

There are various e-Pub readers such as for Windows: Digital Editions, OS X: Calibre/Bookworm, iPhone/iPod Touch/iPad: Stanza, and Linux: Calibre/Bookworm.

E-Book for desktop

One can also see the entire issue as printed here in a 'flip book' version on desktops.

Links are available from as well as Archives pages.

Click on  View as eBook