




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Available at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/hemonc

LETTER TO EDITOR

Outcome of myeloma patients with COVID-19 on active lenalidomide-based therapy: Does lenalidomide protect from severe COVID-19?

Imran K. Tailor^{*}, Nawal F. Alshehry, Syed Z. Zaidi, Mohammed A. Marei, Ibraheem H. Motabi, Mansour Alfayez, Syed Y. Altaf

Department of Adult Hematology and Bone Marrow Transplantation, King Fahad Medical City, Riyadh, Saudi Arabia

Received 26 July 2020; accepted 28 August 2020

To the Editor

Patients with haematological cancers and concurrent coronavirus disease 2019 (COVID-19) reportedly have poor outcomes as indicated by high hospitalisation rates, high case fatality rates in mechanically ventilated patients, and a high mortality rate of $\geq 10\%$ [1–3]. A recent small study reported a high mortality rate among patients with myeloma and concurrent COVID-19 infection [4]. Interestingly, one large study reported that active or recent chemotherapy was not a predictor of severe disease, whereas checkpoint inhibitor therapy and age over 65 years were associated with adverse outcomes [1]. Here we describe the outcomes of three consecutive patients with myeloma and concurrent COVID-19 infection admitted at our tertiary care centre between April and June 2020, whilst on lenalidomide-based therapy, two of whom underwent autologous stem cell transplant, with at least 30 days of follow-up.

Clinical features, laboratory characteristics, and other outcomes of the three patients with myeloma are listed in Table 1.

Outcomes of patients with solid tumours and haematological cancers and concurrent COVID-19 infection are poor with high rates of hospitalisations, higher need for ventilatory support, and increased mortality rates [1–4,6,7]. Our study demonstrated excellent outcomes of three consecutive patients with myeloma whilst on lenalidomide-based therapy, with none of them developing severe disease despite being in an immunocompromised state. Patients with myeloma have an increased risk of developing bacterial (7-fold) and viral infections (10-fold) [5]. In our series, two patients underwent stem cell transplant and two of them had significant comorbidities. Although no conclusions can be drawn, our study raises a very interesting question: does active chemotherapy or immunomodulatory treatment have any protective role against severe COVID-19?

One large study involving over 400 patients with cancer, including haematological cancers, and concurrent COVID-19 infection reported that recent chemotherapy was not associated with adverse outcomes, whereas checkpoint inhibitor

^{*} Corresponding author.

E-mail address: doctorimran77@yahoo.co.uk (I.K. Tailor).

<https://doi.org/10.1016/j.hemonc.2020.08.002>

1658-3876/© 2020 King Faisal Specialist Hospital & Research Centre. Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article as: I. K. Tailor, N. F. Alshehry, S. Z. Zaidi et al., Outcome of myeloma patients with COVID-19 on active lenalidomide-based therapy: Does lenalidomide protect from severe COVID-19?, *Hematol Oncol Stem Cell Ther*, <https://>

Table 1 Clinical and Laboratory Characteristics of Patients.

Characteristic	Patient 1	Patient 2	Patient 3
Age	57	43	49
Sex	Female	Male	Male
Diagnosis	Relapsed light chain myeloma post bortezomib/autologous stem cell transplant/lenalidomide/daratumumab	Ig A myeloma post bortezomib/autologous stem cell transplant/lenalidomide	Newly diagnosed light chain myeloma with plasma cell leukaemia with recent history of acute kidney injury
Previous autologous stem cell transplant	Yes	Yes	No
Comorbidities	Diabetes, hypertension, ischemic heart disease, chronic kidney disease	Learning difficulties, care home resident	None
ECOG performance status at presentation	2	1	1
Therapy at the time of presentation	Lenalidomide, dexamethasone	Lenalidomide, dexamethasone	Lenalidomide, bortezomib, dexamethasone
Active treatment for myeloma at COVID-19 diagnosis or in the past 4 wk	Yes	Yes	Yes
Symptoms	Fever, cough, shortness of breath	Cough	Fever, cough
Chest x-ray	Infiltrates present	Not performed	Normal
Oxygen requirement	2 L/min	None	None
Laboratory findings	Mild anaemia, leukopenia, moderate thrombocytopenia; Normal coagulation/D-dimers, Chemistry Increased CRP, ferritin, and LDH	Not done	Mild anaemia only
Noninvasive ventilation	No	No	No
Mechanical ventilation	No	No	No
Treatment	Antibiotics and supportive care	Supportive care	Supportive care
IVIg therapy in previous 4 wk	Yes	No	No
Hospitalisation (d)	13	No (care home isolation only)	25 (till negative COVID swab)
Time taken for negative COVID-19 from initial positive	14	Not tested	25
Outcome	Recovered	Recovered	Recovered
Follow-up (d)	30	59	83

Note. CRP = C-reactive protein; D = day; ECOG = Eastern Cooperative Oncology Group; IVIG = intravenous immunoglobulin; LDH = lactate dehydrogenase; min = minute; wk = week.

therapy (which acts by harnessing and activating the immune system) was a predictor of poor outcome [11]. Other smaller studies also made similar observation that there was no correlation between active treatment with chemotherapy and outcome among patients with haematological cancers [6]. Acute respiratory distress syndrome (ARDS) is the most common cause of death among patients with COVID-19 and develops secondary to cytokine storm [8,10]. Cytokine storm is characterised by disproportionate response to infection or inflammation inducing significant increase in cytokines, such as interleukin (IL)-1, IL-2, IL-6, granulocyte colony-stimulating factor, interferon δ , and tumour necrosis factor (TNF) α , which interact with the complement system and coagulation system and may lead to disseminated intravascular coagulation, multiorgan failure, and death in some patients [9]. COVID-19 is associated with excessive cytokine levels and ARDS; however, there is no consensus if viral replication and/or cytokine storm are responsible for ARDS [9]. This forms the basis of use of many anti-inflammatory drugs in COVID-19, including tocilizumab (IL-6 inhibitor), Bruton tyrosine kinase inhibitors (e.g., ibrutinib), and Janus kinase-2 inhibitors (e.g., ruxolitinib), which have shown favourable outcomes in small series of patients [8,11]. Dexamethasone, an immunosuppressive agent, has shown to reduce COVID-19-related mortality rate in oxygenated and ventilated patients by 20% and 35%, respectively, in one of the largest randomised controlled trials conducted till date for COVID-19 patients [12].

Lenalidomide, an immunomodulatory agent, is one of the backbones in the treatment of myeloma leading to great outcomes; it is used at various stages of management including at induction, in relapsed setting, and as maintenance therapy [13]. Its immunomodulatory action inhibits the production of proinflammatory cytokines, such as TNF- α , IL-1, IL-6, and IL-12, and elevates the production of anti-inflammatory cytokine IL-10, and this has been implicated in clinical efficacy in myeloma [14]. This particular effect of lenalidomide may be playing a protective role in preventing the catastrophic cytokine storm, thereby dampening the immune system which may well be helpful in patients with COVID-19.

Based on aforementioned observations, although our study is a small series, we hypothesise that lenalidomide may play a protective role in myeloma patients and prevent developing severe form of disease; however, this warrants further testing. Second, it raises another important question: should immunomodulatory drugs such as lenalidomide be continued during active COVID-19 infection? This theory also supports the notion that initiation or continuation of chemotherapy or immunosuppressive therapy should not be delayed or stopped in patients with haematological cancers even if they are deemed to be stable or low risk, lest it should alter overall outcome, just for the fear of pandemic. Moreover, this may have an implication in other autoimmune conditions too, wherein use of immunomodulatory drugs is the norm; stopping a therapy may be associated with adverse outcomes from not only severe COVID-19 but also progression of autoimmune disease.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] Robilotti EV, Babady NE, Mead PA, Rolling T, Perez-Johnston R, Bernardes M, et al. Determinants of COVID-19 disease severity in patients with cancer. *Nat Med* 2020;26(8):1218–23. <https://doi.org/10.1038/s41591-020-0979-0>.
- [2] Yu J, Ouyang W, Chua MLK, Xie C. SARS-CoV-2 transmission in patients with cancer at a tertiary care hospital in Wuhan, China. *JAMA Oncol* 2020;6(7):1108. <https://doi.org/10.1001/jamaoncol.2020.0980>.
- [3] Mehta V, Goel S, Kabarriti R, Cole D, Goldfinger M, Acuna-Villaorduna A, et al. Case fatality rate of cancer patients with COVID-19 in a New York hospital system. *Cancer Discov* 2020;10(7):935–41.
- [4] Dhakal B, D'Souza A, Chhabra S, Hari P. Multiple myeloma and COVID-19. *Leukemia* 2020;34(7):1961–3. <https://doi.org/10.1038/s41375-020-0879-9>.
- [5] Blimark C, Holmberg E, Mellqvist UH, Landgren O, Bjorkholm M, Hultcrantz M, et al. Multiple myeloma and infections: a population-based study on 9253 multiple myeloma patients. *Haematologica* 2015;100:107–13.
- [6] Aries JA, Davies JK, Auer RL, Hallam SL, Montoto S, Smith M, et al. Clinical outcome of coronavirus disease 2019 in haematology patients. *Br J Haematol* 2020;190(2). <https://doi.org/10.1111/bjh.16852>.
- [7] Malard F, Genthon A, Brissot E, van de Wyngaert Z, Marjanovic Z, Ikhlef S, et al. COVID-19 outcomes in patients with hematologic disease. *Bone Marrow Transplant* 2020:1–5. <https://doi.org/10.1038/s41409-020-0931-4>.
- [8] Alijotas-Reig J, Esteve-Valverde E, Belizna C, Selva-O'Callaghan A, Pardos-Gea J, Quintana A, et al. Immunomodulatory therapy for the management of severe COVID-19. Beyond the anti-viral therapy: A comprehensive review. *Autoimmun Rev* 2020;19:102569. <https://doi.org/10.1016/j.autrev.2020.102569>.
- [9] Mangalmurti N, Hunter CA. Cytokine storms: understanding COVID-19. *Immunity* 2020;53(1):19–25. <https://doi.org/10.1016/j.immuni.2020.06.017>.
- [10] Vardhana SA, Wolchok JD. The many faces of the anti-COVID immune response. *J Exp Med* 2020;217:e20200678. <https://doi.org/10.1084/jem.20200678>.
- [11] Treon SP, Castillo JJ, Skarbnik AP, Soumerai JD, Ghobrial IM, Guerrero ML, et al. The BTK inhibitor ibrutinib may protect against pulmonary injury in COVID-19–infected patients. *Blood* 2020;135(21):1912–5. <https://doi.org/10.1182/blood.202006288>.
- [12] Horby P, Lim WS, Emberson J, Mafham M, Bell J. Effect of dexamethasone in hospitalized patients with COVID-19—preliminary report. *MedRxiv* 2020. <https://doi.org/10.1101/2020.06.22.20137273>.
- [13] Rajkumar SV. Multiple myeloma: 2020 update on diagnosis, risk-stratification and management. *Am J Hematol* 2020;95(5):548–67. <https://doi.org/10.1002/ajh.25791>.
- [14] Kotla V, Goel S, Nischal S, Heuck C, Vivek K, Das B, et al. Mechanism of action of lenalidomide in hematological malignancies. *J Hematol Oncol* 2009;2(1):36. <https://doi.org/10.1186/1756-8722-2-36>.