

In-vitro Wound Healing Effect of 15-Hydroxyprostaglandin Dehydrogenase Inhibitor from Plant

Sandeep Karna

National Institute of Horticultural and Herbal Science, Nonsaengmyeong-ro, Iseo-myeon, Wangju-gun, Jeollabuk-do, Korea

Submitted: 30-07-2016

Revised: 08-09-2016

Published: 07-04-2017


ABSTRACT

Background: Prostaglandins (PGs) have short existence *in vivo* because they are rapidly metabolized by NAD⁺-dependent 15-hydroxyprostaglandin dehydrogenase (15-PGDH) to 15-ketoprostaglandins. Inhibition of 15-PGDH causes elevated level of PGE₂ in cellular system. It will be valuable for the therapeutic management of diseases requiring elevated PGE₂ levels, like wound healing. **Objective:** Ninety-eight plant samples were screened for the discovery of potent 15-PGDH inhibitor. Among them, top five plant extracts as potent 15-PGDH inhibitor were chosen to determine PGE₂ release from HaCaT (Keratinocyte cell line) cell line. Finally, top 15-PGDH inhibitor was selected to evaluate *in vitro* wound healing effect on HaCaT scratch model. **Method:** The inhibitory activity for 15-PGDH inhibitors was evaluated using fluorescence spectrophotometer by measuring the formation of NADH at 468 nm following excitation at 340 nm. Cell viability assay and PGE₂ release was evaluated in HaCaT cell line after treatment of 15-PGDH inhibitors. Scratches were made using sterile 200 μL on HaCaT cell and wound-healing effect was evaluated after treatment of 15-PGDH inhibitor. **Results:** 15-PGDH inhibitors elevated PGE₂ levels in concentration-dependent manner. Ethanol extract of *Artocarpus heterophyllus* (EEAH), the most potent 15-PGDH inhibitor (IC₅₀ = 0.62 μg/mL) with least cytotoxicity (IC₅₀ = 670 μg/ml), elevated both intracellular and extracellular PGE₂ levels. EEAH facilitated *in-vitro* wound healing in a HaCaT (Keratinocyte cell line) scratch model. **Conclusion:** EEAH might apply to treat dermal wounds by elevating PGE₂ levels via COX-1 induction and 15-PGDH inhibition.

Key words: Prostaglandins, cyclooxygenase, PGE₂, wound healing, 15-hydroxyprostaglandin dehydrogenase

SUMMARY

- Biological inactivation of 15-PGDH causes elevated level of PGE₂ which will be useful for the management of disease that requires elevated level of PGE₂.



Abbreviations used: 15-PGDH: 15-hydroxyprostaglandin dehydrogenase, COX: Cyclooxygenase, DTT: Dithiothreitol, DMEM: Dulbecco's modified Eagle's media, EEAH: Ethanol extract of *Artocarpus heterophyllus*, MRP4: Multidrug resistance 4, PGs: Prostaglandins, PGT: Prostaglandin transporter, SDS: Sodium dodecylsulfate

Correspondence:

Dr. Sandeep Karna, National Institute of Horticultural and Herbal Science, Nonsaengmyeong-ro, Iseo-myeon, Wangju-gun, Jeollabuk-do, Korea.
E-mail: skarna@gmail.com
DOI: 10.4103/0973-1296.203971

Access this article online

Website: www.phcog.com

Quick Response Code:



INTRODUCTION

Prostaglandins (PGs) are lipid compounds that participate in a variety of physiologic and pathologic processes, and among them, PGE₂ is a major mediator for inflammation.^[1] PGE₂ is formed by PG synthetase from PGH₂ through the cyclooxygenase (COX) pathway. Two types of COX have been identified: (i) COX-1 has been expressed constitutively in various tissues, including stomach, and (ii) COX-2 has been induced by cytokines, growth factors, tumor promoters, and other agents.^[2] Newly synthesized PGE₂ simply diffused and actively extruded by the multidrug resistance 4 (MRP4) from the cells.^[3] Subsequently, EPR receptor is activated followed by pericellular PGE₂ is cleared via re-uptake of PGE₂ by PG transporter (PGT)^[4] and then rapidly metabolized by cytosolic enzyme named NAD⁺-dependent 15-hydroxyprostaglandin dehydrogenase (15-PGDH).^[5] This enzyme is expressed ubiquitously in mammalian tissues and responsible for biologic inactivation of PGE₂ to 15-ketoprostaglandins.^[6]

PGE₂ has been known as an important mediator for bone formation,^[7,8] gastric ulcer healing,^[9,10] and dermal wound healing.^[11,12] Additionally, PGE₂ has been used to treat gastric ulcer in spite of high price and low efficacy.^[13,14] Therefore, PGE₂ elevation using 15-PGDH inhibitor would be valuable for the management disease that required elevated PGE₂, like wound healing. Wound healing is a complicated process in human or animal in which skin or another organ-tissue repair itself after having wound.^[15] In normal skin, epidermis and dermis maintain steady-

state equilibrium to maintain protective barrier against the external environment. Once the protective barrier is broken, the wound-healing process immediately set in motion and complex biochemical events takes place to repair the damage.^[16] The aim of this study was to screen most potent 15-PGDH inhibitor from plant source and evaluate its wound-healing efficacy using *in-vitro* scratch model in HaCaT cell.

MATERIALS AND METHODS

Materials, reagents, and instruments

Plant extracts were purchased from the Korean Plant Extract Bank (Daejeon, Korea). PGE₂, NAD⁺, NADH, glutathione-sepharose 4B, dithiothreitol (DTT), sodium dodecylsulfate (SDS), EDTA, reduced glutathione, mitomycin, and other chemicals and reagents were purchased

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Karna S. *In-vitro* wound healing effect of 15-hydroxyprostaglandin dehydrogenase inhibitor from plant. *Phcog Mag* 2017;13:122-6.

from Sigma (St. Louis, MO, USA). TGF- β 1 was purchased from BioVision (California, USA) and pGEX-2T expression vector from Pharmacia Crop. (New Jersey, USA). cDNA of human 15-PGDH was cloned from human placenta as illustrate earlier.^[17] PGE₂ enzyme immunoassay kits were obtained from Thermo Scientific (Rockford, IL, USA). UV-Vis spectrophotometer (Shimadzu, Japan) was used to obtain UV spectra. Real-time PCR was performed using a Light Cycler 2.0 (Roche, Mannheim, Germany). Scratches were visualized and pictures captured using a transmission electron microscope (Hitachi, Tokyo, Japan).

Sample preparation

Different parts of plants were dried in the dark for 7 days and then grounded to powder. The powder samples were extracted three times with ethanol and extracts were obtained through removal of solvents during evaporations. The concentrated samples were stored at -20°C. One hundred milligram of crude extracts were dissolved in 1 ml of commercial grade ethanol.

Expression and purification of 15-PGDH

pGEX-2T expression vector was used to transform 15-PGDH cDNA plasmid into *E. coli* BL-21 lysS. The cells were grown in LB Broth (1.0 L) containing 50 mg/L ampicillin at 37°C and 230 × g. Isopropyl-1-thio- β -D-galactopyranoside was added to 1 mM when OD₆₀₀ reached 0.8 and cells were allowed to grow for additional 12 h at 25°C. Cells were harvested by centrifugation at 4000g for 20 min at 4°C and pellets were resuspended in 20 mL of cold cell lysis buffer (1× PBS buffer pH 7.4 containing 1 mM EDTA and 0.1 mM DTT) and sonicated (3 × 15 s at 4°C) to disrupt cells. Sonicated pallets were centrifuged at 4000g for 20 min at 4°C and supernatant obtained was applied to a glutathione-sepharose 4B column. The column was rinsed with lysis buffer until OD₂₈₀ reach less than 0.002. 15-PGDH was eluted from column using elution buffer (50 mM Tris-HCl pH 8.0 containing 10 mM reduced glutathione, 1 mM EDTA, and 0.1 mM DTT). The purity of 15-PGDH was examined by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and concentration measured.

15-PGDH assay

The inhibitory activity was evaluated using fluorescence spectrophotometer, measuring formation of NADH at 468 nm following excitation at 340 nm. Reaction mixture was prepared in 2 mL of Tris-HCl buffer (50 mM, pH 7.5) containing 0.1 mM DTT, 0.25 mM NAD⁺, purified enzyme (10 μ g), 21 μ M PGE₂ and various concentration inhibitors. 15-PGDH inhibitory activities were evaluated using standard curve of NADH. Each concentration of inhibitor was assayed in triplicate.

Cell culture, cell viability assay, and determination of PGE₂ release

HaCaT cell line (Keratinocyte cells) was cultured in Dulbecco's modified Eagle's media (DMEM) supplemented with 10% heat inactivated fetal bovine serum (Sigma) and 100 μ g/mL penicillin. Cells were cultured in a 5% CO₂ atmosphere at 37°C. Cell viabilities were evaluated using MTT assay.^[18] HaCaT cells (1 × 10⁴/90 μ L of DMEM medium) were seeded in 96-well plates. After overnight incubation, plant extract samples were treated for 72 h and after that 10 μ L of MTT (5 mg/mL stock solution) was added into each well, incubated for 4 h at 5% CO₂ atmosphere at 37°C. Medium was then removed and 150 μ L of DMSO was added to dissolve Formazan crystals. Absorbance was measured at 540 nm using ELISA microplate reader (Perkin-Elmer, California, USA). Extracellular PGE₂ releases were determined in HaCaT cell line using various concentrations of five most potent plant extracts as a 15-PGDH inhibitor. HaCaT cells were seeded (5 × 10⁵ cells/well) into six-well culture plates in DMEM,

supplemented with fetal bovine serum and antibiotics for overnight under 5% CO₂, 37°C. Various concentrations of plant extract were added into individual well and media collected after 6 h for extracellular determination and cells were harvested for intracellular determination of PGE₂. Concentrations of PGE₂ were determined using PGE₂ enzyme immunoassay kit (Cayman, MI, USA).

Quantitative real-time PCR

Total cellular RNA was isolated from HaCaT cells according to TRI reagent-specific instruction (RNA iso Plus, Takara Bio. Inc., Shiga, Japan). cDNA was synthesized using reagent-specific instruction of Superscript First Strand synthesis system for reverse transcription-PCR (Invitrogen, Carlsbad, CA, USA) from isolated RNAs. Primers used for real-time PCR were as follows: COX-1 forward, 5'-CCTCATGTTTGCCTTCTTGC-3' and reverse, 5'-GGC GGTACATTTCTCCATC-3' COX-2 forward, 5'-GATCTA CCCTCCTCAA-3' and reverse, 5'-GAACAACCTGC TCATCAC-3' MRP4 forward, 5'-AACCTCTAACCGACATTCCTG-3' and reverse, 5'-TCAACATA TTACAGCCACCATC-3' human PGT forward, 5'-GGATGCTGTTTGGAGGAATCCTCA-3' and reverse, 5'-GCACGATCCTGTC TTGCTGAA-3' and β -actin forward, 5'-GACTATGACTTAGTTGCGTTA-3' and reverse, 5'-GTTGAACCTCTACATAC TTCCG-3'. PCR reaction mixture contained 4 μ L of diluted cDNA (1:5), 10 pmole of each forward and reverse primer, 4 mM MgCl₂, and 4 μ L of Fast Starter Mix buffer (dNTPs, SYBR Green dye and Tag polymerase).

In-vitro wound healing

In-vitro scratch assay was performed in HaCaT cell line to examine wound-healing effect as previously reported^[19,20] using EEAH. HaCaT cells were seeded into six-well plates (5 × 10⁵ cells per well) and grown until reached to 80% confluence. After that media was replaced with serum-free DMEM containing mitomycin (10 μ g/mL) and cells were incubated for 2 h to prevent wound proliferation. Plates were extensively washed with PBS and then scratches were made using sterile 200 μ L pipette tip, and cells were re-washed. TGF- β 1 (100 pg/mL) as a positive control or EEAH were added to the medium. Pictures were taken in the same position before and after incubation to document the wound-healing process. Scratch experiments were repeated thrice and representative pictures are included in this study. Scratches were photographed under microscope (×100) immediately after scratching and also after 48 h of incubation at 37°C in 5% CO₂ incubator.

Statistical analysis

Each experiment was performed at least three times and data are expressed as mean \pm SE. Statistical significance was determined using paired Student's *t*-test and *P* value less than 0.05 were considered statistically significant.

RESULTS

15-PGDH inhibitory activity assay, cytotoxic assay, and *In-vitro* PGE₂ determination

15-PGDH inhibitory activities (IC₅₀ μ g/mL) are illustrated in Supplementary [Table S1]. *Glochidione hirsulta* was the most potent inhibitor for 15-PGDH having 0.36 μ g/mL IC₅₀. Five most potent inhibitors were assayed for cytotoxicity. Relative cytotoxicity was determined *in vitro* by anchorage-dependent cells, HaCaT. Result showed that EEAH was more toxic (cytotoxic IC₅₀ 70 μ g/mL) among tested inhibitor. Further, it is necessary to check whether 15-PGDH inhibitors could increase PGE₂ in cellular or not. Most potent inhibitors were treated for 12 h with concentration of 10 and 100 times of IC₅₀ of

Table S1: Plant extracts with potential 15-PGDH-inhibitory activity

Serial No.*	Plant sample	IC ₅₀ (µg/mL)	Serial No.*	Plant Sample	IC ₅₀ (µg/mL)
1	<i>Glochidion hirsulta</i>	0.37	2	<i>Glochidion velutinum</i>	0.51
3	<i>Cinnamomum tetragonum</i>	0.52	4	<i>Memecylon edule</i>	0.57
5	<i>Artocarpus heterophyllus</i>	0.62	6	<i>Allospodias lakonensis</i>	0.64
7	<i>Commersonia batramia</i>	0.66	8	<i>Syzygium bullockii aff.</i>	0.72
9	<i>Syzygium formosum</i>	0.73	10	<i>Choerospondias axillaris</i>	0.84
11	<i>Lumnitzera racemosa</i>	1.31	12	<i>Osbeckia stellata</i>	1.52
13	<i>Trema orientalis</i>	1.79	14	<i>Lithocarpus gymnocarpus</i>	1.92
15	<i>Planchonella obovata</i>	2.68	16	<i>Triumfetta grandidens</i>	2.87
17	<i>Triumfetta bartramia</i>	2.94	18	<i>Mallotus apelta</i>	3.19
19	<i>Baccaurea ramiflora</i>	3.44	20	<i>Sonneratia alba</i>	4.08
21	<i>Daphniphyllum calycinum</i>	4.10	22	<i>Wandlandia paniculata</i>	4.24
23	<i>Mucuna pruriens</i>	4.59	24	<i>Polygonum hydropiper</i>	4.94
25	<i>Macrosolen ochinchinensis</i>	4.94	26	<i>Lindra myrrha</i>	5.47
27	<i>Melastoma sanguineum</i>	5.62	28	<i>Rhodomyrtus tomentosa</i>	7.56
29	<i>Melastoma normale</i>	7.56	30	<i>Catharanthus roseus</i>	8.51
31	<i>Hibiscus tiliaceus</i>	10.07	32	<i>Mallotus floribundus</i>	10.95
33	<i>Maesa tomentella</i>	11.83	34	<i>Machilus velutina</i>	12.42
35	<i>Archidendron poilanei</i>	13.85	36	<i>Breynia indosinensis</i>	17.53
37	<i>Caesalpinia crista</i>	18.20	38	<i>Hopea odorata</i>	19.36
39	<i>Ipomoea congesta</i>	42.86	40	<i>Cleistanthus tonkinensis</i>	50.70
41	<i>Cratoxylum maingayi</i>	330.30	42	<i>Gouania leptostachya</i>	512.54
43	<i>Engelhardia roxburghina</i>	573.70	44	<i>Machilus thunbergii</i>	916.10
45	<i>Euodia lepta aff.</i>	1,170	46	<i>Senna alata</i>	2,000
47	<i>Vitex tripinnata</i>	2,040	48	<i>Philydrum lanugiosum</i>	2,440
49	<i>Mallotus philippinensis</i>	2,980	50	<i>Pandanus humiulis</i>	3,350
51	<i>Aidia cochinchinensis</i>	3,380	52	<i>Actephila excelsa</i>	4,040
53	<i>Aleurites montata</i>	4,080	54	<i>Streblus asper</i>	4,460
55	<i>Ilex triflora</i>	5,310	56	<i>Vernonia cinerea</i>	6,290
57	<i>Grewia paniculata</i>	6,940	58	<i>Eupatorium odoratum</i>	7,490
59	<i>Broussonetia papyrifera</i>	8,660	60	<i>Carica papaya</i>	8,770
61	<i>Rubus cochinchinensis</i>	9,160	62	<i>Helicteres hirsuta</i>	9,470
63	<i>Ludwigia epilobioides</i>	10,050	64	<i>Desmos chinensis</i>	10,120
65	<i>Gomphandra tonkinensis</i>	10,990	66	<i>Ficus heteropleura</i>	11,370
67	<i>Cerbera manghas</i>	12,500	68	<i>Litsea glutinosa</i>	14,350
69	<i>Helixanthera parasitica</i>	14,740	70	<i>Villebrunea tonkinensis</i>	15,220
71	<i>Memecylon umbellatum</i>	15,520	72	<i>Ageratum conyzoides</i>	15,900
73	<i>Parthenium hysterophorus</i>	15,950	74	<i>Spinifex littoreus</i>	16,480
75	<i>Celosia argentea</i>	20,330	76	<i>Acalypha siamensis</i>	20,980
77	<i>Vitex rotundifolia</i>	22,270	78	<i>Bridelia monoica</i>	22,810
79	<i>Ligustrum sinense</i>	23,680	80	<i>Carallia lanceaefolia</i>	24,430
81	<i>Symplocos cochinchinensis</i>	25,220	82	<i>Leucaena leucocephala</i>	29,960
83	<i>Ficus hirta</i>	32,130	84	<i>Litsea cubeba</i>	57,250
85	<i>Ehretia acuminata</i>	104,200	86	<i>Aidia oxyodonta</i>	154,300
87	<i>Eurya cerasifolia</i>	992,500	88	<i>Alocasia odora</i>	NA
89	<i>Amaranthus tricolor</i>	NA	90	<i>Angelonia goyazensis</i>	NA
91	<i>Aporosa tetrapleura</i>	NA	92	<i>Archidendron clypearia</i>	NA
93	<i>Ardisia quinquegona</i>	NA	94	<i>Embelia laeta</i>	NA
95	<i>Gymnema sylvestre</i>	NA	96	<i>Eurya cerasifolia</i>	NA
97	<i>Kibatalia macrophylla</i>	NA	98	<i>Hyptis brevipes</i>	NA

*Potency order of 15-PGDH-inhibitory activity NA: No activity against 15-PGDH

Table 1: Cytotoxic activities of strong 15-PGDH inhibitors and increment of PGE₂ in HaCaT cells

S. No.*	Inhibitor	Cytotoxicity IC ₅₀ (µg/mL)	% Increment of PGE ₂ of the control	
			10 × IC ₅₀	100 × IC ₅₀
1	<i>Glochidion hirsulta</i>	710	50.90	181.84
2	<i>Glochidion velutinum</i>	270	27.75	274.92
3	<i>Cinnamomum tetragonum</i>	535	39.9	375.65
4	<i>Memecylon edule</i>	450	3.55	265.28
5	<i>Artocarpus heterophyllus</i>	70	42.03	923.29

*Potency order of 15-PGDH-inhibitory activity

Table 2: Intracellular and extracellular PGE₂ levels after 12-h treatment of tested samples (10 × IC₅₀ of 15-PGDH) in HaCaT cells

Sample name	Intracellular (pg/μg) (Mean ± SD)	Extracellular (pg/mL) (Mean ± SD)
Control	1.96 ± 0.18	393.67±12.46
<i>Artocarpus heterophyllus</i>	3.85 ± 0.43*	450.96 ± 15.53*

*Statistically significant $P < 0.05$ **Table 3:** Extracellular PGE₂ levels (pg/mL) influenced by EEAH in HaCaT cells during *in vitro* wound healing

Time	Sample	Control
0 h	Control	143.6
	EEAH	143.6
12 h	Control	420.36
	EEAH	854.62
24 h	Control	442.86
	EEAH	885.62
48 h	Control	323.28
	EEAH	773.05



15-PGDH inhibition and percentage increment in PGE₂ were measured using ELISA kit. They increased PGE₂ in a concentration-dependent manner and EEAH increased PGE₂ level significantly [Table 1]. Results confirmed that tested inhibitors increased PGE₂ level *in vitro* cellular system. Further investigation was concentrated on EEAH.

Regulation of COX-1/2, PGT, and MRP4 by EEAH

PGE₂ level in cellular system might depend on the expression level of COX-1/2, MRP4, and PGT. COX-1/2 is responsible for the production of PGE₂ from PGH₂ and MRP4 and PGT plays important role in the transportation of PGE₂. Relatively low concentration of extracellular PGE₂ compared with intracellular PGE₂ was observed after the treatment with EEAH, suggesting that above-mentioned factors would determine PGE₂ regulation in biologic systems [Table 2]. HaCaT cells were treated with EEAH and regulation of COX-1/2, MRP4, and PGT was evaluated using real-time PCR. Real-time PCR assay showed that EEAH increased expression level of COX-1/2, MRP4, and PGT [Figure 1]. Observed data showed that high intracellular PGE₂ concentration in HaCaT cells may be due to the activation of COX-1 and 15-PGDH inhibition by EEAH.

Wound-healing effect of EEAH

In-vitro wound healing study was performed using HaCaT scratch model. Various experiment sets were designed to evaluate the results: no drug treatment as a negative control, TGF-β1 (100 pg/mL) as a positive control, EEAH (6.2 μg/mL) only and in combination with COX-1/2 inhibitor (SC 560 0.5μM and Celecoxib 0.5μM), respectively. The photographs were taken before treatment and after 2 days incubation at 37°C, 5% CO₂. EEAH facilitated wound healing (recovery distance: 60%) as compared with the negative control (recovery distance: 15%). Wound-healing effect of EEAH was comparable with TGF-β1 (recovery distance 81%). Similarly, COX-1 inhibitor (SC560) prevented the would-healing effect of EEAH partially (recovery distance: 37%); however, COX-2 inhibitor (Celecoxib, recovery distance 40%) did not interfere with wound-healing effect of EEAH. These results suggested that COX pathway as supplier of PGE₂ plays important role in wound healing [Figure 2]. During wound healing, we have also monitored elevation level of PGE₂ on time interval and higher PGE₂ concentration was observed during 12–24 h and slightly reduced at 48 hr [Table 3].

**Figure 1:** Effects of EEAH on mRNA expression of COX-1, COX -2, MRP4, and PGT * Statistically significant $P < 0.05$ **Figure 2:** *In vitro* wound-healing effect of EEAH in HaCaT cells. After scratch being made, immediately photographs were taken and widths were measured. Scratched cells were incubated for 2 days in 5% CO₂ incubator, the picture was again captured at the same place and widths were measured. Average width of wound was calculated with longest, medium and shortest length between the both cell populations. Control, no drug treatment; TGF-β1 (100 pg/mL), positive control; EEAH, ethanol extracts of *Artocarpus heterophyllus* (6.2 μg/mL); SC 560 (0.5 μM); C, Celecoxib (0.5 μM)

DISCUSSION

Plant as a source of medicine used for traditional healthcare system for several thousands of years.^[21] New drug development scientists have been focusing their studies on medicinal plants in different parts of the world.^[22,23] There are significant economic benefits in the development of new medicines from plants for the treatment of various diseases.^[24] Twenty-five percent of medicines belong to plants source and their derivatives.^[25] Plants as a source of traditional medicine are used to cure diseases involving skin problems, cold, fever, cough, headache, diarrhea, fertility problems, and toothache.^[26] Therefore, 15-PGDH inhibitor was investigated in this study because of diverse therapeutic values and ample of opportunity to develop new drug from plant source. 15-PGDH negatively regulates PGE₂ levels and activity *in vivo*.^[27,28] In this study, extracellular and intracellular PGE₂ levels were elevated in HaCaT cell line after treatment of 15-PGDH inhibitors Table 1 and

Table 2. The levels of PGE₂ available to cells are dependent on function of COX-1/2, MRP4, and PGT in PGE₂ signaling. EEAH elevate COX-1 mRNA expression significantly, while MRP mRNA expression slightly, which helps to increase availability of PGE₂ levels in HaCaT cell line [Figure 1]. Recently, Kochel and Fulton reported that MRP4, PGT, and 15-PGDH play important role in regulating PGE₂ levels.^[29]

Previously, several scientists have reported wound-healing efficacy of plant extracts.^[30-33] This study investigated the role of PGE₂ during the wound-healing process that was induced by plant extract. Tissue regeneration plays important role during recovery from injury, including wound healing, and PGE₂ is candidate molecule that helps in regeneration of tissue.^[34] After scratching, extracellular medium was collected on different time interval (0, 12, 24, and 48 h) and PGE₂ concentration was measured. When wound being, the activity of PGE₂ was increased because there was increase on extracellular PGE₂ concentration during 12 and 24 h in all the tested samples because of inflammatory and proliferative actions on wound bed. COX-2 expression and PGE₂ production in the wound bed increase as dermal wound healing transitions from scarless to a scarring phenotype.^[12] At a cellular level, PGE₂ was shown to regulate the proliferation of fibroblasts from both wound-healing phenotypes. Hemostasis, inflammatory, proliferative, and remodeling are the major steps of wound healing, where PGE₂ plays an important role.^[35] EEAH plays important role to elevate PGE₂ level in cellular system by inhibiting 15-PGDH that accelerates wound healing.

CONCLUSIONS

15-PGDH inhibitor alone will give better result of wound healing than COX inhibitors because of elevated level of PGE₂. Therefore, inhibition of 15-PGDH with plant extracts will be valuable for the therapeutic management of diseases requiring elevated PGE₂ levels like in wound healing. The clinical efficacy and safety of these plant extracts as well as the purification of active ingredients from these plants remains to be done.

Acknowledgement

Plant samples and 15-PGDH plasmid were kindly gifted by Cheol-Hee Choi, Chosun University, Gwangju, Republic of Korea.

Financial support and sponsorship

Nil.

Conflicts of interest

There is no conflict of interests

REFERENCES

- Griffiths R. Prostaglandins and inflammation. In: Inflammation: basic principles and clinical correlates. Philadelphia, PA: Lippincott Williams and Wilkins 1999;349-60.
- Crofford LJ. COX-1 and COX-2 tissue expression: implications and predictions. J Rheumatol Suppl 1997;49:15-9.
- Schuster VL. Prostaglandin transport Prostaglandins Other Lipid Mediat. 2002;68:633-47.
- Schuster VL. Molecular mechanisms of prostaglandin transport. Annu Rev Physiol 1998;60:221-42.
- Anggard E, Samuelsson B. Smooth muscle stimulating lipids in sheep iris. The identification of prostaglandin F 2a. Prostaglandins and related Factors 21. Biochem Pharmacol 1964;13:281-3.
- Anggard E. The biological activities of three metabolites of prostaglandin E 1. Acta Physiol Scand 1966;66:509-10.
- Li M, Thompson DD, Paralkar VM. Prostaglandin E(2) receptors in bone formation. Int Orthop 2007;31:767-72.
- Keller J, Klamer A, Bak B, Suder P. Effect of local prostaglandin E2 on fracture callus in rabbits. Acta Orthop Scand 1993;64:59-63.
- Hatazawa R, Tanaka A, Tanigami M, Amagase K, Kato S, Ashida Y, et al. Cyclooxygenase-2/

- prostaglandin E2 accelerates the healing of gastric ulcers via EP4 receptors. Am J Physiol Gastrointest Liver Physiol 2007;293:G788-97.
- Shigeta J, Takahashi S, Okabe S. Role of cyclooxygenase-2 in the healing of gastric ulcers in rats. J Pharmacol Exp Ther 1998;286:1383-90.
- Parekh A, Sandulache VC, Singh T, Cetin S, Sacks MS, Dohar JE, et al. Prostaglandin E2 differentially regulates contraction and structural reorganization of anchored collagen gels by human adult and fetal dermal fibroblasts. Wound Repair Regen 2009;17:88-98.
- Wilgus TA, Bergdall VK, Tober KL, Hill KJ, Mitra S, Flavahan NA, et al. The impact of cyclooxygenase-2 mediated inflammation on scarless fetal wound healing. Am J Pathol 2004;165:753-61.
- Fung WP, Karim SM. Effect of 15 (R) 15 methyl prostaglandin E2 on the healing of gastric ulcers: a double-blind endoscopic study. Med J Aust 1976;2:127-8.
- Gibinski K, Rybicka J, Mikos E, Nowak A. Double-blind clinical trial on gastroduodenal ulcer healing with prostaglandin E2 analogues. Gut 1977;18:636-9.
- Nguyen DT, Orgill DP, Murphy GF. The pathophysiologic basis for wound healing and cutaneous regeneration. 2009;In: Biomaterials for treating skin loss Chapter 4 Cambridge/ Boca Raton: Woodhead Publishing (UK/Europe) and CRC Press (US).
- Stadelmann WK, Digenis AG, Tobin GR. Physiology and healing dynamics of chronic cutaneous wounds. Am J Surg 1998;176:26S-38S.
- ENSOR CM, Yang JY, Okita RT, Tai HH. Cloning and sequence analysis of the cDNA for human placental NAD(+)-dependent 15-hydroxyprostaglandin dehydrogenase. J Biol Chem 1990;265:14888-91.
- Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. J Immunol Methods 1983;65:55-63.
- Hintermann E, Bilban M, Sharabi A, Quaranta V. Inhibitory role of alpha 6 beta 4-associated erbB-2 and phosphoinositide 3-kinase in keratinocyte haptotactic migration dependent on alpha 3 beta 1 integrin. J Cell Biol 2001;153:465-78.
- Koivisto L, Jiang G, Hakkinen L, Chan B, Larjava H. HaCaT keratinocyte migration is dependent on epidermal growth factor receptor signaling and glycogen synthase kinase-3alpha. Exp Cell Res 2006;312:2791-805.
- Abu-Rabia A. Urinary diseases and ethnobotany among pastoral nomads in the Middle East. J Ethnobiol Ethnomed 2005;1:4.
- Gazzaneo LR, de Lucena RF, de Albuquerque UP. Knowledge and use of medicinal plants by local specialists in an region of Atlantic Forest in the state of Pernambuco (Northeastern Brazil). J Ethnobiol Ethnomed 2005;1:9.
- Hanazaki N, Tamashiro JY, Leitao-Filho H, Gegossi A. Diversity of plant uses in two Caicaras communities from the Atlantic forest coast, Brazil. Biodivers Conserv 2000;9:597-615.
- Azaizeh H, Fulder S, Khalil K, Said O. Ethnobotanical knowledge of local Arab practitioners in the Middle Eastern region. Fitoterapia 2003;74:98-108.
- Principe P, Riley PA, Slater TF. Cellular thiols in rat liver cell lines possessing different growth characteristics. Cell Biochem Funct 1991;9:125-33.
- Muthu C, Ayyanar M, Raja N, Ignacimuthu S. Medicinal plants used by traditional healers in Kancheepuram district of Tamil Nadu, India. J Ethnobiol Ethnomed 2006;2:43.
- Tai HH, Ensor CM, Tong M, Zhou H, Yan F. Prostaglandin catabolizing enzymes Prostaglandins Other Lipid Mediat. 2002;68-69:483-93.
- Myung SJ, Rerko RM, Yan M, Platzer P, Guda K, Dotson A, et al. 15-Hydroxyprostaglandin dehydrogenase is an *in vivo* suppressor of colon tumorigenesis. Proc Natl Acad Sci USA 2006;103:12098-102.
- Kochel TJ, Fulton AM. Multiple drug resistance-associated protein 4 (MRP4), prostaglandin transporter (PGT), and 15-hydroxyprostaglandin dehydrogenase (15-PGDH) as determinants of PGE₂ levels in cancer. Prostaglandins Other Lipid Mediat 2015;116-117:99-103.
- Shailajan S, Gurjar D. Wound healing activity of *Chrysophyllum cainito* L leaves: evaluation in rats using excision wound model. J Young Pharm 2016;8:96-103.
- Shailajan S, Gurjar D. Wound healing efficacy of *Mimusops elengi* L flowers in albino Wistar rats using excision wound model. Pharmacogn Commn 2016;6:72-9.
- Hasamnis AA, Mohanty BK, Murali krishna, Patil S. Evaluation of wound healing effect of topical phenytoin on excisional wound in albino rats. J Young Pharm 2010;2:59-6.
- Gutierrez RMP, Soils RV. Anti-inflammatory and wound healing potential of *Prosthechea michuacana* in rats. Pharmacog Mag 2009;5:219-25.
- Zhang Y, Desai A, Yang SY, Bae KB, Antczak MI, Fink SP, et al. Tissue regeneration. Inhibition of the prostaglandin-degrading enzyme 15-PGDH potentiates tissue regeneration. Science 2015;348:aaa2340.
- Quinn JV. Tissue adhesives in wound care. 1998;Hamilton, ONB.C Decker, Inc.