

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

rac-2-Iodo-3,4-dihydronaphthalen-1(2*H*)-one

Abdul Rauf Raza,^a M. Nawaz Tahir,^{b*} Ayesha Sultan,^a Muhammad Danish^a and Muhammad Sohail^a

^aDepartment of Chemistry, University of Sargodha, Sargodha, Pakistan, and

^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 17 November 2009; accepted 18 November 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; disorder in main residue; R factor = 0.024; wR factor = 0.059; data-to-parameter ratio = 18.5.

In the title compound, $\text{C}_{10}\text{H}_9\text{IO}$, the asymmetric unit contains two molecules, in which the iodo-bearing six-membered rings adopt envelope conformations [displacements of the flap atoms = 0.419 (3) and 0.431 (3) Å]. In both molecules, the I atoms are disordered over two set of sites in 0.54 (4):0.46 (4) and 0.71 (3):0.29 (3) ratios. In the crystal, the packing features a weak $\text{C}-\text{H}\cdots\pi$ interaction.

Related literature

For a related structure, see: Haddad (1986).



Experimental

Crystal data

$\text{C}_{10}\text{H}_9\text{IO}$
 $M_r = 272.07$

Monoclinic, $P2_1/c$
 $a = 6.115$ (5) Å

$b = 19.658$ (4) Å
 $c = 15.896$ (5) Å
 $\beta = 90.551$ (5)°
 $V = 1910.7$ (17) Å³
 $Z = 8$

Mo $K\alpha$ radiation
 $\mu = 3.31$ mm⁻¹
 $T = 296$ K
 $0.28 \times 0.20 \times 0.18$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.685$, $T_{\max} = 0.717$

19043 measured reflections
4397 independent reflections
3632 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.024$
 $wR(F^2) = 0.059$
 $S = 1.07$
4397 reflections

238 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.39$ e Å⁻³
 $\Delta\rho_{\min} = -0.50$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C5}-\text{H5}\cdots\text{Cg3}$	0.93	2.95	3.700 (4)	139

Cg3 is the centroid of the $\text{C11}-\text{C16}$ ring.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5236).

References

- Bruker (2005). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Haddad, S. F. (1986). *Acta Cryst.* **C42**, 581–584.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o3172 [doi:10.1107/S1600536809049095]

***rac*-2-Iodo-3,4-dihydronaphthalen-1(2*H*)-one**

A. R. Raza, M. N. Tahir, A. Sultan, M. Danish and M. Sohail

Comment

The title compound (I, Fig. 1) is an intermediate for the total synthesis of steroidal hormones. The crystal structures of (II) 2,2-Dibromo-3,4-dihydro-1(2*H*)-naphthalenone (Haddad, 1986) has been published, which seems relevant to (I).

The asymmetric unit of title compound consists of two individual molecules which are clearly racemate. In the molecule having (*S*)-configuration, the I-atom containing ring A (C1/C6/C7—C10) is twisted with maximum puckering amplitude $Q_T = 0.431(3) \text{ \AA}$, whereas in (*R*)-configuration the puckering parameter is $Q_T = 0.419(3) \text{ \AA}$. In two molecules the groups of benzene rings along with two adjacent C-atoms, C (C1—C6/C7/C10) and D (C11—C16/C17/C20) are planar with maximum r. m. s. deviations of 0.0114 and 0.0280 \AA respectively, from the respective mean square planes. The dihedral angle between C/D is 66.83(7) \AA . In the first molecule the I-atom is disordered over two set of sites having occupancy ratio of 0.54(4):0.46(4). Similarly in the other molecule the I-atom is disordered over two set of sites having occupancy ratio of 0.71(3):0.29(3). The molecules are stabilized due to C—H $\cdots\pi$ interactions (Table 1).

Experimental

A solution of I₂ (7.75 g, 30.5 mmol) in CHCl₃ was added as drops to a solution of 1-tetralone (2.198 g, 15.2 mmol) in acetic acid (9.156 g, 0.1526 mol) and refluxed for 28 h. The H₂O (30 ml) was added for partitioning. The reaction mixture was extracted with CHCl₃ (3 \times 15 ml). The combined organic layer was concentrated *in vacuo*, the crude was dissolved in ethyl acetate, washed with 5% Na₂S₂O₃ (2 \times 15 ml), dried over anhydrous Na₂SO₄, filtered, boiled with charcoal, concentrated under reduce pressure and allowed for crystallization, which afforded colourless prisms (89%) of (I).

Refinement

The other H-atoms were positioned geometrically (C—H = 0.93–0.98 \AA) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. View of (I) with the I atoms having greater occupancies. The displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. View of (I) with the I atoms having lesser occupancies. The displacement ellipsoids are drawn at the 30% probability level.

rac-2-Iodo-3,4-dihydronaphthalen-1(2H)-one

Crystal data

$C_{10}H_9IO$	$F(000) = 1040$
$M_r = 272.07$	$D_x = 1.892 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: $-P 2_1/c$	Cell parameters from 4397 reflections
$a = 6.115 (5) \text{ \AA}$	$\theta = 2.4\text{--}27.8^\circ$
$b = 19.658 (4) \text{ \AA}$	$\mu = 3.31 \text{ mm}^{-1}$
$c = 15.896 (5) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 90.551 (5)^\circ$	Prism, colourless
$V = 1910.7 (17) \text{ \AA}^3$	$0.28 \times 0.20 \times 0.18 \text{ mm}$
$Z = 8$	

Data collection

Bruker Kappa APEXII CCD diffractometer	4397 independent reflections
Radiation source: fine-focus sealed tube graphite	3632 reflections with $I > 2\sigma(I)$
Detector resolution: $7.50 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.026$
ω scans	$\theta_{\text{max}} = 27.8^\circ$, $\theta_{\text{min}} = 2.4^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$h = -8 \rightarrow 7$
$T_{\text{min}} = 0.685$, $T_{\text{max}} = 0.717$	$k = -25 \rightarrow 24$
19043 measured reflections	$l = -20 \rightarrow 20$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.024$	H-atom parameters constrained
$wR(F^2) = 0.059$	$w = 1/[\sigma^2(F_o^2) + (0.0273P)^2 + 0.6672P]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
4397 reflections	$(\Delta/\sigma)_{\text{max}} = 0.002$
238 parameters	$\Delta\rho_{\text{max}} = 0.39 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.50 \text{ e \AA}^{-3}$

0 restraints
 Extinction correction: *SHELXL97* (Sheldrick, 2008),
 $F_c^* = kF_c [1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Primary atom site location: structure-invariant direct methods
 Extinction coefficient: 0.00283 (17)

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	U_{iso}^*/U_{eq}	Occ. (<1)
I1A	0.8199 (5)	0.0828 (2)	0.36895 (19)	0.0502 (6)	0.54 (4)
I1B	0.8299 (7)	0.0783 (3)	0.3658 (2)	0.0622 (8)	0.46 (4)
O1	0.6627 (3)	0.25061 (10)	0.31382 (12)	0.0512 (6)	
C1	0.9910 (4)	0.17057 (12)	0.15222 (17)	0.0408 (8)	
C2	0.9965 (5)	0.16572 (14)	0.06450 (19)	0.0573 (10)	
C3	0.8386 (6)	0.19473 (15)	0.01454 (19)	0.0637 (10)	
C4	0.6694 (6)	0.23029 (15)	0.05043 (19)	0.0621 (11)	
C5	0.6596 (4)	0.23677 (13)	0.13679 (17)	0.0473 (9)	
C6	0.8181 (3)	0.20662 (11)	0.18883 (15)	0.0350 (7)	
C7	0.8027 (3)	0.21515 (11)	0.28175 (15)	0.0356 (7)	
C8	0.9707 (4)	0.17888 (13)	0.33563 (17)	0.0439 (8)	
C9	1.1881 (4)	0.16933 (15)	0.2919 (2)	0.0546 (9)	
C10	1.1619 (4)	0.13632 (14)	0.20632 (19)	0.0518 (9)	
I2A	0.2833 (4)	0.51127 (14)	0.4019 (2)	0.0524 (4)	0.71 (3)
I2B	0.2948 (9)	0.5150 (3)	0.3949 (5)	0.0551 (13)	0.29 (3)
O2	0.1573 (3)	0.33920 (10)	0.34826 (14)	0.0590 (7)	
C11	0.4861 (4)	0.42382 (11)	0.19085 (17)	0.0395 (8)	
C12	0.4975 (5)	0.42831 (14)	0.1038 (2)	0.0593 (11)	
C13	0.3390 (7)	0.39932 (18)	0.0525 (2)	0.0735 (13)	
C14	0.1666 (6)	0.36502 (18)	0.0870 (2)	0.0725 (12)	
C15	0.1531 (4)	0.35857 (13)	0.1724 (2)	0.0528 (10)	
C16	0.3121 (3)	0.38740 (11)	0.22589 (16)	0.0366 (7)	
C17	0.2928 (4)	0.37754 (11)	0.31794 (16)	0.0385 (7)	
C18	0.4509 (4)	0.41620 (12)	0.37374 (17)	0.0431 (8)	
C19	0.6704 (4)	0.42740 (14)	0.33282 (19)	0.0504 (9)	
C20	0.6513 (4)	0.45891 (14)	0.24640 (19)	0.0496 (9)	
H2	1.11050	0.14215	0.03943	0.0687*	
H3	0.84562	0.19042	-0.04364	0.0763*	
H4	0.56189	0.24993	0.01653	0.0746*	
H5	0.54622	0.26144	0.16068	0.0568*	

supplementary materials

H8	0.99519	0.20536	0.38710	0.0527*
H9A	1.25810	0.21327	0.28537	0.0654*
H9B	1.28268	0.14126	0.32686	0.0654*
H10A	1.30098	0.13765	0.17757	0.0620*
H10B	1.12210	0.08895	0.21380	0.0620*
H12	0.61398	0.45126	0.07953	0.0712*
H13	0.34957	0.40313	-0.00567	0.0882*
H14	0.05908	0.34619	0.05236	0.0868*
H15	0.03679	0.33471	0.19547	0.0632*
H18	0.47290	0.39074	0.42614	0.0517*
H19A	0.75860	0.45680	0.36852	0.0605*
H19B	0.74537	0.38409	0.32833	0.0605*
H20A	0.61044	0.50634	0.25223	0.0596*
H20B	0.79302	0.45732	0.21957	0.0596*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1A	0.0533 (13)	0.0521 (9)	0.0453 (9)	-0.0069 (5)	0.0019 (5)	0.0139 (8)
I1B	0.084 (2)	0.0428 (9)	0.0599 (13)	-0.0067 (6)	0.0130 (10)	-0.0018 (11)
O1	0.0456 (10)	0.0531 (11)	0.0550 (12)	0.0073 (8)	0.0063 (8)	-0.0161 (9)
C1	0.0450 (12)	0.0273 (12)	0.0504 (16)	-0.0049 (9)	0.0138 (11)	0.0004 (11)
C2	0.0773 (19)	0.0391 (15)	0.0561 (19)	-0.0096 (13)	0.0319 (16)	-0.0049 (13)
C3	0.102 (2)	0.0490 (17)	0.0402 (16)	-0.0163 (17)	0.0091 (16)	0.0025 (14)
C4	0.083 (2)	0.0521 (18)	0.0510 (19)	-0.0057 (15)	-0.0130 (16)	0.0118 (14)
C5	0.0502 (14)	0.0387 (14)	0.0530 (17)	0.0023 (11)	-0.0036 (12)	0.0000 (12)
C6	0.0370 (11)	0.0243 (11)	0.0439 (14)	-0.0035 (9)	0.0054 (10)	-0.0011 (10)
C7	0.0322 (11)	0.0297 (12)	0.0448 (14)	-0.0045 (9)	0.0018 (10)	-0.0046 (10)
C8	0.0442 (13)	0.0389 (14)	0.0485 (15)	-0.0046 (10)	-0.0082 (11)	-0.0064 (11)
C9	0.0346 (12)	0.0499 (16)	0.079 (2)	-0.0019 (11)	-0.0078 (12)	0.0078 (15)
C10	0.0379 (13)	0.0419 (15)	0.076 (2)	0.0064 (10)	0.0160 (13)	0.0039 (14)
I2A	0.0475 (8)	0.0585 (9)	0.0512 (5)	0.0129 (3)	-0.0063 (3)	-0.0177 (5)
I2B	0.077 (3)	0.0322 (19)	0.0562 (13)	0.0131 (8)	0.0055 (13)	-0.0043 (12)
O2	0.0550 (11)	0.0514 (12)	0.0709 (14)	-0.0111 (9)	0.0206 (10)	0.0053 (10)
C11	0.0416 (12)	0.0274 (12)	0.0496 (16)	0.0011 (9)	0.0093 (11)	-0.0039 (10)
C12	0.0745 (19)	0.0488 (17)	0.0551 (19)	-0.0020 (14)	0.0204 (15)	0.0010 (14)
C13	0.105 (3)	0.071 (2)	0.0446 (18)	0.006 (2)	0.0004 (18)	-0.0078 (16)
C14	0.086 (2)	0.064 (2)	0.067 (2)	-0.0004 (18)	-0.0220 (19)	-0.0170 (17)
C15	0.0511 (15)	0.0381 (14)	0.069 (2)	-0.0081 (11)	-0.0073 (13)	-0.0075 (13)
C16	0.0355 (11)	0.0259 (12)	0.0484 (15)	0.0012 (9)	0.0025 (10)	-0.0033 (10)
C17	0.0356 (11)	0.0283 (12)	0.0517 (15)	0.0032 (9)	0.0071 (10)	0.0012 (10)
C18	0.0445 (13)	0.0427 (14)	0.0420 (15)	0.0102 (10)	-0.0023 (11)	0.0008 (11)
C19	0.0341 (12)	0.0504 (16)	0.0665 (19)	0.0033 (10)	-0.0068 (12)	-0.0087 (13)
C20	0.0347 (12)	0.0433 (15)	0.071 (2)	-0.0092 (10)	0.0103 (12)	-0.0039 (13)

Geometric parameters (\AA , $^\circ$)

I1A—C8	2.170 (5)	C9—H9B	0.9700
I1B—C8	2.211 (6)	C10—H10A	0.9700

I2A—C18	2.180 (4)	C10—H10B	0.9700
I2B—C18	2.192 (7)	C11—C12	1.389 (4)
O1—C7	1.220 (3)	C11—C16	1.402 (3)
O2—C17	1.223 (3)	C11—C20	1.503 (4)
C1—C6	1.404 (3)	C12—C13	1.384 (5)
C1—C2	1.398 (4)	C13—C14	1.370 (5)
C1—C10	1.506 (4)	C14—C15	1.367 (5)
C2—C3	1.369 (5)	C15—C16	1.405 (4)
C3—C4	1.377 (5)	C16—C17	1.482 (4)
C4—C5	1.381 (4)	C17—C18	1.511 (4)
C5—C6	1.400 (4)	C18—C19	1.513 (4)
C6—C7	1.490 (4)	C19—C20	1.510 (4)
C7—C8	1.510 (4)	C12—H12	0.9300
C8—C9	1.518 (4)	C13—H13	0.9300
C9—C10	1.514 (4)	C14—H14	0.9300
C2—H2	0.9300	C15—H15	0.9300
C3—H3	0.9300	C18—H18	0.9800
C4—H4	0.9300	C19—H19A	0.9700
C5—H5	0.9300	C19—H19B	0.9700
C8—H8	0.9800	C20—H20A	0.9700
C9—H9A	0.9700	C20—H20B	0.9700
C2—C1—C6	118.2 (2)	C12—C11—C16	118.3 (2)
C2—C1—C10	121.1 (2)	C12—C11—C20	121.1 (2)
C6—C1—C10	120.7 (2)	C16—C11—C20	120.6 (2)
C1—C2—C3	121.8 (3)	C11—C12—C13	121.3 (3)
C2—C3—C4	120.0 (3)	C12—C13—C14	120.3 (3)
C3—C4—C5	119.9 (3)	C13—C14—C15	119.8 (3)
C4—C5—C6	120.8 (2)	C14—C15—C16	121.0 (3)
C1—C6—C5	119.3 (2)	C11—C16—C15	119.3 (2)
C1—C6—C7	121.5 (2)	C11—C16—C17	121.8 (2)
C5—C6—C7	119.21 (19)	C15—C16—C17	118.9 (2)
O1—C7—C6	122.0 (2)	O2—C17—C16	122.1 (2)
O1—C7—C8	120.6 (2)	O2—C17—C18	120.7 (2)
C6—C7—C8	117.38 (18)	C16—C17—C18	117.2 (2)
I1A—C8—C7	105.13 (17)	I2A—C18—C17	104.62 (17)
I1A—C8—C9	112.4 (2)	I2A—C18—C19	112.60 (18)
I1B—C8—C7	106.32 (18)	I2B—C18—C17	105.0 (2)
I1B—C8—C9	109.5 (2)	I2B—C18—C19	109.1 (2)
C7—C8—C9	113.1 (2)	C17—C18—C19	112.7 (2)
C8—C9—C10	112.3 (2)	C18—C19—C20	112.9 (2)
C1—C10—C9	112.9 (2)	C11—C20—C19	113.1 (2)
C1—C2—H2	119.00	C11—C12—H12	119.00
C3—C2—H2	119.00	C13—C12—H12	119.00
C2—C3—H3	120.00	C12—C13—H13	120.00
C4—C3—H3	120.00	C14—C13—H13	120.00
C3—C4—H4	120.00	C13—C14—H14	120.00
C5—C4—H4	120.00	C15—C14—H14	120.00
C4—C5—H5	120.00	C14—C15—H15	119.00
C6—C5—H5	120.00	C16—C15—H15	119.00

supplementary materials

I1A—C8—H8	109.00	I2A—C18—H18	109.00
I1B—C8—H8	111.00	I2B—C18—H18	112.00
C7—C8—H8	109.00	C17—C18—H18	109.00
C9—C8—H8	109.00	C19—C18—H18	109.00
C8—C9—H9A	109.00	C18—C19—H19A	109.00
C8—C9—H9B	109.00	C18—C19—H19B	109.00
C10—C9—H9A	109.00	C20—C19—H19A	109.00
C10—C9—H9B	109.00	C20—C19—H19B	109.00
H9A—C9—H9B	108.00	H19A—C19—H19B	108.00
C1—C10—H10A	109.00	C11—C20—H20A	109.00
C1—C10—H10B	109.00	C11—C20—H20B	109.00
C9—C10—H10A	109.00	C19—C20—H20A	109.00
C9—C10—H10B	109.00	C19—C20—H20B	109.00
H10A—C10—H10B	108.00	H20A—C20—H20B	108.00
C6—C1—C2—C3	-0.3 (4)	C16—C11—C12—C13	1.7 (4)
C10—C1—C2—C3	177.9 (3)	C20—C11—C12—C13	-176.3 (3)
C2—C1—C6—C5	-0.6 (3)	C12—C11—C16—C15	-1.8 (3)
C2—C1—C6—C7	-178.9 (2)	C12—C11—C16—C17	177.3 (2)
C10—C1—C6—C5	-178.8 (2)	C20—C11—C16—C15	176.2 (2)
C10—C1—C6—C7	2.9 (3)	C20—C11—C16—C17	-4.7 (3)
C2—C1—C10—C9	156.0 (2)	C12—C11—C20—C19	-157.1 (2)
C6—C1—C10—C9	-25.8 (3)	C16—C11—C20—C19	25.0 (3)
C1—C2—C3—C4	0.5 (5)	C11—C12—C13—C14	-0.3 (5)
C2—C3—C4—C5	0.2 (5)	C12—C13—C14—C15	-1.0 (5)
C3—C4—C5—C6	-1.1 (4)	C13—C14—C15—C16	0.9 (5)
C4—C5—C6—C1	1.3 (4)	C14—C15—C16—C11	0.6 (4)
C4—C5—C6—C7	179.6 (2)	C14—C15—C16—C17	-178.6 (3)
C1—C6—C7—O1	174.4 (2)	C11—C16—C17—O2	-171.0 (2)
C1—C6—C7—C8	-4.7 (3)	C11—C16—C17—C18	8.1 (3)
C5—C6—C7—O1	-4.0 (3)	C15—C16—C17—O2	8.1 (3)
C5—C6—C7—C8	177.1 (2)	C15—C16—C17—C18	-172.9 (2)
O1—C7—C8—I1A	87.3 (2)	O2—C17—C18—I2A	-89.8 (2)
O1—C7—C8—C9	-149.8 (2)	O2—C17—C18—C19	147.6 (2)
C6—C7—C8—I1A	-93.7 (2)	C16—C17—C18—I2A	91.2 (2)
C6—C7—C8—C9	29.3 (3)	C16—C17—C18—C19	-31.5 (3)
I1A—C8—C9—C10	66.6 (3)	I2A—C18—C19—C20	-65.9 (3)
C7—C8—C9—C10	-52.2 (3)	C17—C18—C19—C20	52.1 (3)
C8—C9—C10—C1	50.1 (3)	C18—C19—C20—C11	-48.6 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C5—H5 \cdots Cg3	0.93	2.95	3.700 (4)	139

Fig. 1




Fig. 2

