

Tumor-derived VEGF modulates hematopoiesis

Yuan Xue^{†1}, Fang Chen^{†2}, Danfang Zhang¹, Sharon Lim¹ and Yihai Cao^{*1}

Address: ¹Department of Microbiology, Tumor and Cell Biology, Karolinska Institute 171 77 Stockholm, Sweden and ²Shanghai Chest Hospital Affiliated to Shanghai Jiaotong University, 200030 Shanghai, PR China

Email: Yuan Xue - yuan.xue@ki.se; Fang Chen - chenfang1975@hotmail.com; Danfang Zhang - danfangzhang@hotmail.com; Sharon Lim - sharon.lim@ki.se; Yihai Cao* - yihai.cao@ki.se

* Corresponding author †Equal contributors

Published: 23 December 2009

Received: 1 September 2009

Journal of Angiogenesis Research 2009, **1**:9 doi:10.1186/2040-2384-1-9

Accepted: 23 December 2009

This article is available from: <http://www.jangiogenesis.com/content/1/1/9>

© 2009 Xue et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

VEGF-induced angiogenesis significantly contributes to tumor growth, invasion and metastasis. However, little is known about its hematopoietic activity during malignant development and progression. Here we show that in a mouse tumor model, tumor-derived VEGF acts as an endocrine-like hormone to induce extramedullary hematopoiesis by targeting distal organs in the host. In tumor-bearing mice, circulating VEGF induced hepatomegaly and splenomegaly owing to vessel dilation, tortuosity and activation of hematopoiesis. Furthermore, VEGFR1 and VEGFR2 were primarily localized in blood vessels rather than hepatocytes or splenocytes, demonstrating that alteration of angiogenic profiles modulates hematopoiesis in these organs. Stimulation of extramedullary hematopoiesis sheds new light on complex biological functions of VEGF and significantly increases our understanding of molecular mechanisms underlying VEGF-induced tumor growth.

Introduction

Malignant tissues produce multiple growth factors and cytokines to induce angiogenesis, which is essential for tumor growth, invasion and metastasis [1,2]. Among tumor-derived angiogenic factors, vascular endothelial growth factor (VEGF) is probably one of the best-characterized molecules. VEGF displays multiple physiological and pathological functions by targeting both vascular and non-vascular systems [3-5]. In developing embryos, deletion of only one allele of the *veg* gene results in severe defects of early embryos that manifest impaired hematopoiesis and collapse of the vascular system [6,7]. In the adult, VEGF is required to maintain the integrity of the vasculature and physiological functions, including endothelial survival, vascular fenestration in several endocrine glands, neurotrophic effects, support of bone marrow hematopoiesis, wound healing, and reproductive

activity [8,9]. To maintain these multiple physiological functions, optimal levels of VEGF expression are required in various tissues and organs [10,11]. When optimal expression levels are altered, VEGF often causes pathological disorders by triggering uncontrolled vascular responses that include pathological vasculogenesis, angiogenesis, and tissue edema. The plasticity features of VEGF expression determine its involvement in a broad spectrum of human diseases including malignant and non-malignant disorders. For example, tissue hypoxia is one of the key factors that elevate VEGF expressions in both tumors and non-malignant disorders [12,13].

In tumors, VEGF is known to significantly contribute to pathological angiogenesis, tortuosity of tumor vasculatures and vasculogenesis, which all together lead to accelerated growth rates of tumors, invasion and metastasis

[14]. In addition to vascular effects, VEGF also mobilizes mononucleic cells and probably endothelial progenitor cells from bone marrow, whereas former enhances tumor inflammation, the latter participates in vasculogenesis [15,16]. VEGF-induced vascular tortuosity and leakiness also provide a structural basis for tumor cell invasion into the circulation system, leading to distal metastasis. In addition to hematogenous metastasis, recent work from our laboratory and others demonstrates that VEGF is also a potent lymphangiogenic factor, which promotes lymphatic metastasis [17-20].

Similar to the prototype member of VEGF, other members in the VEGF family exhibit overlapping and sometimes distinctive biological functions in both physiological and pathological settings, depending on their abilities to activate a subset of VEGF receptors. While VEGFR2 is consistently reported as a functional receptor to transduce angiogenic, vasculogenic and vascular permeability activities, functional properties of VEGFR1 remain controversial [21]. Probably, VEGFR-1 transduces both positive and negative signals in endothelial and non-endothelial cells, depending on the choice of experimental settings [21]. Additionally, high molecular forms of VEGF also bind to neuropilin-1 and -2, which are involved in tumor growth and metastasis [22,23].

Owing to the pivotal role of VEGF in regulation of pathological angiogenesis and tumor growth, several anti-VEGF drugs have been developed for cancer therapy. These include neutralization of the VEGF ligand by antibodies such as bevacizumab and small chemical compounds targeting the receptor signaling pathways such as sorafenib and sunitinib [24]. In general, clinical responses to these drugs in combinations with chemotherapy are very encouraging and they become one of the key components of the first-line therapeutic regimens for various human cancers [25]. In the present study, we provide new evidence that VEGF could induce extramedullary hemetapoiesis in adult tumor-bearing mice. These findings suggest that VEGF may significantly contribute to tumor growth via improvement of hematopoiesis

Methods

Animals and reagents

Female C57Bl/6 mice were anesthetized by Isoflurane (Abbott Scandinavia) before all procedures. The experiments were followed up to 4 weeks. Mice were sacrificed by exposure to a lethal dose of CO₂ followed by cervical dislocation. All animal studies were reviewed and approved by the animal care committee of the North Stockholm Animal Board (Stockholm, Sweden). Antibodies include a rat anti-mouse CD31 monoclonal antibody, a rat anti-mouse erythroid cells (Ter119) monoclonal antibody (BD Pharmingen, San Diego, CA, USA), a rat

anti-mouse VEGFR1 (MF1) antibody, a rat anti-mouse VEGFR2 (DC101) antibody (ImClone Systems Inc., NY, USA), and a rabbit anti-mouse polyclonal VEGFR-2 antibody (kindly provided by Dr. Rolf Brekken at University of Texas Southwestern Medical Center).

Cell culture

Murine T241 fibrosarcoma is transfected with control vector and DNA constructs which stably express human VEGF₁₆₅ as previously described [5,26]. Tumor cell lines were grown and maintained in Dulbecco's Modified Eagles Medium (DMEM) (Sigma) with 10% heat-inactivated fetal bovine serum.

Xenograft tumor model

Approximately 1×10^6 tumor cells were subcutaneously injected into the dorsal region of each mouse. Tumor growth was monitored on a daily basis. When tumors reached approximately 1.0 cm³, the experiments were terminated. Various tissues and organs were dissected and fixed with 4% paraformaldehyde (PFA) overnight, followed by transferring into PBS until further analysis. For some analyses, a portion of each tissue was frozen at -80°C until further use. In blockade experiments, VEGF tumor-bearing mice were randomly divided into two groups (n = 6-8/group) and received VEGFR1 or VEGFR2 blockades as previous described [4,5]. The treatment started at day 2 after tumor implantation. Each mouse in various groups received vehicle, anti-VEGFR1 (MF1, Imclone Inc., 800 µg/mouse) or anti-VEGFR2 (DC101, Imclone Inc., 800 µg/mouse) antibodies twice a week for a total 2-week period.

Histological analysis and immunohistochemistry

Paraffin-embedded tissues including liver, spleen and bone were sectioned in 5 µm thickness and stained with hemtoxylin/eosin (H&E) according to a standard method [5]. Liver sections were stained with an anti-Ter119 antibody as the primary antibody. After extensive washes with PBS, sections were stained with a secondary antibody of HRP-conjugated anti-rat IgG. Positive signals were developed by substrate DAB, followed by capturing under a light microscope (40× magnification).

Cryostat tissue sections of 10 µm thickness were stained according to a standard immunohistochemical procedure [5]. Briefly, sections were incubated with primary antibodies, including anti-CD31, Ter119, VEGFR1 or VEGFR2, overnight at 4 °C. After rigorous washes with PBS for three times, sections were incubated for 1 hour at room temperature with various secondary antibodies, including a goat anti-rat Alexa-555-labeled antibody (1:500), a FITC-labeled rabbit anti-rat IgG (1:300, Vector Laboratories), a Cy5-labeled goat anti-rabbit antibody (1:400, Chemicon International, Temecula, CA, USA),

which were used for either mono- or double-staining. Sections were mounted on glass-slides with Vectashield mounting medium (Vector Laboratories). Positive signals were photographed under a confocal microscopy (20×, Zeiss Confocal LSM510 Microscope).

Results and Discussion

Our recent findings show that tumor-derived VEGF induces a systemic syndrome in mice, manifesting a cancer-associated paraneoplastic syndrome [5]. Using the same tumor model, i.e., murine T241 fibrosarcoma model, we studied the role of VEGF in modulation of hematopoiesis. In a xenograft model, implantation of T241-VEGF tumors in mice led to hepatomegaly and splenomegaly [5]. Histological examination of liver tissues showed that visible hematopoietic islets in liver sections from T241-VEGF tumor-bearing mice but not in liver sections of T241-vector control tumor-bearing mice (Fig. 1). To further validate the identity of hematopoietic islets, liver sections were stained with Ter119, a hematopoietic marker for erythroblasts. Consistent with H&E staining, these morphologically hematopoietic islets exhibited positive signals of Ter119 (Fig. 1). Quantification analysis showed a significant difference between T241-VEGF and T241-vector groups (see ref [5]). Moreover, the sinusoidal hepatic vasculature became highly dilated in livers of T241-VEGF tumor-bearing mice but not in livers of T241-vector tumor-bearing mice (Fig. 1). These findings demonstrate that tumor-derived VEGF significantly contributes hepatic hematopoiesis and the vascular architecture is significantly altered in this organ.

Because extramedullary hematopoiesis often occurs in the spleen of mice, we next examined spleens of tumor-bearing mice. Similar to livers, splenomegaly also existed in T241-VEGF tumor-bearing mice [5]. Histological examination showed that apparent borders between the white pulp (WP) and red pulp (RP) under physiological conditions were vanished and were replaced by a mixture of WP and RP without any distinctive borders throughout the entire spleen of T241-VEGF tumor-bearing mice (Fig. 2). Ter119 staining revealed that active hematopoiesis occurred throughout the entire spleen tissue of T241-VEGF tumor-bearing mice (Fig. 2). In markedly contrast, Ter119 positive signals only present in the RP region in the spleen of T241-vector tumor-bearing mice (Fig. 2). These findings show that tumor-derived VEGF significantly modulates extramedullary hematopoiesis in the adult spleen by expansion of RP areas.

To delineate molecular mechanisms underlying systemic VEGF-induced extramedullary hematopoiesis, VEGFR1 and VEGFR2 were detected in both liver and spleen tissues. Interestingly, expression patterns of VEGFR1 and VEGFR2 were restricted in blood vessels, but not in other

cell types including hepatocytes, splenocytes and stromal cells (Fig. 3). Moreover, VEGF2 positive signals were generally enhanced while VEGFR1 signals were decreased in both spleens and livers of T241-VEGF tumor-bearing mice (Fig. 3). To further distinguish VEGFR1- and VEGFR2-mediated signaling pathways in hematopoiesis, VEGFR1 and VEGFR2 specific blockades were used for treatments. An anti-VEGFR2 specific monoclonal antibody completely restored histological structures of liver and spleen in VEGF tumor-bearing mice, whereas VEGFR1 blockade had no effect in these organs (Fig. 4). Consistent with histological changes, VEGFR2 blockade completely normalized hepatic sinusoidal vasculatures whereas VEGFR1 blockade lacked such an effect. These findings demonstrate that VEGFR2 is a crucial receptor that mediates extramedullary hematopoiesis and tortuosity of vasculatures in these organs.

Although VEGF might directly promote extramedullary hematopoiesis in liver and spleen, it was plausible that VEGF-induced extramedullary hematopoiesis echoes defective bone marrow (BM) hematopoiesis and thus represents a compensatory mechanism of BM deficiency. To test this possibility, we histologically examined BM of VEGF tumor-bearing mice. Markedly, VEGF tumor-bearing mice suffered from a severe defect in BM by losing hematopoietic cells relative to vector control BM (Fig. 5). Interestingly, VEGFR2 blockade could largely restore VEGF-induced BM defects while VEGFR1 blockade was unable to rescue the defective phenotype (Fig. 5). Based on these findings, it is likely that extramedullary hematopoiesis resulted from defective BM hematopoiesis via compensatory activation of this process in liver and spleen.

The data presented in this study provide compelling evidence that tumor-derived VEGF displays a profound effect on the hematopoietic system. It appears that tumor-derived VEGF enters into the circulation and acts on either endothelial cells and/or hematopoietic progenitor cells to modulate hematopoiesis. Although molecular mechanisms underlying VEGF-induced impairment of BM hematopoiesis remain unknown, it is possible that VEGF mobilizes BM hematopoietic stem cells to accumulate in peripheral tissues and organs. Indeed, VEGF has been reported to mobilize BM cells in tumor models and other experimental settings [15,16]. Suppression of BM hematopoiesis would in theory result in a decreased tumor growth rate due to insufficient oxygen supply. In contrast, in our experimental system we have found that VEGF tumors grow at an accelerated rate relative to control tumors [5,26]. There are several possible mechanisms to explain these paradoxical findings: 1) VEGF also exhibits potent angiogenic and vasculogenic activities, which significantly facilitate blood perfusion in tumor tissues; 2)





Figure 1
Hepatic hematopoietic islets and vasculature. Liver sections from T241-VEGF and T241-vector tumor-bearing mice were stained with H&E (A and B), an anti-Ter119 antibody (C and D), or anti-CD31 antibody (E and F). Arrows in panels A, C, and D point to hematopoietic islets. Arrows in E and F point to hepatic sinusoidal microvessels. CV = central vein. HC = hepatocyte. Bar in panels A-D = 100 μm. Bar in panels E and F = 50 μm.

**Figure 2**

Spleno-extramedullary hematopoiesis. Spleen sections from T241-VEGF and T241-vector tumor-bearing mice were stained with H&E (A and B), or were double stained an anti-Ter119 antibody and propidium iodide (PI) (C and D). Dashed lines in panel B encircle white pulp (WP). RP = red pulp. Hematopoietic cells from the erythrocyte lineage were labeled with Ter119 (green) and all cell nuclei were labeled with PI (red). Bar in panel A and B = 100 μ m. Bar in C and D = 50 μ m.

Hypoxia is known to increase tumor invasion and spread. Indeed, necrotic cores are often observed in VEGF tumors relative to control tumors. These findings suggest that VEGF tumors suffer from higher degrees of tissue hypoxia than control tumors, and hypoxia would persuade tumor cells to expand to healthy neighboring tissues; 3) VEGF-mobilized BM-derived inflammatory cells significantly

contribute to tumor growth by producing a range of angiogenic factors and cytokines, and 4) Extramedullary hematopoiesis compensates oxygen supply and supports tumor growth. However, this compensatory mechanism may not sufficiently rescue the defective phenotype of BM hematopoiesis.




Figure 3
Localization of VEGFR1 and VEGFR2 in liver and spleen tissue sections. Liver (A and B) and spleen (C and D) tissue sections from T241-VEGF and T241-vector tumor-bearing mice were double stained with monoclonal antibodies specific for mouse VEGFR1 (red) or VEGFR2 (blue). Yellow arrows point to double positive signals and white arrowheads indicate single positive signals. Bar = 50 μm.

Our findings also show that VEGF targeted therapy has a profound impact on multiple tissues and organs [5]. In the hematopoietic system, anti-VEGF drugs might significantly improve BM hematopoiesis and anemia in patients. However, the distinct functions mediated by VEGFR1 and VEGFR2 should be separated. While VEGFR1 is distributed in hematopoietic cells such as monocytes and neutrophils, VEGFR2 is the key receptor to mediate hematopoietic signals [5]. Consistent with our findings, genetic deletion of *vegfr2* gene in mice results in severe

defects of both vascular and hematopoietic systems in mouse embryos [6,7]. In contrast, deletion of VEGFR1 does not seem to significantly affect hematopoiesis [6,7]. Most currently available drugs such as bevacizumab, sunitinib and sorafenib do not allow us to distinguish the biological functions mediated by VEGFR1 and VEGFR2 [21]. It is unclear if VEGFR2 specific blockade would produce better clinical outcomes than currently available VEGF inhibitors in the clinic. However, it is known that VEGFR1 could also mediate negative signals of angiogen-




Figure 4
Histology and vasculature of liver and spleen treated with VEGFR blockades. Spleen (A-C) and liver (D-F) tissue sections from buffer- (A, D, and G), anti-VEGFR1- (B, E and H), or anti-VEGFR2-(C, F and I) treated T241-VEGF-bearing mice were stained with H&E (A-F) or anti-CD31 (G-I). Arrows in panels of G-I indicate microsinusoidal vasculatures in liver sections. CV = central vein. Bar = 50 µm.

esis under both physiological and pathological conditions [21].

Neutralization of VEGF-induced systemic disorders in cancer patients is probably an important mechanism for cancer therapy although antiangiogenesis is one of the key mechanisms of anti-VEGF drugs. Consistent with this notion, suppression of tumor size has not been positively correlated with clinical benefits of these antiangiogenic drugs [27]. Our current findings provide new evidence that VEGF targets multiple tissues and organs and that

anti-VEGF therapy may have a global impact in cancer patients.

Conclusions

Our findings show that tumor-derived VEGF modulates the hematopoietic system by suppression of BM-hematopoiesis, leading to activation of extramedullary hematopoiesis in liver and spleen. Modulation of hematopoiesis is mediated by VEGFR2 but not by VEGFR1. These results demonstrate that tumor-derived VEGF has a profound impact on multiple tissues and organs and these broad systemic effects might further facilitate tumor growth and




Figure 5

Histological examination of bone marrow. Bone marrows of T241-VEGF (A) and T241-vector (B) tumor-bearing mice were stained with H&E. Bone marrows of buffer- (C), anti-VEGFR1- (D), or anti-VEGFR2-(E) treated T241-VEGF-bearing mice were stained with H&E. Arrows in panels A and C-E point to the residual hematopoietic islets. Bone matrix were encircled with dashed lines. Bar in panel A and B = 100 μ m. Bar in panels C-E = 50 μ m.

metastasis. Additionally, circulating VEGF derived from the malignant tissue also contributes to development of cancer-associated systemic disorders, which significantly jeopardize quality of life and survival of cancer patients. Thus, anti-VEGF therapy may produce more profound beneficial effects in cancer patients by improving hematopoiesis and other systemic functions of cancer patients.

List of abbreviations

VEGF: vascular endothelial growth factor; VEGFR: VEGF receptor; BM: bone marrow; LLC: Lewis lung carcinoma; RP: red pulp; WP: white pulp.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

YX and YC designed research; YX, FC, SL and DZ performed research; YX, FC, SL, DZ and YC analyzed data; YC wrote the paper.

Acknowledgements

This work was supported by the laboratory of YC through research grants from the Swedish Research Council, the Swedish Cancer Foundation, the Karolinska Institute Foundation, the Karolinska gender foundation, the Torsten and Ragnar Söderberg's Foundation, and by European Union Integrated Projects of Angiotargeting Contract 504743 (to YC) and VascuPlug

Contract STRP 013811 (to YC). YC is also a Chang Jiang scholar at the Shandong University, China.

References

- Cao Y: **Antiangiogenic cancer therapy.** *Semin Cancer Biol* 2004, **14**:139-45.
- Cao Y: **Tumor angiogenesis and molecular targets for therapy.** *Front Biosci* 2009, **14**:3962-73.
- Crawford Y, Ferrara N: **VEGF inhibition: insights from preclinical and clinical studies.** *Cell Tissue Res* 2009, **335**:261-9.
- Xue Y, Petrovic N, Cao R, Larsson O, Lim S, Chen S, Feldmann HM, Liang Z, Zhu Z, Nedergaard J, Cannon B, Cao Y: **Hypoxia-independent angiogenesis in adipose tissues during cold acclimation.** *Cell Metab* 2009, **9**:99-109.
- Xue Y, Religa P, Cao R, Hansen AJ, Lucchini F, Jones B, Wu Y, Zhu Z, Pytowski B, Liang Y, Zhong W, Vezzone P, Rozell B, Cao Y: **Anti-VEGF agents confer survival advantages to tumor-bearing mice by improving cancer-associated systemic syndrome.** *Proc Natl Acad Sci USA* 2008, **105**:18513-8.
- Carmeliet P, Ferreira V, Breier G, Pollefeyt S, Kieckens L, Gertsensstein M, Fahrig M, Vandenhoeck A, Harpal K, Eberhardt C, Declercq C, Pawling J, Moons L, Collen D, Risau W, Nagy A: **Abnormal blood vessel development and lethality in embryos lacking a single VEGF allele.** *Nature* 1996, **380**:435-9.
- Ferrara N, Carver-Moore K, Chen H, Dowd M, Lu L, O'Shea KS, Powell-Braxton L, Hillan KJ, Moore MW: **Heterozygous embryonic lethality induced by targeted inactivation of the VEGF gene.** *Nature* 1996, **380**:439-42.
- Ferrara N: **Vascular endothelial growth factor.** *Arterioscler Thromb Vasc Biol* 2009, **29**:789-91.
- Kamba T, McDonald DM: **Mechanisms of adverse effects of anti-VEGF therapy for cancer.** *Br J Cancer* 2007, **96**:1788-95.
- Stockmann C, Doedens A, Weidemann A, Zhang N, Takeda N, Greenberg JL, Cheresch DA, Johnson RS: **Deletion of vascular endothelial growth factor in myeloid cells accelerates tumorigenesis.** *Nature* 2008, **456**:814-8.
- Tam BY, Wei K, Rudge JS, Hoffman J, Holash J, Park SK, Yuan J, Hefner C, Chartier C, Lee JS, Jiang S, Nayak NR, Kuypers FA, Ma L, Sundram U, Wu G, Garcia JA, Schrier SL, Maher JJ, Johnson RS, Yancopoulos GD, Mulligan RC, Kuo CJ: **VEGF modulates erythropoiesis through regulation of adult hepatic erythropoietin synthesis.** *Nat Med* 2006, **12**:793-800.
- Makino Y, Cao R, Svensson K, Bertilsson G, Asman M, Tanaka H, Cao Y, Berkenstam A, Poellinger L: **Inhibitory PAS domain protein is a negative regulator of hypoxia-inducible gene expression.** *Nature* 2001, **414**:550-4.
- Shweiki D, Itin A, Soffer D, Keshet E: **Vascular endothelial growth factor induced by hypoxia may mediate hypoxia-initiated angiogenesis.** *Nature* 1992, **359**:843-5.
- Nagy JA, Chang SH, Dvorak AM, Dvorak HF: **Why are tumour blood vessels abnormal and why is it important to know?** *Br J Cancer* 2009, **100**:865-9.
- Lyden D, Hattori K, Dias S, Costa C, Blaikie P, Butros L, Chadburn A, Heissig B, Marks W, Witte L, Wu Y, Hicklin D, Zhu Z, Hackett NR, Crystal RG, Moore MA, Hajar KA, Manova K, Benezra R, Rafii S: **Impaired recruitment of bone-marrow-derived endothelial and hematopoietic precursor cells blocks tumor angiogenesis and growth.** *Nat Med* 2001, **7**:1194-201.
- Shaked Y, Ciarrocchi A, Franco M, Lee CR, Man S, Cheung AM, Hicklin DJ, Chaplin D, Foster FS, Benezra R, Kerbel RS: **Therapy-induced acute recruitment of circulating endothelial progenitor cells to tumors.** *Science* 2006, **313**:1785-7.
- Bjorn Dahl MA, Cao R, Burton JB, Brakenhielm E, Religa P, Galter D, Wu L, Cao Y: **Vascular endothelial growth factor-a promotes peritumoral lymphangiogenesis and lymphatic metastasis.** *Cancer Res* 2005, **65**:9261-8.
- Cursiefen C, Chen L, Borges LP, Jackson D, Cao J, Radziejewski C, D'Amore PA, Dana MR, Wiegand SJ, Streilein JW: **VEGF-A stimulates lymphangiogenesis and hemangiogenesis in inflammatory neovascularization via macrophage recruitment.** *J Clin Invest* 2004, **113**:1040-50.
- Hong YK, Lange-Asschenfeldt B, Velasco P, Hirakawa S, Kunstfeld R, Brown LF, Bohlen P, Senger DR, Detmar M: **VEGF-A promotes tissue repair-associated lymphatic vessel formation via VEGFR-2 and the alpha1beta1 and alpha2beta1 integrins.** *FASEB J* 2004, **18**:1111-3.
- Nagy JA, Vasile E, Feng D, Sundberg C, Brown LF, Detmar MJ, Lawitts JA, Benjamin L, Tan X, Manseau EJ, Dvorak AM, Dvorak HF: **Vascular permeability factor/vascular endothelial growth factor induces lymphangiogenesis as well as angiogenesis.** *J Exp Med* 2002, **196**:1497-506.
- Cao Y: **Positive and negative modulation of angiogenesis by VEGFR1 ligands.** *Sci Signal* 2009, **2**:re1.
- Caunt M, Mak J, Liang WC, Stawicki S, Pan Q, Tong RK, Kowalski J, Ho C, Reslan HB, Ross J, Berry L, Kasman I, Zlot C, Cheng Z, Le Couter J, Filvaroff EH, Plowman G, Peale F, French D, Carano R, Koch AW, Wu Y, Watts RJ, Tessier-Lavigne M, Bagri A: **Blocking neuropilin-2 function inhibits tumor cell metastasis.** *Cancer Cell* 2008, **13**:331-42.
- Pan Q, Chanthery Y, Liang WC, Stawicki S, Mak J, Rathore N, Tong RK, Kowalski J, Yee SF, Pacheco G, Ross S, Cheng Z, Le Couter J, Plowman G, Peale F, Koch AW, Wu Y, Bagri A, Tessier-Lavigne M, Watts RJ: **Blocking neuropilin-1 function has an additive effect with anti-VEGF to inhibit tumor growth.** *Cancer Cell* 2007, **11**:53-67.
- Cao Y, Zhong W, Sun Y: **Improvement of antiangiogenic cancer therapy by understanding the mechanisms of angiogenic factor interplay and drug resistance.** *Semin Cancer Biol* 2009.
- Collins TS, Hurwitz HI: **Targeting vascular endothelial growth factor and angiogenesis for the treatment of colorectal cancer.** *Semin Oncol* 2005, **32**:61-8.
- Eriksson A, Cao R, Pawliuk R, Berg SM, Tsang M, Zhou D, Fleet C, Tritsarlis K, Dissing S, Le Boulch P, Cao Y: **Placenta growth factor-1 antagonizes VEGF-induced angiogenesis and tumor growth by the formation of functionally inactive PIGF-II/VEGF heterodimers.** *Cancer Cell* 2002, **1**:99-108.
- Ledford H: **Drug markers questioned.** *Nature* 2008, **452**:510-1.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

