

Received: 2022.09.15
Accepted: 2022.09.15
Available online: 2022.09.15
Published: 2022.09.16

Errate: Remimazolam Tosilate Sedation with Adjuvant Sufentanil in Chinese Patients with Liver Cirrhosis Undergoing Gastroscopy: A Randomized Controlled Study

1 YingHao Cao
1 Ping Chi
2 Chen Zhou
1 WenFei Lv
1 ZheFen Quan
1 Fu Shan Xue

1 Department of Anesthesiology, Beijing Friendship Hospital, Capital Medical University, Beijing, PR China
2 Department of Zoology and Physiology, University of Wyoming, Laramie, WY, USA

Corresponding Author: Fu Shan Xue, e-mail: xuefushan@aliyun.com

The results in Table 3 (highlighted) have been corrected to be consistent with the results reported in the text.

Table 3. Primary and secondary outcomes.

		Remimazolam	Propofol	P
Subjects, n		74	74	–
Primary outcome				
Endoscopist satisfaction	Satisfied	67 (90.5%)	57 (77.0%)	0.026
	Fair	6 (8.1%)	13 (17.6%)	0.085
	Unsatisfied	1 (1.4%)	4 (5.4%)	0.367
Secondary outcomes				
Respiratory depression		2 (2.7%)	13 (17.6%)	0.007
Apnea		Nil	3 (4.1%)	0.245
Body movement		6 (8.1%)	17 (23.0%)	0.043
Bradycardia		Nil	4 (5.4%)	0.1021
Hypotension		3 (4.1%)	11 (14.9%)	0.031
Nausea or vomiting		2 (2.7%)	1 (1.4%)	1.000
Somnolence		Nil	3 (4.1%)	0.245
Dizziness		1 (1.4%)	3 (4.1%)	0.621
Fever		Nil	Nil	1.000
Patient satisfaction	Satisfied	74 (100.0%)	72 (97.3%)	0.908
	Fair	Nil	2 (2.7%)	0.497
	Unsatisfied	Nil	Nil	1.000

Reference:

YingHao Cao, Ping Chi, Chen Zhou, WenFei Lv, ZheFen Quan, Fu Shan Xue: Remimazolam Tosilate Sedation with Adjuvant Sufentanil in Chinese Patients with Liver Cirrhosis Undergoing Gastroscopy: A Randomized Controlled Study. Med Sci Monit, 2022; 28: e936580.

DOI: 10.12659/MSM.936580



Full-text PDF: <https://www.medscimonit.com/abstract/index/idArt/938399>