

[Tris(3,5-diphenylpyrazolyl)hydroborato]nickel(II) bromide

David J. Harding,^{a*} Phimpakha Harding^a and Harry Adams^b

^aMolecular Technology Research Unit, Department of Chemistry, Walailak University, Thasala, Nakhon Si Thammarat 80161, Thailand, and ^bDepartment of Chemistry, Faculty of Science, University of Sheffield, Brook Hill, Sheffield S3 7HF, England

Correspondence e-mail: hdavid@wu.ac.th


Received 14 May 2009; accepted 6 June 2009

Key indicators: single-crystal X-ray study; $T = 150\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.028; wR factor = 0.063; data-to-parameter ratio = 12.7.

In the title tris(pyrazolyl)borate (Tp^{Ph_2}) complex, $[\text{NiBr}(\text{C}_{45}\text{H}_{34}\text{BN}_6)]$, the Ni, Br and B atoms lie on a crystallographic threefold axis and a distorted NiN_3Br tetrahedral geometry arises for the metal ion. In the crystal, $\text{C}-\text{H}\cdots(\text{C}=\text{C})$ and $\text{C}-\text{H}\cdots\pi$ interactions help to establish the polar crystal packing.

Related literature

For other Tp^RNiX ($X = \text{Cl}, \text{Br}$) complexes, see: Desrochers *et al.* (2003, 2006); Kunrath *et al.* (2003); Uehara *et al.* (2002); Guo *et al.* (1998); Harding *et al.* (2007). For ionic radius data, see: Shannon (1976).



Experimental

Crystal data

$[\text{NiBr}(\text{C}_{45}\text{H}_{34}\text{BN}_6)]$
 $M_r = 808.21$
Trigonal, $R\bar{3}$
 $a = 12.8227(8)\text{ \AA}$
 $c = 19.327(3)\text{ \AA}$
 $V = 2752.0(5)\text{ \AA}^3$

$Z = 3$
Mo $K\alpha$ radiation
 $\mu = 1.66\text{ mm}^{-1}$
 $T = 150\text{ K}$
 $0.24 \times 0.24 \times 0.21\text{ mm}$

Data collection

Bruker SMART CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 1997)
 $T_{\min} = 0.691$, $T_{\max} = 0.722$

5609 measured reflections
2075 independent reflections
1943 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.063$
 $S = 1.06$
2075 reflections
163 parameters
1 restraint

H-atom parameters constrained
 $\Delta\rho_{\max} = 0.27\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.29\text{ e \AA}^{-3}$
Absolute structure: Flack (1983),
670 Friedel pairs
Flack parameter: 0.020 (8)

Table 1
Selected geometric parameters (\AA , $^\circ$).

Ni1—Br1	2.3523 (6)	Ni1—N1	2.041 (2)
N1—Ni1—N1 ¹	93.11 (8)	N1—Ni1—Br1	123.04 (6)

Symmetry code: (i) $-x + y + 1, -x + 1, z$.

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C5—H5 ⁱ — $\text{Cg}1^{\text{ii}}$	0.95	2.73	3.589 (3)	151

Symmetry code: (ii) $-x + y + \frac{5}{3}, -x + \frac{1}{3}, z - \frac{2}{3}$.

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINT* (Bruker, 1997); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors gratefully acknowledge the Thailand Research Fund (grant No. RMU5080029) and the National Research Council of Thailand (grant No. WU51106) for support of this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2976).

References

- Bruker (1997). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Desrochers, P. J., LeLievre, L., Johnson, R. J., Lamb, B. T., Phelps, A. L., Cordes, A. W., Gu, W. & Cramer, S. P. (2003). *Inorg. Chem.* **42**, 7945–7950.
- Desrochers, P. J., Telszer, J., Zvyagin, S. A., Ozarowski, A., Krzystek, J. & Vicic, D. A. (2006). *Inorg. Chem.* **45**, 8930–8941.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Guo, S., Ding, E., Yin, Y. & Yu, K. (1998). *Polyhedron*, **17**, 3841–3849.
- Harding, D. J., Harding, P., Adams, H. & Tuntulani, T. (2007). *Inorg. Chim. Acta*, **360**, 3335–3340.
- Kunrath, F. A., de Souza, R. F., Casagrande, O. L. Jr, Brooks, N. R. & Young, V. G. Jr (2003). *Organometallics*, **22**, 4739–4743.
- Shannon, R. D. (1976). *Acta Cryst. A* **32**, 751–767.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Uehara, K., Hikichi, S. & Akita, M. (2002). *J. Chem. Soc. Dalton Trans.* pp. 3529–3538.

supplementary materials

Acta Cryst. (2009). E65, m773 [doi:10.1107/S1600536809021606]

[Tris(3,5-diphenylpyrazolyl)hydroborato]nickel(II) bromide

D. J. Harding, P. Harding and H. Adams

Comment

Tris(pyrazolyl)borates are versatile and popular ligands in coordination chemistry with many complexes now known. Despite the C₃ symmetry present in many tris(pyrazolyl)borate ligands few tetrahedral complexes crystallize in space groups containing a C₃ axis (Desrochers *et al.*, 2003, 2006; Kunrath *et al.*, 2003; Uehara *et al.*, 2002). Rare examples which do contain a C₃ axis include [Tp^{Ph₂}NiCl] and [Tp^{Ph₂}Ni(OAc)] despite the later being formally five-coordinate (Guo *et al.*, 1998; Harding *et al.*, 2007). In the following paper we report a further example namely, the title compound, [Tp^{Ph₂}NiBr], (I).

The reaction of NiBr₂.2H₂O with KTp^{Ph₂} readily affords the title complex as a red-purple solid in moderate yield. Crystals were grown by allowing hexanes to diffuse into a concentrated solution of the complex in CH₂Cl₂. The compound crystallizes in the trigonal R3 space group. The structure is shown in Figure 1 while important bond lengths and angles are given in the supporting tables. The geometry around the nickel centre is best described as distorted tetrahedral {N1—Ni—N1ⁱ = 93.11 (8), N1—Ni—Br1 = 123.04 (6)}. The Ni—N bond lengths are very slightly longer by *ca.* 0.01 Å than those found in [Tp^{Ph₂}NiCl] (Guo *et al.*, 1998). A similar difference is observed in the structures of [Tp^{*}NiCl] and [Tp^{*}NiBr] (Desrochers *et al.*, 2003, 2006). The Ni—Br distance is 2.3523 (6) Å, *ca* 0.15 Å longer than the corresponding Ni—Cl distance in [Tp^{Ph₂}NiCl], and consistent with the difference in the bromine and chlorine covalent radii (0.15 Å; Shannon, 1976). Interestingly, the Ni—Br bond length in (I) is significantly longer than that observed for [Tp^{*}NiBr] (2.291 (2) Å). A similar increase, albeit not so marked, is also found between [Tp^{Ph₂}NiCl] and [Tp^{*}NiCl] (Δ Ni—Cl = 0.03 Å) suggesting that the larger Tp^{Ph₂} ligand may be responsible for the longer nickel-halide bond distances.

The crystal packing in the structure of (I) contains several C—H···π interactions between the phenyl rings of neighbouring Tp^{Ph₂} ligands (see Figure 2). The hydrogen atoms H6 and H11 are directed at the π bonds between C1—N1 and C4—C5, respectively {(C1—N1)π···H6 2.657 (3) Å; (C4—C5)π···H11 2.834 (5) Å} while H5 interacts with a phenyl ring (Cg1···H5 2.730 (3) Å; Cg1 is the centroid of ring C10—C15). Similar interactions occur on all three faces of the Tp^{Ph₂} ligand creating a network of triangular columns such that all the [Tp^{Ph₂}NiBr] molecules point in the same direction and the phenyl rings adopt a propeller configuration.

Experimental

NiBr₂.2H₂O (34 mg, 0.14 mmol) was dissolved in THF (5 ml) giving a green solution and then stirred for 5 min. KTp^{Ph₂} (95 mg, 0.15 mmol) was dissolved in THF (5 ml) giving a pale yellow solution. Addition of the KTp^{Ph₂} solution to the Ni solution resulted in a colour change to a red-pink solution. The solution was stirred for 16 hrs. The solution was reduced to dryness, redissolved in fresh THF (2 ml) and filtered through celite. The purple-pink solution was layered with hexanes (10 ml). After two days purple-pink blocks of (I) appeared. These were washed with EtOH (3 x 3 ml) and hexanes (2 x 5

supplementary materials

ml) (59 mg, 54%). $\nu_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ 3059w, 2966w, 2854w (vCH), 2626w (vBH). δ_{H} (300 MHz; CDCl_3 ; SiMe₄), 7.63 (br m, *m*- and *p*-Ph, 18H), 8.53 (br s, *o*-Ph, 6H), 8.86 (br s, *o*-Ph, 6H). UV–Vis $\lambda_{\text{max}}(\text{CH}_2\text{Cl}_2)/\text{nm}$ 318 ($\epsilon/\text{dm}^3\text{mol}^{-1}\text{cm}^{-1}$ 3140), 502 (600), 828 (120), 926 (160). m/z (ESI) 727 [M—Br]⁺. Anal. Calc. for $\text{C}_{45}\text{H}_{34}\text{N}_6\text{BBrNi}[\text{Tp}^{\text{Ph}_2}\text{NiBr}]$: C, 66.87; H, 4.24; N 10.40 Found: C, 66.62; H, 4.29; N, 10.17%

Figures




Fig. 1. The molecular structure of (I) with displacement ellipsoids drawn at the 50% probability level. [Symmetry codes:(i) - $x + y + 1$, - $x + 1$, z ; (ii) - $y + 1$, $x - y$, z].




Fig. 2. The molecular packing in (I) showing the three C—H···π interactions. Only selected atoms are labelled or shown for clarity [Symmetry codes:(i) 4/3 - $x + y$, 2/3 - x , -1/3 + z ; (ii) 4/3 - y , -1/3 + $x - y$, -1/3 + z].

[Tris(3,5-diphenylpyrazolyl)hydroborato]nickel(II) bromide

Crystal data

[NiBr(C ₄₅ H ₃₄ BN ₆)]	$Z = 3$
$M_r = 808.21$	$F_{000} = 1242$
Trigonal, R3	$D_x = 1.463 \text{ Mg m}^{-3}$
Hall symbol: R 3	Mo $K\alpha$ radiation
$a = 12.8227 (8) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 12.8227 (8) \text{ \AA}$	Cell parameters from 2851 reflections
$c = 19.327 (3) \text{ \AA}$	$\theta = 2.8\text{--}30.6^\circ$
$\alpha = 90^\circ$	$\mu = 1.66 \text{ mm}^{-1}$
$\beta = 90^\circ$	$T = 150 \text{ K}$
$\gamma = 120^\circ$	Block, purple–pink
$V = 2752.0 (5) \text{ \AA}^3$	$0.24 \times 0.24 \times 0.21 \text{ mm}$

Data collection

Bruker SMART CCD diffractometer	2075 independent reflections
Radiation source: fine-focus sealed tube	1943 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.037$

Detector resolution: 100 pixels mm⁻¹
 $T = 150\text{ K}$
 φ scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 1997)
 $T_{\min} = 0.691, T_{\max} = 0.722$
 5609 measured reflections

$\theta_{\max} = 27.5^\circ$

$\theta_{\min} = 2.1^\circ$

$h = -16 \rightarrow 16$

$k = -11 \rightarrow 16$

$l = -21 \rightarrow 25$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.063$
 $S = 1.06$
 2075 reflections
 163 parameters
 1 restraint
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0169P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.27\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.29\text{ e \AA}^{-3}$
 Extinction correction: none
 Absolute structure: Flack (1983), 670 Friedel pairs
 Flack parameter: 0.020 (8)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.6667	0.3333	0.732391 (16)	0.02249 (14)
Ni1	0.6667	0.3333	0.61068 (2)	0.01335 (14)
N1	0.75319 (18)	0.26622 (19)	0.55312 (10)	0.0127 (4)
N2	0.75175 (19)	0.29001 (19)	0.48407 (10)	0.0128 (4)
B1	0.6667	0.3333	0.4567 (3)	0.0139 (10)
H1B	0.6667	0.3333	0.3983	0.017*
C1	0.8082 (2)	0.2003 (2)	0.55910 (12)	0.0146 (5)
C2	0.8419 (3)	0.1815 (3)	0.49355 (13)	0.0166 (6)
H2	0.8812	0.1373	0.4829	0.020*

supplementary materials

C3	0.8064 (2)	0.2405 (2)	0.44716 (12)	0.0136 (5)
C4	0.8224 (3)	0.2532 (3)	0.37198 (14)	0.0135 (5)
C5	0.7922 (2)	0.1509 (2)	0.33065 (13)	0.0173 (6)
H5	0.7601	0.0738	0.3515	0.021*
C6	0.8094 (3)	0.1632 (3)	0.25990 (16)	0.0182 (7)
H6	0.7890	0.0942	0.2325	0.022*
C7	0.8557 (2)	0.2743 (3)	0.22844 (13)	0.0201 (6)
H7	0.8668	0.2814	0.1797	0.024*
C8	0.8860 (3)	0.3759 (3)	0.26839 (13)	0.0215 (6)
H8	0.9179	0.4525	0.2469	0.026*
C9	0.8698 (3)	0.3652 (3)	0.33951 (14)	0.0190 (6)
H9	0.8913	0.4350	0.3665	0.023*
C10	0.8239 (2)	0.1510 (3)	0.62468 (13)	0.0163 (6)
C11	0.8460 (2)	0.2131 (3)	0.68680 (14)	0.0185 (6)
H11	0.8564	0.2919	0.6872	0.022*
C12	0.8530 (3)	0.1605 (3)	0.74866 (17)	0.0230 (7)
H12	0.8650	0.2023	0.7912	0.028*
C13	0.8422 (3)	0.0472 (3)	0.74807 (16)	0.0269 (7)
H13	0.8468	0.0113	0.7901	0.032*
C14	0.8246 (3)	-0.0137 (3)	0.68574 (15)	0.0234 (7)
H14	0.8191	-0.0904	0.6849	0.028*
C15	0.8150 (3)	0.0381 (3)	0.62476 (16)	0.0209 (6)
H15	0.8022	-0.0041	0.5824	0.025*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.02665 (19)	0.02665 (19)	0.0142 (2)	0.01332 (9)	0.000	0.000
Ni1	0.0140 (2)	0.0140 (2)	0.0120 (3)	0.00702 (10)	0.000	0.000
N1	0.0107 (10)	0.0149 (11)	0.0121 (9)	0.0061 (9)	-0.0007 (8)	0.0007 (8)
N2	0.0137 (11)	0.0146 (11)	0.0096 (8)	0.0068 (9)	0.0007 (8)	0.0006 (8)
B1	0.0112 (15)	0.0112 (15)	0.019 (2)	0.0056 (8)	0.000	0.000
C1	0.0121 (13)	0.0105 (13)	0.0199 (12)	0.0047 (11)	-0.0016 (10)	0.0005 (10)
C2	0.0186 (14)	0.0165 (15)	0.0191 (16)	0.0122 (13)	-0.0028 (13)	0.0009 (12)
C3	0.0097 (13)	0.0094 (12)	0.0189 (12)	0.0026 (11)	-0.0001 (9)	0.0010 (9)
C4	0.0106 (13)	0.0148 (14)	0.0169 (12)	0.0077 (11)	0.0010 (10)	-0.0012 (11)
C5	0.0171 (15)	0.0124 (14)	0.0221 (13)	0.0071 (12)	0.0005 (11)	0.0031 (11)
C6	0.0191 (16)	0.0155 (15)	0.0234 (16)	0.0113 (13)	-0.0015 (13)	-0.0039 (13)
C7	0.0180 (15)	0.0237 (15)	0.0153 (12)	0.0080 (13)	0.0044 (11)	0.0003 (11)
C8	0.0210 (15)	0.0179 (15)	0.0210 (12)	0.0062 (12)	0.0060 (11)	0.0053 (11)
C9	0.0199 (16)	0.0172 (15)	0.0193 (13)	0.0087 (13)	0.0016 (11)	-0.0029 (11)
C10	0.0089 (13)	0.0166 (14)	0.0215 (13)	0.0049 (11)	0.0001 (10)	0.0043 (11)
C11	0.0136 (14)	0.0155 (14)	0.0233 (13)	0.0050 (12)	-0.0050 (11)	0.0012 (11)
C12	0.0204 (17)	0.0291 (19)	0.0180 (15)	0.0112 (15)	-0.0034 (13)	0.0002 (14)
C13	0.0206 (16)	0.0304 (17)	0.0268 (14)	0.0105 (14)	-0.0022 (13)	0.0126 (13)
C14	0.0206 (16)	0.0195 (16)	0.0333 (16)	0.0124 (13)	-0.0008 (12)	0.0075 (13)
C15	0.0185 (15)	0.0194 (16)	0.0257 (14)	0.0101 (14)	-0.0019 (12)	-0.0011 (12)

Geometric parameters (\AA , $^\circ$)

Ni1—Br1	2.3523 (6)	C5—H5	0.9500
Ni1—N1	2.041 (2)	C6—C7	1.380 (4)
Ni1—N1 ⁱ	2.041 (2)	C6—H6	0.9500
Ni1—N1 ⁱⁱ	2.041 (2)	C7—C8	1.392 (4)
N1—C1	1.350 (3)	C7—H7	0.9500
N1—N2	1.371 (3)	C8—C9	1.387 (4)
N2—C3	1.360 (3)	C8—H8	0.9500
N2—B1	1.544 (3)	C9—H9	0.9500
B1—N2 ⁱ	1.544 (3)	C10—C11	1.389 (4)
B1—N2 ⁱⁱ	1.544 (3)	C10—C15	1.394 (4)
B1—H1B	1.1278	C11—C12	1.397 (4)
C1—C2	1.398 (4)	C11—H11	0.9500
C1—C10	1.475 (3)	C12—C13	1.390 (5)
C2—C3	1.388 (3)	C12—H12	0.9500
C2—H2	0.9500	C13—C14	1.391 (5)
C3—C4	1.465 (4)	C13—H13	0.9500
C4—C9	1.397 (4)	C14—C15	1.388 (4)
C4—C5	1.415 (4)	C14—H14	0.9500
C5—C6	1.381 (4)	C15—H15	0.9500
N1—Ni1—N1 ⁱ	93.11 (8)	C4—C5—H5	120.0
N1—Ni1—N1 ⁱⁱ	93.11 (8)	C7—C6—C5	121.0 (3)
N1 ⁱ —Ni1—N1 ⁱⁱ	93.11 (8)	C7—C6—H6	119.5
N1—Ni1—Br1	123.04 (6)	C5—C6—H6	119.5
N1 ⁱ —Ni1—Br1	123.04 (6)	C6—C7—C8	119.7 (2)
N1 ⁱⁱ —Ni1—Br1	123.04 (6)	C6—C7—H7	120.1
C1—N1—N2	106.99 (19)	C8—C7—H7	120.1
C1—N1—Ni1	141.45 (17)	C9—C8—C7	119.9 (3)
N2—N1—Ni1	111.47 (15)	C9—C8—H8	120.0
C3—N2—N1	109.82 (19)	C7—C8—H8	120.0
C3—N2—B1	127.6 (2)	C8—C9—C4	120.9 (3)
N1—N2—B1	120.3 (2)	C8—C9—H9	119.5
N2—B1—N2 ⁱ	108.89 (19)	C4—C9—H9	119.5
N2—B1—N2 ⁱⁱ	108.89 (19)	C11—C10—C15	118.8 (3)
N2 ⁱ —B1—N2 ⁱⁱ	108.89 (19)	C11—C10—C1	121.9 (3)
N2—B1—H1B	110.0	C15—C10—C1	119.2 (2)
N2 ⁱ —B1—H1B	110.0	C10—C11—C12	120.4 (3)
N2 ⁱⁱ —B1—H1B	110.0	C10—C11—H11	119.8
N1—C1—C2	109.5 (2)	C12—C11—H11	119.8
N1—C1—C10	124.6 (2)	C13—C12—C11	120.1 (3)
C2—C1—C10	125.8 (2)	C13—C12—H12	120.0
C3—C2—C1	106.1 (2)	C11—C12—H12	120.0
C3—C2—H2	127.0	C14—C13—C12	119.8 (3)
C1—C2—H2	127.0	C14—C13—H13	120.1

supplementary materials

N2—C3—C2	107.6 (2)	C12—C13—H13	120.1
N2—C3—C4	122.9 (2)	C15—C14—C13	119.7 (3)
C2—C3—C4	129.5 (2)	C15—C14—H14	120.2
C9—C4—C5	118.4 (2)	C13—C14—H14	120.2
C9—C4—C3	121.7 (3)	C14—C15—C10	121.1 (3)
C5—C4—C3	119.9 (2)	C14—C15—H15	119.4
C6—C5—C4	119.9 (3)	C10—C15—H15	119.4
C6—C5—H5	120.0		


Symmetry codes: (i) $-x+y+1, -x+1, z$; (ii) $-y+1, x-y, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C5—H5 \cdots Cg1 ⁱⁱⁱ	0.95	2.73	3.589 (3)	151

Symmetry codes: (iii) $-x+y+5/3, -x+1/3, z-2/3$.

Fig. 1



supplementary materials

Fig. 2

