



# Moderate Exercise Training Combined With a High-Fat and Sucrose Diet Protects Pancreatic Islet Function in Male C57BL/6J Mice

Katherine Veras<sup>1\*†</sup>, Camila Ferraz Lucena<sup>1</sup>, Julia Goedcke<sup>2</sup>, Fabiana S. Evangelista<sup>3</sup>, Angelo Carpinelli<sup>1</sup> and Carla Roberta de Oliveira Carvalho<sup>1</sup>

<sup>1</sup> Institute of Biomedical Sciences, Department of Physiology and Biophysics, University of São Paulo, São Paulo, Brazil,

<sup>2</sup> Non-Communicable Diseases Research Unit, South African Medical Research Council, Cape Town, South Africa,

<sup>3</sup> School of Arts, Sciences and Humanities, University of São Paulo, São Paulo, Brazil

## OPEN ACCESS

### Edited by:

Emmanuel Ampofo,  
Saarland University Hospital, Germany

### Reviewed by:

Sabrina Grassioli,  
Universidade Estadual do Oeste do  
Paraná, Brazil  
Paul Frank Williams,  
The University of Sydney, Australia

### \*Correspondence:

Katherine Veras  
Katherine.veras@gmail.com

### †Present address:

Katherine Veras,  
School of Health & Human  
Performance, Dublin City University,  
Dublin, Ireland

### Specialty section:

This article was submitted to  
Diabetes: Molecular Mechanisms,  
a section of the journal  
Frontiers in Endocrinology

Received: 22 February 2022

Accepted: 13 April 2022

Published: 20 May 2022

### Citation:

Veras K, Lucena CF, Goedcke J,  
Evangelista FS, Carpinelli A and  
Carvalho CRO (2022) Moderate  
Exercise Training Combined With  
a High-Fat and Sucrose Diet  
Protects Pancreatic Islet Function  
in Male C57BL/6J Mice.  
Front. Endocrinol. 13:881236.  
doi: 10.3389/fendo.2022.881236

Obesity is mainly caused by excess energy intake and physical inactivity, and the number of overweight/obese individuals has been steadily increasing for decades. Previous studies showed that rodents fed westernized diets exhibit endocrine pancreas deterioration and a range of metabolic disorders. This study evaluated the effects of moderated aerobic treadmill exercise training on pancreatic islet cell viability and function in mice consuming a high-fat and sucrose diet. In the present study, 60-day-old male C57BL/6J mice were divided into four groups: control (C), fed a standard diet AIN-93M (3.83 kcal/g; 70% carbohydrate (cornstarch and dextrinized starch were chosen as the major source of carbohydrate for the AIN-93 diet. In addition, a small amount of sucrose), 20% protein (casein), and 10% fat (soybean) with no training (i.e., sedentary); C + training (CTR, fed the standard diet with eight weeks of exercise; high-fat diet + sucrose (HFDS), fed a high fat and sucrose diet (5.2 kcal/g; 20% carbohydrate (cornstarch and dextrinized starch were chosen as the major source of carbohydrate), 20% protein (casein), 60% fat (Lard was chosen as the major source of fat and a small amount of soybean) + 20% sucrose diluted in drinking water with no training; and HFDS + training (HFDSTR). After eight weeks, the HFDS mice displayed increased body weight ( $P < 0.001$ ) and epididymal, inguinal and retroperitoneal adipose tissue mass ( $P < 0.01$ ). These mice also presented insulin resistance ( $P < 0.01$ ), glucose intolerance ( $P < 0.001$ ), impaired glucose-stimulated insulin secretion (GSIS) and were less responsive to the physiological net ROS production induced by glucose stimulus. The HFDS group's pancreatic islet cells were 38% less viable and 59% more apoptotic than those from the C group ( $P < 0.05$ ). The HFDSTR improved glucose tolerance, body mass, insulin sensitivity and GSIS ( $P < 0.05$ ). Furthermore, HFDSTR mice had 53% more viable isolated pancreatic islets cells and 29% fewer apoptotic cells than the HFDS group ( $P < 0.01$ ). Thus, exercise training may slow down and/or prevent adverse metabolic effects associated with consuming a westernized diet.

**Keywords:** high fat and sucrose diet, aerobic exercise training, obesity, pancreatic islets, glucose tolerance, insulin sensitivity

## INTRODUCTION

The rising worldwide prevalence of overweight and obese individuals has been described as a global pandemic (1). Notably, no country has reported a decrease in the prevalence or rate of obesity in the last 33 years (2). It is well known that obesity is mostly caused by excess energy intake from diets rich in saturated fat and sugar (i.e., westernized diet) and reduced energy expenditure such as physical inactivity, a combination frequently observed in modern human societies (3–5).

Obesity can lead to insulin resistance (IR) with compensatory insulin secretion to maintain glucose homeostasis (6). Under chronic conditions, this mechanism impairs pancreatic  $\beta$ -cell function, resulting in glucose intolerance and type 2 diabetes mellitus (T2DM) (7). Extensive studies in rodent models have shown that diets resembling those commonly consumed by humans in modernized societies (i.e., westernized diet) cause endocrine pancreas deterioration and metabolic disorders (8, 9). Moreover, a high-fat diet supplemented with 20% sucrose (HFDS) has been utilized in rodents to study obesity-related morbidities (9–11).

The mechanisms whereby a hypercaloric diet can impair the pancreatic endocrine function, including glucose-stimulated insulin secretion (GSIS), may be partially related to decreased cell viability and antioxidant capacity and/or oxidative stress due to unbalanced reactive oxygen species (ROS) production and elimination (12). Indeed, studies have shown that excessive ROS production promotes apoptosis in insulin-secreting cell lines and pancreatic islets from rodents (13, 14). It has also been reported that long-term HFDS feeding induced pancreatic steatosis and increased oxidative stress and  $\beta$ -cell apoptosis in Bama minipigs (15).

Physical exercise promotes several systemic adaptations, including increased insulin sensitivity and improved glucose homeostasis, that reduce obesity and T2DM risks (16, 17). Studies have shown that the insulin response in peripheral tissues to regular exercise may be decreased in healthy individuals because physical activity attenuates the demand on  $\beta$ -cells (18, 19). Exercise also increases cellular antioxidant capacity, shielding  $\beta$ -cells from oxidative stress-induced dysfunction and death (20). Furthermore, exercise protects rodent and human pancreatic  $\beta$  cells under diabetogenic conditions, enhancing  $\beta$ -cell viability (21–23).

Unfortunately, due to the low cost, easy access, and taste, the consumption of highly processed foods high in fat and sugar is increasing globally (24). Previous work showed that combined exercise strategies to improve cardiorespiratory fitness and weight management could prevent diet-induced obesity and associated complications (25, 26). Indeed, a randomized controlled trial showed that two weeks of cycling exercise training improved  $\beta$ -cell function in prediabetic and T2DM individuals and decreased pancreatic fat independent of baseline glucose tolerance (27). Additionally, another study showed that combining moderate running exercise with cola-drinking ameliorated the metabolic disorders and promoted an approximately 50% increase in pancreatic  $\beta$  cell mass observed in the sedentary cola-drinking group (28).

Most of the findings related to this topic are disease models in which the metabolic disorder was already present (23, 29–31). Therefore, our study aimed to investigate the effects of combining aerobic exercise training with a high-fat and sucrose diet on cell viability and  $\beta$ -cell function in pancreatic islets of male C57BL/6J mice. Despite the reduced caloric intake, there was an increased feeding efficiency ratio (FER) and body mass in the HFDS group. These findings provide insights into preventive strategies that could mitigate the deleterious effects of the western diet on the endocrine pancreas and glucose homeostasis.

## MATERIAL AND METHODS

### Ethical Approval

All experiments were performed following the ethical guidelines for experiments with mice. Before starting the study, the protocol was prospectively approved by the Biomedical Sciences Institute's Animal Health Committee at the University of São Paulo (CEUA). The Brazilian Society of Science in Laboratory Animals (SBCAL) also approved the experimental protocols of this study (protocol #065, page 07 of Book 03).

### Experimental Design

Sixty-day-old male C57BL/6J mice were obtained from our breeding colony at the Institute of Biomedical Sciences at the University of São Paulo. These animals were randomly divided into four groups: control (C), fed a standard diet AIN-93M (3.83 kcal/g; 70% carbohydrate (cornstarch and dextrinized starch were chosen as the major source of carbohydrate for the AIN-93 diet. In addition, a small amount of sucrose), 20% protein (casein), and 10% fat (soybean) (32) with no training (i.e., sedentary); C + training (CTR, fed the standard diet with eight weeks of exercise (**Figure 1**); high-fat diet + sucrose (HFDS), fed a high fat and sucrose diet (5.2 kcal/g; 20% carbohydrate (cornstarch and dextrinized starch were chosen as the major source of carbohydrate), 20% protein (casein), 60% fat (Lard was chosen as the major source of fat and a small amount of soybean) + 20% sucrose diluted in drinking water adapted from (32–34) with no training; and HFDS + training (HFDSTR) (**Figure 1**).

The animals were housed in a temperature ( $22^{\circ}\text{C} \pm 2^{\circ}\text{C}$ ) and humidity (60%) controlled room with a 12h light/dark cycle. The mice had free access to food and water throughout the experimental protocol. Body weight and food and water intake were recorded weekly. Total food, water and energy intake were estimated by calculating the area under the curve (AUC) over 8 weeks. The FER was calculated by dividing the weight gained by kilocalories consumed over 8 weeks (35). At the end of the experimental protocol and 48 hours post-exercise training, mice were exposed to 70%  $\text{CO}_2$  and 30%  $\text{O}_2$  and euthanized by decapitation. Tissue weights of the epididymal, inguinal, and retroperitoneal adipose depots and soleus and gastrocnemius skeletal muscles were recorded and normalized to final body weight, and samples were stored at  $-80^{\circ}\text{C}$ .

## Exercise Intervention

As described in **Figure 1**, one week before starting the training protocol, mice were familiarized with the treadmill by carrying out four mild exercise sessions at 8.3 m/min for 10 min. The exercise capacity of each animal was evaluated through a maximal effort test ( $T_{max}$ ) on a motorized treadmill, starting at 5 m/min and increasing the speed by 1 m/min until exhaustion. The maximal exercise capacity (run-to-exhaustion) was defined as when the mouse stopped running spontaneously (36, 37). After the initial effort test, mice in the CTR and HFDSTR groups were exercise-trained 30 min/day, five times per week at 60% of the peak velocity treadmill running speed [adapted from (38)]. A  $T_{max}$  test was performed at the end of week three to maintain a constant relative intensity throughout the 8-week training period, and the treadmill running speed was adjusted accordingly. Another  $T_{max}$  test was conducted after completing the 8-week training period to assess post-test performance changes (**Figure 1**).

## Insulin Sensitivity, Glucose Tolerance and Homeostasis Model Assessments

After eight weeks and 48 hours after the last training session, the animals were restricted from eating for six hours and then subjected to a 30-min insulin tolerance test (short ITT) ( $n=42$ ) or a glucose tolerance test (GTT) ( $n=40$ ). For the short ITT, blood glucose was measured (Accue check active, Roche) before delivering (zero time point) an intraperitoneal insulin bolus (0.75 U/Kg) and after 5, 10, 15, 20, 25 and 30 minutes. The glucose disappearance rate (Kitt) was calculated based on the slope from a linear regression of glucose concentration from 5 to 25 minutes after insulin administration (39). For the GTT, blood glucose was measured before an intraperitoneal glucose injection (1 mg/g) (zero time point) and after 15, 30, 45, 60, 90, and 120 minutes. The data were presented as glucose area under the curve (AUC)


calculated from 120 min of an intraperitoneal bolus by total area using the trapezoidal rule. The homeostasis model assessment of insulin resistance (HOMA-IR) and the homeostasis model of  $\beta$ -cell function index (HOMA-B) (40) were calculated using fasting glucose and insulin concentrations.

## Fasting Serum Analyses

Forty-eight hours after the last training session, the mice were fasted overnight for 12 hours, decapitated, and then total trunk blood samples were collected ( $n=101$ ). Serum was separated by centrifugation at  $239 \times g$  for 20 min at  $4^{\circ}\text{C}$ , and samples were stored at  $-20^{\circ}\text{C}$ . Serum insulin concentrations were measured using a radioimmunoassay (RIA; PerkinElmer, Boston, MA, USA) with  $^{125}\text{I}$ -labeled insulin (41). The serum triglycerides (TG), total cholesterol (TC) and glucose concentrations were determined using an *Accutrend Plus*<sup>®</sup> device (Roche). The serum aspartate aminotransferase (AST) and alanine aminotransferase (ALT) activities were measured using commercially available kits (Labtest Diagnóstica, Minas Gerais, Brazil). Lipid peroxidation was estimated by quantifying thiobarbituric acid reactive species (TBARS) levels in the serum, as previously described (42).

## Citrate Synthase Activity

Forty-eight hours after the last training session, the maximum citrate synthase (CS) activity was determined in the gastrocnemius muscle ( $\sim 20$  mg) ( $n=31$ ). Briefly, 0.5 mM Tris-HCl extraction buffer and 1.0 mM EDTA pH 7.4 was used for protein extraction, and Tris/aminomethane (100 mM), 5,5'-Dithiobis (2-nitrobenzoic acid) (DTNB; 0.2 mM), acetyl CoA (0.1 mM) and Triton X-100 (0.1% v/v), pH 8.1 was used to measure the enzymatic activity. The reaction was initiated by adding 3  $\mu\text{L}$  of muscle homogenate and 10  $\mu\text{L}$  of oxaloacetic acid (10 mM final concentration) to the samples. The reactions were monitored spectrophotometrically by measuring the absorbance



at 412 nm at 25°C, as previously described (43). The activity was measured by quantifying the complex formed between Coenzyme A (CoA) released by the medium and oxaloacetate DTNB. The maximal enzyme activity was expressed as nanomoles per minute per milligram of protein.

### Liver Lipid Droplet Determination

The livers of 12 mice were fixed in 10% formaldehyde solution for eight hours in individual cassettes to evaluate liver histology. Then, the fixed samples were incubated in 70% alcohol overnight. Next, the samples were dehydrated through a series of baths in 95% alcohol, 100% alcohol and xylene. Following dehydration, the tissue samples were embedded in paraffin at 60°C, and a microtome (Zeiss, Jena, Germany) cut the samples into 5-micron semi-serial slices, which were then stained with hematoxylin and eosin (H&E). The cellular histology was evaluated, and the intracellular liver lipid droplets (LLDs) were quantified. Twenty images from each animal were obtained using a Nikon Eclipse Ti-U microscope at 20× magnification coupled with a Nikon DS-R1 digital camera and NIS-Elements BR 3.1 software. These images were projected onto a high-resolution LCD monitor. The test system contained 270 symmetrically distributed points produced from a transparent sheet placed on the monitor screen. Each point that remained on the LLDs was counted and named EP. Then the EP/P ratio, which corresponds to the amount of LLDs, was calculated (44). The results were expressed as a percentage.

### Pancreatic Islet Isolation

Forty-eight hours after the last training session and following decapitation, the pancreas from 101 mice were collected, and islets were isolated by collagenase digestion (45, 46). Briefly, the pancreas was inflated with Hanks solution containing 0.68 g/L type IV collagenase (Sigma–Aldrich Chemical, St. Louis, MO, USA), excised and maintained at 37°C for 25 min. The digested tissue was harvested, and the islets were hand-picked. Krebs–Henseleit buffer containing 136 mM NaCl, 5.36 mM KCl, 4.17 mM NaHCO<sub>3</sub>, 1.26 mM CaCl<sub>2</sub>·2H<sub>2</sub>O, 0.81 mM MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.3 mM NaHPO<sub>4</sub>·12H<sub>2</sub>O and 0.44 mM KH<sub>2</sub>PO<sub>4</sub> was used for islet isolation and pooling. The pancreatic islet samples were utilized in the cell viability, DNA fragmentation, GSIS and ROS assays.

### Pancreatic Islet Cellular Viability

Pancreatic islets were disrupted by trypsin digestion and gentle pipetting (47). Living, apoptotic and dead cells were identified with the Guava ViaCountR assay (Millipore, Billerica, MA, USA) based on staining the cytoplasm and nucleus with different dyes. One dye is membrane-permeable and stains all nucleated cells, whereas the other is membrane-impermeable and only stains damaged cells, indicating apoptotic and dying cells. The cell suspensions were incubated with the ViaCount reagent for five minutes at room temperature. Data acquisition was performed on a Guava EasyCyte flow cytometer (Millipore, Billerica, MA, USA), and data analysis was performed using the ViaCount software module.

### Islet Cell Apoptosis

Islet dispersed cell apoptosis detection was performed by using the Cell Death Detection ELISA<sup>PLUS</sup> kit (Roche Diagnostics, Mannheim, Germany), according to the manufacturer's instructions. This assay measures the amount of histone-associated DNA fragments (mono- and oligonucleosomes). Herein, batches containing 80 islets per mouse were lysed in 100 µL of lysis buffer, and the cytosolic and nuclear fractions were separated by centrifugation at 200 × g for 10 min. The mean absorbance at 405 nm of each sample was normalized to the DNA content in the nuclear fraction.

### Glucose Stimulated Insulin Secretion (GSIS) and Net Cytosolic ROS Production

GSIS was assessed by incubating five islets in 0.5 mL Krebs–Henseleit buffer containing 0.2% albumin and 5.6 mM glucose (pre-incubation phase) at 30°C for 30 min. After the pre-incubation phase, the islets were incubated in 2.8 mM and 16.7 mM glucose for 60 min (47). Then, 300 µL of the supernatant were stored at -20°C until determining “insulin secretion”. For total “insulin content”, the pancreatic islets were disrupted in a 52 ethanol:17 water:1 hydrochloric acid solution and sonicated. The “insulin secretion” and “insulin content” were measured by radioimmunoassay (RIA) (41). The insulin secretion data (corresponding to groups of five islets per glucose concentration) was expressed as the insulin secretion/insulin content ratio.

Islet superoxide content was evaluated by flow cytometry (Guava EasyCyte, Millipore), monitoring its reaction with dihydroethidium (DHE; Life Technologies, Eugene, Oregon, USA) (48), which generates the fluorescent product 2-hydroxyethidium (2-OH-E<sup>+</sup>) (49). Groups of 25 islets were pre-incubated at 37°C for 30 min in Krebs–Henseleit buffer, pH 7.4 containing 5.6 mM glucose, and then incubated for one hour in Krebs–Henseleit buffer with 0.2% BSA and 2.8 mM or 16.7 mM glucose (47). Next, 50 µL DHE was added, and the samples were incubated in the dark at room temperature for 20 min. The islets were trypsin-digested, gently pipetted up and down and resuspended in 200 µL of RPMI 1640 culture medium at 37°C for two minutes. Afterward, an additional 600 µL of RPMI-1640 medium was added to inactivate the trypsin, as described previously with adaptations (47). The yellow fluorescence was measured using an excitation wavelength of 488 nm, and the emission signal was acquired with a yellow filter (583 nm ± 26 nm). Despite the superoxide radical's short half-life, 5000 events were recorded per group of 25 islets per mouse.

### Statistical Analysis

GraphPad Prism v.9 (GraphPad Software, La Jolla, CA, USA) was used to perform the statistical analyses. All data are expressed as the mean ± standard deviation (SD) using a scatter dot plot that allows the reader to visualize the number of individual samples in each analysis. Two-way ANOVA with Tukey's *post-hoc* test was used to analyze diet and training effects.

For all analyses, a P-value of <0.05 was considered statistically significant.

## RESULTS

### Exercise Training Efficiency

Exercise training, irrespective of the diet, resulted in significant increases in run-to-exhaustion times [ $F = (1, 58) = 39.78$ ,  $P < 0.0001$ ] (**Figure 2A**). The run-to-exhaustion times of the CTR and HFDSTR groups increased by 55.3 and 53.8%, respectively (CTR, week 8 vs. 0: -4.155, 95% CI -6.932 to -1.378,  $P = 0.0004$  and HFDS, week 8 vs. 0: -4.208, 95% CI -7.415 to -1.002,  $P = 0.0028$ ). On the other hand, the C (-1.395, 95% CI -3.879 to 1.089,  $P = 0.6448$ ) and HFDS (-1.413, 95% CI -4.031 to 1.205,  $P = 0.6888$ ) groups were not significantly different (18.7 and 18.4%, respectively) (**Figure 2A**). As shown in **Figure 2B**, CS activity in the gastrocnemius muscle increased by 1.6 and 2-fold in the exercise-trained groups compared to their respective controls without training (CTR vs. C: -0.1251, 95% CI -0.2247 to -0.02541,  $P = 0.0098$ ) and HFDSTR vs. HFDS: -0.1405, 95% CI -0.2513 to -0.02971,  $P = 0.0090$ ).

### Body Weight Changes

The CTR and HFDS animals displayed a 1.8-fold-decrease and 2.1-fold increase, respectively ( $3.5 \pm 1.2$ g and  $14 \pm 3.2$ g) in total body mass gain compared to the C mice [C vs. CTR: 3.0, 95% CI 0.5 to 5.50,  $P < 0.01$ ; C vs. HFDS: -7.4, 95% CI -10 to -4.7,  $P < 0.0001$ ; HFDS vs. HFDSTR: 7.01, 95% CI 4.3 to 9.74,  $P < 0.0001$ ] (**Figure 3A**). Furthermore, the HFDSTR mice had 2-fold decreased the total body mass gain than in HFDS group (HFDS vs. HFDSTR: 7.01, 95% CI 4.3 to 9.74,  $P <$


0.0001). Throughout the eight-week experimental period, the CTR mice gained 0.43 g of b.w./week while the HFDS mice gained 1.75 g/week on average compared to the C and HFDSTR groups which only gained 0.80 and 0.87 g/week ( $P < 0.0001$ , **Figure 3B**).

### Food Intake


As shown in **Figure 3C**, the caloric intake was reduced in both HFDS and HFDSTR by 90 and 85% compared to the C and CTR groups, respectively (C vs. HFDS: 14.60, 95% CI 0.8319 to 28.37,  $P = 0.0343$  and CTR vs. HFDSTR: 27.30, 95% CI 13.96 to 40.64,  $P < 0.0001$ ). Nonetheless, the FER was decreased in CTR mice compared to the C group (C vs. CTR: 0.2556, 95% CI 0.07577 to 0.4355,  $P = 0.0028$ ) (**Figure 3D**). Despite the 14 kcal decrease in the HFDS animals' total caloric intake during the eight-week experiment (C vs. HFDS: -0.6730, 95% CI -0.8578 to -0.4882,  $P < 0.0001$ ) (**Figure 3C**), there was an increase in the FER, an observation that was attenuated in the HFDSTR mice (HFDS vs. HFDSTR: 0.5750, 95% CI 0.3854 to 0.7646,  $P < 0.0001$ ) (**Figure 3D**).

### Tissue Weights and Liver Lipid Droplets Percentages

**Table 1** shows the Tissue Weights. Despite adjusting for body weight, a diet effect ( $P < 0.05$ ) was observed and the HFDS group presented increased epididymal (C vs. HFDS: -2.292, 95% CI -3.487 to -1.097,  $P < 0.0001$ ) inguinal (C vs. HFDS: -1.792, 95% CI -3.349 to -0.2341,  $P = 0.0191$ ) and retroperitoneal (C vs. HFDS: -1.664, 95% CI -2.546 to -0.7828,  $P < 0.0001$ ) WAT weights, while the soleus [ $F(1, 28) = 0.01299$ ,  $P = 0.9101$ ] and gastrocnemius [ $F(1, 32) = 0.06137$ ,  $P = 0.8059$ ] muscle weights were not significantly altered. Additionally, exercise training did not alter the weights of the epididymal [ $F(1, 53) = 6.106$ ,  $P = 0.0167$ ], inguinal [ $F(1, 32) =$



**FIGURE 2** | Effects of endurance training on exercise capacity (run-to-exhaustion time). **(A)** C, CTR, HFDS and HFDSTR (week 0 vs. after week 8). Differences in CTR and HFDSTR between week 0 vs. 8, respectively ( $P < 0.01$ ). **(B)** Citrate Synthase Activity in the gastrocnemius muscle of C, CTR, HFDS and HFDSTR after eight weeks: Endurance training effect ( $P < 0.01$ ). Data are expressed as the mean  $\pm$  SD.



**FIGURE 3** | Effects of eight weeks of exercise training and HFDS diet on metabolic parameters in male C57BL6J mice. **(A)** Total body mass gain after eight weeks. C vs. HFDS, HFDS vs. HFDSTR, CTR vs. HFDSTR ( $P < 0.01$ ). **(B)** Changes in body mass over the eight-week intervention. C vs. CTR ( $P < 0.05$ ), C vs. HFD ( $P < 0.001$ ), HFDS vs. HFDSTR and CTR vs. HFDSTR ( $P < 0.05$ ). **(C)** Total caloric intake; C vs. HFDS, HFDS vs. HFDSTR ( $P < 0.01$ ). **(D)** FER: C vs. CTR, C vs. HFDS, HFDS vs. HFDSTR and CTR vs. HFDSTR ( $P < 0.001$ ). Data were analyzed by two-way ANOVA as described in the Materials and Methods. All data presented as the mean  $\pm$  SD.

**TABLE 1** | Tissue weights and LLDs in hepatocytes following the eight-week intervention in the C, CTR, HFDS and HFDSTR groups.

Variables	N	C	CTR	HFDS	HFDSTR
Epididymal WAT	47	2.15 $\pm$ 0.97	1.67 $\pm$ 0.96	4.43 $\pm$ 1.63*	3.24 $\pm$ 1.43 <sup>#</sup>
Inguinal WAT	36	1.72 $\pm$ 0.80	1.53 $\pm$ 0.88	3.51 $\pm$ 1.81*	2.53 $\pm$ 1.28
Retroperitoneal WAT	35	1.07 $\pm$ 0.43	0.86 $\pm$ 0.38	2.73 $\pm$ 1.02*	2.21 $\pm$ 0.72
Soleus Muscle	32	0.07 $\pm$ 0.007	0.07 $\pm$ 0.007	0.06 $\pm$ 0.02	0.08 $\pm$ 0.02
Gastrocnemius Muscle	36	0.83 $\pm$ 0.31	1.0 $\pm$ 0.16	0.81 $\pm$ 0.10	0.97 $\pm$ 0.15
LLDs	12	ND	ND	37.2 $\pm$ 2.5*	22.5 $\pm$ 5.4 <sup>#§</sup>

Tissues weights are expressed as g/g body mass. LLDs were expressed as a percentage. N indicates the total number of mice used. ND is non-detectable. Symbols represents post-hoc test ( $P < 0.05$ ): \*C vs. HFDS and <sup>#</sup>CTR vs. HFDSTR and <sup>§</sup>HFDS vs. HFDSTR. Data are expressed as the mean  $\pm$  SD.

1.933,  $P = 0.1740$ ] and retroperitoneal [ $F(1, 31) = 2.508, P = 0.1234$ ] tissues or the soleus [ $F(1, 28) = 3.753, P = 0.0629$ ] and gastrocnemius [ $F(1, 32) = 0.06137, P = 0.8059$ ] muscles (**Table 1**).

As expected, there were no detectable (ND) LLDs in the hepatocytes of the C and CTR mice (0.000, 95% CI -6.316 to 6.316,  $P > 0.9999$ ). In contrast, the LLD percentage in the HFDS mice was markedly increased to over 35% (-37.75, 95% CI -44.45 to -31.05,  $P < 0.0001$ ). Notably, the hepatocytes from the HFDSTR group contained 39.5% fewer liver LLDs than those from HFDS mice (15.25, 95% CI 9.134 to 21.37,  $P < 0.0001$ ) (**Table 1**).

### Biochemical Analysis

A summary of the biochemical marker data for each group is presented in **Table 2**. Compared to C, the HFDS group presented higher glucose (-73.98, 95% CI -94.89 to -53.07,  $P < 0.0001$ ), insulin (0.3367, 95% CI 0.07933 to 0.5940,  $P = 0.0068$ ), HOMA-B (-23.54, 95% CI -42.39 to -4.694,  $P < 0.005$ ) and TG (22.02, 95% CI 1.838 to 42.19,  $P < 0.005$ ). Several of these variables including glucose, (61.33, 95% CI 40.69 to 81.96,  $P < 0.0001$ ), insulin (-0.3492, 95% CI -0.6065 to -0.09183,  $P = 0.0049$ ), and HOMA-B (38.70, 95% CI 18.83 to 58.57,  $P < 0.0001$ ) were decreased in the

**TABLE 2** | Serum biochemical profile following the eight-week intervention in the C, CTR, HFDS and HFDSTR groups.

Variables	N	C	CTR	HFDS	HFDSTR
Insulin (ng/dL)	31	0.71 ± 0.12	0.75 ± 0.09	1.05 ± 0.02*	0.7 ± 0.18 <sup>§</sup>
Glucose (mg/dL)	15	171.30 ± 29.63	162.5 ± 19.46	245.3 ± 32.72*	184.0 ± 18.16 <sup>§</sup>
HOMA-IR (UA)	31	2.52 ± 0.18	2.60 ± 0.19	2.80 ± 0.23	2.85 ± 0.30
HOMA-β (UA)	21	67.70 ± 7.12	54.4 ± 11.81	90.3 ± 3.17*	51.6 ± 12.26 <sup>§</sup>
TG (mg/dL)	32	118.42 ± 12.43	111.80 ± 6.50	140.40 ± 13.23*	154.12 ± 22.02 <sup>#</sup>
TC (mg/dL)	31	175.75 ± 3.65	173.66 ± 3.77	176.62 ± 2.38	170.25 ± 6.75
ALT (UI/mL)	18	1.10 ± 0.09	1.10 ± 0.07	1.40 ± 0.28	1.32 ± 0.18
AST (UI/mL)	18	1.10 ± 0.11	1.10 ± 0.06	1.25 ± 0.20	1.25 ± 0.17
AST/ALT	18	1.01 ± 0.05	1.00 ± 0.01	1.11 ± 0.13	1.05 ± 0.01
TBARS (nmol MDA/mL)	32	0.70 ± 0.17	0.63 ± 0.30	0.58 ± 0.40	0.46 ± 0.21


TG, triglycerides; TC, total cholesterol. Differences ( $P < 0.05$ ) were observed in \*C vs. HFDS and <sup>#</sup>HFDS vs. HFDSTR, and <sup>§</sup>HFDS vs. HFDSTR. Data are expressed as the mean ± SD.


HFDSTR group. There were no significant alterations in HOMA-IR [ $F(1, 27) = 0.03797$ ,  $P = 0.8470$ ], TC values [ $F(1, 30) = 0.8323$ ,  $P = 0.3689$ ], ALT [ $F(1, 14) = 0.1409$ ,  $P = 0.7130$ ] and AST [ $F(1, 14) = 8.490e-9$ ,  $P > 0.9999$ ] activities, or TBARS levels [ $F(1, 28) = 0.1480$ ,  $P = 0.7033$ ] between groups.

### Insulin Sensitivity and Glucose Tolerance

The GTT of HFDS mice (**Figure 4A**), also represented by the AUC of glucose in **Figure 4B**, was 61% greater than in the C group (18102, 95% CI -25249 to -10956,  $P < 0.0001$ ) and 55%

greater than the HFDSTR group (16291, 95% CI 9317 to 23265,  $P < 0.0001$ ). The AUCs of glucose for the C and HFDSTR groups were not significantly different (-1812, 95% CI -8958 to 5334,  $P = 0.9030$ ). In contrast, the KITT of the HFDS group was 2.3-fold lower than the C group (3.657, 95% CI 1.214 to 6.100,  $P = 0.0014$ ). Furthermore, as shown in **Figure 4C**, the KITT of the HFDSTR mice was 1.9-fold higher than that of the HFDS mice (-2.507, 95% CI -4.600 to -0.4147,  $P = 0.0133$ ) but was not significantly different from the C group (-1.232, 95% CI to 3.532,  $P = 0.5729$ ).





**FIGURE 5** | Dispersed pancreatic islet after the eight-week intervention in the C, CTR, HFDS and HFDSTR groups. **(A)** Percentage of viable pancreatic cells: C vs. HFDS and HFDS vs. HFDSTR ( $P < 0.05$ ). **(B)** Cellular apoptosis: C vs. HFDS and HFDS vs. HFDSTR ( $P < 0.05$ ). **(C)** DNA fragmentation: C vs. HFDS and HFDS vs. HFDSTR ( $P < 0.05$ ). Data are expressed as the mean  $\pm$  SD.

## Pancreatic Islet Cellular Viability and Islet Cell Apoptosis

As expected, the cellular viability of dispersed pancreatic islets was reduced by 38% in the HFDS group compared to the C group (19.97, 95% CI 4.615 to 35.32,  $P = 0.0071$ ). This effect was attenuated in the HFDSTR mice (-17.17, 95% CI -30.29 to -4.04,  $P = 0.0068$ ) which presented 53% more viable cells vs HFDS (Figure 5A). Conversely, cellular apoptosis (Figure 5B) and DNA fragmentation (Figure 5C) increased 1.6 (-2.7, 95% CI -4.605 to -0.7947,  $P = 0.0042$ ) and 3.9-fold (2.752, 95% CI -5.018 to -0.4859,  $P = 0.0170$ ) in the HFDS mice compared to the C group. Notably, in the HFDSTR mice, there was a 29% reduction in cellular apoptosis (2.108, 95% CI 0.2920 to 3.925  $P = 0.0197$ ), accompanied by a 99% reduction in the DNA fragmentation compared to the HFDS group (3.669, 95% CI 1.403 to 5.935,  $P = 0.0024$ ), and not significantly different from C mice (-0.5913, 95% CI -2.496 to 1.314,  $P = 0.8165$ ) and (0.9167, 95% CI -1.181 to 3.015,  $P = 0.5729$ ), respectively.

## Glucose Stimulated Insulin Secretion (GSIS) and Net Cytosolic ROS Production

The GSIS rate was increased 2-fold with high glucose (16.7 mmol/L) compared to low glucose (2.8 mmol/L) in the C (-0.01210, 95% CI -0.02088 to -0.003325,  $P = 0.0019$ ), CTR (-0.01071, 95% CI -0.01949 to -0.001934,  $P = 0.0080$ ) and HFDSTR (-0.009751, 95% CI -0.01776 to -0.001739,  $P = 0.0082$ ) groups. As expected, the


HFDS group did not display an increase in static insulin secretion induced by the high glucose concentration (0.0001243, 95% CI -0.007293 to 0.007542,  $P > 0.9999$ ) (Figure 6A). Lastly, reduced indirect ROS levels were detected in the high glucose condition compared to the low glucose in the C (78.58, 95% CI 26.26 to 130.9,  $P = 0.0005$ ), CTR (70.29, 95% CI 9.079 to 131.5,  $P = 0.0143$ ) and HFDSTR (85.68, 95% CI 3.135 to 168.2,  $P = 0.0368$ ) groups, but not in the HFDS mice (39.00, 95% CI -2.267 to 80.27,  $P = 0.0764$ ) (Figure 6B).

## DISCUSSION

The present study showed that moderate-intensity exercise training could avoid some of the deleterious effects of a high fat and sucrose diet on murine pancreatic cell viability, glucose tolerance and insulin sensitivity. The benefits of this specific physical exercise regimen were associated with maintaining GSIS and cytosolic ROS levels, consequently improving glucose tolerance and insulin sensitivity.

Despite the HFDS group ingesting fewer calories, the FER was increased, and these animals gained more body and WAT weight than the C group. A classical situation of body mass gain, particularly fatty mass, is related to reduced energetic expenditure and increased energetic intake. Unfortunately, in our study, we could not analyze indirect calorimetry. However, evidence indicates that long-term ingestion of a sugar-rich or





**FIGURE 6** | Effects of eight weeks of exercise training and HFDS diet on pancreatic islets. **(A)** GSIS normalized for total insulin content. Glucose responsiveness (2.8 vs. 16.7mM) in C ( $P < 0.0001$ ), CTR ( $P < 0.0001$ ) and HFDSTR groups ( $P < 0.0001$ ). **(B)** Ethidium bromide content and average fluorescence intensity by DHE in dispersed pancreatic islets after 1-hour of glucose stimulus (2.8 vs. 16.7mM). Glucose effect (2.8 vs. 16.7mM) in C, CTR and HFDSTR ( $P < 0.001$ ). Data are expressed as the mean  $\pm$  SD.

high-fat diet causes severe obesity in rats without hyperphagia (50). On the other hand, another study (51) showed that energy intake was similar between the high-fat and low-fat diet groups due to the HFD group consuming less food. The latter study also showed that energy expenditure was lower in the HFD group than in the low-fat diet group, increasing body weight and fat mass and contributing to insulin resistance. Accordingly, the HFDS mice presented impaired glucose tolerance, higher body fat mass, reduced caloric intake, and insulin sensitivity in the present study.

We found that exercise (CTR vs. C and HFDSTR vs. HFDS) attenuated the observed weight gain and elevated the FER but did not affect the epididymal, inguinal, and retroperitoneal tissue weights. Notably, the average weight of the HFDSTR mice was comparable to the C group. It is plausible that intramuscular and other ectopic fat content could be involved in body weight regulation, insulin sensitivity and glucose tolerance and improved by exercise training (52). Future studies measuring the whole fat and fat-free mass content are needed to investigate this possibility.

The HFDS group displayed increased TG levels and a higher percentage of LLDs than the C group, but the AST and ALT activities remained unchanged. Non-alcoholic fatty liver disease (NAFLD) is characterized by the aberrant intracellular accumulation of lipids (>5%) in hepatocytes (53) and is highly correlated with obesity, insulin resistance and T2DM (54). While exercise training did not change the TG content in HFDSTR mice, it attenuated the hepatic steatosis grade in these animals. In this sense, exercise could delay non-alcoholic steatohepatitis progression.

We also evaluated endocrine pancreas adaptations by monitoring insulin and glucose metabolism in response to a high-fat and sucrose diet and exercise training. The HFDS mice had fewer viable pancreatic islets, impaired GSIS, reduced cytosolic ROS

content, and elevated serum insulin concentrations and HOMA- $\beta$  values, likely due to the excess intake of fatty acids and sugar. Indeed, previous studies have reported that increased  $\beta$ -cell mass over time may lead to a reduced  $\beta$ -cell mass, culminating in  $\beta$ -cell failure (9, 55). Additional studies evaluating the parameters involved in the pancreatic  $\beta$ -cell adaptations are necessary to elucidate further the precise mechanisms involved.

ROS are constitutively produced and removed, driving a redox state and acting as a signal for cell processes, including pancreatic  $\beta$ -cell signaling and control of insulin secretion (56). Physiologically, acutely elevated glucose concentrations reduce net ROS detection, preventing pancreatic cells from these powerful free radicals and oxidants (57). It is important to point out that the redox homeostasis systems of eukaryotic cells are highly compartmentalized (58), and we only measured net cytosolic ROS production and not mitochondrial and non-mitochondrial ROS production and activity. Our findings demonstrated that the HFDS mice lost, at least in part, the GSIS-associated mechanism with no significant alterations in the systemic TBARS levels compared to the C group. However, the pancreatic islets from HFDS fed mice were less responsive to the physiological net ROS production induced by glucose stimulus. Thus, our results indicate that moderate exercise training effectively stimulates protective hormetic responses (59) and ROS maintenance (60) to compensate for the peripheral insulin resistance elicited by the high-fat and sucrose diet.

Once again, we did not measure antioxidant enzyme content, which could be associated with the observed reduction in ROS production and apoptotic protein content in the pancreatic islet of rats after eight weeks of endurance training (60, 61). Importantly, the eight-week exercise training protocol used in the present study attenuated and prevented the high-fat and sucrose diet-induced effects in mice, evidenced by the

maintenance of glucose homeostasis and insulin sensitivity in the HFDSTR animals. These findings are probably associated with more efficient pancreatic islets since exercise increased cell viability and reduced DNA fragmentation and apoptosis, maintaining ROS net content, and preventing GISIS impairment.

Although these results are not necessarily surprising, our study provides valuable insights into how hypercaloric diets and regular aerobic exercise training could prevent or improve obesity-related complications. However, it is unclear if exercise alone can offset the negative effects of foods rich in fat and sugar. Additional studies are required to explore the molecular mechanisms underlying the observed responses and adaptations.

## CONCLUSION

Consistent with the hypothesis of this study, we showed that moderate exercise training performed simultaneously with a hypercaloric diet protected insulin sensitivity, glucose tolerance and pancreatic islets function in male C57BL/6J mice. Furthermore, this study provides experimental evidence demonstrating that moderate aerobic exercise training could attenuate the risks and complications associated with the westernized diet.

## DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## ETHICS STATEMENT

The animal study was reviewed and approved by Biomedical Sciences Institute's Animal Health Committee at the University

## REFERENCES

1. Popkin BM, Adair LS, Ng SW. Global Nutrition Transition and the Pandemic of Obesity in Developing Countries. *Nutr Rev* (2012) 70(1):3–21. doi: 10.1111/j.1753-4887.2011.00456.x
2. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, Regional, and National Prevalence of Overweight and Obesity in Children and Adults During 1980–2013: A Systematic Analysis for the Global Burden of Disease Study 2013. *Lancet* (2014) 384(9945):766–81. doi: 10.1016/S0140-6736(14)60460-8
3. Ishimoto T, Lanaspa MA, Rivard CJ, Roncal-Jimenez CA, Orlicky DJ, Cicerchi C, et al. High-Fat and High-Sucrose (Western) Diet Induces Steatohepatitis That is Dependent on Fructokinase. *Hepatology* (2013) 58(5):1632–43. doi: 10.1002/hep.26594
4. Dixon JB. The Effect of Obesity on Health Outcomes. *Mol Cell Endocrinol* (2010) 316(2):104–8. doi: 10.1016/j.mce.2009.07.008
5. Nurwanti E, Uddin M, Chang JS, Hadi H, Syed-Abdul S, Su EC, et al. Roles of Sedentary Behaviors and Unhealthy Foods in Increasing the Obesity Risk in Adult Men and Women: A Cross-Sectional National Study. *Nutrients* (2018) 10(6). doi: 10.3390/nu10060704
6. Kahn SE. The Relative Contributions of Insulin Resistance and Beta-Cell Dysfunction to the Pathophysiology of Type 2 Diabetes. *Diabetologia* (2003) 46(1):3–19. doi: 10.1007/s00125-002-1009-0

of São Paulo (CEUA). The Brazilian Society of Science in Laboratory Animals (SBCAL) also approved the experimental protocols of this study. The approved protocol number is 065, as stated on Sheet 07 of Book 03.

## AUTHOR CONTRIBUTIONS

KV contributed to the project's conception and experimental design, data collection, analysis, and interpretation, and drafting of and revising of the article. CM and FE performed experiments and collected data. JG interpreted the data and revised the manuscript. AC and CCRO conceived and designed the experiments, analyzed, and interpreted the data and interpretation and revised the manuscript. All authors contributed to the article and approved the submitted version.

## FUNDING

This work was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq number: 159288/2011-8) and Fundação de Apoio à Pesquisa do Estado de São Paulo (FAPESP Number: 2019/20464-8).

## ACKNOWLEDGMENTS

We are grateful for the technical assistance provided by Marilu Mazzaro (ICB/USP) and Mikaely Ribeiro (UMC), Dr. Flávio de Oliveira Pires (EACH/USP) for helping with statistics and editorial comments, Dr. Felipe Natali Almeida, Dr. Lyanne C. C. Araujo, Dr. Marcelo Nicolás Muscará and Dr. Simone Aparecida Teixeira (ICB/USP) for the technical assistance and discussions.

7. Kahn SE, Prigeon RL, Schwartz RS, Fujimoto WY, Knopp RH, Brunzell JD, et al. Obesity, Body Fat Distribution, Insulin Sensitivity and Islet Beta-Cell Function as Explanations for Metabolic Diversity. *J Nutr* (2001) 131(2):354S–60S. doi: 10.1093/jn/131.2.354S
8. Buettner R, Schölmerich J, Bollheimer LC. High-Fat Diets: Modeling the Metabolic Disorders of Human Obesity in Rodents. *Obes (Silver Spring)* (2007) 15(4):798–808. doi: 10.1038/oby.2007.608
9. Burchfield JG, Kebede MA, Meoli CC, Stöckli J, Whitworth PT, Wright AL, et al. High Dietary Fat and Sucrose Results in an Extensive and Time-Dependent Deterioration in Health of Multiple Physiological Systems in Mice. *J Biol Chem* (2018) 293(15):5731–45. doi: 10.1074/jbc.RA117.000808
10. Kowalski GM, Kraakman MJ, Mason SA, Murphy AJ, Bruce CR. Resolution of Glucose Intolerance in Long-Term High-Fat, High-Sucrose-Fed Mice. *J Endocrinol* (2017) 233(3):269–79. doi: 10.1530/JOE-17-0004
11. Yang ZH, Miyahara H, Takeo J, Katayama M. Diet High in Fat and Sucrose Induces Rapid Onset of Obesity-Related Metabolic Syndrome Partly Through Rapid Response of Genes Involved in Lipogenesis, Insulin Signalling and Inflammation in Mice. *Diabetol Metab Syndr* (2012) 4(1):32. doi: 10.1186/1758-5996-4-32
12. Pi J, Collins S. Reactive Oxygen Species and Uncoupling Protein 2 in Pancreatic  $\beta$ -Cell Function. *Diabetes Obes Metab* (2010) 12 Suppl 2:141–8. doi: 10.1111/j.1463-1326.2010.01269.x

13. Roma LP, Bosqueiro JR, Cunha DA, Carneiro EM, Gurgul-Convey E, Lenzen S, et al. Protection of Insulin-Producing Cells Against Toxicity of Dexamethasone by Catalase Overexpression. *Free Radic Biol Med* (2009) 47(10):1386–93. doi: 10.1016/j.freeradbiomed.2009.08.010
14. Newsholme P, Haber EP, Hirabara SM, Rebelato EL, Procopio J, Morgan D, et al. Diabetes Associated Cell Stress and Dysfunction: Role of Mitochondrial and non-Mitochondrial ROS Production and Activity. *J Physiol* (2007) 583(Pt 1):9–24. doi: 10.1113/jphysiol.2007.135871
15. Zhao ZZ, Xin LL, Xia JH, Yang SL, Chen YX, Li K. Long-Term High-Fat High-Sucrose Diet Promotes Enlarged Islets and  $\beta$ -Cell Damage by Oxidative Stress in Bama Minipigs. *Pancreas* (2015) 44(6):888–95. doi: 10.1097/MPA.0000000000000349
16. Zierath JR. Invited Review: Exercise Training-Induced Changes in Insulin Signaling in Skeletal Muscle. *J Appl Physiol* (1985) (2002) 93(2):773–81. doi: 10.1152/jappphysiol.00126.2002
17. Ropelle ER, Pauli JR, Prada PO, de Souza CT, Picardi PK, Faria MC, et al. Reversal of Diet-Induced Insulin Resistance With a Single Bout of Exercise in the Rat: The Role of PTP1B and IRS-1 Serine Phosphorylation. *J Physiol* (2006) 577(Pt 3):997–1007. doi: 10.1113/jphysiol.2006.120006
18. Galbo H, Hedekov CJ, Capito K, Vinten J. The Effect of Physical Training on Insulin Secretion of Rat Pancreatic Islets. *Acta Physiol Scand* (1981) 111(1):75–9. doi: 10.1111/j.1748-1716.1981.tb06707.x
19. Brandt C, Pedersen BK. The Role of Exercise-Induced Myokines in Muscle Homeostasis and the Defense Against Chronic Diseases. *J BioMed Biotechnol* (2010) 2010:520258. doi: 10.1155/2010/520258
20. Calegari VC, Abrantes JL, Silveira LR, Paula FM, Costa JM, Rafacho A, et al. Endurance Training Stimulates Growth and Survival Pathways and the Redox Balance in Rat Pancreatic Islets. *J Appl Physiol* (2012) 112(5):711–8. doi: 10.1152/jappphysiol.00318.2011
21. Paula FMM, Leite NC, Borck PC, Freitas-Dias R, Cnop M, Chacon-Mikahil MPT, et al. Exercise Training Protects Human and Rodent  $\beta$  Cells Against Endoplasmic Reticulum Stress and Apoptosis. *FASEB J* (2018) 32(3):1524–36. doi: 10.1096/fj.201700710R
22. Paula FM, Leite NC, Vanzela EC, Kurauti MA, Freitas-Dias R, Carneiro EM, et al. Exercise Increases Pancreatic  $\beta$ -Cell Viability in a Model of Type 1 Diabetes Through IL-6 Signaling. *FASEB J* (2015) 29(5):1805–16. doi: 10.1096/fj.14-264820
23. Nieuwoudt S, Fealy CE, Foucher JA, Scelsi AR, Malin SK, Pagadala M, et al. Functional High-Intensity Training Improves Pancreatic  $\beta$ -Cell Function in Adults With Type 2 Diabetes. *Am J Physiol Endocrinol Metab* (2017) 313(3):E314–E20. doi: 10.1152/ajpendo.00407.2016
24. Kearney J. Food Consumption Trends and Drivers. *Philos Trans R Soc Lond B Biol Sci* (2010) 365(1554):2793–807. doi: 10.1098/rstb.2010.0149
25. Ortega FB, Ruiz JR, Labayen I, Lavie CJ, Blair SN. The Fat But Fit Paradox: What We Know and Don't Know About it. *Br J Sports Med* (2018) 52(3):151–3. doi: 10.1136/bjsports-2016-097400
26. Church T. Exercise in Obesity, Metabolic Syndrome, and Diabetes. *Prog Cardiovasc Dis* (2011) 53(6):412–8. doi: 10.1016/j.pcad.2011.03.013
27. Heiskanen MA, Motiani KK, Mari A, Saunavaara V, Eskelinen JJ, Virtanen KA, et al. Exercise Training Decreases Pancreatic Fat Content and Improves Beta Cell Function Regardless of Baseline Glucose Tolerance: A Randomised Controlled Trial. *Diabetologia* (2018) 61(8):1817–28. doi: 10.1007/s00125-018-4627-x
28. Otero-Losada M, González J, Müller A, Ottaviano G, Cao G, Azzato F, et al. Exercise Ameliorates Endocrine Pancreas Damage Induced by Chronic Cola Drinking in Rats. *PLoS One* (2016) 11(5):e0155630. doi: 10.1371/journal.pone.0155630
29. Park S, Hong SM, Lee JE, Sung SR. Exercise Improves Glucose Homeostasis That has Been Impaired by a High-Fat Diet by Potentiating Pancreatic Beta-Cell Function and Mass Through IRS2 in Diabetic Rats. *J Appl Physiol* (1985) (2007) 103(5):1764–71. doi: 10.1152/jappphysiol.00434.2007
30. Gomes RM, Tófolo LP, Rinaldi W, Scomparin DX, Grassioli S, Barella LF, et al. Moderate Exercise Restores Pancreatic Beta-Cell Function and Autonomic Nervous System Activity in Obese Rats Induced by High-Fat Diet. *Cell Physiol Biochem* (2013) 32(2):310–21. doi: 10.1159/000354439
31. Moien-Afshari F, Ghosh S, Elmi S, Rahman MM, Sallam N, Khazaei M, et al. Exercise Restores Endothelial Function Independently of Weight Loss or Hyperglycaemic Status in Db/Db Mice. *Diabetologia* (2008) 51(7):1327–37. doi: 10.1007/s00125-008-0996-x
32. Reeves PG, Nielsen FH, Fahey GC. AIN-93 Purified Diets for Laboratory Rodents: Final Report of the American Institute of Nutrition Ad Hoc Writing Committee on the Reformulation of the AIN-76A Rodent Diet. *J Nutr* (1993) 123(11):1939–51. doi: 10.1093/jn/123.11.1939
33. Larqué C, Velasco M, Navarro-Tableros V, Duhne M, Aguirre J, Gutiérrez-Reyes G, et al. Early Endocrine and Molecular Changes in Metabolic Syndrome Models. *IUBMB Life* (2011) 63(10):831–9. doi: 10.1002/iub.544
34. Guimarães VHD, Lelis DF, Oliveira LP, Borém LMA, Guimarães FAD, Farias LC, et al. Comparative Study of Dietary Fat: Lard and Sugar as a Better Obesity and Metabolic Syndrome Mice Model. *Arch Physiol Biochem* (2020), 1–11. doi: 10.1080/13813455.2020.1835986
35. Bernier M, Mitchell SJ, Wahl D, Diaz A, Singh A, Seo W, et al. Disulfiram Treatment Normalizes Body Weight in Obese Mice. *Cell Metab* (2020) 32(2):203–14.e4. doi: 10.1016/j.cmet.2020.04.019
36. Haubold KW, Allen DL, Capetanaki Y, Leinwand LA. Loss of Desmin Leads to Impaired Voluntary Wheel Running and Treadmill Exercise Performance. *J Appl Physiol* (1985) (2003) 95(4):1617–22. doi: 10.1152/jappphysiol.00408.2003
37. Lerman I, Harrison BC, Freeman K, Hewett TE, Allen DL, Robbins J, et al. Genetic Variability in Forced and Voluntary Endurance Exercise Performance in Seven Inbred Mouse Strains. *J Appl Physiol* (1985) (2002) 92(6):2245–55. doi: 10.1152/jappphysiol.01045.2001
38. Falone S, D'Alessandro A, Mirabilio A, Cacchio M, Di Ilio C, Di Loreto S, et al. Late-Onset Running Biphasically Improves Redox Balance, Energy- and Methylglyoxal-Related Status, as Well as SIRT1 Expression in Mouse Hippocampus. *PLoS One* (2012) 7(10):e48334. doi: 10.1371/journal.pone.0048334
39. Bonora E, Targher G, Alberiche M, Bonadonna RC, Saggiani F, Zenere MB, et al. Homeostasis Model Assessment Closely Mirrors the Glucose Clamp Technique in the Assessment of Insulin Sensitivity: Studies in Subjects With Various Degrees of Glucose Tolerance and Insulin Sensitivity. *Diabetes Care* (2000) 23(1):57–63. doi: 10.2337/diacare.23.1.57
40. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis Model Assessment: Insulin Resistance and Beta-Cell Function From Fasting Plasma Glucose and Insulin Concentrations in Man. *Diabetologia* (1985) 28(7):412–9. doi: 10.1007/BF00280883
41. Scott AM, Atwater I, Rojas E. A Method for the Simultaneous Measurement of Insulin Release and B Cell Membrane Potential in Single Mouse Islets of Langerhans. *Diabetologia* (1981) 21(5):470–5. doi: 10.1007/BF00257788
42. Ohkawa H, Ohishi N, Yagi K. Assay for Lipid Peroxides in Animal Tissues by Thiobarbituric Acid Reaction. *Anal Biochem* (1979) 95(2):351–8. doi: 10.1016/0003-2697(79)90738-3
43. SRERE PA. CITRYL-COA. AN SUBSTRATE FOR THE CITRATE-CLEAVAGE ENZYME. *Biochim Biophys Acta* (1963) 73:523–5. doi: 10.1016/0926-6569(63)90146-9
44. Catta-Preta M, Mendonca LS, Fraulob-Aquino J, Aguilá MB, Mandarim-de-Lacerda CA. A Critical Analysis of Three Quantitative Methods of Assessment of Hepatic Steatosis in Liver Biopsies. *Virchows Arch* (2011) 459(5):477–85. doi: 10.1007/s00428-011-1147-1
45. Lacy PE, Kostianovsky M. Method for the Isolation of Intact Islets of Langerhans From the Rat Pancreas. *Diabetes* (1967) 16(1):35–9. doi: 10.2337/diab.16.1.35
46. Carpinelli AR, Curi R, Malaisse WJ. Long-Term Regulation of Pancreatic B-Cell Responsiveness to D-Glucose by Food Availability, Feeding Schedule, and Diet Composition. *Physiol Behav* (1992) 52(6):1193–6. doi: 10.1016/0031-9384(92)90481-G
47. Munhoz AC, Riva P, Simões D, Curi R, Carpinelli AR. Control of Insulin Secretion by Production of Reactive Oxygen Species: Study Performed in Pancreatic Islets From Fed and 48-Hour Fasted Wistar Rats. *PLoS One* (2016) 11(6):e0158166. doi: 10.1371/journal.pone.0158166
48. Bindokas VP, Kuznetsov A, Sreeman S, Polonsky KS, Roe MW, Philipson LH. Visualizing Superoxide Production in Normal and Diabetic Rat Islets of Langerhans. *J Biol Chem* (2003) 278(11):9796–801. doi: 10.1074/jbc.M206913200
49. Zhao H, Kalivendi S, Zhang H, Joseph J, Nithipatikom K, Vásquez-Vivar J, et al. Superoxide Reacts With Hydroethidine But Forms a Fluorescent Product That is Distinctly Different From Ethidium: Potential Implications in

- Intracellular Fluorescence Detection of Superoxide. *Free Radic Biol Med* (2003) 34(11):1359–68. doi: 10.1016/S0891-5849(03)00142-4
50. Oscari LB, Miller WC, Arnall DA. Effects of Dietary Sugar and of Dietary Fat on Food Intake and Body Fat Content in Rats. *Growth* (1987) 51(1):64–73.
  51. Choi MS, Kim YJ, Kwon EY, Ryoo JY, Kim SR, Jung UJ. High-Fat Diet Decreases Energy Expenditure and Expression of Genes Controlling Lipid Metabolism, Mitochondrial Function and Skeletal System Development in the Adipose Tissue, Along With Increased Expression of Extracellular Matrix Remodelling- and Inflammation-Related Genes. *Br J Nutr* (2015) 113(6):867–77. doi: 10.1017/S0007114515000100
  52. Shaw CS, Clark J, Wagenmakers AJ. The Effect of Exercise and Nutrition on Intramuscular Fat Metabolism and Insulin Sensitivity. *Annu Rev Nutr* (2010) 30:13–34. doi: 10.1146/annurev.nutr.012809.104817
  53. (EASL) EAftSotL, (EASD) EAftSoD, (EASO) EAftSoO and EASL-EASD-EASO. Clinical Practice Guidelines for the Management of non-Alcoholic Fatty Liver Disease. *J Hepatol* (2016) 64(6):1388–402. doi: 10.1016/j.jhep.2015.11.004
  54. Younossi ZM, Koenig AB, Abdelatif D, Fazel Y, Henry L, Wymer M. Global Epidemiology of Nonalcoholic Fatty Liver Disease—Meta-Analytic Assessment of Prevalence, Incidence, and Outcomes. *Hepatology* (2016) 64(1):73–84. doi: 10.1002/hep.28431
  55. Paschen M, Moede T, Valladolid-Acebes I, Leibiger B, Moruzzi N, Jacob S, et al. Diet-Induced  $\beta$ -Cell Insulin Resistance Results in Reversible Loss of Functional  $\beta$ -Cell Mass. *FASEB J* (2019) 33(1):204–18. doi: 10.1096/fj.201800826R
  56. Leloup C, Tourrel-Cuzin C, Magnan C, Karaca M, Castel J, Carneiro L, et al. Mitochondrial Reactive Oxygen Species are Obligatory Signals for Glucose-Induced Insulin Secretion. *Diabetes* (2009) 58(3):673–81. doi: 10.2337/db07-1056
  57. Pi J, Bai Y, Zhang Q, Wong V, Floering LM, Daniel K, et al. Reactive Oxygen Species as a Signal in Glucose-Stimulated Insulin Secretion. *Diabetes* (2007) 56(7):1783–91. doi: 10.2337/db06-1601
  58. Roma LP, Jonas JC. Nutrient Metabolism, Subcellular Redox State, and Oxidative Stress in Pancreatic Islets and  $\beta$ -Cells. *J Mol Biol* (2020) 432(5):1461–93. doi: 10.1016/j.jmb.2019.10.012
  59. Kolb H, Eizirik DL. Resistance to Type 2 Diabetes Mellitus: A Matter of Hormesis? *Nat Rev Endocrinol* (2011) 8(3):183–92. doi: 10.1038/nrendo.2011.158
  60. Calegari VC, Zoppi CC, Rezende LF, Silveira LR, Carneiro EM, Boschero AC. Endurance Training Activates AMP-Activated Protein Kinase, Increases Expression of Uncoupling Protein 2 and Reduces Insulin Secretion From Rat Pancreatic Islets. *J Endocrinol* (2011) 208(3):257–64. doi: 10.1530/JOE-10-0450
  61. Medina-Contreras JML, Colado-Velázquez J, Gómez-Viquez NL, Mailloux-Salinas P, Pérez-Torres I, Aranda-Fraustro A, et al. Effects of Topical Capsaicin Combined With Moderate Exercise on Insulin Resistance, Body Weight and Oxidative Stress in Hypoestrogenic Obese Rats. *Int J Obes (Lond)* (2017) 41(5):750–8. doi: 10.1038/ijo.2017.33
- Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.
- Copyright © 2022 Veras, Lucena, Goedcke, Evangelista, Carpinelli and Carvalho. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.