

BMJ Open Determinants of short birth intervals among married women: a cross-sectional study in Karachi, Pakistan

Sidrah Nausheen,¹ Maria Bhura,² Kristy Hackett ,³ Imtiaz Hussain,⁴ Zainab Shaikh,⁵ Arjumand Rizvi,⁴ Uzair Ansari,⁴ David Canning,³ Iqbal Shah,⁶ Sajid Soofi ⁴

To cite: Nausheen S, Bhura M, Hackett K, *et al*. Determinants of short birth intervals among married women: a cross-sectional study in Karachi, Pakistan. *BMJ Open* 2021;**11**:e043786. doi:10.1136/bmjopen-2020-043786

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-043786>).

Received 15 August 2020
Revised 15 February 2021
Accepted 25 February 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Obstetrics & Gynecology, Aga Khan University, Karachi, Pakistan

²Center of Excellence in Women & Child Health, Aga Khan University, Karachi, Pakistan

³Harvard T.H. Chan School of Public Health, Harvard University, Boston, Massachusetts, USA

⁴Pediatrics & Child Health, Aga Khan University, Karachi, Pakistan

⁵Center of Excellence & Child Health, Aga Khan University, Karachi, Pakistan

⁶Global Health and Population, Harvard University T H Chan School of Public Health, Boston, Massachusetts, USA

Correspondence to
Professor Sajid Soofi;
sajid.soofi@aku.edu

ABSTRACT

Introduction Birth spacing is a critical pathway to improving reproductive health. WHO recommends a minimum of 33-month interval between two consecutive births to reduce maternal, perinatal, infant morbidity and mortality. Our study evaluated factors associated with short birth intervals (SBIs) of less than 33 months between two consecutive births, in Karachi, Pakistan.

Methods We used data from a cross-sectional study among married women of reproductive age (MWRA) who had at least one live birth in the 6 years preceding the survey (N=2394). Information regarding their sociodemographic characteristics, reproductive history, fertility preferences, family planning history and a 6-year reproductive calendar were collected. To identify factors associated with SBIs, we fitted simple and multiple Cox proportional hazards models and computed HRs with their 95% CIs.

Results The median birth interval was 25 months (IQR: 14–39 months), with 22.9% (833) of births occurring within 33 months of the index birth. Women's increasing age (25–30 years (aHR 0.63 (0.53 to 0.75)), 30+ years (aHR 0.29, 95% CI 0.22 to 0.39) compared with 20–24 years; secondary education (aHR 0.75, 95% CI 0.63 to 0.88), intermediate education (aHR 0.62, 95% CI 0.48 to 0.80), higher education (aHR 0.69, 95% CI 0.51 to 0.92) compared with no education, and a male child of the index birth (aHR 0.81, 95% CI 0.70 to 0.94) reduced the likelihood of SBIs. Women's younger age <20 years (aHR 1.24, 95% CI 1.05 to 1.24) compared with 20–24 years, and those who did not use contraception within 9 months of the index birth had a higher likelihood for SBIs for succeeding birth compared with those who used contraception (aHR 2.23, 95% CI 1.93 to 2.58).

Conclusion Study shows that birth intervals in the study population are lower than the national average. To optimise birth intervals, programmes should target child spacing strategies and counsel MWRA on the benefits of optimal birth spacing, family planning services and contraceptive utilisation.

BACKGROUND

Birth spacing is integral to improving reproductive health. WHO recommends a minimum 24-month birth-to-pregnancy interval, or a 33-month interval between

Strengths and limitations of this study

- This is the first study that has investigated birth spacing in urban areas of Karachi, Pakistan.
- It is a cross-sectional study that has employed a three-stage random sampling design that is, at cluster level, at household level and at individual level.
- There may be an under-representation of birth intervals because the study did not consider abortions or miscarriages.
- The study only considered births in 6-year calendar time and therefore births occurred before or after this calendar time were considered as no-event.

two consecutive births to reduce the risk of adverse maternal, perinatal and infant health outcomes.¹ Birth spacing is highly influenced by socioeconomic, demographic, cultural and behavioural characteristics.² Short birth-to-birth intervals, also known as, short birth intervals (SBIs) are associated with poor neonatal and infant outcomes,³ including low birth weight,⁴ preterm births,⁵ small-for-gestational age,⁶ neonatal mortality^{7 8} and infant mortality.^{4 9 10} Short birth-to-pregnancy intervals are also associated with a 61% increased risk in neonatal mortality and a 48% increased risk in under-5 mortality if the interval is less than 24 months.¹¹

Similarly, maternal health is negatively impacted by SBIs, where women do not have sufficient time to physically recuperate from their previous pregnancy.¹² Closely spaced pregnancies increase maternal nutrition depletion, resulting in a reduction of the mother's nutritional status.¹³ Birth-to-pregnancy intervals of less than 6 months can significantly increase the odds of maternal mortality by 150% (95% CI 22% to 438%), and are associated with an increased risk of third trimester bleeding, premature rupture of membranes, postpartum endometriosis and anaemia.¹⁴ A systematic review



of studies from Ethiopia found that women with birth-to-pregnancy intervals of less than 2 years were twice at risk of developing anaemia during their next pregnancy since repeated pregnancies tend to deplete a woman's iron stores.¹⁵ However, systematic reviews have reported conflicting and low-quality evidence between maternal health outcomes and SBIs.^{16 17}

Longitudinal data on singleton live births in Bangladesh found that shorter intervals between birth and pregnancy were associated with higher infant and child mortality, and longer birth intervals improved child survival.^{18 19} Several studies have found associations between SBIs and neonatal and infant mortality in both low-income and middle-income countries (LMICs) and high-income countries over time.^{20–25} Systematic reviews and Demographic Health Survey (DHS) analyses have also studied the impact of SBIs on infant mortality, particularly in low-income countries.^{26 27} SBIs are associated with infant morbidity and poor health outcomes in multiple ways, for both the older child as well as the one born after the SBI. Women with closely spaced pregnancy may less likely to attend antenatal care services (which are critical for monitoring pregnancy and identifying complications) because they have other child to take care of.²⁸ Furthermore, lactation may be impaired due to maternal nutritional depletion and they may be unable to provide adequate breastfeeding to their older infant.²⁸ Children who are closely spaced are more likely to compete for resources, such as breastmilk, parental attention and time.^{24 29}

Pakistan has a population of over 216.6 million people in 2019 and is currently the fifth most populous country in the world, with an annual population growth rate of 2.1% and a fertility rate of 3.6 children per woman in 2017.^{30 31} The country possesses a maternal mortality ratio of 276 deaths per 100 000 live births, neonatal mortality of 42 deaths per 1000 live births, and infant mortality at 62 deaths per 1000 live births.^{32–34} The median age at first birth is 22.8 years among married women of reproductive age (MWRA). Moreover, the use of any method of family planning by currently married women is 34%, with 25% using a modern method and 9% using a traditional method of contraception.³⁵ Although Pakistan's median birth interval is 28.2 months according to Pakistan Demographic and Health Survey (PDHS) 2017–2018, 37% of the births occur within 24 months of the preceding birth.³⁵ This statistic is higher among younger women, where women aged 15–19 years have birth intervals which are 12.4 months shorter, on average, than women aged 30–39 years.³⁵

An earlier study across 21 LMICs revealed that Pakistan has one of the highest percentages (60%) of short birth-to-pregnancy intervals (<23 months after birth) with 31% unmet need for spacing and 29% unmet need for limiting.³⁶ The unmet need for spacing and limiting pregnancies in Pakistan is 17%, indicating that several women who want to space or limit pregnancies do not use any method to do so.³⁵ Therefore, opting for family planning and contraceptive use after childbirth can help women

achieve healthy spacing of pregnancies.³⁷ In Pakistan, preference for a male child is deeply entrenched, therefore, couple's wait before moving to subsequent pregnancy is short as long as desired number of son(s) are not born.^{38 39} A recent study from Pakistan has reported that birth intervals of less than 24 or 18 are higher among women without one or more sons.³⁹ Other predictors that contribute to birth intervals in other studies include wealth indices, women's education, maternal age, later start of reproductive years, gender of an index child and parity according to studies conducted in Bangladesh, Iran and Ethiopia.^{40–42} However, there is a lack of data on birth intervals in Pakistan. Our study seeks to explore the socio-economic, demographic and reproductive factors associated with SBIs of less than 33 months using retrospective data from urban populations in Karachi, Pakistan.

METHODS

Study design

This study draws on data from an evaluation of the Willows programme (<https://projects.iq.harvard.edu/willowsimpacteval>), a community-based reproductive health programme that provides family planning information, education and referral through household visits to WRA. The parent study assessed the effect of the Willows programme on modern contraceptive use with an aim to guide future programming for family planning interventions in Pakistan. The current study was a cross-sectional assessment of retrospective data that evaluated the Willows Programme, and took place between August and December 2018.

Study setting and participants

This cross-sectional study was conducted in Korangi Town, PIB Colony, and Dalmia/Shanti Nagar, three periurban municipalities in Karachi, Pakistan. All areas are home to both locals and migrants from within the country, as well as Afghanistan, Bangladesh and Burma (now Myanmar), and have a majority of Muslim population. Women were eligible to participate in the study if they were married, usual household members, spoke at least one of the four commonly spoken languages (Urdu, Pushto, English or Sindhi), were between the ages of 15 and 49 years, and self-reported themselves as fertile.

Sample size and sampling strategy

For a parent study, a sample size of 1836 (~2000) from each area intervention and control area was required assuming an estimated modern contraceptive prevalence rate of below 30% in selected areas, methodology has been described in detail elsewhere.⁴³ A three-stage random sampling design was carried out in STATA using a uniform (0, 1) random number generator with a fixed seed. First, we used geographical information systems technology to construct a sampling frame with distinct area and cluster demarcation of the survey sites, forming 708 clusters in total. Next with a goal of an average of 60

households per cluster, we randomly selected 220 clusters, with 110 clusters from Korangi Town, and 110 clusters from PIB Colony and Dalmia/Shanti Nagar combined. Since PIB colony and Dalmia/Shanti Nagar are smaller in geographical and population size compared with Korangi town, therefore, equal numbers of clusters were selected from Korangi Town and PIB colony and Dalmia/Shanti Nagar combined. Proceeding that, an android application for household listing questionnaire was developed to assess the number of women between 15 and 49 years of age. If more than one WRA lived in a selected household, we randomly selected one from the household.

Data collection

We conducted face to face interviews with eligible women using a structured tablet-based questionnaire on the CommCare application for this survey. The survey questionnaire included a range of topics on women's reproductive health, including information on sociodemographic characteristics of women and their husband, reproductive history, obstetric history, family planning history, fertility preferences and a reproductive calendar of pregnancies, births, terminations and contraceptive use for the preceding 6 years. This study used a month-by-month calendar, similar to those collected in DHS and was based on a 5-year recall period.

Data analysis

Measures and outcomes

Information on birth intervals was analysed using the contraceptive calendar for all participating women. Of all (4336) the randomly selected women, 4193 consented for participation in this retrospective survey. Of these, 2394 women who had given live birth at least once in the 6 years preceding the survey by using the calendar data were included in the analysis, and a total of 1799 MWRA were excluded because they did not give birth to any child in the 6 years preceding the survey, or their pregnancies resulted in abortions or miscarriages. Index births were defined as the birth preceding the birth interval. We assessed the association between birth spacing and socio-demographic characteristics, including woman's age at index birth, woman's education, husband's education, wealth quintiles, ethnicity, sex of the infant, contraceptive uptake within 9 months of the index birth and length of the first birth interval in months. Wealth quintiles were measured in five categories that is, poorest, poorer, middle, richer and richest; however, for this study, we recategorised them into three and created poorest/poorer, middle and richer/richest. In examining the determinants of SBIs, we defined an event as the interval between the index birth and the next birth (live or still-born) of less than 33 months, corresponding to recommendations by the WHO.¹ Women who gave birth after 33 months or those who did not give birth after the index child were considered no-event by the survey as information only until the time of the interview was recorded.

Statistical analysis

A discrete time survival analysis technique was carried out using logistic functional form.⁴⁴ This model specification allows flexible baseline hazard, so there is no need to assume a functional form of the effect of duration. For this analysis, duration between consecutive births divided into two categories using 33 months as a cut-off for SBI. This model specification facilitates the introduction of time-varying covariates in the model and censoring in the data.

We initially performed bivariate analyses to examine the association between explanatory variables and the outcome variable (model A). A multivariable model was adjusted for all covariates (model B). Another multivariable Cox proportional hazard model was fitted by including variables with $p < 0.2$ in the bivariate model (model C) using a backward elimination method, and variables with $p < 0.05$ were retained within the model. HRs and their 95% CIs were computed with statistical significance determined at the 5% level ($p < 0.05$). All analyses account clustering for the sampling design and women level using clustered robust SEs. The model was checked for multicollinearity using variance inflation factor using cut-offs of ≥ 10 . All analyses were performed in STATA V.15.

Patient and public involvement

The public was not involved in the design of the research tools, but they were part of the study. The key findings will be shared with their representatives as part of the dissemination plan at local level.

Informed written consent was obtained from each study participant. Women who were unable to sign provided consent with a thumb impression in the presence of witnesses.

RESULTS

Descriptive results

A total of 4336 MWRA were approached; of those 4193 women consented for participation in this retrospective survey. A total of 1799 MWRA were excluded because of they had no index birth history. Therefore, 2394 women were included in our analysis who had given birth to a total of 3641 children in the 6 years preceding the survey. Of the total births, 833 (22.9%) occurred in less than 33 months of the index birth; and the median birth interval in our study was 25 months (IQR: 14–39 months). Descriptive results for participants are presented in [table 1](#) with median and IQRs for birth intervals in months for each category.

One in three women in our study had achieved secondary education (36.6%), with higher than secondary education being the least common (9.8%) and about one-quarter (28.7%) women had no formal education. Similarly, one in three husbands had achieved secondary education (38.3%) and quarter of them had no education. Majority of our sample (84.3%) were

Table 1 Per cent distribution of sociodemographic characteristics of participants with mean and median birth interval (N=2394)

	n (%) (N=2394)	Birth interval in months	
		Mean	Median (IQR)
Women's education			
None	686 (28.7)	26.8	25 (14–37)
Primary	319 (13.3)	28.1	25 (14–40)
Secondary	877 (36.6)	27.8	25 (14–40)
Intermediate	277 (11.6)	30.0	28 (16–42)
Higher	235 (9.8)	28.1	27 (14–39)
Husband education			
None	642 (26.8)	27.4	25 (14–38)
Primary	220 (9.2)	26.9	25 (14–37)
Secondary	916 (38.3)	27.9	25 (14–40)
Intermediate	317 (13.2)	28.2	26 (14–39)
Higher	299 (12.5)	28.7	26 (15–42)
Age at first marriage (n=2393)			
10–19	1003 (41.9)	28.0	25 (15–40)
20–24	1016 (42.4)	27.4	25 (14–39)
25–29	321 (13.4)	28.5	25 (14–40)
30–45	53 (2.2)	26.9	23 (15–37.5)
Age at first birth			
10–19	657 (27.4)	28.0	25 (15–39)
20–24	1126 (47.0)	27.7	25 (14–39)
25–29	495 (20.7)	27.6	25 (14–39)
30–45	116 (4.8)	28.7	23.5 (14–41)
Wealth quintiles			
Poorest and poorer	641 (26.8)	27.4	25 (14–38)
Middle	889 (37.1)	27.5	25 (14–39)
Richest and richer	864 (36.1)	28.4	26 (15–40)
Ethnicity			
Urdu	1152 (48.1)	28.6	26 (15–40)
Sindhi	197 (8.2)	26.3	22 (14–36)
Punjabi	314 (13.1)	28.4	26 (14–40)
Other	731 (30.5)	26.8	25 (14–37)

married between 10 and 24 years of age, and 47.0% had their first birth between 20 and 24 years of age. About half the respondents (48.1%) belonged to an Urdu-speaking caste. From all index births included in our study, 32.5% were born when their mothers were 20–24 years of age, and 39.9% between 25– and 29 years of age. Majority of women belonged to middle wealth quintile (37.1%), while a similar number belonged to combined two richest

quintiles (36.1%), and a quarter belonged to combined two poorest two quintiles (26.8%) (table 2).

When asked about contraceptive use within 9 months of the index birth, about a quarter (28.3%) of participants did not use contraception (table 2). Among those who used contraception, more than half (68.5%) used modern contraceptive methods, a quarter (25.6%) used traditional methods, and 5.9% used both modern and traditional methods. Women who did not use contraceptive methods had a shorter birth interval (median: 22 months, IQR: 14–35 months) than those who used modern contraceptive methods (median: 26 months, IQR: 14–41 months) or traditional contraceptive methods (median: 28 months, IQR: 16–41 months) (table 2). Birth intervals varied slightly depending on the sex of the index birth. Data reveals that length of succeeding birth interval is shorter when the sex of index child is female, and this puts woman in even greater pressure to try for a male child earlier (table 2).

Predictors of SBI

Bivariate analyses of predictors of SBIs (<33 months) are displayed in table 3. They indicate that women aged 25–30 years and women who were greater than 30 years of age were less likely to have an SBIs compared with those younger than aged 20–24 years. However, the likelihood of SBI was higher among women less than 20 years old compared with women 20–24 years of age. Mothers who received secondary, intermediate, and higher education were also less likely to have SBIs than those who received no formal education. Likewise, husbands who received intermediate and higher were also less likely to have an SBI for the subsequent birth. Couples who did not use contraceptives within 9 months were more likely to have SBIs. SBIs were also associated with the gender of the child born prior to the index birth; wealth quintiles, where those belonging to the middle and richest wealth quintiles were less likely to have SBIs and ethnicity, with those belonging to a Sindhi or other background more likely to have SBIs compared with Urdu-speaking families (table 3).

Two multivariate models were generated, with model B adjusted for all variables and model C adjusted for significant explanatory variables (table 3). When adjusted for all explanatory variables, women who were less than 20 years of age were more likely and those between the ages of 25–29 years and 30+ years were less likely to have SBIs compared with women 20–24 years of age. Similarly, women with secondary, and intermediate education also had fewer SBIs compared with those with no education. Couples who did not use contraception within 9 months of the index birth, and women who were between 25 and 29 years at first birth were more likely to have SBIs and those with more male children were less likely to have shorter birth intervals.

Mother's age, mother's education, contraceptive use within 9 months of index birth, age at first marriage, and gender of child born prior to index birth were fitted into

Table 2 Percent distribution of births with mean and median birth intervals (N=3641)

	n (%) N=3641	Birth interval in months	
		Mean	Median (IQR)
Total		27.8	25 (14–39)
Age of woman at index birth			
<20	244 (6.7)	27.4	25 (15.5–36.5)
20–24	1182 (32.5)	27.2	25 (15–37)
25–30	1453 (39.9)	27.1	25 (14–39)
>30	762 (20.9)	29.9	28 (15–42)
Contraceptive use within 9 months after index birth			
Used	2612 (71.7)	28.8	27 (15–41)
Did not use	1029 (28.3)	25.5	22 (14–35)
Contraceptive methods used within 9 months after index birth			
Modern	1791 (68.5)	28.4	26 (14–41)
Traditional	668 (25.6)	29.8	28 (16–41)
Both	153 (5.9)	29.1	27 (16–40)
Gender of index child*			
Male	1732 (51.9)	29.0	27 (15–41)
Female	1603 (48.1)	27.2	25 (14–38)

*Denominator was 3335 for this variable as some of the children were the index birth.

a discrete time proportional hazards model (model C) and were found to be significantly associated with SBIs. Similar to model A, women younger than 20 years of age had a higher likelihood for SBIs (aHR 1.36, 95% CI 1.07 to 1.73), and women between the ages of 25 and 30 years (aHR 0.63, 95% CI 0.53 to 0.75) and 30+ years (aHR 0.29, 95% CI 0.22 to 0.39) were less likely to have SBIs compared with women 20–24 years of age. Women who had attained secondary (aHR 0.75, 95% CI 0.63 to 0.88), intermediate (aHR 0.62, 95% CI 0.48 to 0.80) and higher education (aHR 0.69, 95% CI 0.51 to 0.92) had fewer SBIs compared with those with no education, and having a male index child resulted in SBIs (aHR 0.81, 95% CI 0.70 to 0.94). Moreover, women's age at index birth also influenced birth intervals; compared with those who were younger than 20 years of age at marriage, those who were 20–24 years at marriage (aHR 1.24, 95% CI 1.05 to 1.24), 25–29 years of age (aHR 1.54, 95% CI 1.18 to 2.02) and those 30–45 years of age (aHR 2.64, 95% CI 1.55 to 4.47) had a higher likelihood of SBIs. Couples who did not use contraception within 9 months of the index birth also had a higher likelihood for SBIs compared with those who used contraception (aHR 2.33, 95% CI 1.93 to 2.58). Kaplan-Meier survival curves depict the probability of SBIs by the various subgroups (figure 1).

DISCUSSION

SBIs are associated with adverse neonatal outcomes and neonatal mortality; and contribute to the burden of disease among neonates in LMICs.⁴⁵ This study evaluated

the predictors of SBIs (<33 months) in urban areas of Karachi, Pakistan, and found that 22.9% of births that occurred within 6 years of the study had a following birth less than 33 months of the index birth. The average median birth interval in our study was 25 months, which is lower than the national median of 29.8 months in urban areas, and lower than the recommended duration.³⁵ This interval is much shorter than study conducted in neighbouring Iran where the median duration between two live births was 39 months,⁴⁰ but higher than a study in rural Uganda where the median birth interval was 22 months.⁴⁶ Another large-scale cross-sectional study in rural Bangladesh found that 24.6% had SBIs of less than 33 months, which is very similar to our results, although our study was conducted in an urban setting.⁴² The median birth interval duration reported in our study is also relatively less than the ones obtained in similar studies conducted in Ethiopia,⁴⁷ Myanmar⁴⁸ and India⁴⁹ with values ranging between 30 and 32.6 months.

In our study, maternal age, education, contraceptive use within 9 months of the index birth and gender of the child prior to the index birth were the strongest predictors of SBIs. Maternal age was a major determinant of all birth intervals in a similar study in Pakistan on the determinants of higher-order birth intervals, where increasing maternal age increased birth intervals.⁵⁰ These findings are also consistent with those reported from Bangladesh, where mother's age at first birth, parity, survival status of the index child, mother's education, place of residence and family composition, that is, having a male child was

Table 3 The discrete time proportional hazard model for predictors of SBI (birth interval <33 months)

	Model A—bivariate		Model B—multivariate (all variables)		Model C—multivariate (reduced)	
	HR (95% CI)	P value	Adjusted HR (95% CI)	P value	Adjusted HR (95% CI)	P value
Woman age at index birth						
<20	1.32 (1.07 to 1.62)	0.009	1.40 (1.09 to 1.80)	0.008	1.36 (1.07 to 1.73)	0.012
20–24	1	.	1	.	1	.
25–30	0.66 (0.57 to 0.77)	<0.0001	0.63 (0.50 to 0.72)	<0.0001	0.63 (0.53 to 0.75)	<0.0001
>30	0.37 (0.30 to 0.47)	<0.0001	0.28 (0.21 to 0.38)	<0.0001	0.29 (0.22 to 0.39)	<0.0001
Study arm						
Control	1		1			
Intervention	0.98 (0.86 to 1.12)	0.757	1.03 (0.88 to 1.20)	0.707		
Mother education						
None	1	.	1	.	1	.
Primary	0.83 (0.67 to 1.02)	0.071	0.82 (0.66 to 1.03)	0.085	0.81 (0.65 to 1.01)	0.057
Secondary	0.79 (0.68 to 0.92)	0.003	0.76 (0.62 to 0.92)	0.005	0.75 (0.63 to 0.88)	0.001
Intermediate	0.62 (0.49 to 0.78)	<0.0001	0.63 (0.47 to 0.85)	0.002	0.62 (0.48 to 0.80)	<0.001
Higher	0.68 (0.52 to 0.89)	0.004	0.70 (0.49 to 0.99)	0.043	0.69 (0.51 to 0.92)	0.011
Husband education						
None	1	.	1	.		
Primary	0.95 (0.74 to 1.21)	0.647	0.95 (0.74 to 1.23)	0.702		
Secondary	0.88 (0.76 to 1.03)	0.114	0.98 (0.82 to 1.17)	0.817		
Intermediate	0.78 (0.63 to 0.98)	0.033	0.95 (0.72 to 1.25)	0.713		
Above	0.68 (0.54 to 0.86)	0.001	0.94 (0.71 to 1.24)	0.653		
Contraceptive use within 9 months after index birth						
Use	1	.	1	.	1	.
Did not use	2.45 (2.14 to 2.81)	<0.0001	2.20 (1.90 to 2.56)	<0.0001	2.23 (1.93 to 2.58)	<0.0001
Age at first marriage						
10–19	1	.	1	.	1	.
20–24	0.96 (0.83 to 1.10)	0.535	1.08 (0.87 to 1.36)	0.481	1.24 (1.05 to 1.47)	0.011
25–29	0.85 (0.69 to 1.04)	0.117	1.06 (0.71 to 1.59)	0.773	1.54 (1.18 to 2.02)	0.002
30–45	0.97 (0.61 to 1.55)	0.889	1.82 (0.79 to 4.19)	0.158	2.64 (1.59 to 4.47)	<0.0001
Age at first birth						
10–19	1	.	1	.		
20–24	0.96 (0.82 to 1.11)	0.563	1.16 (0.92 to 1.46)	0.213		
25–29	0.95 (0.79 to 1.14)	0.585	1.61 (1.12 to 2.32)	0.010		
30–45	0.82 (0.58 to 1.18)	0.287	1.62 (0.79 to 3.34)	0.195		
First marriage and first birth interval (months)	1.00 (1.00 to 1.00)	0.448	1.00 (0.99 to 1.00)	0.51		
Gender of index child						
Female	1	.	1	.	1	.
Male	0.85 (0.73 to 0.98)	0.023	0.81 (0.71 to 0.94)	0.005	0.81 (0.70 to 0.94)	0.004
Wealth quintiles						
Two poorest	1	.	1	.		
Middle	0.80 (0.68 to 0.93)	0.005	0.90 (0.76 to 1.07)	0.229		
Two richest	0.75 (0.64 to 0.88)	<0.001	0.94 (0.77 to 1.14)	0.552		
Ethnicity						

Continued

Table 3 Continued

	Model A—bivariate		Model B—multivariate (all variables)		Model C—multivariate (reduced)	
	HR (95% CI)	P value	Adjusted HR (95% CI)	P value	Adjusted HR (95% CI)	P value
Urdu	1	.	1	.		
Sindhi	1.36 (1.08 to 1.71)	0.01	0.99 (0.79 to 1.31)	0.943		
Punjabi	1.07 (0.88 to 1.32)	0.491	1.08 (0.87 to 1.34)	0.468		
Other	1.25 (1.07 to 1.44)	0.004	0.98 (0.82 to 1.19)	0.872		

A—bivariate analysis.

B—model includes all predictors regardless of their significance in bivariate analysis.

C—the predictors significant at $p < 0.2$ in bivariate analysis considered for adjustment. Parsimonious model selected using backward elimination, $p < 0.05$ considered significant.

SBI, short birth interval.

significantly associated with length of birth intervals.⁵¹ Similarly, in Iran, the current age of women and maternal age at the time of delivery were strongly associated with birth interval duration.^{42 52} Our results correspond with a study in Uganda where SBIs were associated with younger maternal age.^{42 46} The PDHS 2017–2018 also found that younger women had SBIs compared with older women.³⁵

This could possibly be due to the increasing maternal age not only raises concerns for infertility; but also motivate woman to quickly have her desired number of children. In addition, women have more autonomy in making reproductive decisions when they are older.⁴⁶ Moreover, older women are also more likely to have achieved their desired family size, and therefore, have longer birth intervals.⁴⁷




Figure 1 Kaplan-Meier survival estimates.

As expected, women who did not use any contraceptive method 9 months prior to the index birth were also more likely to have SBIs compared with those who used any form of contraception. The results are consistent with the findings from a literature review of 14 studies conducted in developed and developing countries which found the use of contraceptive is protective against SBIs.⁵³ Though, many of the published evidence in this domain from Pakistan is 20 years old^{54–56} studies from Bangladesh and India supported the evidence.^{49–51} Similar findings have been reported from Africa, where lack of contraceptive use was found to be one of the strongest predictors of SBIs in Ethiopia.⁵⁷ We found that women with higher education were less likely to have SBIs, which is in concordance with studies from Bangladesh and Saudi Arabia.^{42–51–58} A study in India found that education and women's autonomy were both strongly associated with longer birth intervals.⁵⁹ An analysis between education and fertility in Indonesia proposed that women who are more educated have a higher likelihood of participating in family planning programmes, using services and using long-acting modern contraceptives since they have more knowledge of birth control methods and utilise them accurately.⁶⁰ Moreover, educated women are likely to marry later and thus limit their reproductive years and number of children.

Another finding of our study was that women who had a male index child had a reduced likelihood of SBIs than those who gave birth to a female child. Parental attitudes and preference for male children in Middle Eastern and South Asian cultures may be the reason for this finding, since male children are typically regarded as economic assets as well as future bread earners for the family.^{61–63} Societal pressures for a woman to demonstrate her fertility and for her to bear a son may be influencing her ability to make decision around the spacing of children and use of contraceptives.^{61–64} A recent study was conducted analyses using three DHS from Pakistan on preferences for male children and its impact on birth intervals. They found a significant impact of son preference on birth intervals during the first two parities, where women who had daughters had significantly shorter subsequent birth intervals compared with those who had more sons.³⁹ Moreover, women with one or more sons were more likely to use contraceptive methods, indicating a strong preference for sons compared with daughters.³⁹ In order to tackle this pervasive desire for male children, gender equality measures, importance of girls and awareness is crucial. This has major policy implications for the family planning programmes which should be questioned for investing more money into motivational campaigns and should have more integrative policies to promote education for girl child, implementation of legislation against discrimination on the grounds of sex, abolition of practices such as dowry and bride prices, and promoting social welfare and social security so a son is no longer considered an asset and security for an old age.

This is one of the first studies to investigate birth spacing in urban areas of Karachi, Pakistan. The study, however, is not without limitations. First, our study was conducted in selected low-income areas of Karachi, Pakistan, and is therefore not representative of the national or the local population. Second, our analyses do not include pregnancies that resulted in abortions or miscarriages, and therefore, when live or stillbirths are preceded by a non-live pregnancy, there could have been an underestimation of the proportion of closely spaced pregnancies. Third, the determinants identified are only for births that occurred within our study period, and it is possible that other variables could have played a role in predicting birth intervals in the participants in our study. Finally, due to the 6-year time frame, children born to women in our study before or after the time period were not included and were therefore counted as no-event. The last-born infant of each woman in the study timeframe was also included as no-event since there was no data for live births after that infant, and this may have introduced an under-representation of the number of SBIs in our study.

CONCLUSION

Optimal birth spacing has the potential to improve maternal, neonatal and infant health outcomes, reduce familial financial burdens and allow parents to provide children with comprehensive care and attention. Our findings suggest that reproductive health interventions should address underlying socioeconomic factors that contribute to SBIs, such as preferences for male child, education and younger MRWA. Family planning should be integrated with other multisectoral programmes such as education, where girls from the early stage should be empowered and given awareness on these issues. Moreover, family planning strategies should not only focus on increasing coverage of services but also to create awareness about optimal birth intervals and interventions to enhance modern contraceptive utilisation behaviours among WRA.

Twitter Imtiaz Hussain @imtiazhilbi and Sajid Soofi @sajidsoofi

Acknowledgements We would like to thank all study participants, including surveyors, data enumerators and team leads.

Contributors SS was the principal investigator for Pakistan. SS, IS and DC conceptualised the idea and designed the study. SN, MB and ZS drafted the manuscript. AR and UA managed and analysed the data. IH and KH were involved in the implementation of project and contributed to the development of monthly, quarterly and final report and manuscript. SS, DC, KH and IS interpreted the data and critically reviewed the manuscript. All authors contributed in manuscript review and approved the final manuscript.

Funding Funding for this study received by a grant from an anonymous donor via Harvard T.H. Chan School of Public Health, USA

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval The study received ethical approval from the Ethical Review Committee (ERC) at the Aga Khan University (AKU) (4964-Ped-ERC-17) and the

Institutional Review Board (IRB) at the Harvard T.H. Chan School of Public Health (IRB17-1864).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. Data are available on reasonable request. The data of the study used for analysis is available on ZENODO <https://doi.org/10.5281/zenodo.4661082>.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Kristy Hackett <http://orcid.org/0000-0002-3969-659X>

Sajid Soofi <http://orcid.org/0000-0003-4192-8406>

REFERENCES

- World Health Organization. *Report of a technical consultation on birth spacing*. Geneva: World Health Organization, 2005.
- Ford K, Huffman SL, Chowdhury AKMA, et al. Birth-interval dynamics in rural Bangladesh and maternal weight. *Demography* 1989;26:425–37.
- Conde-Agudelo A, Rosas-Bermudez A, Castaño F, et al. Effects of birth spacing on maternal, perinatal, infant, and child health: a systematic review of causal mechanisms. *Stud Fam Plann* 2012;43:93–114.
- Zhu B-P, Rolfs RT, Nangle BE, et al. Effect of the interval between pregnancies on perinatal outcomes. *N Engl J Med Overseas Ed* 1999;340:589–94.
- DeFranco EA, Stamilio DM, Boslaugh SE, et al. A short interpregnancy interval is a risk factor for preterm birth and its recurrence. *Am J Obstet Gynecol* 2007;197:264.e1–264.e6.
- Shults RA, Arndt V, Olshan AF, et al. Effects of short interpregnancy intervals on Small-for-Gestational age and preterm births. *Epidemiology* 1999;10:250–4.
- Smith GCS, Pell JP, Dobbie R. Interpregnancy interval and risk of preterm birth and neonatal death: retrospective cohort study. *BMJ* 2003;327:313.
- Klerman LV, Cliver SP, Goldenberg RL. The impact of short interpregnancy intervals on pregnancy outcomes in a low-income population. *Am J Public Health* 1998;88:1182–5.
- Appareddy S, Pryor J, Bailey B. Inter-pregnancy interval and adverse outcomes: evidence for an additional risk in health disparate populations. *J Matern Fetal Neonatal Med* 2017;30:2640–4.
- Marston C. *Report of a WHO technical consultation on birth spacing Geneva Switzerland 13-15 June 2005*. Geneva, Switzerland: WHO, 2006.
- Kozuki N, Walker N. Exploring the association between short/long preceding birth intervals and child mortality: using reference birth interval children of the same mother as comparison. *BMC Public Health* 2013;13 Suppl 3:S6.
- Winkvist A, Rasmussen KM, Habicht JP. A new definition of maternal depletion syndrome. *Am J Public Health* 1992;82:691–4.
- King JC. The risk of maternal nutritional depletion and poor outcomes increases in early or closely spaced pregnancies. *J Nutr* 2003;133:1732S–6.
- Conde-Agudelo A, Belizán JM. Maternal morbidity and mortality associated with interpregnancy interval: cross sectional study. *BMJ* 2000;321:1255–9.
- Kassa GM, Muche AA, Berhe AK, et al. Prevalence and determinants of anemia among pregnant women in Ethiopia; a systematic review and meta-analysis. *BMC Hematol* 2017;17:17.
- Wendt A, Gibbs CM, Peters S, et al. Impact of increasing inter-pregnancy interval on maternal and infant health. *Paediatr Perinat Epidemiol* 2012;26:239–58.
- Dewey KG, Cohen RJ. *Does birth spacing affect maternal or child nutritional status? A systematic literature review*. UK: Oxford, 2007: Vol 3. 151–73.
- Davanzo J, Hale L, Razzaque A, et al. The effects of pregnancy spacing on infant and child mortality in Matlab, Bangladesh: how they vary by the type of pregnancy outcome that began the interval. *Popul Stud* 2008;62:131–54.
- Greenspan A. Family planning's benefits include improved child health and nutrition: new data from Bangladesh. *Asia Pac Pop Policy* 1993;26:1–4.
- Cleland JG, Sathar ZA. The effect of birth spacing on childhood mortality in Pakistan. *Popul Stud* 1984;38:401–18.
- Curtis SL, Diamond I, McDonald JW. Birth interval and family effects on postneonatal mortality in Brazil. *Demography* 1993;30:33–43.
- Millman SR, Cooksey EC. Birth weight and the effects of birth spacing and breastfeeding on infant mortality. *Stud Fam Plann* 1987;18:202.
- Becher H, Müller O, Jahn A. Risk factors of infant and child mortality in rural Burkina Faso. *Bulletin of the World Health Organization* 2004;82:265.
- Molitoris J. Heterogeneous effects of birth spacing on neonatal mortality risks in Bangladesh. *Stud Fam Plann* 2018;49:3–21.
- Gonçalves SD, Moultrie TA. Short preceding birth intervals and child mortality in Mozambique. *Afr J Reprod Health* 2012;16:29–42.
- Dadi AF, Abel Fekadu D. A systematic review and meta-analysis of the effect of short birth interval on infant mortality in Ethiopia. *PLoS One* 2015;10:e0126759.
- Rutstein SO. Effects of preceding birth intervals on neonatal, infant and under-five years mortality and nutritional status in developing countries: evidence from the demographic and health surveys. *Int J Gynaecol Obstet* 2005;89 Suppl 1:S7–24.
- Boerma JT, Bicego GT. Preceding birth intervals and child survival: searching for pathways of influence. *Stud Fam Plann* 1992;23:243–56.
- Ramachandran P. Maternal Nutrition—Effect on fetal growth and outcome of pregnancy. *Nutr Rev* 2002;60:S26–34.
- UNDP Pakistan. *Population growth: implications for human development*. Islamabad, Pakistan: UNDP Pakistan, 2019.
- Population Reference Bureau. International data: Pakistan, 2019. Available: <https://www.prb.org/international/geography/pakistan>
- UN-IGME. Neonatal mortality rate: Pakistan, 2019. Available: <https://data.worldbank.org/indicator/SH.DYN.NMRT>
- UNICEF, WHO, Bank W, Division UNP. *Levels and trends in child mortality: report 2019. estimates developed by the un Interagency group for child mortality estimation*. New York, USA: UNICEF, WHO, Bank W, Division UNP, 2019.
- UNICEF. Maternal and newborn health disparities in Pakistan: key facts. in, 2017. Available: https://data.unicef.org/wp-content/uploads/country_profiles/Pakistan/country%20profile_PAK.pdf
- NIPS, ICF. *Pakistan demographic and health survey 2017-18*. Islamabad, Pakistan, and Rockville, Maryland, USA: NIPS, ICF, 2019.
- Moore Z, Pfitzer A, Gubin R, et al. Missed opportunities for family planning: an analysis of pregnancy risk and contraceptive method use among postpartum women in 21 low- and middle-income countries. *Contraception* 2015;92:31–9.
- Rajan S, Speizer IS, Calhoun LM, et al. Counseling during maternal and infant health visits and postpartum contraceptive use in Uttar Pradesh, India. *Int Perspect Sex Reprod Health* 2016;42:167.
- Qadir F, Khan MM, Medhin G, et al. Male gender preference, female gender disadvantage as risk factors for psychological morbidity in Pakistani women of childbearing age - a life course perspective. *BMC Public Health* 2011;11:745.
- Javed R, Mughal M. Preference for boys and length of birth intervals in Pakistan. *Research in Economics* 2020;74:140–52.
- Fallahzadeh H, Farajpour Z, Emam Z. Duration and determinants of birth interval in Yazd, Iran: a population study. *Iran J Reprod Med* 2013;11:379–84.
- Hailu D, Gulte T. Determinants of short Interbirth interval among reproductive age mothers in Arba Minch district, Ethiopia. *Int J Reprod Med* 2016;2016:1–17.
- de Jonge HCC, Azad K, Seward N, et al. Determinants and consequences of short birth interval in rural Bangladesh: a cross-sectional study. *BMC Pregnancy Childbirth* 2014;14:427.
- Özçelik EA, Rohr J, Hackett K, et al. Applying inverse probability weighting to measure contraceptive prevalence using data from a community-based reproductive health intervention in Pakistan. *Int Perspect Sex Reprod Health* 2020;46:21–33.
- Jenkins SP. Easy estimation methods for discrete-time duration models. *Oxf Bull Econ Stat* 1995;57:129–36.
- Kozuki N, Lee ACC, Silveira MF, et al. The associations of birth intervals with small-for-gestational-age, preterm, and neonatal and infant mortality: a meta-analysis. *BMC Public Health* 2013;13:S3.
- Aleni M, Mbalinda SN, Muhindo R. Birth intervals and associated factors among women attending young child clinic in Yumbe Hospital, Uganda. *Int J Reprod Med* 2020;2020:1–11.
- Yohannes S, Wondafrash M, Abera M, et al. Duration and determinants of birth interval among women of child bearing age in southern Ethiopia. *BMC Pregnancy Childbirth* 2011;11:38.
- Nyein C, Keiwkarnka B, Sillabutra J. Factors affecting the birth spacing among rural pregnant women in Salin township, Myanmar. *Journal of Health Research* 2014;28:165–71.



- 49 Singh SN, Singh SN, Narendra R. Demographic and socio-economic determinants of birth interval dynamics in Manipur: a survival analysis. *Journal of Health and Allied Sciences* 2011;9:3.
- 50 Kamal A, Pervaiz MK. Determinants of higher order birth intervals in Pakistan. *Journal of Statistics* 2012;19:54–82.
- 51 Khan JR, Bari W, Latif AHMM. Trend of determinants of birth interval dynamics in Bangladesh. *BMC Public Health* 2016;16:934.
- 52 Eini-Zinab H, Agha H. Demographic and Socio-Economic determinants of birth interval dynamics in Iran: a hazard function analysis. paper presented at: paper accepted to be appeared as poster at the XXV general population conference of the International Union for the Scientific Study of Population (IUSSP) during 2005.
- 53 Yeakey MP, Muntifering CJ, Ramachandran DV, et al. How contraceptive use affects birth intervals: results of a literature review. *Stud Fam Plann* 2009;40:205–14.
- 54 Kiani MF, Nazli S. Dynamics of birth spacing in Pakistan. *Pak Dev Rev* 1998;27:655,657.
- 55 Rajpoot UA. Correlates of birth interval: fresh evidence from WFS data. *Int J Contemp Sociol* 1995;32:81–96.
- 56 Sathar Z. Birth spacing in Pakistan. *J Biosoc Sci* 1988;20:175–94.
- 57 Tessema GA, Zeleke BM, Ayele TA. Birth interval and its predictors among married women in Dabat district, Northwest Ethiopia: a retrospective follow up study. *Afr J Reprod Health* 2013;17:39–45.
- 58 Rasheed P, Al-Dabal BK. Birth interval: perceptions and practices among urban-based Saudi Arabian women. *East Mediterr Health J* 2007;13:881–92.
- 59 Jatrana S, Pasupuleti SSR. Women's autonomy, education and birth intervals in India. *Asian Popul Stud* 2015;11:172–90.
- 60 Kim J. Women's education and fertility: an analysis of the relationship between education and birth spacing in Indonesia. *Econ Dev Cult Change* 2010;58:739–74.
- 61 Chakraborty N, Sharmin S, Islam MA. Differential pattern of birth intervals in Bangladesh. *Asia Pac Popul J* 1996;11:1–9.
- 62 Hajian-Tilaki KO, Asnafi N, Aliakbarnia-Omrani F. The patterns and determinants of birth intervals in multiparous women in Babol, Northern Iran. *Southeast Asian J Trop Med Public Health* 2009;40:852.
- 63 Saeed S. Towards an explanation of son preference in Pakistan. *Soc Dev Issues*;37:17–36.
- 64 Gray E, Evans A. Parity progression in Australia: what role does sex of existing children play? *Australian Journal of Social Issues* 2005;40:505.