

ORIGINAL ARTICLE

A Study of Awareness and Screening Behavior of Cardiovascular Risk Factors in Patients with Psoriasis and Dermatologists

Min Kyung Lee, Han Su Kim, Eun Byul Cho, Eun Joo Park, Kwang Ho Kim, Kwang Joong Kim

Department of Dermatology, Hallym University College of Medicine, Anyang, Korea

Background: A number of studies have suggested an increased frequency of cardiovascular (CV) diseases in patients with psoriasis. **Objective:** In this study, we assessed the awareness among psoriasis patients and dermatologists in private primary clinics about the increased CV risk linked to psoriasis, and examined the screening behaviors of dermatologists for CV risk factors in psoriasis patients. **Methods:** We distributed questionnaires to dermatologists in primary clinics and to psoriasis patients about their awareness of the increased CV risk factors related with psoriasis. **Results:** One hundred four patients and 50 dermatologists were included. 64.4% of patients and 92% of dermatologists answered that they know about the increased risk of CV diseases in psoriasis patients. However, far fewer dermatologists than expected followed the screening guidelines for CV risk factors. We found that the duration ($p < 0.0001$) and severity ($p < 0.0001$) of psoriasis were related to patient's awareness. A significant correlation between dermatologists' awareness and the number of psoriasis patients they treated each month was also observed ($p < 0.024$). **Conclusion:** This study may help promote the idea that psoriasis patients require education about their increased CV risk factors and that dermatologists require further education about screening practices to detect CV risk in psoriasis patients. (**Ann Dermatol** 27(1) 59~65,

Received July 24, 2014, Revised November 25, 2014, Accepted for publication December 3, 2014

Corresponding author: Kwang Joong Kim, Department of Dermatology, Hallym University Sacred Heart Hospital, 22, Gwanpyeong-ro 170beon-gil, Dongan-gu, Anyang 431-796, Korea. Tel: 82-31-380-3765, Fax: 82-31-386-3761, E-mail: kkj51818@hallym.or.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

2015)

-Keywords-

Awareness, Cardiovascular diseases, Psoriasis

INTRODUCTION

Psoriasis is a chronic immune-mediated inflammatory disease, characterized by scaly plaques or pustules, with a prevalence rate of approximately 1.5% ~ 3%¹⁻³. Recent studies have revealed that systemic inflammatory diseases such as psoriasis, systemic lupus erythematosus, and rheumatoid arthritis are associated with an increased risk of cardiovascular (CV) diseases⁴. In particular, it has been well established that the prevalence of CV diseases and conventional CV risk factors, such as obesity, smoking, diabetes mellitus, hyperlipidemia, and hypertension, is higher in patients with psoriasis than in the general population⁵⁻⁸. Moreover, psoriasis is related to high CV mortality rates, whereas its severity is associated with an increased risk of CV disease⁹⁻¹¹.

Despite previous studies that demonstrated an increased risk of CV disease and the establishment of screening guidelines for CV risk factors in psoriasis patients¹², it is not certain whether psoriasis patients and dermatologists are aware of this risk and whether dermatologists follow the recommended screening guidelines. We assumed that dermatologists in private primary practice have relatively fewer opportunities to obtain knowledge about the increased CV risk factors in psoriasis patients than dermatologists working in university hospitals, and we wondered about the awareness and screening behaviors of dermatologists in private primary clinics as well as about the awareness of psoriasis patients. The aims of this study are to

assess the awareness among psoriasis patients and dermatologists in primary clinics about the increased CV risks linked to psoriasis, and to evaluate the screening behaviors for CV risk factors among dermatologists.

MATERIALS AND METHODS

Subjects and data collection

First, psoriasis patients were randomly selected among outpatients with psoriasis aged ≥20 years who presented to the Department of Dermatology, Hallym University Sacred Heart Hospital. One hundred four patients, with adequate time to receive an explanation about the study, provide consent, and complete questionnaires, were enrolled between June 1 and December 31, 2013. The patients completed questionnaires about their awareness of increased CV risks linked to psoriasis and the screening behaviors in detecting risk factors (Table 1). To evaluate the clinical variables that are related with patients' awareness, a detailed history was obtained from each patient,

including sociodemographic issues and the duration of their psoriasis. Information about the severity of psoriasis, assessed by using the Psoriasis Area Severity Index (PASI), was obtained from the patients' medical records, and the highest score was selected. The patients were questioned about how they obtained the information about increased CV risk factors.

Board-certified, Korean dermatologists practicing in private primary clinics were randomly selected from the Korean Dermatological Association. One hundred questionnaires were distributed through electronic mail. A total of 50 dermatologists completed the questionnaire (Table 2). The questionnaire identified the dermatologists' awareness of the high CV risk factors in psoriasis patients and their screening practices for CV risk factors when treating these patients. To identify the relevant factors associated with their awareness, the dermatologists were asked about the number of patients with psoriasis they treated per month. They were also questioned about their screening methods for CV risk factors.

Table 1. Questionnaire for patients about the awareness of increased cardiovascular (CV) risk in psoriasis patients

1. Please indicate your sex. ① Male ② Female
2. What is your age? _____ years
3. Did/do you have any of the following comorbid diseases?
 ① Hypertension ② Diabetes mellitus ③ Hyperlipidemia ④ Other _____
4. Smoking history
 ① Current smoker (____ pack(s) per day on average, for _____ year(s))
 ② Ex-smoker (____ pack(s) per day on average, for _____ year(s))
 ③ None
5. When were you diagnosed with psoriasis? _____ year(s) ago
6. Please indicate all the following psoriasis treatments until date. (Please indicate all that you are aware of.)
 ① Systemic therapy (cyclosporine, vitamin A, methotrexate)
 ② Topical therapy (vitamin D, steroid)
 ③ Phototherapy (narrow band UVB, PUVA)
 ④ Biologics
 ⑤ Other _____
7. Compared to the general population, psoriasis patients are at greater risk of having CV disease risk factors (such as high blood pressure, diabetes mellitus, obesity, and hyperlipidemia) and CV diseases (such as coronary artery disease, myocardial infarction, and stroke). Are you aware of this fact?
 ① I am fully aware of this. (Please answer questions 7-1 and 7-2.)
 ② I have heard this, but do not know what it means to me. (Please answer questions 7-1 and 7-2.)
 ③ I have not heard this.
- 7-1. Where have you hear about the information in question 7?
 ① Internet ② Media (TV, newspaper, magazine, etc.)
 ③ Doctor ④ Acquaintance ⑤ Other _____
- 7-2. After you learned about the information in question 7, did you make any of the following lifestyle changes?
 ① I started getting regular medical checkups.(blood sugar, cholesterol, blood pressure, body mass index)
 ② I started working out regularly.
 ③ I changed my diet.
 ④ I quit smoking.
 ⑤ I started drinking less.
 ⑥ Other _____

Table 2. Questionnaire for dermatologists on the awareness of increased cardiovascular (CV) risk in psoriasis patients

1. Please indicate your sex.	<input type="radio"/> Male	<input type="radio"/> Female	
2. What is your age? _____ years			
3. Please indicate in which of the following setting you practice medicine.			
① Primary clinic (level 1 medical institution)			
② Hospital (level 2 or 3 medical institution)			
③ Other _____			
4. How long have you practiced medicine as a dermatologist?	<input type="radio"/> <5 years	<input type="radio"/> 5~10 years	
	<input type="radio"/> 10~20 years	<input type="radio"/> >20 years	
5. How many psoriasis patients do you typically examine per month?	<input type="radio"/> 0	<input type="radio"/> 1~5	<input type="radio"/> 6~10
	<input type="radio"/> 11~15	<input type="radio"/> >20	
6. What percentage of your patients have psoriasis?	<input type="radio"/> <20%	<input type="radio"/> 20%~40%	<input type="radio"/> 40%~60%
	<input type="radio"/> 60%~80%	<input type="radio"/> 80%~100%	
7. Compared to the general population, psoriasis patients are at greater risk of having CV disease risk factors (such as high blood pressure, diabetes mellitus, obesity, and hyperlipidemia) and CV diseases (such as coronary artery disease, myocardial infarction, and stroke). Are you aware of this fact?			
① I am fully aware of this. (Please answer question 7-1.)			
② I have heard this, but do not know what it means to me. (Please answer question 7-1.)			
③ I have not heard this.			
7-1. If you are aware of this information, do you explain this to your patients?	<input type="radio"/> Yes	<input type="radio"/> No	
8. Do you screen psoriasis patients for dyslipidemia?	<input type="radio"/> (Please answer question 8-1.)	<input type="radio"/> No	
8-1. Which of the following items do you examine?			
① Medical history			<input type="radio"/> Total cholesterol
④ High-density lipoprotein			<input type="radio"/> Low-density lipoprotein
⑤ Triglyceride			<input type="radio"/> Other _____
9. Do you screen psoriasis patients for obesity (body mass index measurement)?	<input type="radio"/> Yes	<input type="radio"/> No	
10. Do you screen psoriasis patients for diabetes?	<input type="radio"/> Yes (Please answer question 10-1.)	<input type="radio"/> No	
10-1. Which of the following methods do you use? (Please indicate all that apply.)			
① Refer to medical history			
② Random plasma glucose test			
③ Fasting plasma glucose test			
④ Oral glucose tolerance test			
⑤ HbA1c			
11. Do you screen psoriasis patients for blood pressure?			
① Yes (Please answer question 11-1.)			<input type="radio"/> No
11-1. Which of the following method do you use?			
① Refer to medical history			<input type="radio"/> Check blood pressure
12. If you do not perform any of the screening tests mentioned in questions 8~11 for psoriasis patients, why? (Please indicate all that apply.)			
① I did not know it was necessary.			
② Patients' financial circumstances			
③ Lack of time			
④ Lack of facilities or testing equipment			
⑤ Other _____			

Statistical analysis

Data are expressed as mean \pm standard deviation. An independent t-test was used to analyze the data. A χ^2 test was also used to identify the factors that were related to the awareness of increased CV risk in patients with psoriasis. All statistical analyses were performed by using SPSS ver. 13.0 (SPSS Inc., Chicago, IL, USA). p -values <0.05 were considered statistically significant.

Ethics statement

This study received approval from the institutional review board of Hallym University Sacred Heart Hospital on May 10, 2013 (IRB No. 2014-I011). Written informed consent was obtained from all participants.

RESULTS

Demographic data of the patients

A total of 104 patients (49 men, 55 women) with plaque type psoriasis were included in this study (Table 3). The mean age of psoriasis patients was 49.7 ± 12.6 years, and the mean duration of psoriasis was 10.5 ± 8.7 years.

Table 3. Sociodemographic characteristics of the patients (n=104)

Demographics	Value
Age (y)	49.7 ± 12.6
Sex	47.1
Annual household income (\$)	$61,550 \pm 10,300$
Employment status	
Employed	62
Unemployed	13
Student	6
Homemaker	16
Retired	3
Educational level	
High school or less	39
College education	51
Graduate school education	10

Values are presented as mean \pm standard deviation or percentage of total subjects.

Table 4. Comparisons of awareness of psoriasis patients by clinical factor

Variable	n	Unaware of increased CV risk factors (n=37)	Aware of the risk, but not very well (n=39)	Well aware of increased CV risk factors (n=28)	p-value*
Sex					0.792
Male	49	19 (38.8)	17 (34.7)	13 (26.5)	
Female	55	18 (32.7)	22 (40.0)	15 (27.3)	
Age (y)					0.097
20~30	2	2 (100.0)	0 (0)	0 (0)	
31~40	21	6 (28.6)	11 (52.4)	4 (19.0)	
41~50	32	7 (21.9)	12 (37.5)	13 (40.6)	
51~60	27	10 (37.0)	10 (37.0)	7 (25.9)	
61~70	14	6 (42.9)	4 (28.6)	4 (28.6)	
≥ 71	8	6 (75.0)	2 (25.0)	0 (0)	
Duration (y)					0.000
<1	6	2 (33.3)	4 (66.7)	0 (0)	
1~5	35	15 (42.9)	19 (54.3)	1 (2.9)	
5~10	12	7 (58.3)	3 (25.0)	2 (16.7)	
10~20	35	9 (25.7)	9 (25.7)	17 (48.6)	
>20	16	4 (25.0)	4 (25.0)	8 (50.0)	
Severity [†]					0.000
Mild	88	36 (40.9)	35 (39.8)	17 (19.3)	
Moderate to severe	16	1 (6.3)	4 (25.0)	11 (68.8)	

Values are presented as number of subjects (percentage of total). CV: cardiovascular. *p-values <0.05 considered statistically significant.

[†]The patients with a Psoriasis Area Severity Index (PASI) score <10 were considered to have mild psoriasis, whereas those with a PASI score ≥ 10 were considered to have moderate to severe psoriasis.

Awareness among psoriasis patients of their increased CV risk

Only 28 of the 104 (26.9%) patients were well aware of their increased CV risk factors, whereas 39 (37.5%) had heard about the increased CV risk in psoriasis patients. However, these patients did not have a thorough understanding of the risk. The other 37 (35.6%) patients answered that they did not have any idea about their increased risk of CV disease. The way in which patients acquired this knowledge was noted. More than one answer was possible. The most common source of information was mass media (67 patients, 40.3%), followed by the internet (25, 37.3%) and their doctors (18, 26.9%). The patients were also asked whether they had made any efforts to reduce their risk of adverse CV outcome. A total of 53 (79.1%) patients responded that they did, whereas the rest responded that they had made no effort at all. The most common efforts made were exercising and changing dietary habits, accounting for 24 (35.8%) and 21 (31.3%) patients, respectively. Eighteen of the 67 patients (26.9%) received screening such as regular laboratory evaluations, whereas 8 (11.9%) reduced their alcohol consumption, and 6 (9.0%) had quit smoking.

The correlation between patients' awareness and their clinical factors was examined. The mean duration of psor-

iasis was 10.5 ± 8.7 years, and the mean PASI score was 5.7 ± 3.5 . We found that the factors, significantly associated with patients' awareness, were the duration ($p < 0.0001$) and severity ($p < 0.0001$) of psoriasis (Table 4). Among the patients who had psoriasis for < 1 year, none were well aware of their increased CV risk. Eight of the 16 (50.0%) patients who had psoriasis for > 20 years showed the greatest awareness. Although only 17 (19.3%) patients with mild psoriasis were well aware of their increased CV risk, a larger proportion (11 of 16, 68.8%) of patients with moderate to severe psoriasis were well aware of their increased CV risk. There was no sex- or age-related predominance in awareness among the participants of this study.

Awareness among dermatologists of increased CV risk

Forty-six of the 50 (92%) dermatologists were aware that psoriasis patients have a greater CV risk than the normal population. Thirty-six (72%) dermatologists were well aware of the risk, whereas the other 10 (20%) had heard of the increased risk, but did not know much about it. The mean duration of clinical practice was 12.1 ± 9.0 years, and the mean number of psoriasis patients that the dermatologists treated per month was 7.1 ± 5.0 . A significant correlation between dermatologists' awareness and the number of psoriasis patients they treated per month was observed ($p < 0.024$; Table 5). None of the dermatologists who treated no patients with psoriasis were well aware of the increased CV risk factors, whereas the four dermatologists who treated > 20 patients per month recognized the risk factors well. The duration of clinical practice was not significantly related to their awareness.

Despite their awareness, the number of dermatologists who screened psoriasis patients for CV risk factors was far less than expected. Five of the 36 (13.9%) dermatologists

screened patients for dyslipidemia, 17 (47.2%) for obesity, 11 (30.6%) for hypertension, and 10 (27.8%) for diabetes mellitus. The reasons why the dermatologists did not screen for CV risk factors were a lack of screening facilities (50%), lack of time (27.1%), lack of feeling the need to perform screening (20.2%), and lack of patient compliance (12.2%). Four dermatologists answered that they did not know exactly what they had to do to evaluate a patient for CV risk factors.

DISCUSSION

In this study, less than half of the psoriasis patients were well aware of their increased CV risk factors. The results of this study suggest that the severity and duration of psoriasis are associated with the awareness of CV risk factors in psoriasis patients. The severe group with PASI ≥ 10 tended to be more aware of the high CV risk factors related with psoriasis. The results also revealed that patients who had psoriasis for a longer duration recognized their risk better than those who had psoriasis for a shorter duration. Whether duration or severity was correlated with the increased CV risk is controversial; however, the patients' awareness in this study was related to the duration and severity of their psoriasis. We assume that this relation exists because psoriasis patients with longer duration and higher severity are likely to have more opportunities to obtain information about increased CV risk during their long and severe disease course.

We also found that patients acquired their knowledge about the increased CV risk factors more often through non-professional channels than from their doctors. This finding suggests that most dermatologists do not educate their patients about their increased CV risk factors or their

Table 5. Comparisons of dermatologist awareness according to the number of psoriasis patients and duration of clinical practice

Variable	n	Unaware of increased CV risk factors (n=4)	Aware of the risk, but not very well (n=10)	Well aware of increased CV risk factors (n=36)	p-value*
Number of psoriasis patients					0.024
0	2	0 (0)	2 (100.0)	0 (0)	
1~5	17	1 (5.9)	3 (17.6)	13 (76.5)	
6~10	14	1 (7.1)	4 (28.6)	9 (64.3)	
11~15	5	2 (40.0)	1 (20.0)	2 (40.0)	
16~20	8	0 (0)	0 (0)	8 (100.0)	
>20	4	0 (0)	0 (0)	4 (100.0)	
Duration of clinical practice (y)					0.766
<5	9	0 (0)	2 (22.2)	7 (77.8)	
5~10	16	1 (6.3)	4 (25.0)	11 (68.8)	
10~20	18	2 (11.1)	4 (22.2)	12 (66.7)	
>20	7	1 (14.3)	0 (0)	6 (85.7)	

Values are presented as number of subjects (percentage of total). CV: cardiovascular. *p-values < 0.05 considered statistically significant.

need to undergo screening. Less than half of these patients were routinely screened for CV risk factors, such as through regular laboratory evaluations, and we can assume that this result reflects low patient awareness, as this study showed.

On the other hand, most of the dermatologists responded that they know about the increased CV risk factors of psoriasis patients. Increasing numbers of studies about the co-morbidities of psoriasis, especially CV co-morbidities, have been established; this may have contributed to the dermatologists' awareness^{13,14}. This study revealed that dermatologists who treated a greater number of psoriasis patients were more likely to be well aware of the increased CV disease risk and more likely to screen for risk factors. An extensive experience with psoriasis patients has helped dermatologists learn more about their CV risk factors. The duration of clinical practice was not significantly related to the dermatologists' awareness. We can assume that dermatologists who have had a longer practice are likely to have fewer opportunities to obtain the latest knowledge such as on the co-morbidities of psoriasis. However, we can also carefully speculate that dermatologists with a longer duration of practice have more experience treating psoriasis patients, resulting in their increased awareness. These two possible contrasting ideas may be helpful in understanding our data, which showed no significant findings between the dermatologists' awareness and their duration of clinical practice.

Our study also found that the dermatologists did not always screen for CV risk factors, such as hypertension, dyslipidemia, diabetes mellitus, and obesity. Most of the dermatologists responded that they did not screen for CV risk factors because of a lack of facilities. This result is consistent with the fact that obesity and hypertension, which are much easier to check in outpatient clinics, seem to be screened at higher rates than other risk factors. Notably, more than half of the dermatologists who were aware of the high CV risk factors in psoriasis patients did not educate their patients about this risk despite their awareness.

In an effort to reduce CV risk factors in psoriasis patients, the National Psoriasis Foundation recommended the use of the American Heart Association guidelines, and suggested that psoriasis patients be screened for CV risk factors starting at 20 years of age¹⁵. Wohlrab et al.¹⁶ recently developed a checklist that allows dermatologists to detect all relevant factors related to the co-morbidities of patients with moderate to severe psoriasis, and they emphasized the importance of interdisciplinary complementary assessments.

Overall, the findings of this study suggest that although most dermatologists are aware of the greater CV risk fac-

tors and CV disease rates in psoriasis patients, only a small proportion of psoriasis patients are well aware of their risk. Dermatologists should feel a responsibility to educate their patients and screen for CV risk factors on the basis of accurate knowledge about the association between psoriasis and CV diseases. In addition, larger, multi-institutional studies are needed to confirm the factors related to the awareness of increased CV risk factors.

This study has several limitations. First, it used questionnaires distributed by electronic mail to dermatologists, which means the possibility of statistical bias is present. Second, the awareness of increased CV risk factors can differ according to the existence of CV diseases in psoriasis patients. Further studies with matching co-morbidities are needed.

REFERENCES

- Rapp SR, Feldman SR, Exum ML, Fleischer AB Jr, Reboussin DM. Psoriasis causes as much disability as other major medical diseases. *J Am Acad Dermatol* 1999;41:401-407.
- Schön MP, Boehncke WH. Psoriasis. *N Engl J Med* 2005; 352:1899-1912.
- Federman DG, Shelling M, Prodanovich S, Gunderson CG, Kirsner RS. Psoriasis: an opportunity to identify cardiovascular risk. *Br J Dermatol* 2009;160:1-7.
- Griffiths CE, Iaccarino L, Naldi L, Olivier I, Pipitone N, Salvarani C, et al. Psoriasis and psoriatic arthritis: immunological aspects and therapeutic guidelines. *Clin Exp Rheumatol* 2006;24(1 Suppl 40):S72-S78.
- Neumann AL, Shin DB, Wang X, Margolis DJ, Troxel AB, Gelfand JM. Prevalence of cardiovascular risk factors in patients with psoriasis. *J Am Acad Dermatol* 2006;55:829-835.
- Mallbris L, Ritchlin CT, Stähle M. Metabolic disorders in patients with psoriasis and psoriatic arthritis. *Curr Rheumatol Rep* 2006;8:355-363.
- Henseler T, Christophers E. Disease concomitance in psoriasis. *J Am Acad Dermatol* 1995;32:982-986.
- Choi WJ, Park EJ, Kwon IH, Kim KH, Kim KJ. Association between psoriasis and cardiovascular risk factors in Korean patients. *Ann Dermatol* 2010;22:300-306.
- Wakkee M, Thio HB, Prens EP, Sijbrands EJ, Neumann HA. Unfavorable cardiovascular risk profiles in untreated and treated psoriasis patients. *Atherosclerosis* 2007;190:1-9.
- Gelfand JM, Neumann AL, Shin DB, Wang X, Margolis DJ, Troxel AB. Risk of myocardial infarction in patients with psoriasis. *JAMA* 2006 296:1735-1741.
- Mallbris L, Akre O, Granath F, Yin L, Lindelöf B, Ekbom A, et al. Increased risk for cardiovascular mortality in psoriasis inpatients but not in outpatients. *Eur J Epidemiol* 2004;19: 225-230.
- Friedewald VE, Cather JC, Gelfand JM, Gordon KB, Gibbons GH, Grundy SM, et al. AJC editor's consensus: psoriasis and coronary artery disease. *Am J Cardiol* 2008;102:1631-1643.

13. Gisondi P, Farina S, Giordano MV, Girolomoni G. Usefulness of the framingham risk score in patients with chronic psoriasis. *Am J Cardiol* 2010;106:1754-1757.
14. Mehta NN, Azfar RS, Shin DB, Neumann AL, Troxel AB, Gelfand JM. Patients with severe psoriasis are at increased risk of cardiovascular mortality: cohort study using the General Practice Research Database. *Eur Heart J* 2010;31: 1000-1006.
15. Kimball AB, Gladman D, Gelfand JM, Gordon K, Horn EJ, Korman NJ, et al; National Psoriasis Foundation. National Psoriasis Foundation clinical consensus on psoriasis comorbidities and recommendations for screening. *J Am Acad Dermatol* 2008;58:1031-1042.
16. Wohlrab J, Fiedler G, Gerdes S, Nast A, Philipp S, Radtke MA, et al. Recommendations for detection of individual risk for comorbidities in patients with psoriasis. *Arch Dermatol Res* 2013;305:91-98.