

Review Article

PPAR- δ in Vascular Pathophysiology

Nanping Wang

Laboratory of Vascular Molecular Biology and Bioengineering, Institute of Cardiovascular Science and Diabetes Center, Peking University Health Science Center, Beijing 100191, China

Correspondence should be addressed to Nanping Wang, npwang@bjmu.edu.cn

Received 25 July 2008; Accepted 27 October 2008

Recommended by Francine M. Gregoire

Peroxisome proliferator-activated receptors belong to the superfamily of ligand-dependent nuclear receptor transcription factors, which include three subtypes: PPAR- α , β/δ , and γ . PPAR- δ , play important roles in the regulation of cell growth and differentiation as well as tissue wound and repair. Emerging evidence has also demonstrated that PPAR- δ is implicated in lipids and glucose metabolism. Most recently, the direct effects of PPAR- δ on cardiovascular processes such as endothelial function and angiogenesis have also been investigated. Therefore, it is suggested that PPAR- δ may have critical roles in cardiovascular pathophysiology and is a potential target for therapeutic intervention of cardiovascular disorders such as atherosclerosis.

Copyright © 2008 Nanping Wang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

Peroxisome proliferator-activated receptors (PPARs) are members of nuclear receptor/ligand-activated transcription factors superfamily. PPAR subfamily consists of 3 subtypes: PPAR- α , β/δ , and γ . PPARs form heterodimers with a retinoid X receptor (RXR) and bind to specific PPAR-responsive elements (PPREs) to regulate target gene expression. In the absence of specific ligands, the PPAR-RXR heterodimer forms repressive complex with corepressors and histone deacetylases. Upon ligand binding, the receptor undergoes conformational changes that cause the dissociation of repressors, the recruitment of coactivators, and the activation of gene transcription [1]. Recently, extensive studies have been performed to characterize the biological and pathophysiological roles of PPAR- α and γ , which are pharmacological targets of the clinical interventions for dyslipidemia and type 2 diabetes, respectively. Fibrates class of lipid-lowering drugs such as fenofibrate and gemfibrozil are agonists for PPAR- α . Thiazolidinedione class of insulin sensitizers including troglitazone (Rezulin), rosiglitazone (Avandia), and pioglitazone (Actos) are specific ligands for PPAR- γ [2]. PPAR- α , primarily expressed in liver, muscles, heart, and kidney, plays a key role in fatty acid catabolism such as β -oxidation. PPAR- γ is highly expressed in fat, controls adipogenesis, and regulates insulin action. However, relatively little has been known with regards to the function

of PPAR- δ , the only subtype of PPARs that is not a target of current drug. Newly developed synthetic ligands and genetically modified mouse models for PPAR- δ have rapidly advanced our understanding of the important roles of PPAR- δ in tissue development, repair, inflammation, and metabolism [3–7]. Most recently, the direct effects of PPAR- δ on cardiovascular processes such as endothelial function and angiogenesis have also been investigated. In this review, we will focus on the recent advancements regarding the roles of PPAR- δ in vascular pathophysiological processes.

2. MATERIALS AND METHODS

2.1. Gene and protein

PPAR- δ , also known as nuclear hormone receptor 1 (NUC1), PPAR- β , or NR1C2, was first cloned in 1992 [8, 9]. The PPAR- δ gene is mapped to human chromosome 6p21.2-p21.1 and has 11 exons, spanning 35 kilobase-pair [10]. Like other PPARs, PPAR- δ protein has a modular structure consisted of 5 regions: an N-terminal region (A/B), a DNA-binding domain (C), a flexible hinge region (D), ligand-binding domain (E), and a C-terminal region (F). X-ray crystallographic study revealed that PPAR- δ has an exceptionally large ligand binding pocket, which maybe related to the promiscuous accommodation of a large range of mostly amphipathic ligands [11, 12].

2.2. Endogenous and synthetic ligands

Several 14- to 18-carbon saturated fatty acids and 16- to 20-carbon polyunsaturated fatty acids can bind PPAR- δ [13–15]. Naturally occurring or synthetic eicosanoids such as prostaglandin A1 and carbaprostacyclin have been shown to bind and activate PPAR- δ [16]. Very low-density lipoprotein (VLDL) derived has also been demonstrated to activate the PPAR- δ target genes in a receptor-dependent manner [17]. Since these agonists activate PPAR- δ all with affinities at molar range, it raises a question as to whether these are bona fide physiological ligands for PPAR- δ . However, many of the above-mentioned agonists under the physiological or pathological conditions are either released by the vessels, such as PGI₂, or being exposed to vascular endothelium, such as VLDL. It would be intriguing to examine whether PPAR- δ in the vessel wall is activated *in vivo*. In addition, it was recently shown that retinoic acid, a ligand for retinoic acid receptor, also can activate the PPAR- δ with nanomolar affinity without affecting the other two subtypes of PPARs [18]. This finding expanded our understanding of the mechanisms of PPAR- δ activation.

Recently, several synthetic ligands have been reported to selectively activate PPAR- δ . The PPAR- δ agonists reported to date were discovered using several strategies: GW501516 and GW-0742 (GlaxoSmithKline) were optimized from a library of hydrophobic carboxylates [19]; L165461 (Merck) was derived from an *in silico* approach [20]. These derivatives of phenoxyacetic acid are the highly selective PPAR- δ ligands with a nanomolar affinity and 1000-fold selectivity over PPAR- α and - γ . Other PPAR- δ agonists including KD3010 (Kalypsys) and MBX-8025 (Metabolex) are currently in clinical development. On the other hand, the development of these specific agonists has greatly aided the investigation in the biological functions of PPAR- δ . A selective antagonist for PPAR- δ , GSK0660, has also been recently demonstrated as it by itself exhibits inverse agonist activity and competes with agonist in a cellular context [21] (Table 1).

2.3. Effects on the vessel wall

Although nearly ubiquitously expressed with highest levels in placenta, skeletal muscles, and adipose tissue, PPAR δ is also expressed in the vascular cells including endothelial cells [22], smooth muscle cells, and macrophages. Particularly, a number of studies during the past 2 years have demonstrated that PPAR- δ plays direct roles in various basic vascular processes such as apoptosis, survival, angiogenesis, and inflammation.

2.4. Endothelial apoptosis

Vascular endothelium, when unperturbed, is considered to provide a relatively nonadhesive and nonthrombotic interface. This characteristic is likely essential to physiological homeostasis. However, endothelial cells (ECs) can undergo apoptosis *in vitro* in response to a variety of pathophysiological conditions including hypoxia, proinflammatory cytokines, bacterial endotoxins, and atherogenic risk factors

such as homocysteine and lipoproteins [23, 24]. EC apoptosis has been implicated in numerous pathophysiological processes, such as angiogenesis, thrombosis, and atherosclerosis. On the other hand, ECs produce a plethora of bioactive molecules to maintain vascular homeostasis. Among those, prostacyclin (PGI₂) protects ECs from apoptosis. Although PPAR- δ has been previously documented to protect against the hypertonicity-induced apoptosis in renal cells [25] and the growth factor deprivation- or anoikis-induced apoptosis in keratinocytes [6], its role in vascular cells has been recently demonstrated. Liou et al. showed that PGI₂ protects ECs from H₂O₂-induced apoptosis via the action of PPAR- δ . By inducing the expression of its target gene 14-3-3 α , PPAR- δ prevents Bad-triggered apoptosis [26]. Treatment with L165041 or overexpression of PPAR- δ also has a similar effect. In addition, small interfering RNA-mediated knockdown of PPAR- δ abrogated the antiapoptotic effect, suggesting that the antiapoptotic role of PPAR- δ appeared to be specific for and dependent on the endogenous PPAR- δ receptor. Interestingly, gene expression of 14-3-3 ϵ was also induced by PPAR- δ through a PPRE-independent mechanism and an interaction between PPAR- δ and CCAAT/enhancer binding protein (C/EBP) [27].

2.5. Endothelial activation

When exposed to proinflammatory stimuli such as tumor necrosis factor (TNF) or lipopolysaccharide (LPS), normally quiescent endothelium undergoes a phenotypic change, which is characterized by induction of proinflammatory and procoagulant factors such as adhesion molecules and tissue factor. Such a phenotypic conversion, referred to as EC activation, is implicated in a number of proinflammatory diseases including atherosclerosis and thrombosis [28]. PPAR- α and - γ have been previously shown to suppress EC expression of proinflammatory adhesion molecules such as intercellular adhesion molecule-1 (ICAM-1), vascular adhesion molecule-1 (VCAM-1), E-selectin, monocyte chemoattractant protein-1 (MCP-1), and ensuing recruitment of leukocytes [22, 29–32]. However, there has also been evidence that suggests a proinflammatory role of PPAR- α or - γ [33, 34]. Recent studies suggested that PPAR- δ also plays a role in inflammatory processes and atherosclerosis. In macrophages, the PPAR- δ agonist GW0742 inhibited lipopolysaccharide (LPS)-induced expression of proinflammatory genes, such as cyclooxygenase (COX)-2 and inducible nitric oxide synthase (iNOS) [35]. GW0742 reduced atherosclerotic lesions and decreased the expression of MCP-1 and ICAM-1 in the aorta of LDLR^{-/-} mice [36, 37]. Given the beneficial effects of PPAR- δ agonists on lipid profiles, it is likely that PPAR- δ agonists may inhibit endothelial activation by improving dyslipidemia. A direct anti-inflammatory effect was also demonstrated. In EAhy926 cells, Rival et al. found that L-165041, at high a concentration up to 100 μ M, inhibited TNF- α -induced VCAM-1 and MCP-1 expressions [38]. In primary culture of human umbilical vein ECs (HUVECs), specific agonists GW0742 and GW501516 inhibited the TNF- α - or interleukin-1 β -induced expression of adhesion

TABLE 1: Ligands for PPAR- δ .

Ligands	Nature	Affinity	Clinical status
<i>Natural agonists</i>			
Linoleic acid	Dietary fatty acid	μM	
Oleic acid	Dietary fatty acid	μM	
Arachidonic acid	Dietary fatty acid	μM	
Eicosapentaenoic acid	Dietary fatty acid	μM	
Docosahexaenoic acid	Dietary fatty acid	μM	
Prostaglandin A1	Endogenous prostaglandin	μM	
<i>Synthetic agonists</i>			
Carbaprostacyclin	Synthetic stable PGI ₂ analogue	μM	
Iliprost	Prostacyclin analogue	μM	
Compound F	Phenoxyacetic acid derivative	nM	
L165,041	Phenoxyacetic acid derivative	nM	
GW501516	Phenoxyacetic acid derivative	nM	
GW0742	Phenoxyacetic acid derivative	nM	
KD2010	not disclosed	nM	Phase I
MBX-8025	not disclosed	nM	Phase II
<i>Synthetic dual agonists</i>			
Compound 23	Dual agonist for γ and δ	nM	
<i>Synthetic antagonist</i>			
GSK0660	Antagonist for δ	nM	

molecules and the monocyte adhesion to ECs. PPAR- δ agonist induced the gene expression of antioxidative enzymes, such as superoxide dismutase-1, catalase, and thioredoxin, and it decreased reactive oxygen species production in ECs. Unexpectedly, the anti-inflammatory effect not only persisted but it also was further enhanced after the decrease of PPAR- δ expression by siRNA knockdown [39]. Given the evidence that the ligand binding caused the dissociation of the transcription repressor BCL-6 from PPAR- δ and the subsequent association of BCL-6 with the VCAM-1 promoter region, this seemingly paradoxical result could be plausibly interpreted with the previously proposed PPAR- δ /Bcl-6 interaction action mode: the synthetic ligand binds to PPAR- δ and recruits the coactivators to replace the corepressors such as Bcl-6. The released corepressors relocate to repress the transcription of proinflammatory genes such as VCAM-1 and E-selectin and thus contribute to the vascular protection.

Ghosh et al. recently showed that the metabolism of endocannabinoids by the endothelial COX-2 coupled to the prostacyclin synthase activates PPAR- δ , which negatively regulates the expression of tissue factor (TF), the primary initiator of blood coagulation [40]. As COX-2 inhibitors suppressed PPAR- δ activity and induced TF expression, these results may help explaining the prothrombotic adverse effects of the cox-2 inhibitors rofecoxib and valdecoxib [41].

2.6. Angiogenesis

Angiogenesis is referred to the formation of new capillaries from the existing blood vessels. Physiological angiogenesis is involved in wound healing and aerobic exercise, whereas

pathological or therapeutic angiogenesis is implicated in cardiovascular diseases, diabetic complications, inflammatory diseases, and cancers [42]. Recent studies have also linked metabolic homeostasis to angiogenesis and further interrogate the potential effects of PPARs on the angiogenic process [43, 44]. An earlier study demonstrated that the PPAR- δ agonist GW501516 dose-dependently stimulates HUVEC proliferation with increased mRNA expression of vascular endothelial growth factor α and its receptor flt-1 [45]. Later, GW501516 was shown to promote endothelial tube formation on an extracellular matrigel, EC outgrowth in a murine aortic ring model, and increased angiogenesis in the implanted matrigel plug assay in vivo through a PPAR- δ - and VEGF-dependent manner [22]. Most recently, Gaudel et al. found that treatment with GW0742 or muscle-specific overexpression of PPAR- δ promoted angiogenesis in mouse skeletal muscle [46]. Besides, arising from sprouts on existing vessels, vessels also arise from endothelial progenitor cells (EPCs), a process referred to as vasculogenesis [47]. Culminating evidence further suggests that circulating EPCs is capable of stimulating angiogenesis [48]. A recent study showed that the proangiogenic effects of human EPCs are in part dependent on the biosynthesis and release of PGI₂, and subsequent activation of PPAR- δ [48, 49]. Furthermore, functional genomic approach provided evidence that silencing of PPAR- δ in the tumor microenvironment impairs angiogenesis and tumor growth, identifying PPAR- δ as one of a few hub nodes in the angiogenic network [49, 50]. Up-to-date results have been largely consistent and pointed toward a proangiogenic activity of PPAR- δ . In corroboration with this, Müller-Brüsselbach et al. found that the growth of syngeneic PPAR- δ wild-type tumors was

impaired in PPAR- $\delta^{-/-}$ mice, concomitant with a reduced blood flow and hyperplastic vascular structures, suggesting that PPAR- δ maybe required in tumor ECs for the formation of functionally mature vessels [51]. Nevertheless, a full understanding of the specific roles of PPAR- δ in specific scenarios of angiogenesis will be imperative for a safe and rational therapeutic strategy.

2.7. Smooth muscle cells

PPAR- δ is expressed in SMCs and is induced in response to platelet-derived growth factor (PDGF) in SMCs, which involved the phosphatidylinositol 3-kinase/Akt signaling pathway. Initial study using overexpression showed that PPAR- δ increased SMC proliferation, indicating a proliferation-promoting effect in SMCs [52]. However, L-165041, a selective PPAR- δ agonist, inhibited SMC proliferation and migration via inhibition of the PDGF-induced expression of cyclin D1 and cyclin-dependent kinase (CDK) 4 and cell cycle progression [53]. In SMCs, GW501516 increased the expression of transforming growth factor- β 1 (TGF- β 1) and the effect seemed to depend on endogenous PPAR- δ . Subsequently, TGF- β 1 was likely responsible for suppression of the IL1 β -induced expression of MCP-1 and proliferation of SMCs [54]. In rats, administration of L-165041 decreased neointima formation in balloon-injured carotid arteries [53]. Thus, synthetic PPAR- δ agonists appear to have antiproliferative and anti-inflammatory properties in SMCs. This is consistent with previous reports that adenovirus-mediated gene transfer of prostacyclin synthase, which produces the endogenous PPAR- δ ligand PGI₂, inhibited SMCs proliferation and intimal hyperplasia [55–57].

2.8. Macrophages

Macrophage infiltration in vessel wall is known to play important role in atherogenesis. PPAR- δ is expressed in macrophages. During past years, the role of PPAR- δ in macrophage biology has been extensively studied. However, existing results still remain controversial. Oliver Jr. et al. showed that GW501516 in a human monocytic cell line increased the expression of ATP-binding cassette A1 (ABCA1) and Apo AI-mediated cholesterol efflux [58]. However, Vosper et al. found that a different PPAR- δ agonist, compound F, increased lipid accumulation in both human primary macrophages and THP-1 cells. Compound F induced the expression of genes involved in lipid uptake and storage such as class A and B scavenger receptors (SRA and CD36) but repressed key genes involved in lipid metabolism and efflux such as Apo E and cholesterol 27-hydroxylase [59]. In mouse macrophages, neither genetic loss of PPAR- δ nor treatment with the PPAR- δ agonists GW501516 or GW0742 significantly influenced cholesterol efflux or accumulation [36, 60]. Beside the effects on lipid trafficking, PPAR δ agonists have a potent anti-inflammatory effect in macrophages. Welch et al. first demonstrated that, in mouse peritoneal macrophages, PPAR- δ agonist GW0742 inhibited LPS-induced expression of COX-2 and

iNOS [32, 35]. Recently, Barish et al. found that GW501516 in mouse macrophages suppressed the gene induction of MCP-1, -3, -5 by IL-1, interferon- γ (IFN- γ), and phorbol ester. The agonist treatment also inhibited transendothelial migration of THP-1 cells [61]. The anti-inflammatory effects of the agonist was lost in the receptor-deficient macrophages [60]. However, in other cell types such as epithelial cells, eosinophils, neutrophils, and lymphocytes, the PPAR- δ agonist was ineffective in inhibiting inflammatory processes, indicating that the effect is cell-type-specific [62].

2.9. Atherosclerosis

To date, several studies have been reported regarding the roles of PPAR- δ in atherosclerosis in different mouse models with different approaches. Lee et al. transplanted PPAR- δ -null bone marrow progenitor cells into LDL receptor-null (LDLR $^{-/-}$) mice. Unexpectedly, the adoptive transfer of PPAR- δ -null macrophages led to a less severe atherosclerosis, suggesting that endogenous PPAR- δ maybe proatherogenic. Although overexpression or deletion of PPAR- δ in macrophages suggested that PPAR- δ is proinflammatory, the agonist GW501516 decreased MCP-1, seemingly having an opposite effect [60]. To reconcile this contradiction, they postulated an unconventional ligand-dependent transcriptional mechanism, which switches PPAR- δ between a “proinflammatory” and “anti-inflammatory”: in the absence of ligand, PPAR- δ sequesters a transcriptional repressor of inflammatory responses such as Bcl-6, permitting induction of proinflammatory genes; in the presence of ligand, PPAR- δ releases the repressor, which is then free to exert its anti-inflammatory effects. Following this loss-of-function approach, two independent studies examined the effect of the PPAR- δ agonist GW0742 on atherogenesis in high fat and cholesterol-fed LDLR $^{-/-}$ mice and yielded divergent results. In the first study, Li et al. found that GW7842 decreased gene expression of proinflammatory cytokines and adhesion molecules within atherosclerotic lesions but failed to alter the progression of atherosclerosis after 14 weeks of treatment (5 mg $^{-1}$ kg $^{-1}$ day $^{-1}$). In another one, Graham et al. used female LDLR $^{-/-}$ mice fed with a diet that induced moderate levels of hypercholesterolemia and observed that GW0742 reduced the lesion size at a higher dose (60 mg $^{-1}$ kg $^{-1}$ day $^{-1}$) after 10 weeks of treatment [37]. Discrepancy between these two studies may be caused by differences in the levels of hypercholesterolemia and different drug doses used. However, the anti-inflammatory effect was generally consistent in both studies, regardless the different effects on the lesion sizes. It is likely that the anti-inflammatory properties of the PPAR- δ agonists on the vessel wall per se are not sufficient to attenuate the progression of atherosclerotic lesions if it is not corroborated by an efficient improvement of metabolic abnormalities. This notion is supported by the data from recently published results. Most recently, GW501516, which has a potent lipid-modifying capacity, has also been demonstrated to have a clear antiatherosclerotic property in apoE $^{-/-}$ mice. Barish et al. showed that GW501516 significantly reduced atherosclerotic lesions with an increase in HDL level and

a reduced expression of chemokines in the aorta and in macrophages [61]. Furthermore, in a model of angiotensin II-accelerated atherosclerosis (LDLR^{-/-} mice), Takata et al. confirmed the atheroprotective effect of GW0742 [63]. After 4 weeks of treatment, GW0742 at both doses (1 and 10 mg⁻¹ kg⁻¹ day⁻¹) significantly inhibited the Ang II induction of atherosclerosis without altering blood pressure. This beneficial effect was likely mediated via the potent anti-inflammatory property since GW0742 increased vascular expression of Bcl-6, the regulators of G protein-coupled signaling (RGS4 and 5) in the artery and suppressed Ang II-induced activation of p38 and ERK in macrophages. However, the metabolic effect of GW0742 may also have contributed to the atheroprotective outcome because GW0742 significantly reduced plasma levels of insulin, glucose, leptin, and decreased triglycerides [63]. Overall, studies in mouse models suggest that PPAR- δ may have an attractive therapeutic target for the treatment of atherosclerosis.

2.10. Cardiovascular risk factors

In addition to the direct effects on the vessel wall, PPAR- δ also has profound effects on various metabolic parameters associated with cardiovascular diseases such as obesity, dyslipidemia, and insulin resistance.

2.11. Obesity

PPAR- δ deficiency causes embryonic lethality due to a placental defect. Some surviving PPAR- δ null mice had reduced fat mass [3], indicating a role of PPAR- δ in adipogenesis. Transgenic mice specifically expressing VP16-PPAR- δ , a constitutively active receptor, in adipose tissue had a reduced body weight, fat mass, and lower levels of circulating free fatty acids and triglycerides [64]. These animals were less susceptible to high-fat diet-induced obesity. In contrast, PPAR- δ null mice were more prone to weight gain on a high-fat diet. Similarly, GW501516 ameliorated diet-induced obesity [65]. It has been known that PPAR- δ activates genes involved in fatty acid oxidation and energy dissipation, such as carnitine palmitoyltransferase 1 (CPT1), acyl-CoA oxidase (AOX), and long chain acyl-CoA dehydrogenase (LCAD), uncoupling proteins. In addition to the direct effects on obesity, PPAR- δ agonists may also have regulatory effects on adipokine profile. For example, administration of L-165041 in rats increased the expression of visfatin and adiponectin, which are known to improve insulin sensitivity and are vasoprotective, but decreased the production of resistin in visceral adipose tissue [66].

2.12. Dyslipidemia

Increased levels of LDL and triglycerides and decreased HDL in plasma are independent risk factors for atherosclerosis and associated with metabolic syndrome as well. Recent studies have demonstrated that activation of PPAR- δ may modify lipid profile in animal models as well as in human. Oliver Jr. et al. first reported that GW501516 significantly improved dyslipidemia in obese primates with an increase

in HDL and a decrease in LDL cholesterol and triglycerides [58]. The beneficial effect of GW0742 and L-165041 on HDL level was also observed in obese and nonobese mice [67, 68]. In addition to enhancing fatty acid oxidation in muscles, PPAR- δ agonists upregulated expression of ABCA1 in several types of cells, which may lead to an increase in HDL and cholesterol reverse transport. In intestinal cells, it also inhibited gene expression of Niemann-Pick C1-like 1, the key molecule for cholesterol absorption [67]. However, the precise mechanisms underlying these lipid-modifying effects still remain to be elucidated. In a small number of healthy human volunteers, GW501516 has been reported to increase HDL cholesterol level and improved the triglycerides clearance [69].

2.13. Insulin resistance and glucose homeostasis

It has been previously known that, in obese primates, GW501516 declined fasting insulin level [58]. GW501516 treated *ob/ob* mice also showed a significantly improved glucose tolerance and with a lower postprandial levels of plasma glucose and insulin [65]. In cultured myotubes, PPAR- δ agonists were found to directly stimulate glucose uptake independent of insulin action. The agonist-stimulated glucose uptake in myotubes appeared to require AMP-activated protein kinase (AMPK) but not PPAR- δ [70, 71]. However, GW501516 had no acute effect on glucose transport in rat skeletal muscles [72]. Recently, Lee et al. showed that PPAR- δ ^{-/-} mice were glucose intolerant. Euglycemic hyperinsulinemic clamp experiments showed that GW501516 improved insulin sensitivity in multiple tissues including hepatic and peripheral tissues. The agonist suppresses hepatic glucose output and increases glucose disposal [73]. Gene array analysis suggested that PPAR- δ might ameliorate hyperglycemia by increasing glucose flux through the pentose phosphate pathway, which is known to enhance de novo fatty acid synthesis. Thus, it could be a concern whether PPAR- δ improves hyperglycemia at the cost of exacerbation of hepatosteatosis, a problem commonly associated with metabolic syndrome and diabetes. However, recent studies demonstrated that PPAR- δ also had a beneficial effect on hepatosteatosis. In a diet-induced mouse model of nonalcoholic steatohepatitis, GW501516 reduced triglycerides accumulation in the livers. In a most recent study, it was demonstrated that PPAR- δ suppressed hepatic lipogenesis via the induction of insulin-induced gene-1 (*insig-1*) and the inhibition of lipogenic sterol-regulatory element binding protein-1 (SREBP-1) activation. In obese diabetic mice, hepatic overexpression of PPAR- δ ameliorated hepatosteatosis [74]. Since *Insig-1* is a critical regulator of lipid homeostasis, identification of the *insig-1* as a target gene of PPAR- δ may also facilitate our understanding of the profound effects of PPAR- δ activation on adipogenesis and lipid metabolism [75, 76]. Recently, in a double-blind and randomized study, PPAR- δ agonist (10 mg o.d. GW501516) was given to a small number of healthy overweight subjects. The results showed that treatment with GW501516 for 2 weeks significantly reduced liver fat content by 20% without increasing oxidative stress [77].




FIGURE 1: Activation of PPAR- δ may have profound effects on vascular homeostasis and coronary artery diseases. These include both the direct actions in the vessel wall and the indirect effects on multiple cardiovascular risk factors. PPAR- δ gene polymorphisms are also linked to cardiovascular diseases.

2.14. Gene polymorphisms

Skogsberg et al. initially described 4 polymorphisms: -409C/T (in the promoter region), +73C/T (exon 1), +255A/G (exon 3), and +294T/C (exon 4). The +294T/C polymorphism showed a significant association with a metabolic trait. The homozygotes for the C allele had a higher plasma LDL and a tendency toward higher risk of CHD compared with homozygous carriers of the T-allele [78, 79]. In addition, there is a highly significant association between the rare C allele and lower plasma HDL concentrations in the female patients with mixed hyperlipidemia. Associations were also found for the C-allele with coronary heart disease and body mass index (BMI) [80, 81]. Chen et al. also demonstrated that, in “lipoprotein and coronary atherosclerosis study” (LCAS) subjects, the PPAR- δ SNPs were strongly associated with the dyslipidemia, the responses to statin, and the atherosclerotic lesions [82]. Vantinen et al. investigated the effects of the PPAR- δ gene SNPs on tissue glucose uptake and suggested that the SNPs regulate insulin sensitivity primarily in skeletal muscles [83]. The PPAR- δ SNPs were genotyped in type II diabetes subjects and normal control. Although no significant association was detected with the risk of type II diabetes, several SNPs were

associated with fasting plasma glucose and BMI [84]. In addition, the association has been found between the PPAR- δ SNPs and metabolic syndrome, and the association was influenced by dietary fat intake [85].

3. CONCLUSIONS

During the last few years, rapid progress has been made with regards to the roles of PPAR- δ in vascular biology. Emerging evidence supports the notion that activation of PPAR- δ may have profound effects on vascular homeostasis and coronary artery diseases. These include both the direct actions in the vessel wall and the beneficial effects on central metabolic pathways (Figure 1). Importantly, PPAR- δ agonists have an unprecedented role in raising HDL level in animals. In addition to a previously reported function of PPAR- δ in increasing oxidative muscle fibers and running endurance, a most recent study revealed that AMPK and PPAR- δ pathways have synergistic effects in terms of exercise-enhancing capacity [86]. The outcome of currently ongoing clinical trial is awaited to prove its clinical efficacy in the treatment of dyslipidemia. Despite the studies in rodent models point to a vascular-protective effect for PPAR- δ agonists, their efficacies in human coronary

artery diseases remain to be clarified. With regards to the effects of PPAR- δ on tumor angiogenesis and the unsettled role in carcinogenesis, safety issues also call for attention [87, 88]. Clearly, further studies are warranted to explore the roles of PPAR- δ in cardiovascular pathophysiology and to exploit this lipid-sensing receptor as a therapeutic target for metabolic syndrome and its cardiovascular complications.

ACKNOWLEDGMENT

Grant support was received from National Science Foundation of China (30890041, 30821001, and 30670848).

REFERENCES

- [1] B. Desvergne and W. Wahli, "Peroxisome proliferator-activated receptors: nuclear control of metabolism," *Endocrine Reviews*, vol. 20, no. 5, pp. 649–688, 1999.
- [2] J. M. Lehmann, L. B. Moore, T. A. Smith-Oliver, W. O. Wilkison, T. M. Willson, and S. A. Kliewer, "An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor γ (PPAR γ)," *The Journal of Biological Chemistry*, vol. 270, no. 22, pp. 12953–12956, 1995.
- [3] Y. Barak, D. Liao, W. He, et al., "Effects of peroxisome proliferator-activated receptor δ on placentation, adiposity, and colorectal cancer," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99, no. 1, pp. 303–308, 2002.
- [4] J. M. Peters, S. S. T. Lee, W. Li, et al., "Growths, adipose, brain, and skin alterations resulting from targeted disruption of the mouse peroxisome proliferator-activated receptor β/δ ," *Molecular and Cellular Biology*, vol. 20, no. 14, pp. 5119–5128, 2000.
- [5] N. S. Tan, L. Michalik, N. Di-Poi, B. Desvergne, and W. Wahli, "Critical roles of the nuclear receptor PPAR β (peroxisome-proliferator-activated receptor β) in skin wound healing," *Biochemical Society Transactions*, vol. 32, no. 1, pp. 97–102, 2004.
- [6] N. S. Tan, L. Michalik, N. Noy, et al., "Critical roles of PPAR β/δ in keratinocyte response to inflammation," *Genes & Development*, vol. 15, no. 24, pp. 3263–3277, 2001.
- [7] L. Michalik, B. Desvergne, N. S. Tan, et al., "Impaired skin wound healing in peroxisome proliferator-activated receptor (PPAR) α and PPAR β mutant mice," *Journal of Cell Biology*, vol. 154, no. 4, pp. 799–814, 2001.
- [8] C. Dreyer, G. Krey, H. Keller, F. Givel, G. Helftenbein, and W. Wahli, "Control of the peroxisomal β -oxidation pathway by a novel family of nuclear hormone receptors," *Cell*, vol. 68, no. 5, pp. 879–887, 1992.
- [9] A. Schmidt, R. L. Vogel, K. M. Witherup, et al., "Identification of fatty acid methyl ester as naturally occurring transcriptional regulators of the members of the peroxisome proliferator-activated receptor family," *Lipids*, vol. 31, no. 11, pp. 1115–1124, 1996.
- [10] T. Yoshikawa, Z. Brkanac, B. R. Dupont, G.-Q. Xing, R. J. Leach, and S. D. Detera-Wadleigh, "Assignment of the human nuclear hormone receptor, NUC1 (PPARD), to chromosome 6p21.1-p21.2," *Genomics*, vol. 35, no. 3, pp. 637–638, 1996.
- [11] S. A. Fyffe, M. S. Alphey, L. Buetow, et al., "Recombinant human PPAR- β/δ ligand-binding domain is locked in an activated conformation by endogenous fatty acids," *Journal of Molecular Biology*, vol. 356, no. 4, pp. 1005–1013, 2006.
- [12] H. E. Xu, M. H. Lambert, V. G. Montana, et al., "Molecular recognition of fatty acids by peroxisome proliferator-activated receptors," *Molecular Cell*, vol. 3, no. 3, pp. 397–403, 1999.
- [13] B. M. Forman, J. Chen, and R. M. Evans, "Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors α and δ ," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 94, no. 9, pp. 4312–4317, 1997.
- [14] K. Yu, W. Bayona, C. B. Kallen, et al., "Differential activation of peroxisome proliferator-activated receptors by eicosanoids," *The Journal of Biological Chemistry*, vol. 270, no. 41, pp. 23975–23983, 1995.
- [15] G. Krey, O. Braissant, F. L'Horsset, et al., "Fatty acids, eicosanoids, and hypolipidemic agents identified as ligands of peroxisome proliferator-activated receptors by coactivator-dependent receptor ligand assay," *Molecular Endocrinology*, vol. 11, no. 6, pp. 779–791, 1997.
- [16] R. A. Gupta, J. Tan, W. F. Krause, et al., "Prostacyclin-mediated activation of peroxisome proliferator-activated receptor δ in colorectal cancer," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 24, pp. 13275–13280, 2000.
- [17] A. Chawla, C.-H. Lee, Y. Barak, et al., "PPAR δ is a very low-density lipoprotein sensor in macrophages," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 3, pp. 1268–1273, 2003.
- [18] N. Shaw, M. Elholm, and N. Noy, "Retinoic acid is a high affinity selective ligand for the peroxisome proliferator-activated receptor β/δ ," *The Journal of Biological Chemistry*, vol. 278, no. 43, pp. 41589–41592, 2003.
- [19] M. L. Sznajdman, C. D. Haffner, P. R. Maloney, et al., "Novel selective small molecule agonists for peroxisome proliferator-activated receptor δ (PPAR δ)—synthesis and biological activity," *Bioorganic & Medicinal Chemistry Letters*, vol. 13, no. 9, pp. 1517–1521, 2003.
- [20] C. Santini, G. D. Berger, W. Han, et al., "Phenylacetic acid derivatives as hPPAR agonists," *Bioorganic & Medicinal Chemistry Letters*, vol. 13, no. 7, pp. 1277–1280, 2003.
- [21] B. G. Shearer, D. J. Steger, J. M. Way, et al., "Identification and characterization of a selective peroxisome proliferator-activated receptor β/δ (NR1C2) antagonist," *Molecular Endocrinology*, vol. 22, no. 2, pp. 523–529, 2008.
- [22] L. Piqueras, A. R. Reynolds, K. M. Hodivala-Dilke, et al., "Activation of PPAR β/δ induces endothelial cell proliferation and angiogenesis," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 27, no. 1, pp. 63–69, 2007.
- [23] S. Dimmeler, J. Haendeler, and A. M. Zeiher, "Regulation of endothelial cell apoptosis in atherothrombosis," *Current Opinion in Lipidology*, vol. 13, no. 5, pp. 531–536, 2002.
- [24] N. Wang and M. B. Stemerman, "Ref-1 and transcriptional control of endothelial apoptosis," *Circulation Research*, vol. 88, no. 12, pp. 1223–1225, 2001.
- [25] E. Letavernier, J. Perez, E. Joye, et al., "Peroxisome proliferator-activated receptor β/δ exerts a strong protection from ischemic acute renal failure," *Journal of the American Society of Nephrology*, vol. 16, no. 8, pp. 2395–2402, 2005.
- [26] J.-Y. Liou, S. Lee, D. Ghelani, N. Matijevic-Aleksic, and K. K. Wu, "Protection of endothelial survival by peroxisome proliferator-activated receptor- δ mediated 14-3-3 upregulation," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 26, no. 7, pp. 1481–1487, 2006.
- [27] L. Brunelli, K. A. Cieslik, J. L. Alcorn, M. Vatta, and A. Baldini, "Peroxisome proliferator-activated receptor- δ upregulates 14-3-3 ϵ in human endothelial cells via CCAAT/enhancer binding

- protein- β ," *Circulation Research*, vol. 100, no. 5, pp. e59–e71, 2007.
- [28] T. M. Carlos and J. M. Harlan, "Leukocyte-endothelial adhesion molecules," *Blood*, vol. 84, no. 7, pp. 2068–2101, 1994.
- [29] N. Marx, G. K. Sukhova, T. Collins, P. Libby, and J. Plutzky, "PPAR α activators inhibit cytokine-induced vascular cell adhesion molecule-1 expression in human endothelial cells," *Circulation*, vol. 99, no. 24, pp. 3125–3131, 1999.
- [30] K. Goya, S. Sumitani, X. Xu, et al., "Peroxisome proliferator-activated receptor α agonists increase nitric oxide synthase expression in vascular endothelial cells," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 24, no. 4, pp. 658–663, 2004.
- [31] N. Wang, L. Verna, N.-G. Chen, et al., "Constitutive activation of peroxisome proliferator-activated receptor- γ suppresses pro-inflammatory adhesion molecules in human vascular endothelial cells," *The Journal of Biological Chemistry*, vol. 277, no. 37, pp. 34176–34181, 2002.
- [32] V. Pasceri, H. D. Wu, J. T. Willerson, and E. T. H. Yeh, "Modulation of vascular inflammation in vitro and in vivo by peroxisome proliferator-activated receptor- γ activators," *Circulation*, vol. 101, no. 3, pp. 235–238, 2000.
- [33] H. Lee, W. Shi, P. Tontonoz, et al., "Role for peroxisome proliferator-activated receptor α in oxidized phospholipid-induced synthesis of monocyte chemotactic protein-1 interleukin-8 by endothelial cells," *Circulation Research*, vol. 87, no. 6, pp. 516–521, 2000.
- [34] N. Marx, T. Bourcier, G. K. Sukhova, P. Libby, and J. Plutzky, "PPAR γ activation in human endothelial cells increases plasminogen activator inhibitor type-1 expression: PPAR γ as a potential mediator in vascular disease," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 19, no. 3, pp. 546–551, 1999.
- [35] J. S. Welch, M. Ricote, T. E. Akiyama, F. J. Gonzalez, and C. K. Glass, "PPAR γ and PPAR δ negatively regulate specific subsets of lipopolysaccharide and IFN- γ target genes in macrophages," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 11, pp. 6712–6717, 2003.
- [36] A. C. Li, C. J. Binder, A. Gutierrez, et al., "Differential inhibition of macrophage foam-cell formation and atherosclerosis in mice by PPAR α , β/δ , and γ ," *Journal of Clinical Investigation*, vol. 114, no. 11, pp. 1564–1576, 2004.
- [37] T. L. Graham, C. Mookherjee, K. E. Suckling, C. N. A. Palmer, and L. Patel, "The PPAR δ agonist GW0742X reduces atherosclerosis in LDLR $^{-/-}$ mice," *Atherosclerosis*, vol. 181, no. 1, pp. 29–37, 2005.
- [38] Y. Rival, N. Benéteau, T. Taillandier, et al., "PPAR α and PPAR δ activators inhibit cytokine-induced nuclear translocation of NF- κ B and expression of VCAM-1 in EAhy926 endothelial cells," *European Journal of Pharmacology*, vol. 435, no. 2–3, pp. 143–151, 2002.
- [39] Y. Fan, Y. Wang, Z. Tang, et al., "Suppression of pro-inflammatory adhesion molecules by PPAR- δ in human vascular endothelial cells," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 28, no. 2, pp. 315–321, 2008.
- [40] M. Ghosh, H. Wang, Y. Ai, et al., "COX-2 suppresses tissue factor expression via endocannabinoid-directed PPAR δ activation," *The Journal of Experimental Medicine*, vol. 204, no. 9, pp. 2053–2061, 2007.
- [41] R. S. Bresalier, R. S. Sandler, H. Quan, et al., "Cardiovascular events associated with rofecoxib in a colorectal adenoma chemoprevention trial," *The New England Journal of Medicine*, vol. 352, no. 11, pp. 1092–1102, 2005.
- [42] J. Folkman, "Angiogenesis," *Annual Review of Medicine*, vol. 57, pp. 1–18, 2006.
- [43] P. Fraisl, M. Baes, and P. Carmeliet, "Hungry for blood vessels: linking metabolism and angiogenesis," *Developmental Cell*, vol. 14, no. 3, pp. 313–314, 2008.
- [44] P. Carmeliet and M. Baes, "Metabolism and therapeutic angiogenesis," *The New England Journal of Medicine*, vol. 358, no. 23, pp. 2511–2512, 2008.
- [45] R. L. Stephen, M. C. U. Gustafsson, M. Jarvis, et al., "Activation of peroxisome proliferator-activated receptor δ stimulates the proliferation of human breast and prostate cancer cell lines," *Cancer Research*, vol. 64, no. 9, pp. 3162–3170, 2004.
- [46] C. Gaudel, C. Schwartz, C. Giordano, N. A. Abumrad, and P. A. Grimaldi, "Pharmacological activation of PPAR β promotes rapid and calcineur-independent fiber remodeling and angiogenesis in mouse skeletal muscle," *American Journal of Physiology*, vol. 295, no. 2, pp. E297–E304, 2008.
- [47] A. Lutun, G. Carmeliet, and P. Carmeliet, "Vascular progenitors: from biology to treatment," *Trends in Cardiovascular Medicine*, vol. 12, no. 2, pp. 88–96, 2002.
- [48] T. Asahara, T. Murohara, A. Sullivan, et al., "Isolation of putative progenitor endothelial cells for angiogenesis," *Science*, vol. 275, no. 5302, pp. 964–967, 1997.
- [49] T. He, T. Lu, L. V. d'Uscio, C. F. Lam, H. C. Lee, and Z. S. Katusic, "Angiogenic function of prostacyclin biosynthesis in human endothelial progenitor cells," *Circulation Research*, vol. 103, no. 1, pp. 80–88, 2008.
- [50] A. Abdollahi, C. Schwager, J. Kleeff, et al., "Transcriptional network governing the angiogenic switch in human pancreatic cancer," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 31, pp. 12890–12895, 2007.
- [51] S. Müller-Brüsselbach, M. Kömhoff, M. Rieck, et al., "Deregulation of tumor angiogenesis and blockade of tumor growth in PPAR β -deficient mice," *The EMBO Journal*, vol. 26, no. 15, pp. 3686–3698, 2007.
- [52] J. Zhang, M. Fu, X. Zhu, et al., "Peroxisome proliferator-activated receptor δ is up-regulated during vascular lesion formation and promotes post-confluent cell proliferation in vascular smooth muscle cells," *The Journal of Biological Chemistry*, vol. 277, no. 13, pp. 11505–11512, 2002.
- [53] H.-J. Lim, S. Lee, J.-H. Park, et al., "PPAR δ agonist L-165041 inhibits rat vascular smooth muscle cell proliferation and migration via inhibition of cell cycle," *Atherosclerosis*. In press.
- [54] H. J. Kim, S. A. Ham, S. U. Kim, et al., "Transforming growth factor- β 1 is a molecular target for the peroxisome proliferator-activated receptor δ ," *Circulation Research*, vol. 102, no. 2, pp. 193–200, 2008.
- [55] Y. Numaguchi, K. Okumura, M. Harada, et al., "Catheter-based prostacyclin synthase gene transfer prevents in-stent restenosis in rabbit atheromatous arteries," *Cardiovascular Research*, vol. 61, no. 1, pp. 177–185, 2004.
- [56] H. Imai, Y. Numaguchi, M. Ishii, et al., "Prostacyclin synthase gene transfer inhibits neointimal formation by suppressing PPAR δ expression," *Atherosclerosis*, vol. 195, no. 2, pp. 322–332, 2007.
- [57] M. Yamada, Y. Numaguchi, K. Okumura, et al., "Prostacyclin synthase gene transfer modulates cyclooxygenase-2-derived prostanoid synthesis and inhibits neointimal formation in rat balloon-injured arteries," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 22, no. 2, pp. 256–262, 2002.
- [58] W. R. Oliver Jr., J. L. Shenk, M. R. Snaith, et al., "A selective peroxisome proliferator-activated receptor δ agonist promotes reverse cholesterol transport," *Proceedings of the National*

- Academy of Sciences of the United States of America*, vol. 98, no. 9, pp. 5306–5311, 2001.
- [59] H. Vosper, L. Patel, T. L. Graham, et al., “The peroxisome proliferator-activated receptor δ promotes lipid accumulation in human macrophages,” *The Journal of Biological Chemistry*, vol. 276, no. 47, pp. 44258–44265, 2001.
- [60] C.-H. Lee, A. Chawla, N. Urbiztondo, D. Liao, W. A. Boisvert, and R. M. Evans, “Transcriptional repression of atherogenic inflammation: modulation by PPAR δ ,” *Science*, vol. 302, no. 5644, pp. 453–457, 2003.
- [61] G. D. Barish, A. R. Atkins, M. Downes, et al., “PPAR δ regulates multiple proinflammatory pathways to suppress atherosclerosis,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 11, pp. 4271–4276, 2008.
- [62] A. Trifilieff, A. Bench, M. Hanley, D. Bayley, E. Campbell, and P. Whittaker, “PPAR- α and - γ but not - δ agonists inhibit airway inflammation in a murine model of asthma: in vitro evidence for an NF- κ B-independent effect,” *British Journal of Pharmacology*, vol. 139, no. 1, pp. 163–171, 2003.
- [63] Y. Takata, J. Liu, F. Yin, et al., “PPAR δ -mediated antiinflammatory mechanisms inhibit angiotensin II-accelerated atherosclerosis,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 11, pp. 4277–4282, 2008.
- [64] Y.-X. Wang, C.-H. Lee, S. Tjep, et al., “Peroxisome-proliferator-activated receptor δ activates fat metabolism to prevent obesity,” *Cell*, vol. 113, no. 2, pp. 159–170, 2003.
- [65] T. Tanaka, J. Yamamoto, S. Iwasaki, et al., “Activation of peroxisome proliferator-activated receptor δ induces fatty acid β -oxidation in skeletal muscle and attenuates metabolic syndrome,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 26, pp. 15924–15929, 2003.
- [66] K. C. Choi, S. Y. Lee, H. J. Yoo, et al., “Effect of PPAR- δ agonist on the expression of visfatin, adiponectin, and resistin in rat adipose tissue and 3T3-L1 adipocytes,” *Biochemical and Biophysical Research Communications*, vol. 357, no. 1, pp. 62–67, 2007.
- [67] J. N. van der Veen, J. K. Kruit, R. Havinga, et al., “Reduced cholesterol absorption upon PPAR δ activation coincides with decreased intestinal expression of NPC1L1,” *Journal of Lipid Research*, vol. 46, no. 3, pp. 526–534, 2005.
- [68] M. D. Leibowitz, C. Fiévet, N. Hennuyer, et al., “Activation of PPAR δ alters lipid metabolism in db/db mice,” *FEBS Letters*, vol. 473, no. 3, pp. 333–336, 2000.
- [69] D. L. Sprecher, C. Massien, G. Pearce, et al., “Triglyceride: high-density lipoprotein cholesterol effects in healthy subjects administered a peroxisome proliferator activated receptor δ agonist,” *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 27, no. 2, pp. 359–365, 2007.
- [70] D. K. Krämer, L. Al-Khalili, S. Perrini, et al., “Direct activation of glucose transport in primary human myotubes after activation of peroxisome proliferator-activated receptor δ ,” *Diabetes*, vol. 54, no. 4, pp. 1157–1163, 2005.
- [71] D. K. Krämer, L. Al-Khalili, B. Guigas, Y. Leng, P.M. Garcia-Roves, and A. Krook, “Role of AMP kinase and PPAR δ in the regulation of lipid and glucose metabolism in human skeletal muscle,” *The Journal of Biological Chemistry*, vol. 282, no. 27, pp. 19313–19320, 2007.
- [72] S. Terada, S. Wicke, J. O. Holloszy, and D.-H. Han, “PPAR δ activator GW-501516 has no acute effect on glucose transport in skeletal muscle,” *American Journal of Physiology*, vol. 290, no. 4, pp. E607–E611, 2006.
- [73] C.-H. Lee, P. Olson, A. Hevener, et al., “PPAR δ regulates glucose metabolism and insulin sensitivity,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 103, no. 9, pp. 3444–3449, 2006.
- [74] X. Qin, X. Xie, Y. Fan, et al., “Peroxisome proliferator-activated receptor- δ induces insulin-induced gene-1 and suppresses hepatic lipogenesis in obese diabetic mice,” *Hepatology*, vol. 48, no. 2, pp. 432–441, 2008.
- [75] L. J. Engelking, H. Kuriyama, R. E. Hammer, et al., “Over-expression of Insig-1 in the livers of transgenic mice inhibits SREBP processing and reduces insulin-stimulated lipogenesis,” *The Journal of Clinical Investigation*, vol. 113, no. 8, pp. 1168–1175, 2004.
- [76] D. Yabe, R. Komuro, G. Liang, J. L. Goldstein, and M. S. Brown, “Liver-specific mRNA for insig-2 down-regulated by insulin: implications for fatty acid synthesis,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 6, pp. 3155–3160, 2003.
- [77] U. Risérus, D. Sprecher, T. Johnson, et al., “Activation of peroxisome proliferator-activated receptor (PPAR) δ promotes reversal of multiple metabolic abnormalities, reduces oxidative stress, and increases fatty acid oxidation in moderately obese men,” *Diabetes*, vol. 57, no. 2, pp. 332–339, 2008.
- [78] J. Skogsberg, K. Kannisto, T. N. Cassel, A. Hamsten, P. Eriksson, and E. Ehrenborg, “Evidence that peroxisome proliferator-activated receptor delta influences cholesterol metabolism in men,” *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 23, no. 4, pp. 637–643, 2003.
- [79] J. Skogsberg, A. D. McMahon, F. Karpe, A. Hamsten, C. J. Packard, and E. Ehrenborg, “Peroxisome proliferator activated receptor delta genotype in relation to cardiovascular risk factors and risk of coronary heart disease in hypercholesterolaemic men,” *Journal of Internal Medicine*, vol. 254, no. 6, pp. 597–604, 2003.
- [80] J. Aberle, I. Hopfer, F. U. Beil, and U. Seedorf, “Association of the T+294C polymorphism in PPAR δ with low HDL cholesterol and coronary heart disease risk in women,” *International Journal of Medical Sciences*, vol. 3, no. 3, pp. 108–111, 2006.
- [81] J. Aberle, I. Hopfer, F. U. Beil, and U. Seedorf, “Association of peroxisome proliferator-activated receptor delta +294T/C with body mass index and interaction with peroxisome proliferator-activated receptor alpha L162V,” *International Journal of Obesity*, vol. 30, no. 12, pp. 1709–1713, 2006.
- [82] S. Chen, N. Tsybouleva, C. M. Ballantyne, A. M. Gotto Jr., and A. J. Marian, “Effects of PPAR α , γ and δ haplotypes on plasma levels of lipids, severity and progression of coronary atherosclerosis and response to statin therapy in the lipoprotein coronary atherosclerosis study,” *Pharmacogenetics*, vol. 14, no. 1, pp. 61–71, 2004.
- [83] M. Vääntinen, P. Nuutila, T. Kuulasmaa, et al., “Single nucleotide polymorphisms in the peroxisome proliferator-activated receptor δ gene are associated with skeletal muscle glucose uptake,” *Diabetes*, vol. 54, no. 12, pp. 3587–3591, 2005.
- [84] H. D. Shin, B. L. Park, L. H. Kim, et al., “Genetic polymorphisms in peroxisome proliferator-activated receptor δ associated with obesity,” *Diabetes*, vol. 53, no. 3, pp. 847–851, 2004.
- [85] J. Robitaille, D. Gaudet, L. Pérusse, and M.-C. Vohl, “Features of the metabolic syndrome are modulated by an interaction between the peroxisome proliferator-activated receptor-delta -87T>C polymorphism and dietary fat in French-Canadians,” *International Journal of Obesity*, vol. 31, no. 3, pp. 411–417, 2007.

-
- [86] V. A. Narkar, M. Downes, R. T. Yu, et al., “AMPK and PPAR δ agonists are exercise mimetics,” *Cell*, vol. 134, no. 3, pp. 405–415, 2008.
- [87] R. A. Gupta, D. Wang, S. Katkuri, H. Wang, S. K. Dey, and R. N. DuBois, “Activation of nuclear hormone receptor peroxisome proliferator-activated receptor- δ accelerates intestinal adenoma growth,” *Nature Medicine*, vol. 10, no. 3, pp. 245–247, 2004.
- [88] D. Wang, H. Wang, Y. Guo, et al., “Crosstalk between peroxisome proliferator-activated receptor δ and VEGF stimulates cancer progression,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 103, no. 50, pp. 19069–19074, 2006.