

Complete Genome of *Geobacter pickeringii* G13^T, a Metal-Reducing Isolate from Sedimentary Kaolin Deposits

Jonathan P. Badalamenti, Daniel R. Bond

Department of Microbiology and BioTechnology Institute, University of Minnesota, Saint Paul, Minnesota, USA

We used PacBio sequencing to assemble the genome of the pristine freshwater isolate *Geobacter pickeringii* G13^T into a single 3,618,700-bp circular chromosome polished to 99.999% accuracy (quality value [QV], 50). This isolate shares several features with other *Geobacter* spp., including genes for degradation of aromatics and an abundance of multiheme *c*-type cytochromes.

Received 12 January 2015 Accepted 16 January 2015 Published 5 March 2015

Citation Badalamenti JP, Bond DR. 2015. Complete genome of *Geobacter pickeringii* G13^T, a metal-reducing isolate from sedimentary kaolin deposits. *Genome Announc* 3(2): e00038-15. doi:10.1128/genomeA.00038-15.

Copyright © 2015 Badalamenti and Bond. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Daniel R. Bond, dbond@umn.edu.

Geobacter spp. play important roles in coupling organic matter degradation to metal reduction in sediment and subsurface environments. Genome sequencing efforts have primarily focused on isolates from aquatic sediments, subsurface bioremediation sites, and electricity-producing bioreactors (1–4). *Geobacter pickeringii*, a freshwater isolate recovered from a pristine, uncontaminated Fe(III)-rich kaolin deposit, is of interest for its wide repertoire of reported electron donors and acceptors used for anaerobic respiration, including the rare ability to utilize methanol (5).

Closed assemblies are available for 7 *Geobacter* isolates in GenBank, and these provide important structural information typically lacking in fragmented assemblies. For single-contig assembly of *G. pickeringii*, we leveraged Pacific Biosciences long-read sequencing to span repetitive elements without relying on reads from complementary platforms (6). Approximately 10 μg of genomic DNA from enzymatically lysed, stationary-phase *G. pickeringii* cells was prepared for continuous long reads using a 20-kb insert target and electrophoretically size selected with a 4-kb cutoff using Blue Pippin (Sage Science). Reads were captured as 120-min movies from 5 SMRT cells (P4-C2 chemistry) on a PacBio RS II sequencer (Mayo Clinic Bioinformatics Core, Rochester, MN).

Raw reads (~1 Gbp total) were filtered to remove SMRT bell adapters and short (<100 bp) and low quality reads (<80% accuracy) using SMRT Analysis v. 2.1 within the Amazon EC2 Elastic Compute Cloud. *De novo* assembly was performed with ~30× coverage of self-corrected long reads with a length cutoff of 10,183 bp as determined automatically with HGAP (7). Contig lengths from the first assembly were summed to estimate genome size and rerun HGAP, producing a single contig with self-overlapping ends. Overlaps were manually trimmed and the circularized contig was subjected to two successive rounds of polishing with Quiver using the entire read set (250× coverage) to remove remaining indels (7) and produce a single, 3,618,700-bp circular chromosome with uniform coverage, G+C content of 62.27%, and consensus concordance of 99.999% (quality value [QV], 50). The chromosome was manually reoriented to begin at the replication origin as predicted using Ori-Finder (8) before annotation via the NCBI Prokaryotic Annotation Pipeline.

Despite being isolated from an uncontaminated environment, *G. pickeringii* contains a full suite of genes for anaerobic oxidation of aromatic compounds and a putative *hgcAB* gene cluster (GPICK_04815—GPICK_04820) responsible for mercury methylation (9). *G. pickeringii* also contains a complete tricarboxylic acid (TCA) cycle, 60 predicted histidine kinases, and 56 multiheme *c*-type cytochromes, similar to *Geobacter sulfurreducens* (72) and *Geobacter metallireducens* (60). Long-read assembly successfully resolved 3 rRNA gene operons and a near perfect duplication of a 7.4-kb heterodisulfide reductase gene cluster. In support of the *G. pickeringii* methanol utilization phenotype (5), genes linked to methanol oxidation in *Rhodococcus erythropolis* N9T-4 (10) occurred as a cluster in *G. pickeringii* (GPICK_06655—GPICK_06695) upstream of a biosynthetic pathway predicted to synthesize an electron transfer cofactor (mycofactocin) used in methanol oxidation (11).

Nucleotide sequence accession numbers. The assembly and annotation have been deposited in GenBank under accession number CP009788. Raw PacBio reads and base modification data have been deposited to the NCBI Sequence Read Archive under accession number SRX796422.

ACKNOWLEDGMENTS

This work was supported by the Minnesota Environment and Natural Resources Trust Fund.

We thank Karl Oles (Mayo Clinic Bioinformatics Core) for performing library preparation and sequencing and Jason Smith and Scott Binford (Pacific Biosciences) for assistance with experimental design and data analysis.

REFERENCES

1. Aklujkar M, Krushkal J, DiBartolo G, Lapidus A, Land ML, Lovley DR. 2009. The genome sequence of *Geobacter metallireducens*: features of metabolism, physiology and regulation common and dissimilar to *Geobacter sulfurreducens*. *BMC Microbiol* 9:109. <http://dx.doi.org/10.1186/1471-2180-9-109>.
2. Aklujkar M, Young ND, Holmes D, Chavan M, Risso C, Kiss HE, Han CS, Land ML, Lovley DR. 2010. The genome of *Geobacter bemidjensis*, exemplar for the subsurface clade of *Geobacter* species that predominate in

- Fe(III)-reducing subsurface environments. *BMC Genomics* 11:490. <http://dx.doi.org/10.1186/1471-2164-11-490>.
3. Butler JE, Young ND, Lovley DR. 2010. Evolution of electron transfer out of the cell: comparative genomics of six *Geobacter* genomes. *BMC Genomics* 11:40. <http://dx.doi.org/10.1186/1471-2164-11-40>.
 4. Yi H, Nevin KP, Kim B-C, Franks AE, Klimes A, Tender LM, Lovley DR. 2009. Selection of a variant of *Geobacter sulfurreducens* with enhanced capacity for current production in microbial fuel cells. *Biosens Bioelectron* 24:3498–3503. <http://dx.doi.org/10.1016/j.bios.2009.05.004>.
 5. Shelobolina ES, Nevin KP, Blakeney-Hayward JD, Johnsen CV, Plaia TW, Krader P, Woodard T, Holmes DE, Vanpraagh CG, Lovley DR. 2007. *Geobacter pickeringii* sp. nov., *Geobacter argillaceus* sp. nov. and *Pelosiinus fermentans* gen. nov., sp. nov., isolated from subsurface kaolin lenses. *Int J Syst Evol Microbiol* 57:126–135. <http://dx.doi.org/10.1099/ijs.0.64221-0>.
 6. Koren S, Harhay GP, Smith TP, Bono JL, Harhay DM, Mcvey SD, Radune D, Bergman NH, Phillippy AM. 2013. Reducing assembly complexity of microbial genomes with single-molecule sequencing. *Genome Biol* 14:R101. <http://dx.doi.org/10.1186/gb-2013-14-9-r101>.
 7. Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.
 8. Gao F, Zhang C-T. 2008. Ori-finder: a Web-based system for finding *oriCs* in unannotated bacterial genomes. *BMC Bioinformatics* 9:79. <http://dx.doi.org/10.1186/1471-2105-9-79>.
 9. Parks JM, Johs A, Podar M, Bridou R, Hurt RA, Smith SD, Tomanicek SJ, Qian Y, Brown SD, Brandt CC, Palumbo AV, Smith JC, Wall JD, Elias DA, Liang L. 2013. The genetic basis for bacterial mercury methylation. *Science* 339:1332–1335. <http://dx.doi.org/10.1126/science.1230667>.
 10. Ohhata N, Yoshida N, Egami H, Katsuragi T, Tani Y, Takagi H. 2007. An extremely oligotrophic bacterium, *Rhodococcus erythropolis* N9T-4, isolated from crude oil. *J Bacteriol* 189:6824–6831. <http://dx.doi.org/10.1128/JB.00872-07>.
 11. Haft DH. 2011. Bioinformatic evidence for a widely distributed, ribosomally produced electron carrier precursor, its maturation proteins, and its nicotinoprotein redox partners. *BMC Genomics* 12:21. <http://dx.doi.org/10.1186/1471-2164-12-21>.