

## Review Article

# Sphingosine-1-Phosphate Transporters as Targets for Cancer Therapy

Masayuki Nagahashi,<sup>1</sup> Kazuaki Takabe,<sup>2</sup> Krista P. Terracina,<sup>2</sup> Daiki Soma,<sup>1</sup>  
Yuki Hirose,<sup>1</sup> Takashi Kobayashi,<sup>1</sup> Yasunobu Matsuda,<sup>3</sup> and Toshifumi Wakai<sup>1</sup>

<sup>1</sup> Division of Digestive and General Surgery, Niigata University Graduate School of Medical and Dental Sciences, 1-757 Asahimachi-dori, Chuo-Ku, Niigata 951-8510, Japan

<sup>2</sup> Division of Surgical Oncology, Department of Surgery, Virginia Commonwealth University School of Medicine and the Massey Cancer Center, 1200 E. Broad Street, Richmond, VA 23219, USA

<sup>3</sup> Department of Medical Technology, Niigata University Graduate School of Health Sciences, 2-746 Asahimachi-dori, Chuo-Ku, Niigata 951-8518, Japan

Correspondence should be addressed to Masayuki Nagahashi; [mnagahashi@med.niigata-u.ac.jp](mailto:mnagahashi@med.niigata-u.ac.jp)

Received 22 May 2014; Accepted 26 June 2014; Published 15 July 2014

Academic Editor: Yoshinori Marunaka

Copyright © 2014 Masayuki Nagahashi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Sphingosine-1-phosphate (S1P) is a pleiotropic lipid mediator that regulates cell survival, migration, the recruitment of immune cells, angiogenesis, and lymphangiogenesis, all of which are involved in cancer progression. S1P is generated inside cancer cells by sphingosine kinases then exported outside of the cell into the tumor microenvironment where it binds to any of five G protein coupled receptors and proceeds to regulate a variety of functions. We have recently reported on the mechanisms underlying the “inside-out” signaling of S1P, its export through the plasma membrane, and its interaction with cell surface receptors. Membrane lipids, including S1P, do not spontaneously exchange through lipid bilayers since the polar head groups do not readily go through the hydrophobic interior of the plasma membrane. Instead, specific transporter proteins exist on the membrane to exchange these lipids. This review summarizes what is known regarding S1P transport through the cell membrane via ATP-binding cassette transporters and the spinster 2 transporter and discusses the roles for these transporters in cancer and in the tumor microenvironment. Based on our research and the emerging understanding of the role of S1P signaling in cancer and in the tumor microenvironment, S1P transporters and S1P signaling hold promise as new therapeutic targets for cancer drug development.

## 1. Introduction

It is well recognized that the tumor microenvironment (TME) plays a key role in cancer progression and metastasis [1–3]. Tumors influence the surrounding microenvironment through the release of extracellular signals, such as cytokines, chemokines, and lipid mediators [4–7]. These bioactive molecules secreted from cancer cells and noncancer components in the TME, such as blood vessels, lymphatic vessels, and inflammatory cells, are considered to be potential therapeutic targets.

Sphingosine-1-phosphate (S1P) has emerged as a new player in the TME and cancer progression during the last decade [4]. S1P generated inside cells is exported outside of

cells into the TME where it stimulates specific S1P receptors on the cell surface (Figure 1). This “inside-out” signaling of S1P is considered to play a fundamental role in cancer progression [8]. To date, several S1P transporters have been identified [9–13]. Here, we review the “inside-out” signaling of S1P with a focus on S1P transporters. We will discuss the importance of S1P signaling in cancer and the possibility of targeting S1P transporters for cancer treatment.

## 2. “Inside-Out” Signaling of S1P

S1P is a pleiotropic lipid mediator that regulates cell survival, proliferation, migration, angiogenesis, lymphangiogenesis,




FIGURE 1: “Inside-out” signaling of sphingosine-1-phosphate (SIP). SIP is generated from Sph (sphingosine) by SphK1 (sphingosine kinase 1) in the cytosol of cancer cells and exported via ABC (ATP-binding cassette) transporters or Spns2 (spinster 2) outside of cells (tumor microenvironment). SIP stimulates specific S1P receptors (S1PR1-5) to promote numerous cellular functions, such as cell proliferation, migration, angiogenesis, and lymphangiogenesis.

and the recruitment of immune cells (Figure 1), making it involved in many physiological and pathological conditions including immune function, inflammation, and cancer [14–19]. SIP is generated from sphingosine inside the cells by two sphingosine kinases (SphK1 and SphK2) [20–23]. The balance between the levels of SIP and its metabolic precursors of ceramide and sphingosine has been regarded as a rheostat that could determine whether a cell proliferates or dies [24]. SIP generated within cells is exported out from cells via S1P transporters on the cell membrane, which will be described in detail later in this review. SIP outside of the cells can stimulate any of five specific G protein coupled receptors (S1PR1-5) [10, 12], with each S1P receptor coupled to different G proteins, which regulate activation or inhibition of the downstream intracellular signaling involved in various cellular functions (Figure 2).

Cancer cells and each type of cell in the TME, such as inflammatory cells and endothelial cells, express different combinations of S1P receptors, which contribute to each cellular function regulated by SIP. For example, S1PR1 is important for B and T lymphocyte egression from secondary lymphatic organs, such as lymph nodes [25]. In endothelial cells, S1PR1 and S1PR2 are known to play an important role in vascular development [26–29]. Stimulation of S1PR1 and/or S1PR3 often promotes cell proliferation and migration in normal and cancer cells, while S1PR2 may inhibit the

signaling that promotes cell proliferation and migration [30–32]. Altogether, this “inside-out” signaling of SIP plays a pivotal role in cancer cells and in the TME by stimulating the S1P receptors on each type of cell [33, 34].

In addition to its “inside-out” signaling, SIP is also known to have a variety of intracellular functions. Intracellular SIP produced by SphK1 can bind to TRAF2 (tumor necrosis factor receptor-associated factor 2) and function as a cofactor required for its E3 ubiquitin ligase activity and consequently, Lys-63-linked polyubiquitination of RIP1 (receptor-interacting protein 1) and NF- $\kappa$ B (nuclear factor kappa-light-chain-enhancer of activated B cells) activation [35]. Similarly, it has been shown that SIP enhances cIAP2 (cellular inhibitor of apoptosis 2) mediated K63-linked polyubiquitination of IRF-1 (interferon regulatory factor-1), which is essential for IL-1-induced production of chemokines CXCL10 and CCL5 [36]. Furthermore, SIP produced by SphK2 in the nucleus acts as an endogenous inhibitor of specific histone deacetylases (HDAC1 and HDAC2), thereby regulating gene transcription, including that of the cyclin dependent kinase inhibitor p21 [37].

The relative levels of SIP within body fluids and tissues are important to a variety of physiologic processes [25, 38]. Levels of SIP are maintained by SIP synthesis and degradation, which create an SIP gradient within the tissue [39]. SIP is dephosphorylated to regenerate sphingosine by S1P




FIGURE 2: Downstream signaling pathways of sphingosine-1-phosphate (S1P) receptors. S1P is a ligand for the five different specific G protein coupled receptors: S1PR1/EDG1; S1PR2/EDG5; S1PR3/EDG3; S1PR4/EDG6; and S1PR5/EDG8. Each S1P receptor is coupled to different G proteins, which regulate activation or inhibition of the indicated downstream signaling pathways, such as adenylyl cyclase-cyclic AMP, AC; extracellular signal-regulated kinase, ERK; phospholipase C, PLC; phosphatidylinositol 3-kinase, PI3K; the small GTPases of the Rho family; and Jun amino terminal kinase, JNK. Only a few examples of these pathways are illustrated.

phosphatases (SPPs) and/or lipid phosphate phosphatases (LPPs). S1P is also irreversibly degraded to hexadecenal and ethanolamine phosphate by S1P lyase (SPL). The current consensus is that trafficking of immune cells is controlled by this S1P gradient. For example, in the blood and lymph, S1P levels are relatively high, but in secondary lymphatic tissue, such as lymph nodes and the thymus, S1P is maintained at very low levels [40]. Importantly, any inhibition of S1P signaling, and therefore altering of this S1P gradient, results in alteration of immune cell trafficking. S1P also regulates vascular integrity. In the plasma, a decrease in the amount of S1P causes increased vascular permeability, likely due to loss of signaling through S1PR1 on endothelial cells [41–43]. S1P through its tissue gradients, intracellular functions, and “inside-out” signaling is important in both physiologic and pathologic processes.

### 3. Export of S1P via ATP-Binding Cassette (ABC) Transporters and Spinster 2 (Spns2) Transporter

Until recently, the process through which S1P produced inside cells by the two SphKs reaches its receptors on the cell surface remained obscure. Membrane lipids, including S1P, do not spontaneously exchange through the lipid bilayers of

the plasma membrane since the polar head groups do not readily go through the hydrophobic interior. Though there are many transporter proteins on the membrane for lipid exchange [9], sphingosine is known to spontaneously translocate without the aid of a transporter across intracellular membranes when added to cells or produced intracellularly [9].

Studies from several laboratories, including ours, have suggested the involvement of ABC transporters in the export of S1P from various types of cells *in vitro* [10]. S1P has been shown to be exported from mast cells via ABCC1 (also known as multidrug resistant protein 1; MRP1) [9], from astrocytes via ABCA1 [44], from endothelial cells via ABCA1 and ABCC1 [45], and from thyroid carcinoma cells via ABCC1 [46]. Using pharmacological and molecular approaches, we demonstrated that ABCC1 and ABCG2 (also known as breast cancer resistance protein; BCRP) are involved in estradiol-mediated transport of S1P and dihydro-S1P out of MCF-7 human breast cancer cells [12]. S1P is exported from erythrocytes and platelets by other transporters in the ABC transporter family [47, 48]. In erythrocytes, S1P is exported by an ATP-dependent and vanadate- and glyburide-sensitive transporter [47], while in platelets, S1P export requires an extracellular stimulus such as thrombin and is exported through two independent transporters, a  $Ca^{2+}$ -dependent transporter and an ATP-dependent glyburide-sensitive transporter [48, 49].

Collectively, these studies suggest that members of the large family of ABC transporters are responsible for export of SIP in various types of cells; however, in studies using mice with ABC transporter deficiencies, including animals with knockout of ABCA1, ABCA7, and ABCG1, SIP levels and related functions have been found to be unaltered [45], indicating the existence of compensatory mechanisms with other transporters.

Spns2, a member of the MFS (major facilitator superfamily) that does not have a typical ATP binding motif, has been recently discovered to export SIP from cells [11, 50–52]. Spns2 was identified independently by two groups, both of whom revealed that it transports SIP through observations in zebrafish. They showed that a mutation in Spns2 caused abnormal development resulting in *cardia bifida* (two hearts) [53] and that the phenotype of the Spns2 mutation is rescued by providing exogenous SIP [11]. The *cardia bifida* phenotype in the Spns2 knockout zebrafish was the clue that linked Spns2 to SIP, since the same phenotype was seen in S1PR2 knockout zebrafish [11, 53]. We have also shown that Spns2 can export endogenous SIP and dihydro-SIP from cells [13]. In addition, human Spns2 can transport several SIP analogues, including phosphorylated FTY720 [54]. Importantly, Spns2 was the first SIP transporter discovered to be physiologically functional *in vivo*, in contrast to the ABC transporters [54].

It has been suggested that Spns2 is important for vascular development [29]. We have observed that lymph nodes from Spns2 knockout mice have aberrant lymphatic sinuses that appear collapsed, with reduced numbers of lymphocytes [13]. Our data suggest that Spns2 is an SIP transporter *in vivo* that plays a role in regulation of levels of SIP not only in the blood, but also in the lymph nodes and lymphatic fluid, influencing lymphocyte trafficking and lymphatic vessel network organization [13]. The recent finding that blood endothelial cells purified from the aorta of Spns2-deficient mice are unable to release SIP [52] seems to support a strong role for Spns2 in SIP regulation. The levels of SIP in the plasma of Spns2 knockout mice have been observed to be decreased to 60% of that of wild type mice with endothelial cells contributing 40% of the total plasma SIP [54]. There was no difference between Spns2 knockout mice and wild type in SIP release activity in both erythrocytes and platelets, showing that a disruption of Spns2 does not affect the SIP release from erythrocytes or platelets [52]. Furthermore, bone marrow reconstitution studies revealed that Spns2 was not involved in SIP release from blood cells and suggested a role for Spns2 in other cells [51]. Consistent with these data, specific deletion of Spns2 on endothelial cells has been shown to result in a lack of lymphocyte egress, mimicking observations in global Spns2-knockout mice. These data suggest that Spns2 functions in endothelial cells, not blood cells, to establish the SIP gradient required for T and B cells to egress from their respective lymphoid organs [51].

#### 4. Targeting SIP Transporters for Treatment of Cancer Patients

There have been an increasing number of studies, implicating roles for SIP in different stages of cancer progression in

subtypes of both adult and pediatric malignancies. SphKs and SIP signaling have been suggested to also have a role in acquisition of drug resistance [55, 56]. There is a growing body of literature, with several clinical and pathological reports revealing the importance of SphK1 on cancer metastasis and prognosis [57–61]. Previous clinical studies have shown that SphK1 is overexpressed in human breast cancer and its expression correlates with poor patient outcomes [58, 62].

Studies linking the molecular interactions of SIP signaling with other oncogenic pathways, such as Ras, STAT3 (signal transducer and activator of transcription 3), NF- $\kappa$ B, and estrogen signaling, have been published. K-Ras mutations are known to increase the production of SIP in a SphK1-dependent manner and expression of the K-Ras oncogene leads to plasma membrane localization of SphK1. The RAS/RAF (rapidly accelerated fibrosarcoma)/MEK (mitogen-activated protein kinases)/ERK (extracellular signal-regulated kinases) pathway likely mediates this process, as constitutively active B-Raf or MEK are capable of activating SphK1 [63]. SIP produced by upregulated SphK1 in tumor cells activates S1PR1, which has been shown to lead to activation of STAT3 [34]. SIP is also involved in the activation of NF- $\kappa$ B, thereby regulating the transcription of the proinflammatory cytokines TNF- $\alpha$  (tumor necrosis factor-alpha) and IL-6 (interleukin-6) [34]. Estradiol is known to also stimulate SphK1 activation and the release of SIP, through which estradiol is capable of activating the S1PR3, resulting in EGFR transactivation in a matrix metalloprotease-dependent manner [31]. Altogether, these molecular interactions of SIP signaling with other oncogenic pathways suggest the importance of SIP signaling in cancer.

Communication among tumor cells, the host microenvironment, and inflammatory cells via systemic SIP regulates metastasis. SIP generated in cancer cells is secreted into tissue interstitial fluid in the body and affects the TME by altering immune cells, evoking inflammation, and inducing angiogenesis and lymphangiogenesis thereby promoting cancer metastasis [34, 64]. We have recently measured SIP levels within tumor interstitial fluid and found that significantly higher levels of SIP compared with the interstitial fluid of normal tissue and that inhibiting the “inside-out” signaling of SIP by FTY720 (known as fingolimod) significantly decreased SIP levels in the tumor interstitial fluid (unpublished data). Altogether, the “inside-out” signaling of SIP with overexpressed SphK1 and SIP transporters plays an important role in cancer progression through its effects on the TME.

It has recently been reported that a specific pharmacological inhibitor of SphK1 had no effect on cell proliferation *in vitro* [65]. This result generated some arguments that SIP, the product of SphK1, may not be an ideal anticancer target. On the other hand, targeting SIP signaling leads to a significant suppression of cancer progression *in vivo* [64], especially under conditions in which cancer is associated with inflammation [34]. Considering the importance of “inside-out” signaling of SIP within the TME through promotion of angiogenesis and lymphangiogenesis [4, 66, 67], the effects of targeting SIP signaling may only be adequately assessed *in vivo*.



ABC transporters were originally described as multidrug resistance genes and have been shown to be overexpressed in various solid and hematological cancers [68]. Expression of ABC transporters has been correlated with resistance to chemotherapies and poor prognosis for patients with certain types of cancer [68]. ABCB1, also known as multidrug resistant gene 1 (MDR1), has been targeted in a number of clinical trials that failed to demonstrate significant benefit [69]. Since it has been suggested that several ABC transporters are involved in secretion of S1P from stromal, endothelial, and cancer cells, it is possible that these transporters may play an important role in the pathological processes regulated by S1P and may worsen the biology of cancer cells. Considering that ABCB1 does not have a role for S1P transport, targeting ABCB1 and/or G2 may be a more promising treatment option for cancer patients. In contrast to the ABC transporters, with some of them originally described as multidrug resistant genes, the roles of Spns2 in cancer remain unknown.

Spns2, a 549 amino acids protein, belongs to the MFS transporter family as determined based on its predicted amino acid sequence [49]. The crystal structure of Spns2 and the precise mechanism of S1P transport via this transporter are still under investigation. S1P import by this transporter also remains to be described. To elucidate the role of Spns2 in cancer, we are currently in the midst of a process of developing gene targeting techniques utilizing nanoparticles to downregulate spns2 expression in cancer cells. How effective this may be in light of other transporters that contribute to S1P export, particularly *in vivo* setting, is yet to be determined. Further investigation is needed to clarify the function and role of Spns2 in normal physiological conditions and in the pathological condition of cancer.

The ABC transporter family and Spns2 are found in various types of cells and have different roles in each type of cell, so that it is possible that targeting a specific transporter may present some difficulties for cancer therapy for patients. In addition to targeting S1P transporters, a possible strategy for cancer treatment involves targeting S1P itself using a monoclonal antibody. In an animal model, neutralization of systemic S1P using Sphingomab, an anti-S1P monoclonal antibody, was shown to suppress lung metastasis [70]. The humanized version of Sphingomab, Sonpicipuzumab/ASONEP [71], has finished phase I trials and has recently entered phase II efficacy and safety studies for the treatment of renal cell carcinoma and age-related macular degeneration [72]. Targeting S1P itself may be one of the promising strategies for controlling cancer progression and metastasis.

## 5. Conclusion

S1P promotes a variety of intracellular and extracellular biological functions and is transported from inside cells through several members of the ABC transporter family and the enigmatic Spns2 transporter. Once through the plasma membrane, S1P exerts its “inside-out” signaling in the context of the TME and promotes cancer progression. Targeting S1P transporters and S1P signaling in cancer progression is a promising direction for development of the next generation of cancer therapeutics.

## Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

## Acknowledgment

This work was supported by NIH Grants (R01CA160688 to Kazuaki Takabe and T32CA085159-10 to Krista P. Terracina).

## References

- [1] D. Hanahan and L. M. Coussens, “Accessories to the crime: functions of cells recruited to the tumor microenvironment,” *Cancer Cell*, vol. 21, no. 3, pp. 309–322, 2012.
- [2] D. Hanahan and R. A. Weinberg, “Hallmarks of cancer: the next generation,” *Cell*, vol. 144, no. 5, pp. 646–674, 2011.
- [3] H. Fang and Y. A. DeClerck, “Targeting the tumor microenvironment: from understanding pathways to effective clinical trials,” *Cancer Research*, vol. 73, no. 16, pp. 4965–4977, 2013.
- [4] K. Takabe and S. Spiegel, “Export of sphingosine-1-phosphate and cancer progression,” *The Journal of Lipid Research*, 2014.
- [5] H. Wiig, O. Tenstad, P. O. Iversen, R. Kalluri, and R. Bjerkvig, “Interstitial fluid: the overlooked component of the tumor microenvironment?” *Fibrogenesis & Tissue Repair*, vol. 3, p. 12, 2010.
- [6] H. Haslene-Hox, E. Oveland, K. C. Berg et al., “A new method for isolation of interstitial fluid from human solid tumors applied to proteomic analysis of ovarian carcinoma tissue,” *PLoS ONE*, vol. 6, no. 4, Article ID e19217, 2011.
- [7] H. Wiig and M. A. Swartz, “Interstitial fluid and lymph formation and transport: physiological regulation and roles in inflammation and cancer,” *Physiological Reviews*, vol. 92, no. 3, pp. 1005–1060, 2012.
- [8] G. T. Kunkel, M. MacEyka, S. Milstien, and S. Spiegel, “Targeting the sphingosine-1-phosphate axis in cancer, inflammation and beyond,” *Nature Reviews Drug Discovery*, vol. 12, no. 9, pp. 688–702, 2013.
- [9] P. Mitra, C. A. Oskeritzian, S. G. Payne, M. A. Beaven, S. Milstien, and S. Spiegel, “Role of ABCB1 in export of sphingosine-1-phosphate from mast cells,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 103, no. 44, pp. 16394–16399, 2006.
- [10] R. H. Kim, K. Takabe, S. Milstien, and S. Spiegel, “Export and functions of sphingosine-1-phosphate,” *Biochimica et Biophysica Acta*, vol. 1791, no. 7, pp. 692–696, 2009.
- [11] A. Kawahara, T. Nishi, Y. Hisano, H. Fukui, A. Yamaguchi, and N. Mochizuki, “The sphingolipid transporter Spns2 functions in migration of zebrafish myocardial precursors,” *Science*, vol. 323, no. 5913, pp. 524–527, 2009.
- [12] K. Takabe, R. H. Kim, J. C. Allegood et al., “Estradiol induces export of sphingosine 1-phosphate from breast cancer cells via ABCB1 and ABCG2,” *The Journal of Biological Chemistry*, vol. 285, no. 14, pp. 10477–10486, 2010.
- [13] M. Nagahashi, E. Y. Kim, A. Yamada et al., “Spns2, a transporter of phosphorylated sphingoid bases, regulates their blood and lymph levels, and the lymphatic network,” *The FASEB Journal*, vol. 27, no. 3, pp. 1001–1011, 2013.
- [14] A. Olivera and S. Spiegel, “Sphingosine-1-phosphate as second messenger in cell proliferation induced by PDGF and FCS mitogens,” *Nature*, vol. 365, no. 6446, pp. 557–560, 1993.

- [15] K. Takabe, S. W. Paugh, S. Milstien, and S. Spiegel, "Inside-out signaling of sphingosine-1-phosphate: therapeutic targets," *Pharmacological Reviews*, vol. 60, no. 2, pp. 181–195, 2008.
- [16] D. Shida, K. Takabe, D. Kapitonov, S. Milstien, and S. Spiegel, "Targeting SphK1 as a new strategy against cancer," *Current Drug Targets*, vol. 9, no. 8, pp. 662–673, 2008.
- [17] N. J. Pyne and S. Pyne, "Sphingosine 1-phosphate and cancer," *Nature Reviews Cancer*, vol. 10, no. 7, pp. 489–503, 2010.
- [18] S. Spiegel and S. Milstien, "The outs and the ins of sphingosine-1-phosphate in immunity," *Nature Reviews Immunology*, vol. 11, no. 6, pp. 403–415, 2011.
- [19] W. C. Huang, M. Nagahashi, K. P. Terracina, and K. Takabe, "Emerging role of sphingosine-1-phosphate in inflammation, cancer, and lymphangiogenesis," *Biomolecules*, vol. 3, no. 3, pp. 408–434, 2013.
- [20] H. Liu, D. Chakravarty, M. Maceyka, S. Milstien, and S. Spiegel, "Sphingosine kinases: a novel family of lipid kinases," *Progress in Nucleic Acid Research and Molecular Biology*, vol. 71, pp. 493–511, 2002.
- [21] M. Maceyka, S. G. Payne, S. Milstien, and S. Spiegel, "Sphingosine kinase, sphingosine-1-phosphate, and apoptosis," *Biochimica et Biophysica Acta*, vol. 1585, no. 2-3, pp. 193–201, 2002.
- [22] V. E. Nava, J. P. Hobson, S. Murthy, S. Milstien, and S. Spiegel, "Sphingosine kinase type 1 promotes estrogen-dependent tumorigenesis of breast cancer MCF-7 cells," *Experimental Cell Research*, vol. 281, no. 1, pp. 115–127, 2002.
- [23] H. Liu, R. E. Toman, S. K. Goparaju et al., "Sphingosine kinase type 2 is a putative BH3-only protein that induces apoptosis," *Journal of Biological Chemistry*, vol. 278, no. 41, pp. 40330–40336, 2003.
- [24] O. Cu villier, G. Pirianov, B. Kleuser et al., "Suppression of ceramide-mediated programmed cell death by sphingosine-1-phosphate," *Nature*, vol. 381, no. 6585, pp. 800–803, 1996.
- [25] J. G. Cyster and S. R. Schwab, "Sphingosine-1-phosphate and lymphocyte egress from lymphoid organs," *Annual Review of Immunology*, vol. 30, pp. 69–94, 2012.
- [26] A. B. Shoham, G. Malkinson, S. Krief et al., "SIP<sub>1</sub> inhibits sprouting angiogenesis during vascular development," *Development*, vol. 139, no. 20, pp. 3859–3869, 2012.
- [27] K. Gaengel, C. Niaudet, K. Hagikura et al., "The sphingosine-1-phosphate receptor *s1pr1* restricts sprouting angiogenesis by regulating the interplay between VE-cadherin and VEGFR2," *Developmental Cell*, vol. 23, no. 3, pp. 587–599, 2012.
- [28] B. Jung, H. Obinata, S. Galvani et al., "Flow-regulated endothelial SIP receptor-1 signaling sustains vascular development," *Developmental Cell*, vol. 23, no. 3, pp. 600–610, 2012.
- [29] K. Mendelson, T. Zygmunt, J. Torres-Vázquez, T. Evans, and T. Hla, "Sphingosine 1-phosphate receptor signaling regulates proper embryonic vascular patterning," *The Journal of Biological Chemistry*, vol. 288, no. 4, pp. 2143–2156, 2013.
- [30] H. M. Rosenfeldt, J. P. Hobson, M. Maceyka et al., "EDG-1 links the PDGF receptor to Src and focal adhesion kinase activation leading to lamellipodia formation and cell migration," *The FASEB Journal*, vol. 15, no. 14, pp. 2649–2659, 2001.
- [31] O. Sukocheva, C. Wadham, A. Holmes et al., "Estrogen transactivates EGFR via the sphingosine 1-phosphate receptor Edg-3: the role of sphingosine kinase-1," *Journal of Cell Biology*, vol. 173, no. 2, pp. 301–310, 2006.
- [32] S. K. Goparaju, P. S. Jolly, K. R. Watterson et al., "The SIP2 receptor negatively regulates platelet-derived growth factor-induced motility and proliferation," *Molecular & Cellular Biology*, vol. 25, no. 10, pp. 4237–4249, 2005.
- [33] M. Nagahashi, S. Ramachandran, O. M. Rashid, and K. Takabe, "Lymphangiogenesis: a new player in cancer progression," *World Journal of Gastroenterology*, vol. 16, no. 32, pp. 4003–4012, 2010.
- [34] J. Liang, M. Nagahashi, E. Y. Kim et al., "Sphingosine-1-phosphate links persistent STAT3 activation, chronic intestinal inflammation, and development of colitis-associated cancer," *Cancer Cell*, vol. 23, no. 1, pp. 107–120, 2013.
- [35] S. E. Alvarez, K. B. Harikumar, N. C. Hait et al., "Sphingosine-1-phosphate is a missing cofactor for the E3 ubiquitin ligase TRAF2," *Nature*, vol. 465, no. 7301, pp. 1084–1088, 2010.
- [36] K. B. Harikumar, J. W. Yester, and M. J. Surace, "K63-linked polyubiquitination of transcription factor IRF1 is essential for IL-1-induced production of chemokines CXCL10 and CCL5," *Nature Immunology*, vol. 15, no. 3, pp. 231–238, 2014.
- [37] N. C. Hait, J. Allegood, M. Maceyka et al., "Regulation of histone acetylation in the nucleus by sphingosine-1-phosphate," *Science*, vol. 325, no. 5945, pp. 1254–1257, 2009.
- [38] S. R. Schwab and J. G. Cyster, "Finding a way out: Lymphocyte egress from lymphoid organs," *Nature Immunology*, vol. 8, no. 12, pp. 1295–1301, 2007.
- [39] M. Serra and J. D. Saba, "Sphingosine 1-phosphate lyase, a key regulator of sphingosine 1-phosphate signaling and function," *Advances in Enzyme Regulation*, vol. 50, no. 1, pp. 349–362, 2010.
- [40] S. R. Schwab, J. P. Pereira, M. Matloubian, Y. Xu, Y. Huang, and J. G. Cyster, "Immunology: lymphocyte sequestration through SIP lyase inhibition and disruption of SIP gradients," *Science*, vol. 309, no. 5741, pp. 1735–1739, 2005.
- [41] E. Camerer, J. B. Regard, I. Cornelissen et al., "Sphingosine-1-phosphate in the plasma compartment regulates basal and inflammation-induced vascular leak in mice," *Journal of Clinical Investigation*, vol. 119, no. 7, pp. 1871–1879, 2009.
- [42] M. J. Lee, S. Thangada, K. P. Claffey et al., "Vascular endothelial cell adherens junction assembly and morphogenesis induced by sphingosine-1-phosphate," *Cell*, vol. 99, no. 3, pp. 301–312, 1999.
- [43] A. Mendoza, B. Bréart, W. D. Ramos-Perez et al., "The transporter *Spns2* is required for secretion of lymph but not plasma sphingosine-1-phosphate," *Cell Reports*, vol. 2, no. 5, pp. 1104–1110, 2012.
- [44] K. Sato, E. Malchinkhuu, Y. Horiuchi et al., "Critical role of ABCA1 transporter in sphingosine 1-phosphate release from astrocytes," *Journal of Neurochemistry*, vol. 103, no. 6, pp. 2610–2619, 2007.
- [45] Y. M. Lee, K. Venkataraman, S. I. Hwang, D. K. Han, and T. Hla, "A novel method to quantify sphingosine 1-phosphate by immobilized metal affinity chromatography (IMAC)," *Prostaglandins and Other Lipid Mediators*, vol. 84, no. 3-4, pp. 154–162, 2007.
- [46] N. Bergelin, T. Blom, J. Heikkilä et al., "Sphingosine kinase as an oncogene: autocrine sphingosine 1-phosphate modulates ML-1 thyroid carcinoma cell migration by a mechanism dependent on protein kinase C- $\alpha$  and ERK1/2," *Endocrinology*, vol. 150, no. 5, pp. 2055–2063, 2009.
- [47] N. Kobayashi, A. Yamaguchi, and T. Nishi, "Characterization of the ATP-dependent sphingosine 1-phosphate transporter in rat erythrocytes," *The Journal of Biological Chemistry*, vol. 284, no. 32, pp. 21192–21200, 2009.
- [48] N. Kobayashi, T. Nishi, T. Hirata et al., "Sphingosine 1-phosphate is released from the cytosol of rat platelets in a carrier-mediated manner," *Journal of Lipid Research*, vol. 47, no. 3, pp. 614–621, 2006.

- [49] T. Nishi, N. Kobayashi, Y. Hisano, A. Kawahara, and A. Yamaguchi, "Molecular and physiological functions of sphingosine 1-phosphate transporters," *Biochimica et Biophysica Acta: Molecular and Cell Biology of Lipids*, vol. 1841, no. 5, pp. 759–765, 2013.
- [50] Y. Hisano, N. Kobayashi, A. Kawahara, A. Yamaguchi, and T. Nishi, "The sphingosine 1-phosphate transporter, SPNS2, functions as a transporter of the phosphorylated form of the immunomodulating agent FTY720," *The Journal of Biological Chemistry*, vol. 286, no. 3, pp. 1758–1766, 2011.
- [51] S. Fukuhara, S. Simmons, S. Kawamura et al., "The sphingosine-1-phosphate transporter Spns2 expressed on endothelial cells regulates lymphocyte trafficking in mice," *Journal of Clinical Investigation*, vol. 122, no. 4, pp. 1416–1426, 2012.
- [52] Y. Hisano, N. Kobayashi, A. Yamaguchi, and T. Nishi, "Mouse SPNS2 functions as a sphingosine-1-phosphate transporter in vascular endothelial cells," *PLoS ONE*, vol. 7, no. 6, Article ID e38941, 2012.
- [53] N. Osborne, K. Brand-Arzamendi, E. A. Ober et al., "The spinster homolog, two of hearts, is required for sphingosine 1-phosphate signaling in zebrafish," *Current Biology*, vol. 18, no. 23, pp. 1882–1888, 2008.
- [54] Y. Hisano, T. Nishi, and A. Kawahara, "The functional roles of SIP in immunity," *Journal of Biochemistry*, vol. 152, no. 4, pp. 305–311, 2012.
- [55] A. Salas, S. Ponnusamy, C. E. Senkal et al., "Sphingosine kinase-1 and sphingosine 1-phosphate receptor 2 mediate Bcr-Abl1 stability and drug resistance by modulation of protein phosphatase 2A," *Blood*, vol. 117, no. 22, pp. 5941–5952, 2011.
- [56] O. Cuvillier, I. Ader, P. Bouquerel, L. Brizuela, C. Gstalder, and B. Malavaud, "Hypoxia, therapeutic resistance, and sphingosine 1-phosphate," *Advances in Cancer Research*, vol. 117, pp. 117–141, 2013.
- [57] J. R. van Brooklyn, C. A. Jackson, D. K. Pearl, M. S. Kotur, P. J. Snyder, and T. W. Prior, "Sphingosine kinase-1 expression correlates with poor survival of patients with glioblastoma multiforme: roles of sphingosine kinase isoforms in growth of glioblastoma cell lines," *Journal of Neuropathology and Experimental Neurology*, vol. 64, no. 8, pp. 695–705, 2005.
- [58] E. Ruckhäberle, A. Rody, K. Engels et al., "Microarray analysis of altered sphingolipid metabolism reveals prognostic significance of sphingosine kinase 1 in breast cancer," *Breast Cancer Research and Treatment*, vol. 112, no. 1, pp. 41–52, 2008.
- [59] W. Li, C. P. Yu, J. T. Xia et al., "Sphingosine kinase 1 is associated with gastric cancer progression and poor survival of patients," *Clinical Cancer Research*, vol. 15, no. 4, pp. 1393–1399, 2009.
- [60] S. Q. Liu, Y. J. Su, M. B. Qin, Y. B. Mao, J. A. Huang, and G. D. Tang, "Sphingosine kinase 1 promotes tumor progression and confers malignancy phenotypes of colon cancer by regulating the focal adhesion kinase pathway and adhesion molecules," *International Journal of Oncology*, vol. 42, no. 2, pp. 617–626, 2013.
- [61] S. Pyne, J. Edwards, J. Ohotski, and N. J. Pyne, "Sphingosine 1-phosphate receptors and sphingosine kinase 1: novel biomarkers for clinical prognosis in breast, prostate, and hematological cancers," *Frontiers in Oncology*, vol. 2, article 168, 2012.
- [62] Y. Zhang, Y. Wang, Z. Wan, S. Liu, Y. Cao, and Z. Zeng, "Sphingosine kinase 1 and cancer: a systematic review and meta-analysis," *PLoS ONE*, vol. 9, no. 2, Article ID e90362, 2014.
- [63] C. R. Gault, S. T. Eblen, C. A. Neumann, Y. A. Hannun, and L. M. Obeid, "Oncogenic K-Ras regulates bioactive sphingolipids in a sphingosine kinase 1-dependent manner," *The Journal of Biological Chemistry*, vol. 287, no. 38, pp. 31794–31803, 2012.
- [64] M. Nagahashi, S. Ramachandran, E. Y. Kim et al., "Sphingosine-1-phosphate produced by sphingosine kinase 1 promotes breast cancer progression by stimulating angiogenesis and lymphangiogenesis," *Cancer Research*, vol. 72, no. 3, pp. 726–735, 2012.
- [65] M. E. Schnute, M. D. McReynolds, T. Kasten et al., "Modulation of cellular S1P levels with a novel, potent and specific inhibitor of sphingosine kinase-1," *Biochemical Journal*, vol. 444, no. 1, pp. 79–88, 2012.
- [66] K. Takabe, A. Yamada, and O. M. Rashid, "Twofer anti-vascular therapy targeting sphingosine-1-phosphate for breast cancer," *Gland Surgery*, vol. 1, no. 2, pp. 80–83, 2012.
- [67] T. Aoyagi, M. Nagahashi, A. Yamada, and K. Takabe, "The role of sphingosine-1-phosphate in breast cancer tumor-induced lymphangiogenesis," *Lymphatic Research and Biology*, vol. 10, no. 3, pp. 97–106, 2012.
- [68] G. D. Kruh and M. G. Belinsky, "The MRP family of drug efflux pumps," *Oncogene*, vol. 22, no. 47, pp. 7537–7552, 2003.
- [69] L. Amiri-Kordestani, A. Basseville, K. Kurdziel, A. T. Fojo, and S. E. Bates, "Targeting MDR in breast and lung cancer: discriminating its potential importance from the failure of drug resistance reversal studies," *Drug Resistance Updates*, vol. 15, no. 1–2, pp. 50–61, 2012.
- [70] S. Ponnusamy, S. P. Selvam, S. Mehrotra et al., "Communication between host organism and cancer cells is transduced by systemic sphingosine kinase 1/sphingosine 1-phosphate signalling to regulate tumour metastasis," *EMBO Molecular Medicine*, vol. 4, no. 8, pp. 761–775, 2012.
- [71] R. A. Sabbadini, "Sphingosine-1-phosphate antibodies as potential agents in the treatment of cancer and age-related macular degeneration," *British Journal of Pharmacology*, vol. 162, no. 6, pp. 1225–1238, 2011.
- [72] S. Schwalm, J. Pfeilschifter, and A. Huwiler, "Targeting the sphingosine kinase/sphingosine 1-phosphate pathway to treat chronic inflammatory kidney diseases," *Basic and Clinical Pharmacology and Toxicology*, vol. 114, no. 1, pp. 44–49, 2014.