CORRECTION Open Access

## Correction to: Comparative genomics of *Alternaria* species provides insights into the pathogenic lifestyle of *Alternaria brassicae* – a pathogen of the *Brassicaceae* family



Sivasubramanian Rajarammohan<sup>1,2</sup>, Kumar Paritosh<sup>3</sup>, Deepak Pental<sup>3</sup> and Jagreet Kaur<sup>1\*</sup>

Correction to: BMC Genomics (2019) 20:1036 https://doi.org/10.1186/s12864-019-6414-6

Following the publication of this article [1], the authors reported that the captions of Figs. 2 and 3 were published in the incorrect order, whereby they mismatch with their corresponding images. The figures are reproduced in the correct sequence with the correct captions in this Correction article. The original article has been corrected.

## Author details

<sup>1</sup>Department of Genetics, University of Delhi, South Campus, New Delhi 110021, India. <sup>2</sup>Present Address: National Agri-Food Biotechnology Institute, Mohali, India. <sup>3</sup>Centre for Genetic Manipulation of Crop Plants, University of Delhi South Campus, New Delhi, India.

Published online: 28 January 2020

## Reference

 Rajarammohan, et al. BMC Genomics. 2019;20:1036. https://doi.org/10.1186/ s12864-019-6414-6.


The original article can be found online at https://doi.org/10.1186/s12864-019-6414-6

Full list of author information is available at the end of the article




<sup>\*</sup> Correspondence: jagreet@south.du.ac.in

<sup>&</sup>lt;sup>1</sup>Department of Genetics, University of Delhi, South Campus, New Delhi



**Fig. 2** Whole-genome alignments of *A. alternata* PN1 and PN2 with *A. brassicae*. **a** Circos plot showing macrosynteny of *A. alternata* PN1 and PN2 with *A. brassicae* across all contigs except the dispensable contigs (ABRSC11, scaffold13,17,18,19), **b** and **c** Syntenic dotplots of *A. brassicae* with *A. alternata* PN1 and PN2



**Fig. 3** Phylogenetic tree of Alternaria species with S. lycopersici as an outgroup. The tree was constructed using 29 single copy orthologs, which had the highest phylogenetic signal as calculated in Mirlo. Branch support values from 1000 bootstrap replicates are shown