



Microarray Analysis of Genes Involved with Shell Strength in Layer Shell Gland at the Early Stage of Active Calcification

Zhangguo Liu^{1,2,a}, Qi Zheng^{3,a}, Xueyu Zhang⁴ and Lizhi Lu^{5,*}

¹ The Nurturing Station for the State Key Laboratory of Subtropical Silviculture, Zhejiang A & F University, Lin'an, Zhejiang, 311300, China

ABSTRACT: The objective of this study was to get a comprehensive understanding of how genes in chicken shell gland modulate eggshell strength at the early stage of active calcification. Four 32-week old of purebred Xianju hens with consistent high or low shell breakage strength were grouped into two pairs. Using Affymetrix Chicken Array, a whole-transcriptome analysis was performed on hen's shell gland at 9 h post oviposition. Gene ontology enrichment analysis for differentially expressed (DE) transcripts was performed using the web-based GOEAST, and the validation of DE-transcripts was tested by qRT-PCR. 1,195 DE-transcripts, corresponding to 941 unique genes were identified in hens with strong eggshell compared to weak shell hens. According to gene ontology annotations, there are 77 DE-transcripts encoding ion transporters and secreted extracellular matrix proteins, and at least 26 DE-transcripts related to carbohydrate metabolism or post-translation glycosylation modification; furthermore, there are 88 signaling DE-transcripts. GO term enrichment analysis suggests that some DE-transcripts mediate reproductive hormones or neurotransmitters to affect eggshell quality through a complex suite of biophysical processes. These results reveal some candidate genes involved with eggshell strength at the early stage of active calcification which may facilitate our understanding of regulating mechanisms of eggshell quality. (**Key Words:** Chicken, Microarray, Differentially Expressed Genes, Eggshell Strength)

INTRODUCTION

The chicken eggshell is a porous bioceramic container which protects the egg against physical damage and microbial contamination. Avian eggshell consists of the innermost bilayered membranes, a calcified layer composed of a mamillary and pallasade layer, and the outermost cuticle. The calcified layer consists of both inorganic minerals and extracellular matrix. It is well known that the shell mineral amount (thickness or density) is the main

factor contributing to the mechanical properties of the eggshell (Ahmed et al., 2005). However, the organic matrix, although its content in the calcified layer is only 2 to 3.5%, is of great importance to the deposition of bicarbonate and calcium ions, and to eggshell strength by controlling calcite crystal nucleation, growth, size and orientation (Greenfield et al., 1984).

The organic matrix in the calcified layer is comprised of a complex suite of components. In the acid soluble part of chicken eggshell matrix, 520 proteins have been identified (Mann et al., 2006), including several abundant proteins such as ovalbumin (Hincke, 1995), ovotransferrin (Gautron et al., 2001b), lysozyme (Hincke et al., 2000), osteopontin (Pines et al., 1995), sialoprotein (Solomon, 1999), clusterin (Mann et al., 2003), ovocleidin-17 (Hincke et al., 1995), ovocleidin-23 (Mann, 1999), ovocleidin-116 (Carrino et al., 1997), ovocalyxin-32 (Gautron et al., 2001a) and ovocalyxin-36 (Gautron et al., 2007). Many of the above components have been reported to undergo various post-translation modifications, which allow them to be effective chelators for interacting with the inorganic materials (Veis, 1989; Reyes-Grajeda et al., 2004; Mann et al., 2007), or to

* Corresponding Author: Lizhi Lu. Tel: +86-571-86406682, Fax: +86-571-86406682, E-mail: Lulizhibox@163.com

² College of Animal Science, Zhejiang A. & F. University, Lin'an, Zhejiang, 311300, China.

³ Department of Biology, 104 Lynch Laboratory, University of Pennsylvania, Philadelphia, PA 19104, USA.

⁴ Institute of Poultry Science, Chinese Academy of Agricultural Science, Yangzhou, Jiangsu, 225003, China.

⁵ Institute of Animal Husbandry and Veterinary Science, Zhejiang Provincial Academy of Agricultural Science, Hangzhou, Zhejiang, 310021, China.

^a These authors contributed equally to this work.

Submitted Jul. 18, 2012; Accepted Sept. 15, 2012; Revised Oct. 8, 2012

mediate protein–protein interactions to facilitate the assembly of the organic matrix (Lakshminarayanan et al., 2002; Ney et al., 2006).

It has been demonstrated that some genes in hen oviduct are associated with eggshell formation, whose expression is dependent on mechanical strain (Pines et al., 1995; Lavelin et al., 1998; Lavelin et al., 2002). It is proposed that some genes may function as crucial modulators for eggshell quality through regulating signal transduction, ion transportation, expression or modification of organic components, and many other processes. However, despite the importance of eggshell strength in the poultry industry, very few transcriptome-wide studies regarding this trait have been published to date (Yang et al., 2007; Dunn et al., 2009; Jonchère et al., 2010).

It is well documented that various parts of the avian eggshell are formed in specific regions of the oviduct as the egg passes through them. During the laying sequence, about 4 h after previous oviposition, the next egg arrives at and will take about 1h to pass through the white isthmus, in which the bilayered shell membranes are built around the egg. Then the egg enters the initial part of the shell gland, the red isthmus (tubular shell gland), and stays there for about 5 h to form mammillary knobs (Reyes-Grajeda et al., 2004). Finally the egg reaches the uterus (the main part of shell gland) and stays for an additional 15 h to form the palisade layer (Creger et al., 1976). It is known that the mamillary layer is the base of calcite crystal nucleation and crystal growth, and the palisade layer is the main part of the calcified shell, both of which affect global eggshell quality (Reyes-Grajeda et al., 2004; Jonchère et al., 2010).

In this study, we focused on the shell gland (uterus tissue near the red isthmus) at about 9 h post oviposition (corresponding to the early stage of active calcification, or to the transition stage from mammillary knob formation to construction of the palisade layer), and identified differentially expressed genes (DE-genes) in the layers with high shell strength compared to those with weak eggshell. Our results provide insight into the candidate genes involved in the mamillary layer formation and calcification that is crucial to the mechanical properties of avian eggshells.

MATERIALS AND METHODS

Animal treatments

Ninety purebred Xianju hens (a widely-bred Chinese indigenous chicken breed) of 28 weeks old were individually housed in laying cages. Birds were maintained under a cycle of 16 h light and 8 h dark. All birds were fed *ad libitum* with water and a mash layer diet (165 g protein, 35 g Ca, 11.29 MJ ME/kg, as recommended by NRC of China, 2004).

After 10 d of adaptation for hens, the oviposition time of each egg was initiated to be observed and recorded, then egg weight and shape index (length/width) were measured immediately. Following strength testing, the egg content was discarded and the shell was washed, dried at room temperature and weighed. Shell thickness without membranes was measured with a digital micrometer. Shell index ($\text{g}/100 \text{ cm}^2$) (Sauveur, 1988) was calculated as $I = (C/S) \times 100$, in which C is the weight of shell with membranes, S is the shell surface (cm^2) with $S = 4.68 \times P^{2/3}$ where P = egg weight (g). All above measurements were consecutively carried out daily for 16 d.

Finally, 2 groups of 2 hens with consistent high or low shell breakage strength were found. The differences between the eggshell properties of the selected 4 hens were analyzed by One-way ANOVA variance analysis in SPSS statistic software.

The four hens of interest were humanely sacrificed about 9 h after the previous oviposition. It is of note that all of the sacrificed hens had eggs in their uteruses (Figure 1A). The fat was removed from the uterus tissues near red isthmus and the tissues were then frozen in liquid nitrogen immediately and stored at -80°C . The animal treatments were approved by the Commission for Animal Welfare of Zhejiang A&F University.




Figure 1. Eggs and forming eggshells obtained from hen shell glands at the moment of tissue sampling. Figure 1A: Eggs obtained from the shell glands of hens when sampling uterus tissues. A-I is the egg from #35 hen, A-II from #19 hen, A-III from #80 hen, A-IV from #40 hen; these hens were all slaughtered at about 9 h after previous oviposition (P.O.). While A-V and A-VI eggs are from another 2 hens culled at 11.5 to 12 h after P.O., respectively. Figure 1B: The forming “shell” sampled at different stage of eggshell formation. B-I is the “shell” from the hen slaughtered at 4 h after P.O.; B-II is the “shell” of above egg from #40 hen. B-III is the “shell” sampled at 12 h post oviposition.

Measurement of eggshell strength

After egg weight and shape index measurements, the uncracked fresh eggs were individually placed lengthways with its blunt end upward in the FHK testing machine (Fujihara Co., Tokyo, Japan), and the vertical pressure was increasingly loaded upon the eggshell until the eggshell cracked and the eggshell strength was recorded as the maximum load (kgf).

RNA preparation

About 500 mg of the tissue of the uterus near red isthmus, including the mucosa, muscularis and outer serosa was powdered under liquid nitrogen. The total RNA was extracted using the RNAiso Plus Mini Kit (TaKaRa, Dalian, P.R. China) according to the manufacturer's instructions. RNA concentration and purity were measured by a NanoDrop spectrophotometer (NanoDrop[®] ND-1000, NanoDrop Technologies, DE).

Microarray hybridization and image acquisition

Microarray analysis was performed by the Bioassay Laboratory of CapitalBio Corporation (CapitalBio Co., Beijing, China). Briefly, the RNA integrity was firstly assessed using a Bioanalyzer (Agilent Technologies, Cheshire, UK), then 2 µg of total RNA was used for reverse transcription and biotin-labeled cRNA synthesis according to the manufactures' instructions, and finally subjected to microarray hybridization. The Affymetrix GeneChip[®] Chicken Genome Array (Affymetrix, Santa Clara, CA, USA) was used in this study, which contains 38,535 probesets corresponding to >28,000 chicken genes. Following 16 h of hybridization, the arrays were immediately washed, stained and scanned using Affymetrix[®] GeneChip[®] scanner 3000 (Affymetrix, Santa Clara, CA, USA), and the image files were processed into raw CEL intensity files using GeneChip Operating Software (GCOS version 1.2).

Pre-processing and normalization of microarray data

The raw intensity files generated by GCOS were imported and processed by R with Bioconductor packages. The total RNA quality was firstly verified statistically again by plotting the 5'-3' hybridization signal trends across all target transcripts. Then the microarray intensity was processed into transcript expression by the Affymetrix MAS5.0 method implemented in the R package, a procedure including background normalization, PM/MM probe correction, expression summarization and constant normalization on probeset level.

Identification of DE-transcripts

To identify DE-transcripts, the 4 array samples were first grouped into two pairs of high vs. low eggshell

strength according to eggshell property differences of the hens (see results). According to Cheuk and Cheng (2011), Affymetrix platform is relatively precise and sensitive in detecting signals, the DE-transcripts were identified as those with fold-change ≥ 2 in either of the two pairs of comparison and a statistical significant difference between high strength and low strength samples ($p < 0.05$, Welch t-test). It is of note that the log-odds values (Lods) of expression fold-change were used in the analysis; therefore, the DE-transcripts always have an absolute Lods value no less than 1 ($|Lods| \geq 1$).

Gene ontology enrichment analysis

Gene ontology enrichment analysis for DE-transcripts was performed using the web-based GOEAST (Zheng and Wang, 2008) Affymetrix analysis tool, with FDR cut-off of 0.05 using Yekutieli's FDR adjustment method.

Validation of differential expression by qRT-PCR experiments

Twenty-one DE-transcripts, with a fold-change ranging from low to high, were selected for further validation with qRT-PCR experiments; and all two groups of microarray samples were tested. Glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*) was used as the internal reference in all the PCR experiments. The primer sequences for qRT-PCR experiments can be found in Table 1.

To begin, total RNA was individually reverse transcribed with the SYBR[®] PrimeScript[™] RT-PCR kit II (TaKaRa, Dalian, China) according to the manufacturer's instructions. Then above RT-PCR kit was further used for fluorescence detection on an ABI Prism 7500 Sequence Detection System (Applied Biosystems, USA). All samples were analyzed in triplicates.

Dissociation curve analysis was conducted to ensure that a single PCR product with appropriate size was amplified in each reaction. On the other hand, the examination of PCR efficiency was performed based on LinRegPCR program (12.X) (Ramakers et al., 2003; Ruijter et al., 2009) to ensure internal and target transcript primers were amplified with similar efficiency.

The differential expression levels (Log₂ units) were calculated using the equation $\text{Log}_2 \text{ units (high versus low)} = -\Delta\Delta\text{Ct}$, where $\Delta\Delta\text{Ct} = (\text{Ct}^{\text{th}} - \text{Ct}^{\text{ih}}) - (\text{Ct}^{\text{tl}} - \text{Ct}^{\text{il}})$. Ct is the threshold cycle number when the amount of amplified product reaches a stable threshold. Ctth and Ct^{ih} represented the Ct of target transcript and internal reference transcript of "high eggshell strength sample", respectively. Correspondingly, Ct^{tl} and Ct^{il} represented the Ct of target transcript and inner-reference transcript of "low eggshell strength sample", respectively.

Table 1. Descriptions of specific primers used for real-time RT-PCR

Gene symbol	Accession no.	Forward primer (5'-3')	Reverse primer (5'-3')	Amplicon (bp)
<i>ACYP2</i>	XM_419292	CGGCTCGCTCAAGTCGGTGG	GGCCCTGAACTTGGCCCCGTC	152
<i>AMDHD1</i>	XM_416158	GCACTGGGAAGTGCATTTGCCA	TCTTCCGTGGCCTTCCTGGTGT	175
<i>ATP6VIA</i>	NM_204974	TGCAACATGGCAGGTGCTGCT	TGCCAGGCCCCAGTTCCACT	187
<i>CA5B</i>	XM_414195	CAGCTTGGCCACCTGCACTCC	ACACGTCGCTGGGTCTGCTAGCT	175
<i>CHST3</i>	NM_205121	TGATGGCCACCACACGCACC	CTGCAGCACGTCGCGGTACA	170
<i>COL12A1</i>	NM_205021	AGGCGAGTCTTCCCCGACGG	GCGCTGTCTCATGTCTGCC	171
<i>CRABP1</i>	NM_001030539	CGCCCCGCCATGCCTAACTT	AACTGGTCCCCGTCCTGGCG	161
<i>CRYBB1</i>	NM_204180	ACCTGGCGGACTGCGGGTT	CGGTAGCTGCTGGACCAGGTG	151
<i>EXOC6B</i>	XM_420892	AACCCACCACAGCCCTCGT	TGGCTGTTGATGAGGCCGCG	149
<i>FGB</i>	XM_420369.2	GCTGCTCCTGCTGCTCCTGC	GTGCCACGGGCCTGAGTGTG	155
<i>GAS2L3</i>	XM_416172	GGAGTAGTGTGGCAGTCCTGC	CCTGGGCCGTGTCTGGGAGT	193
<i>GIT2</i>	NM_204206	TCGCTTGCCATGCCGTGAGG	GCAACGTGGAGCGGGGTGTT	168
<i>MANIA2</i>	XM_416490	ACGTGGACACCAGCAAGGGGG	TCCTTTGCCTCTCCAGGGCCTTT	148
<i>NDST4</i>	XM_420638	CGAGCAGCTTCCCTCATCCCCAA	TGCCAGGGGCTTGACGTAA	156
<i>NPY</i>	NM_204587	GAGGACGCTCCCGCAGAGGA	TCGAAGGGTCTTCAAACCGGGA	175
<i>OC416916</i>	XM_415207	TGGAGGTGGAGCACAAACATCTGC	CCACCGAGCACACAGCCAGAAA	200
<i>PLCXD1</i>	NM_001128637	CCTGGCCTGCAGGAATTTTGATGG	AGCCACGCTGCCACATGGTC	137
<i>RCJMB04_34k20</i>	NM_001031112	GGACAGGCGGGCGAGAGAGT	TGGTGGTAAACGCACGCTGA	126
<i>SLC8A1</i>	XM_415002	CGTGTGTTGTGGCACTGGGGACA	ATGGCCGCGATGGACCAAGC	159
<i>TBXAS1</i>	XM_416334	TGTGTGGTGTGGGACAGCGT	ATACAGCCACGGGGTCTCTGCT	188
<i>WDR72</i>	XM_425069	GGCTGTTATCAGGGGGCCAGGA	GCACACGCAGCACACTACGC	161
<i>GAPDH</i>	NM_204305	GGGCTGCTAAGGCTGTGGGG	TCAGGGGCCATCAGCAGCA	177

RESULTS

Eggshell quality of hens under study

Among 90 tested hens, only 6 of them laid eggs at a similar laying rate with consistent high eggshell strength (defined as ≥ 4.5 kgf) or low strength (defined as ≤ 3.5 kgf). Two of these 6 hens were sacrificed at about 11.5 to 12 h post oviposition, but the eggshells collected from the shell glands showed more calcification extent than expected (Figure 1A and 1B). To focus on the initial stage of active calcification, we decided to use the uterus tissues near red isthmus from the remaining 4 hens, namely #19, #35, #40 and #80, at about 9 hr post oviposition. The eggs and eggshells harvested from these 4 hens at the moment of tissue sampling are shown in Figure 1A and Figure 1B. The eggshell breaking strength is also shown in Table 2 for these 4 hens, with consistent high (#19 and #40) or low (#35 and #80) eggshell quality.

To eggs from #19 and #35 hens, the differences of both

shell strength and shell weight were very significant ($p < 0.01$, t-test), but there was no significant difference ($p > 0.05$) for other eggshell quality metrics, such as shell thickness or shell index (Table 2). On the other hand, the differences of all of above eggshell metrics between eggs from #40 and #80 hens were very significant ($p < 0.01$) (Table 2). To get rigorous microarray data, therefore, we grouped the #19 with #35 hens due to the similarity of some of eggshell mechanical properties of the paired individuals; while #40 and #80 hens were also grouped as another pair.

Differentially expressed transcripts

The expression level of all probesets in 4 array samples were analyzed, and 1,195 DE-transcripts between uterus samples with high shell strength and low shell strength were identified. These DE-transcripts correspond to 941 unique genes. Among them, 407 genes were up-regulated in high strength samples comparing to low strength samples, and the other 534 genes were down-regulated. The expression

Table 2. Parameters related to eggshell quality of hens in this study

Hen	shell strength (kgf)	shell thickness (mm)	shell index (g/100 cm ²)	shell weight (g)	egg weight (g)	shape index
#40	5.17±0.40 ^A	0.367±0.016 ^A	8.07±0.28 ^A	4.297±0.144 ^a	38.46±1.52 ^C	1.274±0.024 ^B
#19	4.75±0.21 ^A	0.328±0.012 ^B	7.18±0.25 ^B	3.968±0.195 ^b	40.72±1.26 ^B	1.322±0.030 ^A
#35	2.99±0.71 ^B	0.324±0.023 ^B	6.97±0.68 ^B	4.363±0.426 ^a	48.60±1.38 ^A	1.278±0.032 ^{AB}
#80	2.54±0.69 ^B	0.272±0.022 ^C	6.25±0.71 ^C	3.451±0.417 ^c	40.58±1.06 ^B	1.313±0.052 ^{AB}

Values are from eggs laid by each hen of interest during the period of observation. Distinct capital letters in the same column indicate parameters between hens with a significant difference ($p < 0.01$), and distinct small letters indicate the significant difference is at level $p < 0.05$.




Figure 2. Heatmap and dendrogram of differentially expressed transcripts (DE-transcripts). Each cell represents the (normalized) gene expression value for given DE-transcript (row) in the specified sample (column). Cell colors indicate gene expression level: red: highly expressed; yellow: medium expression; green: lowly expressed. Row-side and column-side dendrogram represent the hierarchical clustering of DE-transcript expression for different transcripts or samples, respectively. Clustering is based on “complete-linkage” method using Euclidean-distance.

profile of all 1,195 DE-transcripts is shown in the heatmap in Figure 2. As shown in the heatmap, samples #19 with #40 and #35 with #80 were grouped as clusters among different samples, consistent with the similarity of their eggshell quality.

According to gene ontology annotations, the DE-transcripts are involved in a variety of biological processes. The most prominent DE-transcripts were found related to the following processes: signal transduction (88 DE-transcripts), ion transport and extracellular matrix organization (77 DE-transcripts), carbohydrate metabolism and protein modification (26 DE-transcripts) (Table 3).

Furthermore, avian calcified eggshell is a biomaterial composed of calcium salt and special ECM. The ECM is mainly comprised of collagens, glycoproteins and proteoglycans. Among the DE-transcripts, *COL8A2*, *COL12A1*, *COL13A1*, *LOC424798*, *LAMA2*, *LAMA4*, *LAMB4*, and *LAMC1* may be related to extracellular matrix formation; while *CHST3*, *GALNTL1*, *NDST4*, *LARGE*, *POFUT2*, *RCJMB04_28123*, and *MANIA2* are all localized in the endoplasmic reticulum or Golgi apparatus, and likely mediate the processes of carbohydrate metabolism, or posttranslation glycosylation modification.

Gene ontology (GO) term enrichment of DE-transcripts

It is of note that although many DE-transcripts were found related to various biological processes according to their ontology annotations, they are not necessarily correlated to the eggshell quality, due to random noise or other non-specific confounding factors commonly existing in microarray or other high-throughput experiments. Therefore, using web-based GOEAST (Zheng and Wang, 2008) we further identified significantly enriched GO terms among all the DE-transcripts. According to biology processes or molecular functions, the enriched GO terms can be roughly classified into several groups (Tables 4 and 5).

A group of processes are involved in reproductive hormone regulation, which contain Somatotropin secreting cell differentiation (GO:0060126), adenohipophysis development (GO:0021984), and response to estradiol stimulus (GO:0032355) (Table 4).

Table 3. i) DE-transcripts related with signaling, ion transportation, extracellular matrix protein, and carbohydrate metabolism or post-translation glycosylation modification

Gene symbol	Transcript ID	Log2 units (strong VS weak)	p-value	Category
<i>LOC771699</i>	XM_001234946	3.853	0.0004	signaling
<i>SH3PXD2A</i>	XM_421741	3.410	0.0397	signaling
<i>LOC429955</i>	XM_427511	3.296	0.0034	signaling
<i>PDCL2</i>	XM_420702	3.100	0.0115	signaling
<i>LOC430487</i>	XM_428042	3.055	0.0074	signaling
<i>SEMA3G</i>	XM_414289	2.933	0.0213	signaling
<i>RHOBTB2</i>	XM_001232709	2.905	0.0069	signaling
<i>RXFP1</i>	XM_420385	2.795	0.0205	signaling
<i>PIK3C2B</i>	XM_417956	2.777	0.0194	signaling
<i>OR10A7</i>	XM_425093	2.768	0.0067	signaling
<i>NPY</i>	NM_205473	2.401	0.0329	signaling
<i>PDE8B</i>	XM_425218	2.401	0.0498	signaling
<i>GREM2</i>	XM_419552	2.284	0.0423	signaling

Table 3. ii) DE-transcripts related with signaling, ion transportation, extracellular matrix protein, and carbohydrate metabolism or post-translation glycosylation modification (Continued)

Gene symbol	Transcript ID	Log2 units (strong VS weak)	p-value	Category
<i>SPAG9</i>	XM_420098	2.215	0.0423	signaling
<i>LOC396365</i>	NM_205400	2.166	0.0009	signaling
<i>MAPKBP1</i>	XR_026772	2.157	0.0187	signaling
<i>OXTR</i>	NM_001031569	2.128	0.0134	signaling
<i>STC2</i>	XM_414534	2.120	0.0135	signaling
<i>MPP3</i>	XM_418108	1.921	0.0491	signaling
<i>GREM1</i>	NM_204978	1.903	0.0220	signaling
<i>C20orf32</i>	XM_417499	1.889	0.0117	signaling
<i>CRABP1</i>	NM_001030539	1.778	0.0115	signaling
<i>C1orf107</i>	NM_001031051	1.770	0.0411	signaling
<i>RGS9</i>	XM_415685	1.657	0.0177	signaling
<i>ARHGEF12</i>	XM_417890	1.437	0.0014	signaling
<i>LTBP3</i>	XM_426444	1.433	0.0409	signaling
<i>NGEF</i>	NM_001010841	1.390	0.0307	signaling
<i>PDE1A</i>	XM_421969	1.363	0.0127	signaling
<i>CRHBP</i>	XM_424801	1.355	0.0159	signaling
<i>SRGAP1</i>	NM_001080101	1.335	0.0426	signaling
<i>PDE9A</i>	XM_416748	1.196	0.0475	signaling
<i>NPF2FR2</i>	NM_001034825	1.124	0.0283	signaling
<i>WNT4</i>	NM_204783	1.114	0.0180	signaling
<i>FGD4</i>	XM_416365	1.099	0.0121	signaling
<i>SOCS2</i>	NM_204540	1.033	0.0377	signaling
<i>TOB1</i>	NM_001001467	1.015	0.0347	signaling
<i>RND3</i>	XM_422158	0.887	0.0094	signaling
<i>ARHGAP28</i>	XM_419140	-4.485	0.0231	signaling
<i>WNT3</i>	NM_204675	-3.902	0.0422	signaling
<i>LOC428961</i>	NM_001142671	-3.469	0.0058	signaling
<i>VAV2</i>	NM_204142	-3.397	0.0232	signaling
<i>TBC1D20</i>	XM_001235014	-3.154	0.0389	signaling
<i>HTT</i>	XM_420822	-3.086	0.0065	signaling
<i>P2RY2</i>	XM_425667	-3.079	0.0294	signaling
<i>GRAP2</i>	XM_001234081	-2.967	0.0402	signaling
<i>INPP4A</i>	XM_416886	-2.749	0.0130	signaling
<i>ITSN1</i>	XM_416715	-2.744	0.0139	signaling
<i>EDNRB2</i>	NM_204120	-2.644	0.0317	signaling
<i>FGB</i>	XM_420369	-2.631	0.0194	signaling
<i>RXFP3</i>	XM_429217	-2.616	0.0048	signaling
<i>LOC420403</i>	XM_418509	-2.531	0.0049	signaling
<i>SSTR3</i>	NM_001024583	-2.501	0.0203	signaling
<i>GPR39</i>	NM_001080105	-2.376	0.0087	signaling
<i>GPR97</i>	XM_413998	-2.356	0.0230	signaling
<i>FGF14</i>	NM_204777	-2.322	0.0175	signaling
<i>SFRP4</i>	XM_418831	-2.204	0.0162	signaling
<i>LOC430333</i>	XM_001235474	-1.880	0.0153	signaling
<i>GARNLI</i>	XM_421244	-1.850	0.0477	signaling
<i>RAPH1</i>	XM_421961	-1.842	0.0035	signaling
<i>C14orf138</i>	XM_421460	-1.756	0.0012	signaling
<i>LOC421876</i>	XM_419893	-1.750	0.0333	signaling
<i>NLE1</i>	XM_415857	-1.735	0.0463	signaling
<i>CHRM2</i>	NM_001030765	-1.721	0.0310	signaling

Table 3. iii) DE-transcripts related with signaling, ion transportation, extracellular matrix protein, and carbohydrate metabolism or post-translation glycosylation modification (Continued)

Gene symbol	Transcript ID	Log2 units (strong VS weak)	p-value	Category
<i>LOC768958</i>	XM_001232128	-1.622	0.0278	signaling
<i>RASL10B</i>	XM_001233673	-1.546	0.0473	signaling
<i>SIPA1L2</i>	XM_419564	-1.545	0.0020	signaling
<i>LOC769317</i>	XM_001231944	-1.531	0.0287	signaling
<i>PLXDC2</i>	XM_418613	-1.503	0.0155	signaling
<i>SPOCK1</i>	XM_414622	-1.491	0.0375	signaling
<i>CSF2RB</i>	XM_001234608	-1.468	0.0498	signaling
<i>LOC429163</i>	XM_426718	-1.434	0.0220	signaling
<i>PLXNC1</i>	XM_416143	-1.398	0.0103	signaling
<i>RCJMB04_19g9</i>	XM_419989	-1.339	0.0019	signaling
<i>PLXNA1</i>	XM_414370	-1.305	0.0467	signaling
<i>TSPAN5</i>	XM_420654	-1.277	0.0142	signaling
<i>LOC431251</i>	NM_001127171	-1.226	0.0414	signaling
<i>ANXA10</i>	XM_001233661	-1.176	0.0441	signaling
<i>RCJMB04_18c11</i>	NM_001012909	-1.153	0.0436	signaling
<i>SPRED2</i>	XM_419341	-1.124	0.0146	signaling
<i>MOBK1A</i>	XM_420601	-1.116	0.0026	signaling
<i>ALS2CL</i>	XR_026875	-1.038	0.0119	signaling
<i>MPP1</i>	NM_001007917	-1.036	0.0424	signaling
<i>FGF12</i>	NM_204888	-1.016	0.0006	signaling
<i>GNA13</i>	XM_415686	-1.007	0.0195	signaling
<i>ARL10</i>	XM_414552	-0.975	0.0492	signaling
<i>ADRA2B</i>	XM_425203	-0.885	0.0415	signaling
<i>CCKAR</i>	NM_001081501	-0.868	0.0352	signaling
<i>RCJMB04_3n15</i>	NM_001030902	-0.815	0.0373	signaling
<i>TBC1D24</i>	XM_001232296	4.629	0.0023	IT
<i>SCN9A</i>	XM_422021	3.745	0.0047	IT
<i>KCNT2</i>	XM_426614	3.077	0.0015	IT
<i>LOC395893</i>		3.030	0.0392	IT
<i>NIPAL4</i>	XM_414566	2.796	0.0163	IT
<i>ATP6V0A4</i>	NM_001080102	2.790	0.0397	IT
<i>GRIN2B</i>	XM_416204	2.781	0.0291	IT
<i>KCNJ1</i>	XM_425795	2.363	0.0011	IT
<i>POR</i>	XM_415768	2.294	0.0047	IT
<i>KCNK2</i>	XM_001234269	2.270	0.0447	IT
<i>KIRREL3</i>	XR_026874	2.143	0.0193	IT
<i>GABRG2</i>	NM_205345	1.985	0.0176	IT
<i>SLC4A1</i>	NM_205522	1.794	0.0041	IT
<i>NDUFA7</i>	XM_418185	0.995	0.0464	IT
<i>JPH3</i>	XM_414192	0.925	0.0193	IT
<i>CACNA2D1</i>	XM_001231265	0.852	0.0429	IT
<i>EFCAB5</i>	XM_415833	-3.593	0.0342	IT
<i>SPATA22</i>	XM_001235167	-3.590	0.0176	IT
<i>RCJMB04_1f1</i>	NM_001031133	-3.197	0.0328	IT
<i>LOC428404</i>	XM_425965	-3.177	0.0442	IT
<i>ATP13A3</i>	XM_422709	-3.059	0.0065	IT
<i>GABRB2</i>	XM_001232377	-2.963	0.0472	IT
<i>SLC8A1</i>	NM_001079473	-2.694	0.0342	IT
<i>CACNA2D3</i>	XM_414338	-2.622	0.0100	IT
<i>SERINC5</i>	XM_424762	-2.615	0.0129	IT

Table 3. iv) DE-transcripts related with signaling, ion transportation, extracellular matrix protein, and carbohydrate metabolism or post-translation glycosylation modification (Continued)

Gene symbol	Transcript ID	Log2 units (strong VS weak)	p-value	Category
<i>LOC421866</i>	XR_027148	-2.613	0.0307	IT
<i>LOC425295</i>	XM_423073	-2.586	0.0212	IT
<i>LOC772391</i>	XM_001235535	-2.573	0.0163	IT
<i>KCNK17</i>	XM_419477	-2.506	0.0047	IT
<i>KCTD16</i>	XM_425217	-2.363	0.0076	IT
<i>CNNM1</i>	XM_421703	-2.309	0.0446	IT
<i>KCNJ5</i>	XM_417864	-2.138	0.0324	IT
<i>RCJMB04_11e10</i>	NM_001030630	-2.102	0.0226	IT
<i>GRIN3A</i>	XM_001232181	-1.680	0.0355	IT
<i>ATP6V1A</i>	NM_204974	-1.202	0.0065	IT
<i>RCJMB04_16a12</i>	NM_001031305	-1.030	0.0261	IT
<i>CNGA3</i>	NM_205221	-0.962	0.0479	IT
<i>P2RX4</i>	NM_204291	-0.882	0.0252	IT
<i>MEGF10</i>	XM_424719	3.910	0.0398	EM
<i>FAT2</i>	XM_414584	3.892	0.0052	EM
<i>SDK2</i>	NM_204538	2.905	0.0097	EM
<i>NRXN3</i>	XM_421297	2.643	0.0139	EM
<i>LAMA4</i>	XM_419780	2.569	0.0322	EM
<i>NTNG1</i>	XM_001231446	2.004	0.0328	EM
<i>CRTAC1</i>	NM_001080211	1.930	0.0132	EM
<i>LAMC1</i>	NM_204166	1.680	0.0257	EM
<i>LAMB4</i>	XM_001232877	1.642	0.0161	EM
<i>PPFIA1</i>	XM_421074	1.583	0.0166	EM
<i>CLDN20</i>	XM_001232002	1.438	0.0101	EM
<i>CDH9</i>	XM_001231501	1.296	0.0311	EM
<i>CHAD</i>	XM_416236	1.268	0.0240	EM
<i>LOC396026</i>	NM_205128	1.239	0.0403	EM
<i>PCDH21</i>	NM_001001759	1.174	0.0187	EM
<i>EPDR1</i>	XM_418830	1.158	0.0469	EM
<i>CPNE8</i>	XM_001231388	1.121	0.0219	EM
<i>COL12A1</i>	NM_205021	0.996	0.0003	EM
<i>PKP2</i>	XM_416362	0.983	0.0447	EM
<i>CD72</i>	NM_205052	0.855	0.0263	EM
<i>NINJ2</i>	XM_416382	-4.039	0.0409	EM
<i>OTOF</i>	XM_420015	-3.980	0.0016	EM
<i>COL13A1</i>	XM_001232218	-3.260	0.0006	EM
<i>SVEP1</i>	XM_424917	-2.967	0.0250	EM
<i>OTOP1</i>	XM_420790	-2.830	0.0263	EM
<i>GPNMB</i>	XM_425991	-2.771	0.0327	EM
<i>PKP1</i>	XM_419240	-2.349	0.0324	EM
<i>LAMA2</i>	XM_419746	-2.336	0.0000	EM
<i>COL8A2</i>	XM_425780	-2.295	0.0170	EM
<i>FNBP4</i>	XM_424260	-2.279	0.0486	EM
<i>EGFL6</i>	XM_416835	-2.268	0.0392	EM
<i>CDH18</i>	XM_426046	-1.955	0.0090	EM
<i>CLDN8</i>	XM_425544	-1.598	0.0288	EM
<i>RCJMB04_34k20</i>	NM_001031112	-1.214	0.0085	EM
<i>SRPX</i>	XM_416781	-1.123	0.0244	EM
<i>DLG1</i>	XM_422701	-1.084	0.0171	EM
<i>FBLN1</i>	NM_204165	-0.981	0.0206	EM

Table 3. v) DE-transcripts related with signaling, ion transportation, extracellular matrix protein, and carbohydrate metabolism or post-translation glycosylation modification (Continued)

Gene symbol	Transcript ID	Log2 units (strong VS weak)	p-value	Category
<i>F13A1</i>	NM_204685	-0.949	0.0234	EM
<i>FREM1</i>	XM_424932	-0.910	0.0057	EM
<i>MEGF10</i>	XM_424719	3.910	0.0398	EM
<i>MGAT4C</i>	XM_425447	3.007	0.0497	GM or CM
<i>CHST3</i>	NM_205121	2.839	0.0372	GM or CM
<i>EDEM3</i>	XM_422293	2.539	0.0179	GM or CM
<i>LARGE</i>	NM_001004404	2.071	0.0429	GM or CM
<i>GFPT2</i>	XM_424573	1.919	0.0173	GM or CM
<i>GALNTL1</i>	XM_001231964	1.895	0.0452	GM or CM
<i>WDR77</i>	NM_001030916	1.805	0.0445	GM or CM
<i>NDST3</i>	XM_426325	1.403	0.0121	GM or CM
<i>B3GALT1</i>	XM_426584	1.254	0.0483	GM or CM
<i>OGDHL</i>	XM_421503	1.144	0.0022	GM or CM
<i>MAN1A2</i>	XM_416490	-4.700	0.0087	GM or CM
<i>POFUT2</i>	XM_421892	-2.804	0.0156	GM or CM
<i>LOC772154</i>	XM_001235329	-2.798	0.0006	GM or CM
<i>KLB</i>	XM_423224	-2.335	0.0380	GM or CM
<i>TRIM7.2</i>	NM_001099354	-1.986	0.0041	GM or CM
<i>LOC771361</i>	XM_001234647	-1.433	0.0369	GM or CM
<i>RCJMB04_28123</i>	NM_001039316	-1.324	0.0034	GM or CM
<i>NDST4</i>	XM_420638	-1.239	0.0261	GM or CM
<i>PFKM</i>	NM_204223	-1.170	0.0460	GM or CM
<i>NUP153</i>	XM_418937	-1.109	0.0378	GM or CM
<i>GPD1L</i>	XM_418763	-0.947	0.0383	GM or CM
<i>PHKA2</i>	XM_416811	-0.926	0.0326	GM or CM
<i>B3GNTL1</i>	XM_415599	-0.883	0.0109	GM or CM
<i>PMM1</i>	XM_416228	-0.787	0.0490	GM or CM
<i>MMP11</i>	XM_001232776	2.209	0.0391	GM or CM
<i>ST3GAL4</i>	XM_417860	-1.043	0.0094	GM or CM

IT represents ion/proton transporter, EM represents extracellular matrix, GM represents post-translation glycosylation modification, and CM represents carbohydrate metabolism.

As shown in Table 4 and Table 5, many DE-transcripts are involved in signal transduction, such as GO terms purinergic nucleotide receptor activity (GO:0001614), nucleotide receptor activity (GO:0016502), purinergic receptor activity (GO:0035586), transmembrane signaling receptor activity (GO:0004888), and negative regulation of BMP signaling pathway (GO:0030514). Among them, GO:0004888 dominantly contains 33 transcripts encoding signal receptors, and these receptors could be further classified into several subgroups: *OXTR*, *LOC431251* and *SSTR3* belong to reproductive hormone receptors; *CHRM2*, *ADRA2B*, *P2RX4*, *P2RY2*, *EDNRB2*, *GABRB2*, *GABRG2*, *LOC428961* and *NPFFR2* function as receptors mediating neurotransmitters or neuropeptide; *GRIN2B* and *GRIN3A* could modulate the efficiency of synaptic transmission; *NTRK1* and *NTRK2* belong to the receptor tyrosine kinase (RTK) family, and are involved with neurotrophin (GO:0005030 - neurotrophin receptor activity; and GO:0043121 - neurotrophin binding) (Table 5).

Besides various enriched molecular function shown above, many biophysical processes are also found to be enriched among the DE-transcripts, including a series of processes and subgroups (Tables 4 and 5). GO:0003951 (NAD⁺ kinase activity) modulate the metabolism or redox in cell (Table 5). Enrichment of GO:0009409 (response to cold) may reflect the fact the rearing condition of experimental hens was in the winter at room temperature about 2 to 10°C. GO:0046209 (nitric oxide metabolic process) may regulate vascular or smooth muscle relaxation or other functions. GO:0002028 is involved in ion transportation. While the subgroup processes of muscular development and activity include skeletal muscle fiber development (enrichments of GO:0048741, GO:0048747 and GO:0055002) and striated muscle contraction regulation (enrichments of GO:0055117 and GO:0006942). It is of note that there is almost no striated muscle in avian uterus except smooth muscle. However, the chicken genome project was completed in 2004, and the functional

Table 4. Enriched gene ontology (GO) terms revealed from identified DE-transcripts according to biological_process ontology

Group	GOID	Term	p	Gene symbol or representative public ID	
Reproductive hormone synthesis and regulation	GO:0060126	Somatotropin secreting cell differentiation	0.013	<i>OTX2, WNT4</i>	
	GO:0021984	Adenohypophysis development	0.044	<i>OTX2, WNT4</i>	
Signal transduction	GO:0032355	Response to estradiol stimulus	0.044	<i>SOCS2, AREGB</i>	
	GO:0030514	Negative regulation of BMP signaling pathway	0.030	<i>TOB1, GREM1</i>	
Biophysical processes	GO:0048741	Skeletal muscle fiber development	0.012	<i>SLC23A2, CHAT</i>	
	GO:0015074	DNA integration	0.012	<i>LOC770294, LOC770705, ENS-3</i>	
	GO:0055117	Regulation of cardiac muscle contraction	0.013	<i>P2RX4, NKX2-5</i>	
	GO:0009409	Response to cold	0.018	<i>IL4, SLC27A1</i>	
	GO:0048747	Muscle fiber development	0.022	<i>SLC23A2, CHAT</i>	
	GO:0046209	Nitric oxide metabolic process	0.024	<i>P2RX4, CPS1</i>	
	GO:0007586	Digestion	0.030	<i>PGA5, PRSS2, LOC396365</i>	
	GO:0015849	Organic acid transport	0.033	<i>SLC23A2, OCA2, LOC770309, SLC7A14, SLC27A1</i>	
	GO:0046942	Carboxylic acid transport	0.033	<i>SLC23A2, OCA2, LOC770309, SLC7A14, SLC27A1</i>	
	GO:0055002	Striated muscle cell development	0.034	<i>SLC23A2, CHAT, TTN, NKX2-5</i>	
	GO:0006942	Regulation of striated muscle contraction	0.037	<i>P2RX4, NKX2-5</i>	
	GO:0002028	Regulation of sodium ion transport	0.044	<i>NKX2-5, NEDD4L</i>	
	Reproductive biophysical processes	GO:0060748	Tertiary branching involved in mammary gland duct morphogenesis	0.009	<i>WNT4, AR</i>
		GO:0060745	Mammary gland branching involved in pregnancy	0.013	<i>WNT4, AR</i>
GO:0060562		Epithelial tube morphogenesis	0.019	<i>DEAF1, WNT3, GREM1, WNT4, NKX2-5, HOXA11, AR, AREGB</i>	
GO:0060444		Branching involved in mammary gland duct morphogenesis	0.020	<i>WNT4, AR, AREGB</i>	
GO:0009994		Oocyte differentiation	0.024	<i>WNT4, GDF9</i>	
GO:0048599		Oocyte development	0.024	<i>WNT4, GDF9</i>	
GO:0060603		Mammary gland duct morphogenesis	0.033	<i>WNT4, AR, AREGB</i>	
GO:0060135		Maternal process involved in female pregnancy	0.037	<i>WNT4, AR</i>	

GOID represents the identifiers, and Term represents term definitions for Gene Ontology term entities. p: p-value of significance (Welch t-test).

gene database of *G. gallus* remains incomplete, some ontology annotations of DE-genes may refer to mammalian homologs, which may account for our results. The genes related to muscular cell contraction are likely to modulate the mobility of uterus to facilitate egg rotation and calcification (Johnson, 1986; Jonchère et al., 2010). Similarly, there is no digestion process in the uterus, three genes in GO:0007586 (digestion process), *PGA5* (an

aspartic acid protease, which is involved in ovulation (Peluffo et al., 2011)), *PRSS2* and *LOC396365* (preprogastrin), are likely to promote the maturation of secretory extracellular proteins or regulate the secretion of uterus glands and mobility of uterus.

The final group of reproductive biophysical processes also includes several subgroups of processes (Table 4). Epithelial tube morphogenesis (GO:0060562) may regulate

Table 5. Enriched gene ontology (GO) terms revealed from identified DE-transcripts according to molecular function ontology

Group	GOID	Term	p	Gene symbol or representative public ID
Signal transduction	GO:0005030	Neurotrophin receptor activity	0.013	<i>NTRK1, NTRK2</i>
	GO:0001614	Purinergic nucleotide receptor activity	0.017	<i>P2RX4, P2RY2, ENSGALG00000012080</i>
	GO:0016502	Nucleotide receptor activity	0.017	<i>P2RX4, P2RY2, ENSGALG00000012080</i>
	GO:0043121	Neurotrophin binding	0.024	<i>NTRK1, NTRK2</i>
	GO:0035586	Purinergic receptor activity	0.026	<i>P2RX4, P2RY2, ENSGALG00000012080</i>
	GO:0004888	Transmembrane signaling receptor activity	0.049	<i>OXTR, LOC431251, SSTR3, CHRM2, ADRA2B, P2RX4, P2RY2, EDNRB2, GABRB2, GABRG2, LOC428961, NPFFR2, GRIN2B, GRIN3A, NTRK1, NTRK2, EPHB6, DDR2, TMPRSS6, PCSK5, CCKAR, IFNAR2, CSF1R, TLR5, OR10A7, LOC768958, LOC769317, LOC777484, GPR39, GPR97, ENSGALG00000017405, ENSGALG00000017093, ENSGALG00000012080.</i>
Biophysical processes	GO:0003951	NAD+ kinase activity	0.013	<i>C5orf33, NADK</i>
	GO:0005319	Lipid transporter activity	0.049	<i>ATP11C, ATP8A2, ATP8B3, APOB, LOC769564, SLC27A1</i>

GOID represents the identifiers, and Term represents term definitions for gene ontology term entities. p: p-value of significance (Welch t-test).

the development of uterus glands (tubular epithelial glands). Oocyte development subgroup contains oocyte differentiation (GO:0009994) and oocyte development (GO:0048599). Female pregnancy subgroup contains enrichments of GO:0060135, GO:0060745, GO:0060748, GO:0060444 and GO:0060603.

Overall, laying is an avian reproductive behavior, and eggshell calcification is regulated by relative reproductive hormones and neurotransmitters, which may finally affect eggshell quality through a complex suite of biophysical reactions.

Confirmation of DE-transcripts by qRT-PCR

21 DE-transcripts (9 up-regulated and 12 down-regulated) were chosen for validation using qRT-PCR experiments, and the four microarray samples were tested in pairs for #19 vs #35 and #40 vs #80, respectively. As shown in Figure 3, 16 out of the 21 tested transcripts (76%) were confirmed by qRT-PCR experiments, though the absolute fold-change values are slightly different. The remaining transcripts, *CRYBB1, EXOC6B, LOC416916, MANIA2* and *CHST3*, showed inconsistent differential expression between qRT-PCR and microarray experiments.

GAPDH, CHST3, GALNTL1, NDST4, LARGE, SPI, RHOBTB2, and *WDR72* were selected to examine the PCR efficiency. The results showed the PCR efficiency of these genes ranged from 86.8% to 94.2%, and the PCR efficiency of inner reference (*GAPDH*) and other genes seemed nearly

similar.

DISCUSSION

Laying is regarded as avian reproductive behavior, which is regulated by reproductive hormones and neurotransmitters. The chicken oviduct has been extensively used as a model to study hormonal induction of protein synthesis (Khuong and Jeong, 2011). Under the control of steroid hormones or neurotransmitters, the tubular gland epithelial cells synthesize and secrete a great variety of proteins to form egg white and eggshell when egg passes through the oviduct (Mann et al., 2006). In this paper, 1,195 DE-transcripts have been identified to be related with eggshell strength. GOEAST analysis further identify some significantly enriched GO terms, and the enriched GO terms suggest that some DE-transcripts mediate reproductive hormones or neurotransmitters to affect eggshell quality (Tables 4 and 5).

Both terms GO:0060126 and GO:0021984 are involved in reproductive hormone regulation, and share two genes, *OTX2* and *WNT4*.

Otx2 is a paired-like homeodomain transcription factor, which can mediate GnRH (gonadotropin releasing hormone) signaling (Kelley et al., 2000). Functional studies revealed that *Otx2* is required as early as gastrulation for neural induction, and even for brain development (Rhinn et al., 1998). However, *Otx2* is also of importance for




Figure 3. Real-time RT-PCR validation of microarray data. Expression levels of the first 12 transcripts (*AMDHD1*, *ATP6V1A*, *CRYBB1*, *FGB*, *GAS2L3*, *GIT2*, *MAN1A2*, *NDST4*, *PLCXD1*, *RCJMB04-34k20*, *SLC8A1* and *TBXAS1*) were down-regulated in microarray experiment, and the last 9 transcripts (*ACYP2*, *CA5B*, *CHST3*, *COL12A1*, *CRABP1*, *EXOC6B*, *LOC416916*, *NPY* and *WDR72*) were up-regulated in microarray experiment. Among the above transcripts, five transcripts (*CRYBB1*, *MAN1A2*, *CHST3*, *EXOC6B* and *LOC416916*) failed to be verified by real-time RT-PCR.

neurogenesis and cellular proliferation in multiple other tissues (Layman et al., 2011).

As a member of the WNT family, Wnt4 is a secreted glycoprotein signaling molecule and involved in paracrine signaling (Diaz et al., 2011). Wnt4 is critical for female sex determination and differentiation (Chen et al., 2011). In the female, Wnt4 is positively involved in ovarian development; while in the male mutated *WNT4* will result in aberrant testis development (Diaz et al., 2011; Barrionuevo et al., 2012). On the other hand, Wnt4 is also potent to regulate the development of the female reproductive tract (Franco et al., 2011). Furthermore, *WNT4* is expressed postnatally in ovarian follicles and corpora lutea, and its expression increases in response to gonadotropin (Hsieh et al., 2002). Wnt4 mediates follicle development and fertility by regulating the expression of genes involved in steroidogenesis, prostaglandin biosynthesis, tissue remodeling, and angiogenesis (Hsieh et al., 2002; Boyer et al., 2010). Moreover, excluding its reproductive contributions, *WNT4* is also tightly associated with bone strength (Zmuda et al., 2011).

Our results also show that some DE-transcripts are involved in signal transduction (Tables 4 and 5), among which, *NTRK1*, *NTRK2*, *P2RX4*, and *P2RY2* are overlapped in multiple enriched GO terms (Table 5).

Ntrk1, also named TrkA, and Ntrk2 TrkB, are two members of the neurotrophic tyrosine kinase receptor (NTRK) family. These kinases are membrane-bound

receptors mediating various functions of neurotrophins, such as cell survival, migration, outgrowth of axons and dendrites, synaptogenesis, remodeling of synapses, and synaptic transmission (Ohira and Hayashi, 2009). So far, several neurotrophins have been well studied, such as nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), neurotrophin 3 (NT-3), and NT-4/5. NTKRs are high affinity receptors of neurotrophins. TrkA mediates the biological response of NGF, while BDNF and NT-4/5 are the preferred ligands for TrkB (Ohira and Hayashi, 2009). Additionally, NTKRs also play roles in some biomaterials. NT-4 may modulate proliferation and differentiation of the dental epithelium and promote production of the enamel matrix via the TrkB-MAPK pathway (Yoshizaki et al., 2008).

Both *P2RX4* and *P2RY2* are purinergic receptors. Purinergic receptors are subdivided into metabotropic *P2Y* receptors and ionotropic *P2X* receptors. *P2Y* receptors are coupled to G-protein and trigger inositol 1,4,5-triphosphate (IP₃)-induced intracellular Ca²⁺ release following activation of phospholipase C, while *P2X* receptors are ligand-gated ion channels. *P2RX4* will be discussed later, while *P2RY2* performs a dominant role in calcium signaling during osteoblast differentiation (Nishii et al., 2009). It is known that extracellular ATP, UTP, and PPi can strongly block the mineralization of matrix nodules, while this potent inhibition of bone formation is mediated by *P2RY2* (Orriss et al., 2007). Furthermore, *P2RY2* is also involved in

inhibition of intercellular communication between osteoblasts (Hoebertz et al., 2003).

At present, there are at least three cDNA microarray studies globally investigating the gene expression in chicken shell gland (Yang et al., 2007; Dunn et al., 2009; Jonchère et al., 2010), but the overlap among the DE-genes from these studies is not plentiful. Different animals, tissue samples or treatment methods may partially account for this problem.

Yang et al. (2007) harvested uterus tissues at 2 h post oviposition, and screened out 34 known genes in the shell glands between hens with low and high egg productive rates. This study and our data share a single gene *CALD1* (caldesmon 1) (Figure 4). *CALD1* is a ubiquitous actin and calmodulin binding protein, and functions as a substrate for mitogen-activated protein kinase (Childs et al., 1992) or as serine and threonine kinases (Sutherland et al., 1994).

Dunn et al. (2009) identified 266 DE-genes in shell glands from 25-week old mature hens comparing to 12-week old juveniles from high and low bone quality lines, respectively. The tissues sampled when eggs passed through the oviducts but not in shell glands. Three DE-genes are also found in our data: *NADK* (NAD kinase), *LOC422993* (Similar to interferon-induced membrane protein Leu-13/9-27), and *LAMP3* (lysosomal-associated membrane protein 3) (Figure 4), suggesting potential crucial function of these genes in not only early stage of eggshell calcification but also other stages of eggshell formation.

Jonchère et al. (2010) used the 40-week old hens at 18 h

post oviposition (corresponding to the rapid phase of calcification), and identified 469 DE-known genes in uterus versus both white isthmus, and magnum. There are 7 genes consistently identified in their study and our data, such as *P2RX4* (purinergic receptor P2X, ligand-gated ion channel, 4), *FSTL1* (follistatin-like 1), *TUBGCP4* (Tubulin, gamma complex associated protein 4), *WDR77* (WD repeat domain 77), *RCJMB04_6g16* (microtubule-associated protein 1 light chain 3 beta), *PWP1* (PWP1 homolog in *S. cerevisiae*) and *SGK1* (serum/glucocorticoid regulated kinase 1) (Figure 4).

On the other hand, three additional DE-genes in our data were previously found in the acid soluble part of chicken eggshell organic matrix (Mann et al., 2006). These three genes, *FSTL1* (follistatin-like 1), *CAMK2D* (calcium/calmodulin-dependent protein kinase (CaM kinase) II delta) and *KRT75* (keratin 75) (Figure 4), could reflect potential interaction of eggshell calcification and organic matrix formation.

Among these overlapping DE-genes, both *P2RX4* (Jonchère et al., 2010) and *NADK* (Yang et al., 2007) are also present in our enriched GO terms (Tables 4 and 5), and *FSTL1* (Mann et al., 2006; Jonchère et al., 2010) occurs in more than three relative studies.

P2RX4 is one member of the P2X receptors (P2RX). P2RX are ionotropic ATP-gated ion channels conducting Ca^{2+} inflow (Fodor et al., 2009), with high capability of Ca^{2+} permeabilities corresponding to at least 100-fold those of Na^{+} (Burnashev, 1998). In chondrogenic mesenchymal




Figure 4. MA-plot of all microarray tested chicken-genes. X-axis: the average normalized expression values across all 4 (strong+weak) eggshell samples (in \log_2 scale); Y-axis: the log-odds ratio between the average expression values of strong vs. weak eggshell samples; grey dots: nondifferentially expressed genes (non-DE genes); red dots: differentially expressed genes (DE-genes) identified in this study; big blue dots: DE-genes (*CALD1*, *NADK*, *LOC422993*, *LAMP3*, *P2RX4*, *FSTL1*, *TUBGCP4*, *WDR77*, *RCJMB04_6g16*, *PWP1* and *SGK1*) reported in previous studies.

cells, P2X4 receptors could conduct Ca^{2+} inflow to elevate intracellular Ca^{2+} levels, and finally promoting extracellular matrix production (Fodor et al., 2009). Eggshell calcification requires considerable ion transportation, especially Ca^{2+} , and various extracellular matrix synthesis and secretion, whether and how P2RX4 channels regulate these processes requires further studies.

NAD kinases (NADKs) are a family of enzymes transferring a phosphate group from ATP to NAD to generate and maintain the cellular NADP pool (Pollak et al., 2007). It is reported, that during development of placenta, the expression level of *NADK* appears drastically elevated (Lerner et al., 2001).

Fstl1 is a secreted glycoprotein belonging to the BM-40/SPARC/osteonectin family containing both calcium-binding domain and Follistatin-like domain (Hambrock et al., 2004). As a mesenchymal factor, Fstl1 is critical for oviduct development, and determines the differentiation of secretory epithelial cells and ciliated epithelial cells in the oviduct (Umezu et al., 2010). This means Fstl1 may modulate chicken endometrium development during eggshell formation. However, Fstl1 is also present in the organic part of eggshells (Mann et al., 2006), and Jonchère et al. (2010) propose it may be a uterine antiprotease

IMPLICATIONS

Above all, using Affymetrix Chicken Array, 1,195 DE-transcripts were identified in the shell gland between “high shell strength” and “low shell strength” hens, which represent 941 unique known genes. According to gene ontology annotations, these transcripts are involved in a wide range of biological processes; the most prominent DE-transcripts relate to signal transduction, metabolism, extracellular matrix, or ion transport and homeostasis, and so on. Furthermore, Gene Ontology (GO) term enrichment of DE-transcripts suggests that avian eggshell calcification is likely to be regulated by relative reproductive hormones and neurotransmitters, which may finally affect eggshell quality through a complex suite of biophysical processes.

ACKNOWLEDGEMENTS

This work was supported by National Natural Science Foundation of China (grant no. 30700567) and Zhejiang Provincial Natural Science Foundation of China (grant no. LY12C17002).

REFERENCES

- Ahmed, A. M. H., A. B. Rodriguez-Navarro, M. L. Vidal, J. Gautron, J. M. Garcia-Ruiz and Y. Nys. 2005. Changes in eggshell mechanical properties, crystallographic texture and in matrix proteins induced by moult in hens. *Br. Poult. Sci.* 46:268-279.
- Barrionuevo, F. J., M. Burgos, G. Scherer and R. Jiménez. 2012. Genes promoting and disturbing testis development. *Histol. Histopathol.* 27:1361-1383.
- Burnashev, N. 1998. Calcium permeability of ligand-gated channels. *Cell Calcium.* 24:325-332.
- Boyer, A., É. Lapointe, X. Zheng, R. G. Cowan, H. Li, S. M. Quirk, F. J. DeMayo, J. S. Richards, and D. Boerboom. 2010. WNT4 is required for normal ovarian follicle development and female fertility. *FASEB J.* 24:3010-3025.
- Carrino, D. A., J. P. Rodriguez and A. I. Caplan. 1997. Dermatan sulfate proteoglycans from the mineralized matrix of the avian eggshell. *Connect. Tissue Res.* 36:175-193.
- Chen, B., P. Suo, B. Wang, J. Wang, L. Yang, S. Zhou, Y. Zhu, X. Ma and Y. Cao. 2011. Mutation analysis of the WNT4 gene in Han Chinese women with premature ovarian failure. *Reprod. Biol. Endocrinol.* 9:75.
- Cheuk, B. L. and S. W. Cheng. 2011. Differential expression of elastin assembly genes in patients with Stanford Type A aortic dissection using microarray analysis. *J. Vasc. Surg.* 53:1071-1078.
- Childs, T. J., M. H. Watson, J. S. Sanghera, D. L. Campbell, S. L. Pelech and A. S. Mak. 1992. Phosphorylation of smooth muscle caldesmon by mitogen-activated protein (MAP) kinase and expression of MAP kinase in differentiated smooth muscle cells. *J. Biol. Chem.* 267:22853-22859.
- Creger, C. R., H. Phillips and J. T. Scott. 1976. Formation of an eggshell. *Poult. Sci.* 55:1717-1723.
- Diaz, F. J., K. Anthony and A. N. Halfhill. 2011. Early avian follicular development is characterized by changes in transcripts involved in steroidogenesis, paracrine signaling and transcription. *Mol. Reprod. Dev.* 78:212-223.
- Dunn, I. C., P. W. Wilson, Z. Lu, M. M. Bain, C. L. Crossan, R. T. Talbot and D. Waddington. 2009. New hypotheses on the function of the avian shell gland derived from microarray analysis comparing tissue from juvenile and sexually mature hens. *Gen. Comp. Endocrinol.* 163:225-232.
- Fodor, J., C. Matta, T. Juhász, T. Oláh, M. Gönczi, Z. Szűgyártó, P. Gergely, L. Csernoch and R. Zákány. 2009. Ionotropic purinergic receptor P2X4 is involved in the regulation of chondrogenesis in chicken micromass cell cultures. *Cell Calcium.* 45:421-430.
- Franco, H. L., D. Dai, K. Y. Lee, C. A. Rubel, D. Roop, D. Boerboom, J. W. Jeong, J. P. Lydon, I. C. Bagchi, M. K. Bagchi and F. J. Demayo. 2011. WNT4 is a key regulator of normal postnatal uterine development and progesterone signaling during embryo implantation and decidualization in the mouse. *FASEB J.* 25:1176-1187.
- Gautron, J., M. T. Hincke, K. MANN, M. Panhéleux, M. Bain, M. D. McKee, S. E. Solomon and Y. Nys. 2001a. Ovocalyxin-32, a novel chicken eggshell matrix protein: Isolation, amino acid sequencing, cloning and immunocytochemical localization. *J. Biol. Chem.* 276:39243-39252.
- Gautron, J., M. T. Hincke, M. Panhéleux, J. M. Garcia-Ruiz, T. Boldicke and Y. Nys. 2001b. Ovotransferrin is a matrix protein of the hen eggshell membranes and basal calcified layer. *Connect. Tissue Res.* 42:255-267.
- Gautron, J., E. Murayama, A. Vignal, M. Morisson, M. D. McKee,

- S. Réhault, V. Labas, M. Belghazi, M. L. Vidal, Y. Nys and M. T. Hincke. 2007. Cloning of Ovocalyxin-36, a novel chicken eggshell protein related to lipopolysaccharide-binding proteins (LBP), bactericidal permeability-increasing proteins (BPI), and plunc family proteins. *J. Biol. Chem.* 282:5273-5286.
- Greenfield, E. M., D. C. Wilson and M. A. Crenshaw. 1984. Iontropic nucleation of calcium carbonate by molluscan matrix. *Amer. Zool.* 24:925-932.
- Hambrock, H. O., B. Kaufmann, S. Müller, F. G. Hanisch, K. Nose, M. Paulsson, P. Maurer and U. Hartmann. 2004. Structural characterization of TSC-36/Flik: Analysis of two charge isoforms. *J. Biol. Chem.* 279:11727-11735.
- Hincke, M. T. 1995. Ovalbumin is a component of the chicken eggshell matrix. *Connect. Tissue Res.* 31:227-233.
- Hincke, M. T., J. Gautron, M. Panhéleux, J. Garcia-Ruiz, M. D. McKee and Y. Nys. 2000. Identification and localization of lysozyme as a component of eggshell membranes and eggshell matrix. *Matrix Biol.* 19:443-453.
- Hincke, M. T., C. P. Tsang, M. Courtney, V. Hill and R. Narbaitz. 1995. Purification and immunochimistry of a soluble matrix protein of the chicken eggshell (Ovocleidin 17). *Calcif. Tissue Int.* 56:578-583.
- Hsieh, M., M. A. Johnson, N. M. Greenberg and J. S. Richards. 2002. Regulated expression of Wnts and Frizzleds at specific stages of follicular development in the rodent ovary. *Endocrinology.* 143:898-908.
- Johnson, A. L. 1986. Reproduction in the female. In: *Avian Physiology*. 4th ed. (Ed. P. D. Sturkie). Springer-Verlag, New York.
- Jonchère, V., S. Réhault-Godbert, C. Hennequet-Antier, C. Cabau, V. Sibut, L. A. Cogburn, Y. Nys and J. Gautron. 2010. Gene expression profiling to identify eggshell proteins involved in physical defense of the chicken egg. *BMC Genomics.* 11:57-75.
- Kelley, C. G., G. Lavorgna, M. E. Clark, E. Boncinelli and P. L. Mellon. 2000. The Otx2 homeoprotein regulates expression from the gonadotropin-releasing hormone proximal promoter. *Mol. Endocrinol.* 14:1246-1256.
- Khuong, T. T. and D. K. Jeong. 2011. Adipogenic differentiation of chicken epithelial oviduct cells using only chicken serum. *In Vitro Cell. Dev. Biol. Anim.* 47:609-614.
- Lakshminarayanan, R., R. M. Kini and S. Valiyaveettil. 2002. Investigation of the role of ansocalcin in the biomineralization in goose eggshell matrix. *Proc. Natl. Acad. Sci. U.S.A.* 99:5155-5159.
- Lavelin, I., N. Meiri, M. Einat, O. Genina and M. Pines. 2002. Mechanical strain regulation of the chicken glypican-4 gene expression in the avian eggshell gland. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 283:R853-R861.
- Lavelin, I., N. Yarden, S. Ben-Bassat, A. Bar and M. Pines. 1998. Regulation of osteopontin gene expression during egg shell formation in the laying hen by mechanical strain. *Matrix Biol.* 17:615-623.
- Layman, W. S., E. A. Hurd and D. M. Martin. 2011. Reproductive dysfunction and decreased GnRH neurogenesis in a mouse model of CHARGE syndrome. *Hum. Mol. Genet.* 20:3138-3150.
- Lerner, F., M. Niere, A. Ludwig and M. Ziegler. 2001. Structural and functional characterization of human NAD kinase. *Biochem. Biophys. Res. Commun.* 288:69-74.
- Mann, K. 1999. Isolation of a glycosylated form of the chicken eggshell protein ovocleidin and determination of the glycosylation site. *Alternative glycosylation/phosphorylation at an N-glycosylation sequon.* *FEBS Lett.* 463:12-14.
- Mann, K., J. Gautron, Y. Nys, M. D. McKee, T. Bajari, W. J. Schneider and M. T. Hincke. 2003. Disulfide-linked heterodimeric clusterin is a component of the chicken eggshell matrix and egg white. *Matrix Biol.* 22:397-407.
- Mann, K., B. Maček and J. V. Olsen. 2006. Proteomic analysis of the acid-soluble organic matrix of the chicken calcified eggshell layer. *Proteomics.* 6:3801-3810.
- Mann, K., J. V. Olsen, B. Maček, F. Gnad and M. Mann. 2007. Phosphoproteins of the chicken eggshell calcified layer. *Proteomics.* 7:106-115.
- Nishii, N., N. Nejime, C. Yamauchi, N. Yanai, K. Shinozuka and T. Nakabayashi. 2009. Effects of ATP on the intracellular calcium level in the osteoblastic TBR31-2 cell line. *Biol. Pharm. Bull.* 32:18-23.
- Nys, Y., J. Gautron, J. M. Garcia-Ruiz and M. T. Hincke. 2004. Avian eggshell mineralization: biochemical and functional characterization of matrix proteins. *C. R. Palevol.* 3:549-562.
- Ohira, K. and M. Hayashi. 2009. A new aspect of the TrkB signaling pathway in neural plasticity. *Curr. Neuropharmacol.* 7:276-285.
- Orriss, I. R., J. C. Utting, A. Brandao-Burch, K. Colston, B. R. Grubb, G. Burnstock and T. R. Arnett. 2007. Extracellular nucleotides block bone mineralization *in vitro*: Evidence for dual inhibitory mechanisms involving both P2Y2 receptors and pyrophosphate. *Endocrinology* 148:4208-4216.
- Peluffo, M. C., M. J. Murphy, S. T. Baughman, R. L. Stouffer and J. D. Hennebold. 2011. Systematic analysis of protease gene expression in the rhesus macaque ovulatory follicle: metalloproteinase involvement in follicle rupture. *Endocrinology* 152:3963-3974.
- Pines, M., V. Knopov and A. Bar. 1995. Involvement of osteopontin in egg shell formation in the laying chicken. *Matrix Biol.* 14:765-771.
- Pollak, N., M. Niere and M. Ziegler. 2007. NAD kinase levels control the NADPH concentration in human cells. *J. Biol. Chem.* 282:33562-33571.
- Reyes-Grajeda, J. P., A. Moreno and A. Romero. 2004. Crystal structure of ovocleidin-17, a major protein of the calcified Gallus gallus eggshell. *J. Biol. Chem.* 279:40876-40881.
- Ramakers, C., J. M. Ruijter, R. H. Deprez, A. F. Moorman. 2003. Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci. Lett.* 339:62-66.
- Rhinn, M., A. Dierich, W. Shawlot, R. R. Behringer, M. Le Meur and S. L. Ang. 1998. Sequential roles for Otx2 in visceral endoderm and neuroectoderm for forebrain and midbrain induction and specification. *Development.* 125:845-856.
- Ruijter, J. M., C. Ramakers, W. M. Hoogaars, Y. Karlen, O. Bakker, M. J. van den Hoff, A. F. Moorman. 2009. Amplification efficiency: linking baseline and bias in the analysis of quantitative PCR data. *Nucleic Acids Res.* 37:e45-e45.
- Sauveur, B. and M. Reviere. 1988. *Reproduction des volailles et production d'œufs.* INRA Editions, Paris.
- Solomon, S. E. 1999. Gordon Memorial Lecture. An egg ist ein ei,

- es un huevo, est un oeuf. *Br. Poult. Sci.* 40:5-11.
- Sutherland, C., B. S. Renaux, D. J. Mckay and M. P. Walsh. 1994. Phosphorylation of caldesmon by smooth-muscle casein kinase II. *J. Muscle Res. Cell Motil.* 15:440-456.
- Umezu, T., H. Yamanouchi, Y. Iida, M. Miura and Y. Tomooka. 2010. Follistatin-like-1, a diffusible mesenchymal factor determines the fate of epithelium. *Proc. Natl. Acad. Sci. USA.* 107:4601-4606.
- Veis, A. 1989. Chemical and biochemical perspectives. in: *Biom mineralization* (Ed. S. Mann, J. Webb and R. J. P. Williams) pp. 189. VCH, Weinheim, New York.
- Yang, K. T., C. Y. Lin, J. S. Liou, Y. H. Fan, S. H. Chiou, C. W. Huang, C. P. Wu, E. C. Lin, C. F. Chen, Y. P. Lee, W. C. Lee, S. T. Ding, W. T. Cheng and M. C. Huang. 2007. Differentially expressed transcripts in shell glands from low and high egg production strains of chickens using cDNA microarrays. *Anim. Reprod. Sci.* 101:113-124.
- Yoshizaki, K., S. Yamamoto, A. Yamada, K. Yuasa, T. Iwamoto, E. Fukumoto, H. Harada, M. Saito, A. Nakasima, K. Nonaka, Y. Yamada and S. Fukumoto. 2008. Neurotrophic factor neurotrophin-4 regulates ameloblastin expression via full-length TrkB. *J. Biol. Chem.* 283:3385-3391.
- Zheng, Q. and X. J. Wang. 2008. GOEAST: a web-based software toolkit for Gene Ontology enrichment analysis. *Nucleic Acids Res.* 36:358-363.
- Zmuda, J. M., L. M. Yerges-Armstrong, S. P. Moffett, L. Klei, C. M. Kammerer, K. Roeder, J. A. Cauley, A. Kuipers, K. E. Ensrud, C. S. Nestlerode, A. R. Hoffman, C. E. Lewis, T. F. Lang, E. Barrett-Connor, R. E. Ferrell, E. S. Orwoll and Osteoporotic Fractures in Men (MrOS) Study Group. 2011. Genetic analysis of vertebral trabecular bone density and cross-sectional area in older men. *Osteoporos Int.* 22:1079-1090.