

# Pneumopericardium as a complication of pericardiocentesis


Andreas S. Triantafyllis MD PhD, Tatiana Zamfir MD, Nestoras Kontogiannis MD

■ Cite as: *CMAJ* 2023 February 6;195:E193-4. doi: 10.1503/cmaj.221137

A 48-year-old man with trisomy 21 was transferred to our cardiology service because of acute dyspnea. At presentation, his blood pressure was 70/50 mmHg, with a heart rate of 130 beats/min and an oxygen saturation of 88%. Physical examination showed muffled heart sounds, clear lungs and distention of the jugular veins. An echocardiogram showed a large circumferential pericardial effusion.

We performed fluoroscopy-guided pericardiocentesis, under local anesthesia, through the sub-xiphoid approach and inserted a pigtail catheter (8Fr). Drainage of 1000 mL of transudate resulted in rapid clinical improvement. A chest radiograph at 24 hours showed air entrapment between the cardiac silhouette and the pericardium, without signs of pneumothorax (Figure 1). A repeat echocardiogram showed bright echogenic spots swirling in the pericardial cavity, impairing visualization of cardiac structures (video Appendix 1, available at [www.cmaj.ca/lookup/doi/10.1503/cmaj.221137/tab-related-content](http://www.cmaj.ca/lookup/doi/10.1503/cmaj.221137/tab-related-content)). We made the diagnosis of pneumopericardium. Because he was hemodynamically stable, we treated him conservatively and the pericardial air gradually resolved. Diagnostic work-up to identify a cause of his pleural effusion was negative; therefore, we considered the pericardial effusion to be idiopathic.

Pneumopericardium after pericardiocentesis is an uncommon complication that results from air insertion through the pericardial drainage or formation of a direct pleuropericardial communication. The diagnosis is made by chest radiography, computed tomography or echocardiography.<sup>1</sup> Differential diagnosis includes pneumomediastinum, in which air does not typically surround the bottom of the heart and is not confined solely to the heart, but can extend to the superior mediastinum and neck.<sup>2</sup> In our case, pneumopericardium was likely induced by coughing in combination with an unsealed system. Avoidance of vigorous inhalation or coughing, confirmation of proper drain position (i.e., no side holes outside the pericardial space), closure of all drainage system connections and application of surgical gel to the skin around the drain to create an airtight seal are essential to prevent this complication.<sup>3</sup> Pneumopericardium is



**Figure 1:** Chest radiograph of a 48-year-old man after pericardiocentesis, showing air entrapment (asterisk) between the cardiac silhouette and the thin, radiolucent line of the pericardium (arrows).

usually self-limiting and requires no specific therapy. However, the development of cardiac tamponade necessitates repeat pericardiocentesis or surgical decompression.<sup>1</sup>

## References

1. Choi WH, Hwang YM, Park MY, et al. Pneumopericardium as a complication of pericardiocentesis. *Korean Circ J* 2011;41:280-2.
2. Brander L, Ramsay D, Dreier D, et al. Continuous left hemidiaphragm sign revisited: a case of spontaneous pneumopericardium and literature review. *Heart* 2002;88:e5.
3. Narins CR, Lee J, Cole M, et al. Pneumopericardium following pericardiocentesis. *Am J Med* 2016;129:e181-2.

**Competing interests:** None declared.

This article has been peer reviewed.

The authors have obtained patient consent.

**Affiliation:** Department of Cardiology, Asklepeion General Hospital, Athens, Greece

**Content licence:** This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY-NC-ND 4.0) licence, which permits use, distribution and reproduction in any medium, provided that the original publication is properly cited, the use is noncommercial (i.e., research or educational use), and no modifications or adaptations are made. See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

**Correspondence to:** Andreas Triantafyllis, [andreas.triantafyllis@gmail.com](mailto:andreas.triantafyllis@gmail.com)



A video of the echocardiogram of a 48-year-old man after pericardiocentesis, showing bright echogenic spots swirling in the pericardial cavity and impairing visualization of cardiac structures, is in Appendix 1, available at [www.cmaj.ca/lookup/doi/10.1503/cmaj.221137/tab-related-content](http://www.cmaj.ca/lookup/doi/10.1503/cmaj.221137/tab-related-content)

Clinical images are chosen because they are particularly intriguing, classic or dramatic. Submissions of clear, appropriately labelled high-resolution images must be accompanied by a figure caption. A brief explanation (300 words maximum) of the educational importance of the images with minimal references is required. The patient's written consent for publication must be obtained before submission.