


BMJ Open Investigating the risks of cardiovascular disease among premenopausal women using oral contraceptive: a protocol for a systematic review and meta-analysis

Oyesanmi A Fabunmi ,^{1,2} Phiwayinkosi V Dlodla,³ Siphamandla R Ngcobo,¹ Bongani B Nkambule ¹

To cite: Fabunmi OA, Dlodla PV, Ngcobo SR, *et al.* Investigating the risks of cardiovascular disease among premenopausal women using oral contraceptive: a protocol for a systematic review and meta-analysis. *BMJ Open* 2023;**13**:e071118. doi:10.1136/bmjopen-2022-071118

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-071118>).

Received 16 December 2022
Accepted 03 January 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Human Physiology, School of Laboratory Medicine and Medical Sciences (SLMMS), College of Health Sciences, University of KwaZulu-Natal, Durban, South Africa

²Human Physiology, College of Medicine, Ekiti State University, Ado Ekiti, Nigeria

³Biomedical Research and Innovation Platform, South African Medical Research Council, Tygerberg, South Africa

Correspondence to

Professor Bongani B Nkambule; nkambuleb@ukzn.ac.za

ABSTRACT

Introduction The use of oral contraceptives (OCs) is linked to an increased risk of cardiovascular diseases (CVDs) in women of reproductive age. CVD remain one of the top causes of death worldwide, with at least three-quarters of deaths occurring in low-income and middle-income nations. The impact of various types of combined oral contraceptive (COC) on several modifiable risk factors associated with CVDs in premenopausal women is inconsistent regardless of genetic mutations. The aim of this systematic review will be to provide a comprehensive synthesis of the available evidence on the impact of COC usage on modifiable risk factors associated with CVDs and assess ethnic and geographic disparities in the reported prevalence of CVD.

Methods and analysis This systematic review protocol was prepared in accordance with the Preferred Reporting Items for Systematic reviews and Meta-Analyses Protocols 2015 statement. An extensive search on the Embase, MEDLINE and Cochrane Library will be conducted from inception until. Two reviewers will independently screen for eligible studies using a predefined criterion. The risk of bias and quality of included studies will be assessed using the modified Downs and Black's checklist. Whereas the overall quality of included studies will be assessed using the Grading of Recommendations, Assessment, Development and Evaluation assessment tool.

Ethics and dissemination This is a review of existing studies and will not require ethical approval. The findings will be disseminated through peer-reviewed publication. The use of OC and the risk of CVDs including arterial and venous thrombosis remain a major concern among women of reproductive age. Thus, given the impact of COCs on the risk variables linked with CVDs, this review may provide an insight and assistance during COC use.

PROSPERO registration number CRD42020216169.

INTRODUCTION

Women are exposed to estrogen fluctuations throughout their life which can either endogenous or exogenous in nature.¹ The use of exogenous combined oral contraceptives (COCs) remains one of the most commonly used modern methods of birth control due to their high efficacy and safety profile in

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This review will be a comprehensive synthesis of available evidence on the link between oral contraceptive (OC) usage and the risk of cardiovascular diseases (CVDs).
- ⇒ A comprehensive search strategy will be used to identify and retrieve articles relevant to our research question.
- ⇒ This will be the first systematic review and meta-analysis to assess the geographic disparities in the prevalence of CVD linked to OC usage.
- ⇒ The main limitation will be the heterogeneity of the studies available in terms of clinical, methodological, type of study and reported outcomes.

premenopausal women.² The prevalence of COC use in women of reproductive age is around 16% representing around 151 million women worldwide, and in developed countries this accounts for over 30%³⁻⁴, despite certain risk identified to be associated with the usage.^{2,5}

The use of oral contraceptives (OCs) is associated with an increased risk of cardiovascular events^{6,7} such as coronary heart disease, arterial and venous thrombosis, ischaemic or haemorrhagic stroke, and myocardial infarction among others.⁸ Although, these cardiovascular events appear to differ by geographic region, age and gender.^{9,10} The risk for CVD increases when other modifiable cardiovascular factors, such as smoking, obesity, dyslipidaemia, diabetes mellitus, haemostatic disorders and hypertension, are considered.¹¹

Notably, available data regarding the impact of COC on modifiable risk factors associated with cardiovascular diseases (CVDs) in women of reproductive age in different population settings are inconsistent.¹²⁻¹⁴ For instance, findings from an epidemiological study reported changes in levels of systolic blood pressure and diastolic blood pressure

following the administration of OCs.¹³ In contrast, the study by Kharbanda *et al*, observing the cardiovascular effects of COCs in adolescents, reported no changes in blood pressure.¹²

Furthermore, findings from previous studies investigating the impact of COC on haemostatic parameters are also inconsistent.^{5 15} Nevertheless, the impact of COC on the modifiable risk factors associated with CVDs is usually attributed to the direct influence of the oestrogen component, which is known to impact the vascular wall and stimulate endothelial dysfunction, as well as altering the coagulation system.^{2 16} The impact of the oestrogen appears to be countered by the progesterone component, which may be dependent on the dose and duration of use.^{2 17} Prior studies have also revealed the effect of COCs on metabolic parameters, such as changes in lipid profiles,^{14 18} and insulin sensitivity, all of which play a role in the development of a prothrombotic condition.^{19 20}

In spite of high efficacy and safety profile attributed to OCs, data available regarding the impact on the occurrence of atherosclerosis, thrombosis, vasomotion and arrhythmogenesis among others are conflicting and insufficient.²¹ More so, evidence are currently limited as regard the role of geographical areas and ethnicity on the association between the use of OC and risk of CVDs,²² although emerging evidence suggest that ethnic differences can influence CVD risk even at a young age, with pronounced differences in some modifiable risk factors.⁹ The justification for exploring the effect of OCs on cardiovascular events is simply due to the fact that the prolong use of COC and risk of CVD in women of reproductive age still remain an important issue despite premenopausal women having low risk.^{2 23} Thus, determining the impact of COC usage on certain modifiable risk factors associated with CVDs, will help to provide insight and guidance in making an informed decision at each contraceptive consultation (initial and follow-up) in different population settings. Therefore, the aim of this systematic review and meta-analysis will be to provide a comprehensive synthesis of the available evidence on the impact of OC usage on modifiable risk factors associated with the occurrence of CVDs and to further assess the role of ethnic and geographic disparities in the reported prevalence of CVD in women on OCs.

Objectives

The objectives of this study are as follows:

1. To determine the prevalence of CVDs in premenopausal women using OC.
2. To determine the effect of OC on vascular and cellular markers of coagulation in association with immune cell activation in premenopausal women on OC.

METHODS

The systemic review protocol was prepared according to the Preferred Reporting Items for Systemic review and Meta-Analysis Protocol.

Patient and public involvement

There will be no patient and public involvement in the design, interpretation or dissemination of the findings.

Participants

The participants of the systematic review will mainly include healthy adolescent and adult premenopausal women between the age of 18 and 45 years who are not obese and non-smokers.

Exposure

This systematic review will include studies that report on the use of second-generation, third-generation and fourth-generation type of COC.

Comparators

The comparator will include healthy adolescent and adult women who not OCs.

Inclusion and exclusion criteria

This systematic review and meta-analysis will include cross-sectional, cohort, case-control studies and randomised control trials. Studies reporting on the effect of COC usage as form of hormonal therapy or as a method of contraception on the risk factors associated with CVD in healthy premenopausal women who are non-obese, and non-smokers will be included. There will be no language restriction. Reviews, books, letters to editors and studies will be excluded.

Outcomes

The study outcomes will include the following.

Primary outcome

Prevalence or incidence of cardiovascular events.

Surrogate outcome

1. Vascular and cellular markers of coagulation and endothelia dysfunction (endothelin 1, D-dimer, NO, fibrinogen and proinflammatory cytokines (TNF- α , IL-6, C-reactive protein).
2. Traditional cardiovascular risk variables (BMI, systolic and diastolic blood pressure, lipid profile, glucose intolerance and hyperinsulinaemia).

Information sources

A search strategy will be developed using medical subject headings (MeSH) words for MEDLINE, EMBASE and Google Scholar database. The electronic database search will be augmented by searching the Cochrane Central Register of Clinical trials. Papers published in both commercial and in non-commercially operated databases (grey literature) will also be considered. In addition, the reference list of the selected studies will be scanned to identify relevant literature.

Search strategy

The search strategy will be developed using MeSH terms and keywords related to OCs, CVD and premenopausal (online supplemental file). The keywords and

MeSH terms will include OC pills, birth control pills or OCs or contraceptives, premenopausal women, CVD, hypertension, or coronary heart disease.

Study selection

The screening of studies will be conducted by two independent authors (OAF and SRN) to avoid inconsistency in terms of eligibility of studies to enhance objectivity and prevent mistakes. At the initial stage, studies will be screened by the titles, abstracts, keywords and synonyms, then followed by the identification of the full-text articles. Should discrepancies arise (PD) will screen such studies, and consensus will be reached through discussion.

Data management

The Mendeley desktop reference manager (V.1.19.4) will be used to manage retrieved studies, archiving of relevant and excluded studies with reasons. Importantly, reference lists of included studies will be screened to confirm that no relevant studies are left out. Eligible will then be subjected to data collection, critical appraisal, risk and quality evaluation.

Data items and collection process

Relevant data items will be extracted using a structured form depending on the type of study design. To avoid errors during data entry from selected studies, two authors (OAF and SRN) will independently perform this process. Should discrepancies arise, the third and fourth authors (PVD) will be invited for arbitration. The author and year of publication, the country, population (sample size), study design, types, dosage and duration of contraceptive usage, plasma levels of cellular and vascular markers of coagulation, endothelia dysfunction and inflammation or immune activation, lipid profiles, glucose profiles, body mass index (BMI) and haemodynamic indices of hormonal contraceptive users and non-users will be extracted. In case of insufficient data, the main authors of studies will be contacted to obtain enough information.

Risk of bias in individual studies

The risk of bias and quality of included studies will be assessed using the modified Downs and Black's checklist.²⁴ Two reviewers (OAF and BBN) will make independent judgements based on the domains of the tool: reporting bias (10 items), external validity (3 items), internal validity (6 items) and selection bias (7 items). The scores will be rated as excellent (25–26), good (20–24), moderate (14–19), poor (11–13) and very poor (<10). In case of disagreements, PVD will be consulted to arbitrate.

Data synthesis

Review Manager (RevMan) V.5.3 (The Nordic Cochrane Centre, The Cochrane Collaboration, 2014) will be used for statistical analysis. Clinical and methodological heterogeneity assessment will be conducted

first which will then be followed by an assessment of statistical heterogeneity. The χ^2 and I^2 statistic tests will be used to determine the level of heterogeneity across the included studies. An I^2 value of >25 will be considered as moderate or substantial heterogeneity.²⁵

If studies are similar with regard to (participants, intervention, comparisons and outcomes), and more so, information extracted from this sufficient number of studies included are homogeneous, we will conduct a fixed-effect meta-analysis using R statistical Software (The R foundation for statistical computing, Vienna, Austria). A random-effects model will be employed where studies show a substantial level of heterogeneity. Furthermore, a subgroup analysis base on the reported outcome and meta-regression will be conducted to assess the role of geographical areas and ethnicity in association with the use of COC and risk of CVD. The levels of inter-rater agreement will be assessed using Cohen's kappa²⁶ and the funnel plots will be used to assess publication bias.

Quality cumulative evidence

The quality of evidence for primary outcomes will also be evaluated using the grading of recommendations assessment, development and evaluation tool.²⁷ The findings will be summarised and presented in the summary of findings table.

Sensitivity analysis

The sources of statistical heterogeneity will be assessed by performing a sensitivity analysis and excluding studies with high risk of bias based on the risk of bias assessment.

Ethics and dissemination

This is a review of existing studies and will not require ethical approval. The findings will be disseminated through peer-reviewed publication. The use of OC and the risk of CVDs including arterial and venous thrombosis remain a significant concern among premenopausal women. Thus, given the impact of COCs on the risk variables associated with CVDs, this review may provide an insight and assistance during COC use.

Twitter Oyesanmi A Fabunmi @Sormaytion and Bongani B Nkambule @Nkambuleb

Contributors OAF and BBN conceptualised and designed the study. OAF drafted the protocol. SRN and PVD reviewed the drafted the protocol. All authors wrote and approved the final manuscript. BBN is the guarantor of the review.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting or dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; peer reviewed for ethical and funding approval prior to submission.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been

peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Oyesanmi A Fabunmi <http://orcid.org/0000-0002-3005-331X>

Bongani B Nkambule <http://orcid.org/0000-0001-8846-1992>

REFERENCES

- Gialeraki A, Valsami S, Pittaras T, et al. Oral contraceptives and HRT risk of thrombosis. *Clin Appl Thromb Hemost* 2018;24:217–25.
- Williams JS, MacDonald MJ. Influence of hormonal contraceptives on peripheral vascular function and structure in premenopausal females: a review. *Am J Physiol Heart Circ Physiol* 2021;320:H77–89.
- Christin-Maitre S. History of oral contraceptive drugs and their use worldwide. *Best Pract Res Clin Endocrinol Metab* 2013;27:3–12.
- Population Division UN. Contraceptive use by method 2019: data booklet; 2019: 25.
- Hee L, Kettner LO, Vejtorp M. Continuous use of oral contraceptives: an overview of effects and side-effects. *Acta Obstet Gynecol Scand* 2013;92:125–36.
- Nisenbaum MG, Melo NRde, Giribela CRG, et al. Effects of a contraceptive containing drospirenone and ethinyl estradiol on blood pressure and autonomic tone: a prospective controlled clinical trial. *Eur J Obstet Gynecol Reprod Biol* 2014;175:62–6.
- Khialani D, Rosendaal F, Vlieg Avanh. Hormonal contraceptives and the risk of venous thrombosis. *Semin Thromb Hemost* 2020;46:865–71.
- Raps M, Rosendaal F, Ballieux B, et al. Resistance to APC and SHBG levels during use of a four-phasic oral contraceptive containing dienogest and estradiol valerate: a randomized controlled trial. *J Thromb Haemost* 2013;11:855–61.
- Virani SS, Alonso A, Aparicio HJ, et al. Heart disease and stroke Statistics-2021 update: a report from the American heart association. *Circulation* 2021;143:E254–743.
- Centers for Disease Control and Prevention. Data and statistics on venous thromboembolism, 2020. Available: <https://www.cdc.gov/ncbddd/dvt/data.html> [Accessed 27 Oct 2021].
- Rohla M, Weiss TW. Metabolic syndrome, inflammation and atherothrombosis. *Hamostaseologie* 2013;33:283–94.
- Kharbada EO, Parker ED, Sinaiko AR, et al. Initiation of Oral Contraceptives and Changes in Blood Pressure and Body Mass Index in Healthy Adolescents. *J Pediatr* 2014;165:1029–33.
- Park H, Kim K. Associations between oral contraceptive use and risks of hypertension and prehypertension in a cross-sectional study of Korean women. *BMC Womens Health* 2013;13:1–7.
- Farahmand M, Ramezani Tehrani F, Rostami Dovom M, et al. The impact of oral contraceptives on cardiometabolic parameters. *J Endocrinol Invest* 2016;39:277–83.
- Stocco B, Fumagalli HF, Franceschini SA. Comparative study of the effects of combined oral contraceptives in hemostatic variables: an observational preliminary study. *Med* 2015;94.
- Heidarzadeh Z, Asadi B, Saadatnia M, et al. The Effect of Low-dose Combined Oral Contraceptive Pills on Brachial Artery Endothelial Function and Common Carotid Artery Intima-Media Thickness. *Journal of Stroke and Cerebrovascular Diseases* 2014;23:675–80.
- Hirschberg AL. Sex hormones, appetite and eating behaviour in women. *Maturitas* 2012;71:248–56.
- Bakesiima R, Byakika-Kibwika P, Tumwine JK, et al. Dyslipidaemias in women using hormonal contraceptives: a cross sectional study in Mulago Hospital family planning clinic, Kampala, Uganda. *BMJ Open* 2018;8:e022338.
- Paneni F, Beckman JA, Creager MA, et al. Diabetes and vascular disease: pathophysiology, clinical consequences, and medical therapy: Part I. *Eur Heart J* 2013;34:2436–43.
- Vazzana N, Ranalli P, Cucurullo C, et al. Diabetes mellitus and thrombosis. *Thromb Res* 2012;129:371–7.
- Zakharova MY, Meyer RM, Brandy KR, et al. Risk factors for heart attack, stroke, and venous thrombosis associated with hormonal contraceptive use. *Clin Appl Thromb Hemost* 2011;17:323–31.
- Afshari M, Alizadeh-Navaei R, Moosazadeh M. Oral contraceptives and hypertension in women: results of the enrolment phase of Tabari cohort study. *BMC Womens Health* 2021;21.
- Mohamed ABO, Al-Ama N, Al Kreathy H, et al. Oral contraceptive types in relation to ABO blood groups among Saudi women of different reproductive age groups and impact on venous thromboembolism. *Clin Appl Thromb Hemost* 2020;26:107602962096605.
- Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* 1998;52:377–84.
- Imrey PB. Limitations of meta-analyses of studies with high heterogeneity. *JAMA Netw Open* 2020;3:e1919325.
- McHugh ML. Interrater reliability: the kappa statistic. *Biochem Med* 2012;22:276–82.
- Ryan R, Hill S. Supporting implementation of Cochrane methods in complex communication reviews: resources developed and lessons learned for editorial practice and policy. *Health Res Policy Sys* 2019;17.