# Comments on: Comparison of an aspheric monofocal intraocular lens with the new generation monofocal lens using defocus curve

#### Dear Editor,

As we all are aware that due to rapid evolution and technical advancements in the past decade, ophthalmic science has come a long way. The same can be seen in the advancement of various IOL designs and models. After the launch of aspherical IOLs in 2001, there have been only minor improvements in the optical performance of monofocal IOLs. With the introduction of the TECNIS Eyhance IOL, model ICB00,<sup>[1]</sup> from Johnson & Johnson Vision, the changes can be rapidly appreciated in near future. We read the article by Yangzes *et al.*<sup>[2]</sup> in the December 2020 issue of IJO and we must congratulate the authors for taking a leap forward and compare two recently launched newer aspheric monofocal IOL's. However, we have a few important observations and suggestions to make.

Firstly, the authors have mentioned the study as "observational consecutive case series". But since all patients underwent IOL implantation and it was done from September to November 2019, the better terminology will be Prospective, interventional study. Can the authors throw some light over this for the benefit of the readers.

Secondly, it is not clear how the patients were allocated in two groups? The first group had 71 patients and the second group had 45 patients. What rationale did the authors use to divide the patients? This needs clarification. Ideally, the patients could have been randomized to two equal groups to avoid bias and compare the IOLs on similar grounds.

Thirdly, the exclusion criteria are missing. It would be really interesting to know what all patients were excluded from the study.

Lastly, was contrast sensitivity (CS)<sup>[3]</sup> was taken into account for the two IOL's? This could have been really interesting to know to compare the two IOL CS under mesopic and photopic conditions catering to patient satisfaction and final outcome.

#### Acknowledgements

Aravind Eye Hospital and Post Graduate Institute of Ophthalmology, Pondicherry.

Financial support and sponsorship Nil.

## **Conflicts of interest**

There are no conflicts of interest

## Bharat Gurnani, Kirandeep Kaur<sup>1</sup>

Consultant Cornea and Refractive Services, <sup>1</sup>Consultant Pediatric and Squint Services, Aravind Eye Hospital and Post Graduate Institute of Ophthalmology, Pondicherry, India

Correspondence to: Dr. Bharat Gurnani, Consultant Cornea and Refractive Services, Aravind Eye Hospital and Post Graduate Institute of Ophthalmology, Cuddalore Main Road, Thavalukuppam, Pondicherry - 605 007, India. E-mail: drgurnanibharat25@gmail.com

## References

- 1. Tognetto D, Cecchini P, Giglio R. Surface profiles of new-generation IOLs with improved intermediate vision. J Cataract Refract Surg 2020;46:902-6.
- Yangzes S, Kamble N, Grewal S. Comparison of an aspheric monofocal intraocular lens with the new generation monofocal lens using defocus curve. Indian J Ophthalmol 2020;68:3025-9.
- Haaskjold E, Allen ED, Burton RL, Webber SK, Sandvig KU, Jyrkkiö H, et al. Contrast sensitivity after implantation of diffractive bifocal and monofocal intraocular lenses. J Cataract Refract Surg 1998;24:653-8.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code:	Website:
	www.ijo.in
	<b>DOI:</b> 10.4103/ijo.IJO_305_21

**Cite this article as:** Gurnani B, Kaur K. Comments on: Comparison of an aspheric monofocal intraocular lens with the new generation monofocal lens using defocus curve. Indian J Ophthalmol 2021;69:1348.

© 2021 Indian Journal of Ophthalmology | Published by Wolters Kluwer - Medknow