

# Novel Genetic Variant in *HUWE1*

## Prenatal and Postnatal Neuroimaging Phenotype

Mario Tortora, MD, Elisa Cattaneo, MD, Luigina Spaccini, MD, Maria Iascone, PhD, Barbara Scelsa, MD, Alessia Micalizzi, MD, Antonio Novelli, PhD, Mariano Lanna, MD, Andrea Righini, MD, Pierangelo Veggiotti, PhD, and Chiara Doneda, MD

### Correspondence

Dr. Tortora  
mario.tortora@gmail.com

*Neurol Genet* 2024;10:e200169. doi:10.1212/NXG.000000000200169

## Abstract

### Objectives

To provide a comprehensive description of neuroradiologic findings in a patient with a probable pathogenic variant of *HUWE1*, particularly in relation to pontine and cerebellar hypoplasia.

### Methods

We first report prenatal and postnatal neuroradiologic phenotype of a female patient carrying a *HUWE1* likely pathogenic variant and discuss its function.

### Results

An ultrasound shows borderline ventriculomegaly, rotated cerebellar vermis, and dysgenetic corpus callosum. An MR study identifies a short, thin corpus callosum, falcine sinus persistence, reduced cerebellar vermis size, wide inferior IV ventricle, and reduced pontine bulging.

### Discussion

*HUWE1* is a gene encoding an E3 ubiquitin ligase protein involved in nervous system development, function, and disease. The mechanisms by which *HUWE1* gene affects nervous system are still largely unclear, but a growing body of literature described disease-causing variants in this gene. This report may help prenatal diagnostic experts in consider also this entity, especially when dealing with pontine and cerebellar hypoplasia findings.

## Introduction

*HUWE1* is a gene encoding an E3 ubiquitin ligase protein, involved in cell growth and death regulation, tumor development, DNA damage response, CNS function, and inflammasome activation.<sup>1-3</sup> Gene copy number variants and missense mutations have been reported in individuals with syndromic and nonsyndromic X-linked intellectual disability, possibly associated with autism, and can cause minor dysmorphic CNS or facial features, detected through MRI.<sup>3-6</sup> Most patients previously described have no available neuroimaging data, and most who underwent brain MR imaging had a normal examination.<sup>5,7</sup>

We report a fetal and postnatal case in which a novel genetic variant in *HUWE1* gene was identified (c. 273 G>C, p.Met91Ile), aiding prenatal diagnostic experts in recognizing this entity, particularly in cases of pontine and cerebellar hypoplasia.

---

From the Department of Pediatric Radiology and Neuroradiology (M.T., A.R., C.D.); Clinical Genetics Unit (E.C.), Department of Pediatrics; Clinical Genetics Unit (L.S.), Department of Obstetrics and Gynecology, "Vittore Buzzi" Children's Hospital - ASST Fatebenefratelli-Sacco, Milan; Medical Genetics Laboratory (M.I.), Hospital Papa Giovanni XXIII, Bergamo; Pediatric Neurology Unit (B.S.), Department of Pediatrics, "Vittore Buzzi" Children's Hospital - ASST Fatebenefratelli-Sacco, Milan; Translational Cytogenomics Research Unit (A.M., A.N.), IRCCS Bambino Gesù Pediatric Hospital, Rome; Fetal Therapy Unit "U. Nicolini" (M.L.), Department of Obstetrics and Gynecology, Buzzi Children's Hospital, Milan; and Pediatric Neurology Unit (P.V.), "Vittore Buzzi" Children's Hospital, Department of Biomedical and Clinical Sciences "L. Sacco", University of Milan, Italy.

Go to [Neurology.org/NG](https://www.neurology.org/NG) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Copyright © 2024 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.


## Case Description

A 43-year-old woman with 3 previous abortions, no consanguinity, no perinatal infections, or known teratogenic risk medications was referred to our maternal fetal unit. An ultrasound examination at 20 weeks showed borderline ventriculomegaly, rotated and hypoplastic cerebellar vermis and dysgenetic corpus callosum. Fetal MRI at 21 weeks identified a short and thin corpus callosum with mild ventriculomegaly (atrium: 11 mm), falcine sinus persistence; cerebellar vermis was reduced in size compared with normal parameters (superior-inferior diameter: 7 mm vs 8 mm; anterior-posterior diameter: 4 mm vs 5 mm); inferior part of the IV ventricle was wide because of mild cranial rotation of the vermis; reduced pontine bulging was evident too<sup>8</sup> (Figures 1 and 2). NGS panel genes related to prenatal phenotype (cerebellar hypoplasia and corpus callosum hypoplasia) was performed on genomic DNA by using the Twist Custom Panel Kit clinical exome—Twist Bioscience according to the manufacturer's protocol on an NovaSeq6000 platform (Illumina). The reads were aligned to human genome build GRCh37/UCSC hg19. The BaseSpace pipeline and the GeneX software (LifeMap Sciences, Inc.) were used for the variants' calling and annotating, respectively. Putative disease-associated variants were filtered for an allele frequency <1% in Genome Aggregation Database (gnomAD) and were checked based on the public databases (dbSNP), Exome Aggregation Consortium (ExAC) and gnomAD. The variants were evaluated by VarSome and categorized in accordance with the ACMG recommendations.<sup>9</sup> Variants were examined for a minimum depth coverage of 30X and Qscore (minimum threshold of 30) and visualized by the Integrative Genome Viewer. NGS-based trio test allowed to identify a novel de novo heterozygous missense variant c.273G>C (p.Met911Ile) in the *HUWE1* gene (NM\_031407)

(eFigure 1). The missense change is not present in the reference population database (dbSNP), ExAC, and gnomAD and is not previously described in literature. It involves a highly conserved residue across species and is predicted to be deleterious (scaled CADD score 21.70) and can be classified a likely pathogenic variant (class 4 according to the ACMG criteria). We did not perform the X inactivation study on the amniocytes. The couple opted for continuation of pregnancy. Fetal MR imaging, repeated at 28 weeks, revealed dysmorphic ventricular system because of the abnormal corpus callosum, thin and short. Biometric parameters of the cerebellar vermis, cranially rotated, appeared at lower limits for GA (SI diameter: 13 mm; AP diameter: 9 mm). Reduced pontine bulging was detected with smaller cranio-caudal diameter of the pons (Figure 1). The III-trimester of gestation was also complicated by severe polyhydramnios which required amniotic fluid drainage.


The patient was born at 38 + 1 weeks by elective cesarean section. Her birth parameters were as follows: birth weight 3,240 g (72° centile); length 47 cm (22° centile); occipito-frontal circumference 32 cm (12° centile). Apgar was 6/8 at 1st minute and 5th minute. Mild dysmorphic features were observed (thin upper lip, deep set eyes, and epicanthic folds). Neonatal MR imaging, at 4 days of life, confirmed pontocerebellar hypoplasia with significant cranial rotation of the cerebellar vermis and medulla oblongata gross appearance, reduced evidence of anterior fissure, dysgenesis of corpus callosum with reduction of AP diameter, and thickness of the medial and posterior portions. In addition, an occipital median skull defect at the level of tentorium insertion as an intracranial lipoma (Figure 2). At birth, heart ultrasound showed pulmonary hypertension with an estimated pulmonary artery pressure of 60–65 mm Hg. At 1 month, pulmonary pressure had turned

**Figure 1** Supratentorial Abnormalities in Fetal MRI



Midsagittal balance (A, E), coronal (B, C, F, G), and axial (D, H) T2-weighted Ssh sections from 21-week and 28-week prenatal study, respectively, showing thin and short corpus callosum (arrows), lateral ventricles dysmorphism, smaller and partially upward rotated cerebellar vermis (arrowheads), reduced pontine bulging (asterisks), persistent falcine sinus (empty arrow), and mild ventriculomegaly only in the first examination.

**Figure 2** Infratentorial Abnormalities in Fetal MRI




Midsagittal T2-weighted Ssh sections of our patient from 21-week (A) and 28-week (B) prenatal study showing smaller and partially upward rotated cerebellar vermis, reduced pontine bulging (dashed line), gross appearance of the medulla oblongata with poor representation of the groove at the border of the pons. Midsagittal T2-weighted Ssh sections of a normal fetus from 21-week (C) and 28-week (D) prenatal study showing, by comparison, normal appearance of the structures of the posterior cranial fossa.

into normal parameters. Auditory brainstem response was normal, while ophthalmologic evaluation showed a mild atrophy of optic papilla. The child developed hyperopia and strabismus at 6 months. While neurologic evaluation at birth was normal, the neurologic follow-up revealed hypotonia and neurodevelopmental impairment. She started crawling at 15 months and began walking with support at 20 months. No

seizures were reported, and subsequent EEG did not show any epileptic anomalies. Social interaction was initially reduced, but she currently attends daycare with improved interaction with peers. Expressive language is limited to one word, but she is able to point at objects. Standardized neurodevelopmental tests (Griffiths III) showed moderate intellectual disability (GQ74). As soon as she began to grow, it was noted a laryngospasm due

**Figure 3** Neonatal MRI Findings



Midsagittal T1-weighted and T2-weighted (A, E), coronal T2-weighted (B, C, F, G), and axial T2-weighted and T1-weighted (D, H); T2-weighted sections from 4 days and 9 months of age postnatal study, respectively, confirming previous findings as regard to corpus callosum (arrows) and ventricular dysmorphism and proportionally more evident vermian (arrowheads) and pontine (asterisks) hypoplasia. Gross appearance of the medulla oblongata and an occipital median skull defect at the level of tentorium insertion (empty arrows), with a T1-hyperintense linear structure that runned deep down to the torcular region, as an intracranial lipoma, are evident too. Moderate enlargement of frontal periencephalic CSF spaces was also noted in the second examination.

to food aspiration. The baby with persistent oropharyngeal dysphagia was fed through a nasogastric tube and percutaneous endoscopic gastrostomy at 3 months. Oral and swallowing skills are improving, but most food intake is through PEG. At 20 months, the baby's weight and length parameters were below normal, and she started pneumologic follow-up because of pulmonary dysventilation and food aspiration consequences. Several skeletal abnormalities were described, including bilateral rib dysmorphism with partial fusion defects (I and II ribs on the left and VIII and IX ribs on the right), overlapping toes, partial fusion of C2-C3 metameris, and butterfly vertebra at C4. The MR imaging follow-up at 8 months confirmed cranial findings (Figure 3) and excluded occult dysraphism.

## Discussion

The mechanisms by which *HUWE1* gene regulates neuronal function and development are still largely unclear, but a growing body of literature described disease-causing variants in this gene, especially among intellectual disability affected patients.<sup>2,3</sup> We first report extensive description of prenatal and postnatal neuroradiologic phenotype of a female patient carrying a *HUWE1* likely pathogenic variant and discuss its function. In literature, there is no article on genotype-phenotype correlation regarding this *HUWE1* variant. The role of *HUWE1* in brain and cerebellar development has been reported<sup>3,10-12</sup> with correlations to both neural tube defects and medulloblastoma development. The largest case series of patients was described by Moortgat et al.<sup>13</sup>

In addition to intellectual disability, minor face anomalies (long face, a short nose with a broad nasal tip, deep set eyes with epicanthic folds, blepharophimosis, thin upper lip, and full lower lip), and minor SNC anomalies, are described global hypotonia, behavioral disorder (autism spectrum disorder; stereotypies, hyperactivity), skeletal anomalies (short stature, small hands and feet, craniosynostosis, scoliosis, contractures of knees), postnatal microcephaly, and seizure. Less frequently are reported hearing loss, sleep disorder, and feeding difficulties (constipation).<sup>13,14</sup>

Among the described *HUWE1*-related neuroimaging alterations there are undefined focal cortical malformations; thin corpus callosum; delayed myelination; ventriculomegaly; microcephaly and macrocephaly; isolated cerebellar atrophy; hippocampi of rounded and medialized shape; splenium thickening of corpus callosum; filum terminalis lipoma; unspecific hypoxic-ischemic damage.

In our case, the cerebellar vermis and pontine involvement was one of the key features of the prenatal and postnatal assessment. *HUWE1* plays a key role in the differentiation of cerebellar granule neuron precursors (CGNPs) and the maturation of Bergmann glia during cerebellar development.<sup>10</sup> *HUWE1* rules the interaction between the Bergmann glial scaffold and granule neurons, which is essential for granule neuron migration and IGL formation. Mutations in *HUWE1* can lead to abnormal

cerebellar development, misaligned Purkinje cells, disruption of CGNPs differentiation, and disorganization of cerebellar architecture, potentially causing abnormal cell differentiation and tumorigenic states like medulloblastoma.<sup>12</sup> Transcriptional abnormalities in cerebellar formation may directly or indirectly affect brainstem (pons) development. Although initial studies on "cultured ES cells"<sup>15</sup> and *Drosophila* have been performed, still nothing is known about how a *HUWE1* mutation may result in pathologic conditions on cerebellar development and function. Our patient's posterior median skull defect and intracranial lipoma indicate the role of X chromosome genes in neural tube defects through p53 signaling process alteration<sup>11</sup>; however, the impact of *HUWE1* missense mutation on neural tube closure at the tissue morphogenesis remains unexplored. The short and thin corpus callosum may be due to altered axon development because *HUWE1* influences axon termination or branching in flies and worms.<sup>3</sup>

In conclusion, this report highlights *HUWE1* possible role in CNS development and may help prenatal diagnostic experts in consider also this entity, especially when dealing with pontine and cerebellar hypoplasia findings.

## Study Funding

The authors report no targeted funding.

## Disclosure

The authors report no relevant disclosures. Go to [Neurology.org/NG](https://www.neurology.org/NG) for full disclosures.

## Publication History

Received by *Neurology: Genetics* March 8, 2024. Accepted in final form May 8, 2024. Submitted and externally peer reviewed. The handling editor was Editor Stefan M. Pulst, MD, Dr med, FAAN.

## Appendix Authors

Name	Location	Contribution
<b>Mario Tortora, MD</b>	Department of Pediatric Radiology and Neuroradiology, "Vittore Buzzi" Children's Hospital - ASST Fatebenefratelli-Sacco, Milan, Italy	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; analysis or interpretation of data
<b>Elisa Cattaneo, MD</b>	Clinical Genetics Unit, Department of Pediatrics, "Vittore Buzzi" Children's Hospital - ASST Fatebenefratelli-Sacco, Milan, Italy	Major role in the acquisition of data
<b>Luigina Spaccini, MD</b>	Clinical Genetics Unit, Department of Obstetrics and Gynecology, "Vittore Buzzi" Children's Hospital - ASST Fatebenefratelli-Sacco, Milan, Italy	Major role in the acquisition of data
<b>Maria Iascone, PhD</b>	Medical Genetics Laboratory, Hospital Papa Giovanni XXIII, Bergamo, Italy	Major role in the acquisition of data



## Appendix (continued)

Name	Location	Contribution
<b>Barbara Scelsa, MD</b>	Pediatric Neurology Unit, Department of Pediatrics, "Vittore Buzzi" Children's Hospital - ASST Fatebenefratelli-Sacco, Milan, Italy	Major role in the acquisition of data
<b>Alessia Micalizzi, MD</b>	Translational Cytogenomics Research Unit, IRCCS Bambino Gesù Pediatric Hospital, Rome, Italy	Major role in the acquisition of data
<b>Antonio Novelli, PhD</b>	Translational Cytogenomics Research Unit, IRCCS Bambino Gesù Pediatric Hospital, Rome, Italy	Major role in the acquisition of data
<b>Mariano Lanna, MD</b>	Fetal Therapy Unit "U. Nicolini", Department of Obstetrics and Gynecology, Buzzi Children's Hospital, Milan, Italy	Major role in the acquisition of data
<b>Andrea Righini, MD</b>	Department of Pediatric Radiology and Neuroradiology, "Vittore Buzzi" Children's Hospital - ASST Fatebenefratelli-Sacco, Milan, Italy	Analysis or interpretation of data
<b>Pierangelo Veggiotti, PhD</b>	Pediatric Neurology Unit, "Vittore Buzzi" Children's Hospital, Department of Biomedical and Clinical Sciences "L. Sacco", University of Milan, Italy	Analysis or interpretation of data
<b>Chiara Doneda, MD</b>	Department of Pediatric Radiology and Neuroradiology, "Vittore Buzzi" Children's Hospital - ASST Fatebenefratelli-Sacco, Milan, Italy	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; analysis or interpretation of data

## References

- Huibregtse JM, Scheffner M, Beaudenon S, Howley PM. A family of proteins structurally and functionally related to the E6-AP ubiquitin-protein ligase. *Proc Natl Acad Sci U S A*. 1995;92(7):2563-2567. doi:10.1073/pnas.92.7.2563
- Qi L, Xu X, Qi X. The giant E3 ligase *HUWE1* is linked to tumorigenesis, spermatogenesis, intellectual disability, and inflammatory diseases. *Front Cell Infect Microbiol*. 2022;12:905906. doi:10.3389/fcimb.2022.905906
- Giles AC, Grill B. Roles of the *HUWE1* ubiquitin ligase in nervous system development, function and disease. *Neural Dev*. 2020;15(1):6. doi:10.1186/s13064-020-00143-9
- Friez MJ, Brooks SS, Stevenson RE, et al. *HUWE1* mutations in Juberg-Marsidi and Brooks syndromes: the results of an X-chromosome exome sequencing study. *BMJ Open*. 2016;6(4):e009537. doi:10.1136/bmjopen-2015-009537
- Froyen G, Corbett M, Vandewalle J, et al. Submicroscopic duplications of the hydroxysteroid dehydrogenase HSD17B10 and the E3 ubiquitin ligase *HUWE1* are associated with mental retardation. *Am J Hum Genet*. 2008;82(2):432-443. doi:10.1016/j.ajhg.2007.11.002
- Orivoli S, Pavlidis E, Cantalupo G, et al. Xp11.22 microduplications including *HUWE1*: case report and literature review. *Neuropediatrics*. 2016;47(1):51-56. doi:10.1055/s-0035-1566233
- Santos-Rebouças CB, de Almeida LG, Belet S, et al. Novel microduplications at Xp11.22 including *HUWE1*: clinical and molecular insights into these genomic rearrangements associated with intellectual disability. *J Hum Genet*. 2015;60(4):207-211. doi:10.1038/jhg.2015.1
- Garel C. The role of MRI in the evaluation of the fetal brain with an emphasis on biometry, gyration and parenchyma. *Pediatr Radiol*. 2004;34(9):694-699. doi:10.1007/s00247-004-1249-x
- Richards S, Aziz N, Bale S, et al; ACMG Laboratory Quality Assurance Committee. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424. doi:10.1038/gim.2015.30
- D'Arca D, Zhao X, Xu W, Ramirez-Martinez NC, Iavarone A, LaSorella A. *Huwe1* ubiquitin ligase is essential to synchronize neuronal and glial differentiation in the developing cerebellum. *Proc Natl Acad Sci U S A*. 2010;107(13):5875-5880. doi:10.1073/pnas.0912874107
- Delbridge ARD, Kueh AJ, Ke F, et al. Loss of p53 causes Stochastic Aberrant X-chromosome inactivation and female-Specific neural tube defects. *Cell Rep*. 2019;27(2):442-454.e5. doi:10.1016/j.celrep.2019.03.048
- Vriend J, Ghavami S, Marzban H. The role of the ubiquitin proteasome system in cerebellar development and medulloblastoma. *Mol Brain*. 2015;8(1):64. doi:10.1186/s13041-015-0155-5
- Moortgat S, Berland S, Aukrust I, et al. *HUWE1* variants cause dominant X-linked intellectual disability: a clinical study of 21 patients. *Eur J Hum Genet*. 2018;26(1):64-74. doi:10.1038/s41431-017-0038-6
- Ibarluzea N, Hoz AB, Villate O, et al. Targeted Next-Generation sequencing in patients with Suggestive X-linked intellectual disability. *Genes (Basel)*. 2020;11(1):51. doi:10.3390/genes11010051
- Zhao X, Heng JI, Guardavaccaro D, et al. The HECT-domain ubiquitin ligase *Huwe1* controls neural differentiation and proliferation by destabilizing the N-Myc oncoprotein. *Nat Cell Biol*. 2008;10(6):643-653. doi:10.1038/ncb1727