



# No-Touch vs. Conventional Radiofrequency Ablation Using Twin Internally Cooled Wet Electrodes for Small Hepatocellular Carcinomas: A Randomized Prospective Comparative Study

Yun Seok Suh<sup>1, 2\*</sup>, Jae Won Choi<sup>1, 2\*</sup>, Jeong Hee Yoon<sup>1, 2</sup>, Dong Ho Lee<sup>1, 2</sup>, Yoon Jun Kim<sup>3, 4</sup>, Jeong Hoon Lee<sup>3, 4</sup>, Su Jong Yu<sup>3, 4</sup>, Eun Ju Cho<sup>3, 4</sup>, Jung Hwan Yoon<sup>3, 4</sup>, Jeong Min Lee<sup>1, 2, 5</sup>

Departments of <sup>1</sup>Radiology and <sup>3</sup>Internal Medicine, Seoul National University Hospital, Seoul, Korea; <sup>2</sup>Department of Radiology, Seoul National University College of Medicine, Seoul, Korea; <sup>4</sup>Department of Internal Medicine and Liver Research Institute, Seoul National University College of Medicine, Seoul, Korea; <sup>5</sup>Institute of Radiation Medicine, Seoul National University Medical Research Center, Seoul, Korea

**Objective:** This study aimed to compare the efficacy between no-touch (NT) radiofrequency ablation (RFA) and conventional RFA using twin internally cooled wet (TICW) electrodes in the bipolar mode for the treatment of small hepatocellular carcinomas (HCC).

**Materials and Methods:** In this single-center, two-arm, parallel-group, prospective randomized controlled study, we performed a 1:1 random allocation of eligible patients with HCCs to receive NT-RFA or conventional RFA between October 2016 and September 2018. The primary endpoint was the cumulative local tumor progression (LTP) rate after RFA. Secondary endpoints included technical conversion rates of NT-RFA, intrahepatic distance recurrence, extrahepatic metastasis, technical parameters, technical efficacy, and rates of complications. Cumulative LTP rates were analyzed using Kaplan-Meier analysis and the Cox proportional hazard regression model. Considering conversion cases from NT-RFA to conventional RFA, intention-to-treat and as-treated analyses were performed.

**Results:** Enrolled patients were randomly assigned to the NT-RFA group (37 patients with 38 HCCs) or the conventional RFA group (36 patients with 38 HCCs). Among the NT-RFA group patients, conversion to conventional RFA occurred in four patients (10.8%, 4/37). According to intention-to-treat analysis, both 1- and 3-year cumulative LTP rates were 5.6%, in the NT-RFA group, and they were 11.8% and 21.3%, respectively, in the conventional RFA group ( $p = 0.073$ , log-rank). In the as-treated analysis, LTP rates at 1 year and 3 years were 0% and 0%, respectively, in the NT-RFA group and 15.6% and 24.5%, respectively, in the conventional RFA group ( $p = 0.004$ , log-rank). In as-treated analysis using multivariable Cox regression analysis, RFA type was the only significant predictive factor for LTP (hazard ratio = 0.061 with conventional RFA as the reference, 95% confidence interval = 0.000–0.497;  $p = 0.004$ ). There were no significant differences in the procedure characteristics between the two groups. No procedure-related deaths or major complications were observed.

**Conclusion:** NT-RFA using TICW electrodes in bipolar mode demonstrated significantly lower cumulative LTP rates than conventional RFA for small HCCs, which warrants a larger study for further confirmation.

**Keywords:** Hepatocellular carcinoma; Radiofrequency ablation; No-touch technique; Randomized controlled trial

**Received:** January 29, 2021 **Revised:** July 12, 2021 **Accepted:** July 13, 2021

This study was supported by a research grant from RF Medical Co., Ltd. (Seoul, Korea, No. 06-2016-2370).

\*These authors contributed equally to this work.

**Corresponding author:** Jeong Min Lee, MD, PhD, Department of Radiology, Seoul National University College of Medicine, Institute of Radiation Medicine, Seoul National University Medical Research Center, 101 Daehak-ro, Jongno-gu, Seoul 03080, Korea.

• E-mail: [jmlshy2000@gmail.com](mailto:jmlshy2000@gmail.com)

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Radiofrequency ablation (RFA) is currently recommended as an intended curative treatment for very early or early stage hepatocellular carcinoma (HCC) according to several of the most prominent guidelines for the management of HCC [1-3]. However, although RFA is widely accepted as an effective treatment option for small HCCs in nonsurgical candidates, it has a significantly higher rate of local tumor progression (LTP) compared to surgical resection, limiting its use in patients with resectable HCCs [4-7]. Several studies have demonstrated that there are no significant differences in long-term therapeutic outcomes between RFA and surgery, albeit with high LTP in RFA [6,8,9]. However, in patients with LTP, the required number of interventional procedures for controlling recurrent tumors was significantly higher than that in patients without LTP to obtain a similar overall survival outcome [10]. Thus, it is clinically valuable to lower LTP rates with RFA, and for this purpose, the creation of a 5–10 mm ablative margin around the target tumor is necessary, which may not always be possible with RFA [11].

In 2016, Seror et al. [12] reported that no-touch RFA (NT-RFA) using multi-bipolar electrodes resulted in 94% 5-year LTP-free survival, an excellent therapeutic outcome for HCCs within the Milan criteria. Thereafter, several retrospective studies have also reported significantly better LTP rates of NT-RFA using multi-bipolar RF technology than conventional RFA using monopolar RF technology [13-16]. However, a prospective cohort study by Hirooka et al. [17] reported that NT-RFA using multi-bipolar electrodes showed no differences in LTP rates, compared with conventional RFA using monopolar electrodes, despite having significantly better intrasubsegmental tumor recurrence-free survival [17]. Furthermore, in previous studies [12-17], there were differences not only in RFA techniques but also in the number and types of RF electrodes and RF energy delivery modes (multipolar vs. monopolar). Thus, at present, it is not yet certain whether the better LTP rates of NT-RFA were due to the oncologically favorable features of NT-RFA, different efficacy of RF equipment, or differing background clinical characteristics [13-15,17]. Therefore, to answer the question of whether NT-RFA can indeed provide better therapeutic efficacy, we conducted a randomized prospective study comparing NT-RFA with conventional RFA using the same RF ablation system and electrodes for the treatment of small HCCs.

## MATERIALS AND METHODS

This single-center, two-arm, parallel-group, prospective randomized controlled study was approved by the Institutional Review Board of Seoul National University Hospital (IRB No.1604-136-758, NCT 02806076). All participants provided written informed consent for enrollment in the study. This study was financially supported by the RF Medical Co., Ltd. However, the authors had complete control of patient enrollment, data collection, and analysis at all times, without any input from the funding source.

### Study Participants

From October 2016 to September 2018, we recruited participants who met the following eligibility criteria: 1) age 20 to 85 years, 2) Child-Pugh class A, 3) treatment-naive HCC or new HCC that developed more than 2 years after initial curative treatment for HCC, and 4) tumor size  $\leq 2.5$  cm. The 2.5 cm size limit was selected based on the results of a previous study, which showed that the ideal inter-electrode distance for the creation of a spherical ablation zone with bipolar electrodes was 3 cm [18].


The exclusion criteria were as follows: 1)  $> 2$  HCCs or diffuse type, 2) tumors adjacent to the portal or hepatic veins with  $< 5$  mm proximity, 3) invisible tumor on ultrasonography (US)-MR/CT fusion technique, 4) presence of macrovascular invasion or extrahepatic metastasis (EM), or 5) severe coagulopathy or bleeding tendency (platelet count  $\leq 50000/\text{mm}^3$  or international normalized ratio (INR) prolongation  $\geq 50\%$ ). HCC was diagnosed based on its typical hallmarks on CT or the Korean guidelines for HCC management [19].

### Group Assignment

Participants underwent a 1:1 random assignment to the NT-RFA or conventional-RFA group by stratified randomization and block randomization (Fig. 1). Randomization was stratified by the size of the tumor ( $< 2$  cm or 2–2.5 cm), number of tumors (single or two), and type of HCC (treatment-naive or recurrent). The randomization process was performed using a web-based randomization service managed by our institution's medical research collaboration center. Study participants and those assessing outcomes were blinded to group assignments.

### RFA Procedure

One radiologist with 20 years of experience in



**Fig. 1. Patient selection flow and group allocation.** HCC = hepatocellular carcinoma, NT = no-touch, RFA = radiofrequency ablation

percutaneous RFA performed all RFA procedures with assistance from one radiology fellow or resident, on an inpatient basis, under conscious sedation. For RFA procedure planning, pre-procedural CT or MRI studies were reviewed, followed by real-time US-CT/MR fusion imaging (Easy fusion, Samsung Medison) for localization of the target lesion and evaluation of technical feasibility [20]. The detailed RFA procedure of conventional RFA with tumor puncture using twin internally cooled wet (TICW) electrodes (CWTN-T, RF Medical) is described in Supplement.


NT-RFA : bipolar RFA was performed using TICW electrodes and the same generator unit used in the conventional RFA group. For the NT-RFA technique, the two tines of TICW electrodes were inserted into the perimeter of the index tumor (generally 3–5 mm from the margin of the target tumor) at an inter-electrode distance of 2–3 cm, depending on tumor size, under the guidance of real-time US-CT/MR fusion imaging [14,15,17,21]. An example case is shown in Figure 2. However, if there was no safe route for insertion of electrodes into the target tumor using the NT technique under multimodality fusion imaging, we converted patients from the NT-RFA group to the conventional RFA group. Furthermore, if the electrode incidentally passed through the tumor, it was recorded and regarded as an unintentional protocol change into conventional RFA, and the conversion

rates of NT-RFA to conventional RFA were calculated. An example of a converted case is shown in Figure 3.

### Evaluation of Procedure and Follow-Up


After the completion of RFA procedures, multiphasic contrast-enhanced CT studies were immediately conducted to assess ablation size, post-procedural complications, and technical success based on the reporting criteria suggested by the International Working Group on Image-guided Tumor Ablation [22]. Technical success of RFA was defined as complete coverage of the target tumor by the ablation zone on immediate follow-up CT or MRI [22] (Fig. 2). Post-procedural complications were also evaluated according to the guidelines of the Society of Interventional Radiology [23]. A detailed evaluation of the ablation zone and sufficiency of the margin (> 5 mm) is described in Supplement. Any residual enhancement at the ablation margin was considered to be an unablated residual tumor [24], and patients underwent additional ablation on the same day.

After RFA, contrast-enhanced CT or MRI was performed one month after RFA, and every 3 months to detect LTP, intrahepatic distant recurrence (IDR), and EM during the initial 2 years, after which the interval for CT or MRI was adjusted according to the clinician’s decision in the range of 3–6 months [22]. Technical efficacy was defined as



**Fig. 2. Underlying hepatitis C virus related liver cirrhosis patient with HCC.**

**A, B.** Contrast-enhanced CT imaging revealed a 1.0 cm arterial enhancing nodule with weak wash out on segment 5 of the liver, suggesting an HCC (arrows). **C.** Under real-time ultrasonography-CT fusion imaging guidance, a hypoechoic nodule was correlated to the arterial enhancing nodule. **D.** Two electrodes were inserted outside of the target tumor without tumor puncture with an inter-electrode distance of 1.69 cm. **E.** On the portal phase of immediate follow-up CT, complete ablation of the target tumor with sufficient margin was shown to be achieved (arrows). **F.** On 2-year follow-up CT, there was no evidence of local recurrence. HCC = hepatocellular carcinoma



**Fig. 3. Underlying non-hepatitis B, non-hepatitis C liver cirrhosis patient with HCC.**

**A, B.** On gadoteric acid-enhanced MR imaging, a 2.3 cm arterial enhancing nodule with an HBP defect located in segment 3 of the liver was observed, suggesting an HCC (arrows). **C.** Under real-time ultrasonography-CT fusion imaging guidance, a high echogenic nodule with a low echogenic rim was correlated to the arterial enhancing nodule. **D.** There was no safe route to puncture outside of the target tumor owing to insufficient peritumoral parenchyma (< 5 mm). Therefore, we converted to conventional RFA from no-touch RFA. **E.** On the portal phase of immediate follow-up CT, complete ablation of the target tumor with a sufficient margin was shown to be achieved (arrows). **F.** On 2-year follow-up CT, there was no evidence of local recurrence. Note that the axial plane is slightly oblique, compared to previous images. HCC = hepatocellular carcinoma, RFA = radiofrequency ablation

complete coverage of the target tumor by the ablation zone, assessed by imaging at one-month follow up. LTP was defined as newly appearing tumor foci at the margin of the ablation zone after achieving treatment success in a follow-up study with contrast-enhanced CT or MRI [22].

### Outcomes

The primary endpoint was to compare the cumulative incidence of LTP between the NT-RFA and conventional RFA groups. In addition, we determined the NT-RFA to conventional RFA conversion rates and evaluated the cumulative incidence of LTP in study patients according

to the received RFA techniques (as-treated analysis). As secondary endpoints, we also compared IDR, EM, and the following procedural characteristics between the two groups: technical parameters, technical success, technical efficacy, and rates of complications.

### Statistical Analysis

The sample size was approximated to be 55 patients in each group, but based on interim analysis, the study was terminated early in 78 patients (details in Supplement). Statistical evaluation was performed for the intention-to-treat analysis according to the initially assigned group and the as-treated analysis according to the actual treatment method [25]. Demographic factors and technical parameters were analyzed using the chi-squared test or Fisher's exact test for categorical variables and the independent *t* test for continuous variables. Technical success, technique efficacy, and LTP rates were analyzed using per-nodule data, based on the as-treated analysis. The Kaplan-Meier method and log-rank test were used for survival analysis between the two groups. Cox proportional

hazards regression analysis was performed to evaluate predictors of LTP (details in Supplement). Variables with a *p* value < 0.2, on univariable analysis, were included in the multivariable analysis. Statistical significance was set at *p* < 0.05. All statistical analyses were conducted using IBM SPSS Statistics for Windows, Version 25.0 (IBM Corp.) and R Statistical Software (version 4.0.3, R Foundation for Statistical Computing). Our institutional Medical Research Collaborating Center supported us in carrying out the statistical methodology.

## RESULTS

### Participant Enrollment

Between July 2016 and September 2018, we screened 78 potentially eligible patients who were referred for RFA for study enrollment. Five patients were excluded for the following reasons: poor visibility despite fusion imaging (*n* = 1), refusal to submit informed consent (*n* = 2), and poor liver function (*n* = 2). Seventy-three participants with 76 HCCs were enrolled and randomly assigned to the

**Table 1. Characteristics of Patient Groups for Intention-to-Treat Analysis**

Category	Conventional RFA (36 Patients with 38 HCCs)	NT-RFA (37 Patients with 38 HCCs)	<i>P</i>
Sex, M:F	27:11	27:11	> 0.999
Age, year	62.5 ± 7.7	66.1 ± 11.8	0.118
Origin of liver cirrhosis			
Hepatitis B virus-related	24	27	0.464
Hepatitis C virus-related	8	7	0.773
Alcoholism	4	7	0.328
Tumor number			0.692
One	34	35	
Two	2	2*	
Tumor location			0.761
Subcapsular	6	7	
Central	32	31	
Type of tumor			0.312
Naïve HCC	23	28	
Recurrent HCC	13	9	
Tumor size, cm	1.63 ± 0.48	1.74 ± 0.45	0.230
Lab			
Total bilirubin, mg/dL	0.80 ± 0.58	0.81 ± 0.36	0.944
Prothrombin time, INR	1.07 ± 0.12	1.06 ± 0.09	0.751
Albumin, g/dL	3.97 ± 0.48	3.96 ± 0.42	0.920
Platelet, 10 <sup>3</sup> /μL	123.8 ± 54.2	134.4 ± 55.9	0.406
Alpha fetoprotein, ng/mL	236.0 ± 1269.3	48.8 ± 131.8	0.382
Follow-up period, months	30.6 ± 13.3	31.3 ± 11.4	0.664

Data are patient number or average ± standard deviation. \*In one patient, only one of the two tumors was treated because of a poor sonic window. HCC = hepatocellular carcinoma, INR = international normalized ratio, NT = no-touch, RFA = radiofrequency ablation

two groups. Four participants in the NT-RFA group were converted to conventional RFA. Therefore, the intention-to-treat analysis involved 36 conventional RFA participants with 38 HCCs vs. 37 NT-RFA participants with 38 HCCs, and the as-treated analysis comprised 40 conventional RFA participants with 42 HCCs vs. 33 NT-RFA participants with 34 HCCs (Fig. 1). There were no statistically significant differences in demographic factors and tumor characteristics between the groups (Tables 1, 2).

### Procedure Characteristics

Procedure-related technical parameters according to intention-to-treat analysis are presented in Table 3. There were no significant differences in ablation size (Dmax, Dmin, Dv), ablation volume, effective ablation volume, and technical parameters, including procedure time, ablation time, and total energy in both groups (Table 3).

Among the NT-RFA group, conversion to conventional RFA was performed in four patients (10.8%, 4/37). The main reasons for conversion in these four patients were

the lack of a safe access route and subcapsular location with insufficient peritumoral parenchyma (< 5 mm) (Fig. 3). According to the as-treated analysis, 76.1% (32/42) of patients achieved sufficient ablation margin (> 5 mm) in the conventional RFA group, while NT-RFA group patients achieved a higher rate of sufficient ablation margin (88.2%, 30/34), albeit without statistical significance ( $p = 0.178$ ). Technical success was achieved in all cases. The technique efficacy rates of the conventional and NT-RFA groups at the 1-month follow-up were 100% (42/42) and 94.1% (32/34), respectively ( $p = 0.111$ ). The mean follow-up period was  $30.9 \pm 12.3$  months (median: 33.2 months).

### Procedure-Related Complications

In the NT-RFA group, there were no procedure-related deaths or major complications. There was only one major complication of bleeding during the immediate follow-up CT in the conventional RFA group, which had spontaneously stopped on follow-up hepatic angiogram. There were also no significant differences between the two groups in minor

**Table 2. Characteristics of Patient Groups for as-Treated Analysis**

Category	Conventional RFA (40 Patients with 42 HCCs)	NT-RFA (33 Patients with 34 HCCs)	<i>P</i>
Sex, M:F	29:12	25:9	0.668
Age, year	63.4 ± 7.9	65.5 ± 12.2	0.392
Causes of liver cirrhosis			
Hepatitis B virus-related	27	24	0.561
Hepatitis C virus-related	8	7	0.867
Alcoholism	4	7	0.173
Tumor number			
One	38	31	0.916
Two	2	2*	
Tumor location <sup>†</sup>			
Subcapsular	6	7	0.468
Central	36	27	
Type of tumor			
Naïve HCC	27	24	0.668
Recurrent HCC	13	9	
Tumor size, cm	1.65 ± 0.49	1.71 ± 0.44	0.606
Lab			
Total bilirubin, mg/dL	0.80 ± 0.56	0.81 ± 0.37	0.897
Prothrombin time, INR	1.07 ± 0.12	1.06 ± 0.09	0.601
Albumin, g/dL	3.94 ± 0.48	3.97 ± 0.47	0.881
Platelet, 10 <sup>3</sup> /μL	122.5 ± 52.0	137.3 ± 58.0	0.246
Alpha fetoprotein, ng/mL	215.2 ± 1207.5	52.7 ± 139.4	0.452
Follow-up period, months	30.7 ± 13.3	31.3 ± 11.4	0.850

Data are patient number or average ± standard deviation. \*In one patient, only one of the two tumors was treated because of a poor sonic window, <sup>†</sup>Tumor location was classified as a subcapsular or central tumor, based on whether it was within 1 mm from the liver capsule [34]. HCC = hepatocellular carcinoma, INR = international normalized ratio, NT = no-touch, RFA = radiofrequency ablation

complications, based on the as-treated analysis (n = 1, NT-RFA group; n = 2, conventional RFA group; p = 0.673). Minor thermal injuries were reported in three of the cases in post-procedure CT: two cases in the gallbladder (one in each group) and one case in the stomach (conventional RFA group). None of the patients required additional medical treatment.


**Outcome-LTP, IDR, and EM**

In the intention-to-treat analysis, the estimated 1-year and 3-year cumulative incidences of LTP were both 5.6% in the NT-RFA group, and 11.8% and 21.3%, respectively, in the conventional RFA group. The two groups did not differ significantly in LTP rates (log-rank test, p = 0.073) (Fig. 4A). In the as-treated analysis, the estimated 1-year and 3-year cumulative incidences of LTP were both 0% in

**Table 3. Comparison of Technical Parameters between the Two Groups according to Intention-to-Treat Analysis**

Category	Conventional RFA (36 Patients with 38 HCCs)	NT-RFA (37 Patients with 38 HCCs)	P
Power, W	29.4 ± 6.6	26.9 ± 4.9	0.071
Current, A	0.32 ± 0.05	0.30 ± 0.05	0.190
Impedance, Ω	100.9 ± 46.8	93.34 ± 23.19	0.374
Energy, kcal	13.5 ± 8.0	13.1 ± 6.9	0.856
Ablation time, min	12.08 ± 5.06	12.12 ± 5.10	0.973
Dmax, cm	48.7 ± 8.0	50.6 ± 12.2	0.420
Dmin, cm	33.9 ± 5.7	35.6 ± 6.6	0.218
Dv, cm	54.0 ± 13.4	54.0 ± 13.7	0.983
Vab, mL*	48.7 ± 20.9	55.0 ± 31.3	0.306
Veff, mL†	22.1 ± 11.4	26.1 ± 14.4	0.180
Dmin/Dmax	0.70 ± 0.10	0.72 ± 0.14	0.425
Procedure time, min	46.4 ± 12.8	48.8 ± 12.7	0.431

Data are average ± standard deviation. \*Vab - ablation volume (V = π/6 × Dmax × Dmin × Dv), †Veff - effective ablation volume (V = π/6 × Dmin<sup>3</sup>). HCC = hepatocellular carcinoma, NT = no-touch, RFA = radiofrequency ablation



**Fig. 4. Comparison of cumulative incidences of local recurrence between the two groups (A) at intention-to-treat analysis and (B) as-treated analysis.** \*Graphs were obtained using Kaplan-Meier survival curves, and p values were calculated using the log-rank test. LTP = local tumor progression, RFA = radiofrequency ablation

**Table 4. Univariable and Multivariable Cox Regression Analyses in as-Treated Analysis of Risk Factors for Local Tumor Progression**

Characteristics	Univariable	Multivariable		
	<i>P</i>	Hazard Ratio	95% CI	<i>P</i>
Sex	0.762			
Age, year	0.844			
Tumor size, cm	0.328			
Tumor number	0.541			
Treatment history	0.337			
Tumor location	0.194	0.65	0.12–6.08	0.668
Total bilirubin, mg/dL	0.423			
Prothrombin time, INR	< 0.001	268.0	0.44–2.7 × 10 <sup>5</sup>	0.086
Albumin, g/dL	0.016	0.82	0.16–3.80	0.795
Platelet, 10 <sup>3</sup> /μL	0.204			
Alpha fetoprotein, ng/mL	0.224			
Ablation volume, mL	0.440			
RFA type				
NT-RFA vs. conventional RFA as reference	0.002	0.061	0–0.497	0.004

Penalized maximum likelihood estimation was used in predictable factors with separation. CI = confidence interval, INR = international normalized ratio, NT = no-touch, RFA = radiofrequency ablation

**Table 5. Cumulative Incidences of Recurrence according to Intention-to-Treat and as-Treated Analysis**

Outcome Parameter	Conventional RFA (%)			NT-RFA (%)			<i>P</i>	
	1 Year	2 Year	3 Year	1 Year	2 Year	3 Year		
Intention-to-treat analysis	LTP	11.8	18.1	21.3	5.6	5.6	5.6	0.073
	IDR	31.1	40.5	43.8	26.9	38.8	56.6	0.818
	EM	2.9	9.3	9.3	0	9.0	15.5	0.731
As-treated analysis*	LTP	15.6	24.5	24.5	0	0	0	0.004
	IDR	33.1	44.3	47.2	24.1	34.4	53.7	0.746
	EM	2.6	11.2	17.1	0	6.7	6.7	0.380

\*Among the NT-RFA group patients (n = 37), conversion to conventional RFA was performed in 4 patients (10.8%, 4/37) due to a lack of a safe access route and subcapsular location with insufficient peritumoral parenchyma (< 5 mm). EM = extrahepatic metastases, IDR = intrahepatic distant recurrence, LTP = local tumor progression, NT = no-touch, RFA = radiofrequency ablation

the NT-RFA group, and 15.6% and 24.5%, respectively, in the conventional RFA group. The two groups differed significantly in LTP rates (log-rank test, *p* = 0.004) (Fig. 4B).

On multivariable Cox regression analysis in the as-treated analysis, RFA type was the only predictive factor for LTP (hazard ratio = 0.061, 95% confidence interval = 0.000–0.497 with conventional RFA as the reference; *p* = 0.004). Other predictable factors did not show statistical significance at the 5% level (Table 4). As for IDR and EM, there was no significant difference between the two groups in both analyses. The cumulative incidences of these analyses are summarized in Table 5.

## DISCUSSION

In this prospective randomized controlled trial, NT-RFA using TICW electrodes demonstrated better LTP than

conventional RFA using tumor puncture. According to the as-treated analysis, the estimated 1- and 3-year cumulative incidences of LTP were 0% in the NT-RFA group, which was significantly lower than the 15.6% and 24.5% in the conventional RFA group (*p* = 0.004). In addition, the complication rate of NT-RFA (3.3%, 1/33) was shown to be similar to that of conventional tumor puncture RFA (7.5%, 3/40). Indeed, our study results are in good agreement with several previous retrospective studies [13,14,16,26] and a prospective cohort study [17] reporting lower LTP rates of NT-RFA with bipolar electrodes compared to conventional RFA with a monopolar electrode. However, while previous studies utilized different numbers and types of RF electrodes, as well as different RF energy delivery modes on both NT-RFA and conventional RFA groups, our study used the same RF equipment and RF delivery mode for both groups, suggesting that the better LTP rates of NT-RFA



were due to the oncologically favorable features of the NT-RFA technique. Based on our study results, NT-RFA may be preferentially used for the treatment of small HCCs  $\leq 2.5$  cm compared to conventional RFA with tumor puncture.

Of note, although NT-RFA showed better LTP rates than conventional RFA, there was no significant difference in ablation zone size between the two groups. These results regarding ablation size could be attributed to the fact that if necessary to acquire optimal ablation margin, additional cycles of RFA were performed after repositioning of the electrode(s) in both groups. Therefore, the better LTP results of NT-RFA could be explained by several factors. First, the higher energy deposition in the peritumoral ablative margin, related to the geometry between the tumor and electrodes, in the NT-RFA group, might be an important contributing factor. According to a previous multicenter study [27], NT-RFA showed better LTP than standard monopolar RFA for HCCs located near large vessels, because more homogenous and extensive tissue necrosis and ablation zones were achieved beyond the macroscopic tumor boundary by placing the needles away from the tumor periphery. Second, blockage of drainage vessels from the target tumor may have been achieved in the early period of the ablation procedure, which may have blocked the dispersal of tumor cells into the drainage portal bloodstream, and may also be less susceptible to the heat sink effect [17,28]. Third, the centripetal thermal conduction from the periphery to the center of the index tumor may provide a theoretical advantage of less elevation of intratumoral pressure compared to conventional RFA [13,17]. Tanaka et al. [29] and Ueki et al. [26] also reported that intrahepatic metastasis of HCC occurred more frequently when intratumoral pressure increased after ethanol injection, resulting in a lower risk of track seeding along the electrode [30].

In our study, four patients (10.8%, 4/37) were converted from the NT-RFA technique to the conventional RFA technique. This represents the technical difficulty of NT-RFA compared to conventional tumor puncture RFA, as NT-RFA necessitates the insertion of multiple electrodes outside the target tumor with ideal geometry to create an ablative margin around the tumor [13,31]. The insertion of two electrodes around the tumor, at equidistance, while avoiding vital structures, can be technically challenging under the guidance of the US. In that regard, real-time fusion guidance or CT guidance might be more advantageous than US guidance. We found that real-time US-CT/MR fusion

guidance was very useful for placing electrodes in an ideal peritumoral position as it can provide better conspicuity of the target tumor as well as adjacent vascular structures or organs compared with B-mode US guidance [20,32,33]. Thus, to lower the technical difficulty of NT-RFA, additional modifications of RF energy delivery modes, such as combined bipolar and monopolar modes, could be used for NT-RFA. In this regard, further studies on the ideal RF delivery mode for NT-RFA, especially for larger tumors ( $\geq 3$  cm), are warranted. Moreover, NT-RFA had great difficulty in placing the electrodes in the surrounding peritumoral liver parenchyma for subcapsular located tumors, especially near the hepatic angle or hepatic dome portion, and this was the main cause of conversion of NT-RFA to conventional RFA.

Our study has several limitations that warrant mention. First, the small number of study patients was a major limitation. Because we had found a significant difference in LTP rates between NT-RFA and conventional RFA, we chose to terminate our clinical trial earlier than originally planned. Further studies with larger populations are warranted. Second, our short-term follow-up duration after RFA (mean of 36 months) was not sufficient to assess overall survival, and there were no significant differences between the two groups in IDM and EM. However, since the main purpose of our study was to evaluate the efficacy of the two RFA techniques in LTP, three years of follow-up should be sufficient to evaluate LTP after RFA. Third, the number of patients who had received RFA as the first treatment for de novo HCC was limited. Therefore, we could not perform a fair comparison of NT-RFA and conventional RFA in terms of the overall survival of the two groups after RFA. Finally, our study results were achieved by an experienced operator during percutaneous tumor ablation. Therefore, considering the technical difficulty and complexity of the NT-RFA procedure, it is not clear whether the favorable results of NT-RFA in the current study could be reproducible with less experienced operators. However, with the technical development of real-time US fusion imaging, NT-RFA for small HCCs ( $< 2.5$  cm) with favorable locations can be applied to less experienced operators. Despite these limitations, we believe that this study is meaningful, as it is a randomized prospective study comparing NT-RFA and conventional RFA using the same RF equipment and RF methodology, which demonstrated that NT-RFA provided lower LTP at similar safety levels.

In conclusion, NT-RFA using TIWC electrodes in bipolar mode demonstrated significantly lower cumulative LTP rates

than conventional RFA for small HCCs, which warrants a larger study for further confirmation.

## Supplement

The Supplement is available with this article at <https://doi.org/10.3348/kjr.2021.0319>.

## Conflicts of Interest

Jeong Min Lee received a research grant from RF Medical Co., Ltd. Other authors have no potential conflicts of interest to disclose.

## Acknowledgments

We thank Chris Woo, BA for his English editing.

## Author Contributions

Conceptualization: Yun Seok Suh, Jae Won Choi, Jeong Min Lee. Data curation: Yun Seok Suh, Jae Won Choi, Jeong Min Lee. Formal analysis: Yun Seok Suh, Jae Won Choi, Jeong Min Lee. Funding acquisition: Jeong Min Lee. Investigation: Jae Won Choi, Jeong Min Lee. Methodology: Jae Won Choi, Jeong Min Lee. Project administration: Jae Won Choi, Jeong Min Lee. Resources: Yun Seok Suh, Jae Won Choi, Jeong Min Lee. Software: Yun Seok Suh, Jae Won Choi, Jeong Min Lee. Supervision: Jeong Hee Yoon, Dong Ho Lee, Jeong Min Lee. Validation: Yun Seok Suh, Jeong Hee Yoon, Dong Ho Lee, Jeong Min Lee. Visualization: Yun Seok Suh, Jeong Min Lee. Writing—original draft: Yun Seok Suh, Jeong Min Lee. Writing—review & editing: all authors.

## ORCID iDs

Yun Seok Suh

<https://orcid.org/0000-0002-7004-5349>

Jae Won Choi

<https://orcid.org/0000-0002-5937-7238>

Jeong Hee Yoon

<https://orcid.org/0000-0002-9925-9973>

Dong Ho Lee

<https://orcid.org/0000-0001-8983-851X>

Yoon Jun Kim

<https://orcid.org/0000-0001-9141-7773>

Jeong Hoon Lee

<https://orcid.org/0000-0002-0315-2080>

Su Jong Yu

<https://orcid.org/0000-0001-8888-7977>

Eun Ju Cho

<https://orcid.org/0000-0002-2677-3189>

Jung Hwan Yoon

<https://orcid.org/0000-0002-9128-3610>

Jeong Min Lee

<https://orcid.org/0000-0003-0561-8777>

## REFERENCES

1. European Association for the Study of the Liver. EASL clinical practice guidelines: management of hepatocellular carcinoma. *J Hepatol* 2018;69:182-236
2. Korean Liver Cancer Association (KLCA), National Cancer Center (NCC). 2018 Korean Liver Cancer Association-National Cancer Center Korea practice guidelines for the management of hepatocellular carcinoma. *Korean J Radiol* 2019;20:1042-1113
3. Heimbach JK, Kulik LM, Finn RS, Sirlin CB, Abecassis MM, Roberts LR, et al. AASLD guidelines for the treatment of hepatocellular carcinoma. *Hepatology* 2018;67:358-380
4. Ng KKC, Chok KSH, Chan ACY, Cheung TT, Wong TCL, Fung JYY, et al. Randomized clinical trial of hepatic resection versus radiofrequency ablation for early-stage hepatocellular carcinoma. *Br J Surg* 2017;104:1775-1784
5. Chen MS, Li JQ, Zheng Y, Guo RP, Liang HH, Zhang YQ, et al. A prospective randomized trial comparing percutaneous local ablative therapy and partial hepatectomy for small hepatocellular carcinoma. *Ann Surg* 2006;243:321-328
6. Feng K, Yan J, Li X, Xia F, Ma K, Wang S, et al. A randomized controlled trial of radiofrequency ablation and surgical resection in the treatment of small hepatocellular carcinoma. *J Hepatol* 2012;57:794-802
7. Xu XL, Liu XD, Liang M, Luo BM. Radiofrequency ablation versus hepatic resection for small hepatocellular carcinoma: systematic review of randomized controlled trials with meta-analysis and trial sequential analysis. *Radiology* 2018;287:461-472
8. Lee DH, Kim JW, Lee JM, Kim JM, Lee MW, Rhim H, et al. Laparoscopic liver resection versus percutaneous radiofrequency ablation for small single nodular hepatocellular carcinoma: comparison of treatment outcomes. *Liver Cancer* 2021;10:25-37
9. Kang TW, Kim JM, Rhim H, Lee MW, Kim YS, Lim HK, et al. Small hepatocellular carcinoma: radiofrequency ablation versus nonanatomic resection—propensity score analyses of long-term outcomes. *Radiology* 2015;275:908-919
10. Lee DH, Lee JM, Lee JY, Kim SH, Yoon JH, Kim YJ, et al. Radiofrequency ablation of hepatocellular carcinoma as first-line treatment: long-term results and prognostic factors in 162 patients with cirrhosis. *Radiology* 2014;270:900-909
11. Ahmed M, Brace CL, Lee FT Jr, Goldberg SN. Principles of and advances in percutaneous ablation. *Radiology* 2011;258:351-369

12. Seror O, N'Kontchou G, Nault JC, Rabahi Y, Nahon P, Ganne-Carrié N, et al. Hepatocellular carcinoma within milan criteria: no-touch multibipolar radiofrequency ablation for treatment—Long-term results. *Radiology* 2016;280:611-621
13. Hocquet A, Aubé C, Rode A, Cartier V, Sutter O, Manichon AF, et al. Comparison of no-touch multi-bipolar vs. monopolar radiofrequency ablation for small HCC. *J Hepatol* 2017;66:67-74
14. Kawamura Y, Ikeda K, Fujiyama S, Hosaka T, Kobayashi M, Saitoh S, et al. Potential of a no-touch pincer ablation procedure that uses a multipolar radiofrequency ablation system to prevent intrasubsegmental recurrence of small and single hepatocellular carcinomas. *Hepatol Res* 2017;47:1008-1020
15. Seror O, N'Kontchou G, Van Nhieu JT, Rabahi Y, Nahon P, Laurent A, et al. Histopathologic comparison of monopolar versus no-touch multipolar radiofrequency ablation to treat hepatocellular carcinoma within Milan criteria. *J Vasc Interv Radiol* 2014;25:599-607
16. Chai Y, Li K, Zhang C, Chen S, Ma K. The short-term efficacy of no-touch radiofrequency ablation in treating cirrhosis-based small hepatocellular carcinoma. *BMC Cancer* 2019;19:497
17. Hirooka M, Hiraoka A, Ochi H, Koizumi Y, Michitaka K, Joko K, et al. Prospective cohort trial to confirm the efficacy of no-touch radio frequency ablation. *J Gastroenterol Hepatol* 2019;34:567-574
18. Lee JM, Han JK, Kim SH, Lee JY, Shin KS, Choi BI. An ex-vivo experimental study on optimization of bipolar radiofrequency liver ablation using perfusion-cooled electrodes. *Acta Radiol* 2005;46:443-451
19. Korean Liver Cancer Study Group (KLCSG), National Cancer Center, Korea (NCC). 2014 Korean Liver Cancer Study Group-National Cancer Center Korea practice guideline for the management of hepatocellular carcinoma. *Korean J Radiol* 2015;16:465-522
20. Ahn SJ, Lee JM, Lee DH, Lee SM, Yoon JH, Kim YJ, et al. Real-time US-CT/MR fusion imaging for percutaneous radiofrequency ablation of hepatocellular carcinoma. *J Hepatol* 2017;66:347-354
21. Kim TH, Choi HI, Kim BR, Kang JH, Nam JG, Park SJ, et al. No-touch radiofrequency ablation of VX2 hepatic tumors in vivo in rabbits: a proof of concept study. *Korean J Radiol* 2018;19:1099-1109
22. Ahmed M, Solbiati L, Brace CL, Breen DJ, Callstrom MR, Charboneau JW, et al. Image-guided tumor ablation: standardization of terminology and reporting criteria--a 10-year update. *Radiology* 2014;273:241-260
23. Sacks D, McClenny TE, Cardella JF, Lewis CA. Society of Interventional Radiology clinical practice guidelines. *J Vasc Interv Radiol* 2003;14:S199-S202
24. Shropshire EL, Chaudhry M, Miller CM, Allen BC, Bozdogan E, Cardona DM, et al. LI-RADS treatment response algorithm: performance and diagnostic accuracy. *Radiology* 2019;292:226-234
25. Ten Have TR, Normand SL, Marcus SM, Brown CH, Lavori P, Duan N. Intent-to-treat vs. non-intent-to-treat analyses under treatment non-adherence in mental health randomized trials. *Psychiatr Ann* 2008;38:772-783
26. Ueki T, Sakaguchi S, Miyajima Y, Hatono N, Tohara K, Nakabayashi S, et al. Usefulness of tumor pressure as a prognostic factor in cases of hepatocellular carcinoma where the diameter of the tumor is 3 cm or less. *Cancer* 2002;95:596-604
27. Mohkam K, Dumont PN, Manichon AF, Jouvet JC, Bousset L, Merle P, et al. No-touch multibipolar radiofrequency ablation vs. surgical resection for solitary hepatocellular carcinoma ranging from 2 to 5 cm. *J Hepatol* 2018;68:1172-1180
28. Mohkam K, Rode A, Mabrut JY. Influence of ablation technique on treatment failure for perivascular hepatocellular carcinoma. *J Hepatol* 2018;69:750-751
29. Tanaka T, Yamanaka N, Oriyama T, Furukawa K, Okamoto E. Factors regulating tumor pressure in hepatocellular carcinoma and implications for tumor spread. *Hepatology* 1997;26:283-287
30. Otsuka Y, Kaneko H. Laparoscopic liver resection in the treatment of HCC with liver cirrhosis: would it provide superiority to conventional open hepatectomy? *Hepatobiliary Surg Nutr* 2017;6:356-358
31. Yoon JH, Lee JM, Woo S, Hwang EJ, Hwang I, Choi W, et al. Switching bipolar hepatic radiofrequency ablation using internally cooled wet electrodes: comparison with consecutive monopolar and switching monopolar modes. *Br J Radiol* 2015;88:20140468
32. Lee MW. Fusion imaging of real-time ultrasonography with CT or MRI for hepatic intervention. *Ultrasonography* 2014;33:227-239
33. Lee DH, Lee JM. Recent advances in the image-guided tumor ablation of liver malignancies: radiofrequency ablation with multiple electrodes, real-time multimodality fusion imaging, and new energy sources. *Korean J Radiol* 2018;19:545-559
34. Kang TW, Lim HK, Lee MW, Kim YS, Rhim H, Lee WJ, et al. Long-term therapeutic outcomes of radiofrequency ablation for subcapsular versus nonsubcapsular hepatocellular carcinoma: a propensity score matched study. *Radiology* 2016;280:300-312