














Original article:

**INHIBITION OF ANGIOTENSIN II TYPE 1 RECEPTOR BY
CANDESARTAN REDUCES TUMOR GROWTH AND AMELIORATES
FIBROSIS IN COLORECTAL CANCER**

Ehsan Tabatabai^{1,2#} , Majid Khazaei^{3,4#} , Fereshteh Asgharzadeh³ , Seyedeh Elnaz Nazari³, Neda Shakour^{5,6} , Hamid Fiuji⁴, Aghigh Ziaemehr⁴, Asma Mostafapour⁴ , Mohammad Reza Parizadeh⁴ , Mohammad Nouri^{1,2} , Seyed Mahdi Hassanian^{4,7} , Farzin Hadizadeh^{5,8} , Gordon A Ferns⁹ , Mohammad Rahmati^{1,2*} , Farzad Rahmani^{10*} , Amir Avan^{4,11#} 

¹ Stem Cell Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

² Department of Clinical Biochemistry, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

³ Department of Physiology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

⁴ Metabolic Syndrome Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

⁵ Department of Medicinal Chemistry, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

⁶ Student Research Committee, Mashhad University of Medical Sciences, Mashhad, Iran

⁷ Department of Medical Biochemistry, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

⁸ Biotechnology Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran

⁹ Brighton & Sussex Medical School, Division of Medical Education, Falmer, Brighton, Sussex BN1 9PH, UK

¹⁰ Iranshahr University of Medical Sciences, Iranshahr, Iran

¹¹ Medical Genetics Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

These authors contributed equally as first authors.

* **Corresponding authors:** Farzad Rahmani, PhD, Iranshahr University of Medical Sciences, Iranshahr, Iran. E-mail: Rahmany.farzad@gmail.com
Mohammad Rahmati, PhD, Department of Clinical Biochemistry, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran. Stem Cell Research Center, Tabriz University of Medical Sciences, Tabriz, Iran E-mail: Rahmatibio@gmail.com

<http://dx.doi.org/10.17179/excli2021-3421>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

ABSTRACT

Colorectal cancer (CRC) is an important cause of cancer-related mortality. Aberrant activation of the renin-angiotensin system (RAS) is reported to be associated with poor clinical outcomes in patients with CRC. This study

was designed to explore the anti-tumor effects of the angiotensin receptor blocker Candesartan either alone or in combination with 5-FU in *in vitro* and *in vivo* models of CRC. The cytotoxic effects of Candesartan were assessed using the MTT assay in two colorectal cancer cell lines (CT-26 and SW-480). To investigate the potential regulatory role of Candesartan on tumor growth, apoptosis, and migration, the expression levels of Cyclin D1, Survivin, MMP3, MMP9, and E-cadherin mRNAs were evaluated. The oxidant/antioxidant balance was also examined by determining the levels of MDA, thiols, SOD, and CAT. We used a xenograft model of colon cancer to investigate the effects of Candesartan alone, or in combination with 5-FU, on tumor growth following histological staining (Hematoxylin & Eosin and Masson trichrome staining) and biochemical studies as well as gene expression analyses by RT-PCR and western blotting. Candesartan suppressed tumor cell proliferation and migration by modulating Cyclin D1, MMP3/9, and E-cadherin. Treatment with Candesartan either alone, or in combination with 5-FU decreased tumor size in the mouse model, and also increased the level of oxidative markers MDA and reduced CAT, SOD, and thiols. Histological evaluation showed that Candesartan increased tumor necrosis, reduced tumor density and attenuated collagen deposition reducing tumor fibrosis in tumor xenograft. Candesartan, an inhibitor of the RAS, when used in combination with 5-FU reduced tumor growth by inhibiting fibrosis and inducing ROS production, supporting further clinical studies on this therapeutic approach for treatment of CRC.

Keywords: Candesartan, colorectal cancer, renin-angiotensin system

INTRODUCTION

Colorectal cancer (CRC) is one of the most common and aggressive malignancies with a high mortality globally. Despite recent improvements in treatment modalities or identification of novel diagnostic biomarkers, the overall survival of CRC patients treated with standard chemotherapeutic regimens remains extremely poor (Enofe et al., 2020; Wong et al., 2021). Thus, it is necessary to identify novel alternative therapeutics for treatment and management of CRC.

There is growing evidence suggesting a possible association between cancer incidence and the use of angiotensin II receptor type 1 (AT1R) blockers, that are used in the management of hypertension and cardiovascular disease. AT1R have been shown to be upregulated in various cancer cells including melanoma, glioblastoma, prostate, renal, bladder, breast, and pancreatic cancer (Miyajima et al., 2002; Du et al., 2012; Renziehausen, 2017; Renziehausen et al., 2019; Afsar et al., 2021; Arjmand et al., 2020; Asgharzadeh et al., 2020a). The overexpression of AT1R was reported to be associated with the malignant features of tumors by inducing angiotensin II. Angiotensin II is the main effector of the renin-angiotensin system, and has a prominent role in renal and vascular homeostasis. Additionally, the renin-angiotensin system has also been shown to regulate

various aspects of tumor growth and development including: cell proliferation, angiogenesis, inflammation, and apoptosis. Further studies indicate that angiotensin II exerts its tumorigenic effects through binding to the AT1R and angiotensin II receptor type II (AT2R) (Childers, 2015; Pei et al., 2017; Ishikane and Takahashi-Yanaga, 2018).

AT1R blockers, such as candesartan, losartan, and valsartan are commonly used as anti-hypertensive drugs in patients with cardiovascular diseases (Minatoguchi et al., 2013; Abraham et al., 2015). There is growing evidence that suggests that the suppression of angiotensin II receptor signaling by AT1R blockers can inhibit tumor progression in several types of cancers including renal, colon, breast, bladder, and prostate cancer (Plastowska-Ciesielska et al., 2013; Arrieta et al., 2015; Asgharzadeh et al., 2018, 2020a; Hashemzahi et al., 2021a, b). Candesartan has been reported to inhibit cancer cell growth and development in multiple human tumors when used alone or in combination with other anticancer drugs (Molteni et al., 2006; Fan et al., 2016; Ishikane and Takahashi-Yanaga, 2018). In support of these findings, it has been shown that administration of candesartan, prevents metastasis of renal cancer in a rat allograft model (Miyajima et al., 2002). Moreover, treatment with AT1R blockers has been reported to be associated with longer progres-

sion free survival and overall survival in patients with advanced pancreatic cancer and colon cancer (Mc Menamin et al., 2012; Nakai et al., 2013). Taken together, these findings suggest an important role of suppressing AT1R in patient with colon cancer and led us to investigate whether an AT1R blocker may inhibit CRC cell growth either alone or in combination with other chemotherapeutic drugs.

Here we investigated the therapeutic potential and the molecular mechanism of actions of candesartan in the CRC progression by using cellular system and animal model, for a better understanding and hence a better management of the disease.

MATERIAL AND METHODS

Drugs and chemicals

Candesartan and 5-FU were purchased from Cayman Chemical (Ann Arbor, MI). Roswell Park Memorial Institute (RPMI) 1640 Medium, fetal bovine serum (FBS), penicillin (50 IU/ml) and streptomycin (50 µg/ml) were purchased from Gibco (Gaithersburg, MD). Other laboratory chemicals were obtained from Sigma-Aldrich (Zwijndrecht, The Netherlands).

Cell culture

Two different CRC cell lines (CT-26 and SW-480) were used for the experiments. The cells were purchased from Pasteur Institute (Tehran, Iran). Cells were grown in RPMI 1640 medium supplemented with 10 % FBS (Gibco, USA) and 1 % penicillin/streptomycin at 37.0 °C in 5 % CO₂ humidified air.

Growth inhibition (MTT) assay

The MTT tetrazolium colorimetry assay was used to evaluate the cytotoxicity of candesartan on the CRC cells. Briefly, the CT-26 and SW-480 cells were seeded in a 96-well plate and treated with different concentrations of candesartan for 24 hours. After cell treatment, 20 µL of the MTT solution (5 mg/ml) was transferred to each well and plates were incubated at 37 °C for 4 hours to investigate

the viable cells can produce formazan crystals. Then, the medium was replaced by DMSO (100 µL) to solubilize the formazan crystals formed by living cells. The optical density was then measured at 545-630 nm using a micro plate reader (TECAN, Austria) (Amerizadeh et al., 2018). Mean results were reported from three independent experiments carried out at least in triplicate.

Animal experiment

Eight-week-old inbred BALB/c mice were obtained from the Pasteur Institute (Tehran, Iran). The CT-26 cells (2×10^6 cells) were subcutaneously injected into the right flank of each mouse. Next, the mice were divided into 4 groups: group 1 (control animals), group 2 were treated with CRC standard treatment, 5-FU (5 mg/kg every other day). Mice in group 3 were treated with candesartan (20 µM) and group 4 was treated with a 5-FU/ candesartan combination. Treatment was started approximately 2 weeks after injection when the tumor volume reached 80-100 mm³. The animal tumors were carefully monitored and measured every other day. The tumor volume (V) was calculated according to the formula $V = AB^2/2$, where A was the length of the major axis and B was the length of the minor axis. At day 21, mice were sacrificed and colon tumors were collected for further examination by trichrome and hematoxylin and eosin staining methods (Amerizadeh et al., 2018; Hashemzahi et al., 2021b). All animal procedures were conducted according to the guidelines established by the Ethics Committee of Mashhad University of Medical Sciences.

Quantitative Real-Time Polymerase Chain Reaction (qRT-PCR)

Total RNA was extracted from the treated cells (7×10^5 cells/well, in 6-well plates) using the RNeasy® mini kit (Qiagen GmbH, Hilden, Germany). Quantification and quality control of total RNA was performed in triplicate by a NanoDrop2000 spectrophotometer (Thermo Fisher Scientific, Inc.). Then, RNAs were reverse-transcribed using the PrimeScript™ RT reagent kit (TaKaRa Holdings,

Inc., Kyoto, Japan) (Afshari et al., 2019; Rahmani et al., 2019). Also, quantitative RT-PCR analysis was performed using RealQ Plus 2X MasterMix Green-without Rox™ (Amplicon, Stenhuggervej, Denmark). Next, quantitative RT-PCR was performed using specific primers for Cyclin D1, Survivin, E-cadherin, MMP-3, MMP-9, and GAPDH (Table 1) that were obtained from Macrogen (Macrogen Co., Seoul, Korea). The cDNA amplification was performed by using the LightCycler 96 real-time PCR system (Roche Applied Science, Pleasanton, CA, USA). Also, gene expression data were normalized against GAPDH. The 2- $\Delta\Delta C_t$ method was used to analyze the relative expression of target genes. The primer sequences (forward and reverse) are listed in Table 1.

Quantification of apoptosis

The regulatory effect of candesartan on CRC cell apoptosis was studied by flow cytometric analysis using Annexin V/PI apoptosis assay kit (Cayman Chemical, Ann Arbor, MI) (Ghanaatgar-Kasbi et al., 2019). In brief, CT-26 and SW-480 cells were exposed to candesartan (20 μ M) for 24 hours and the population of apoptotic cells were detected by flow cytometry according to manufacturer's instruction. Each experiment was performed in triplicate.

Histopathological staining

The CRC tumor tissues were fixed in formaldehyde, and subsequently embedded into paraffin blocks. The paraffin-embedded blocks were then sectioned to a thickness of approximately 5-7 μ m and multiple tissue sections were mounted onto a glass microscope slide. Following de-paraffinization, mounted tissue sections were stained with hematoxylin and eosin and Masson's trichrome and reviewed by at least 2 independent researchers (Rahmani et al., 2020; Soleimani et al., 2020).

Migration assay

The migration assay was performed as described previously (Hashemzahi et al., 2018). Cells were treated with candesartan and cellular migration was evaluated using the Leica DMI300B (Leica) at different time intervals (0, 16, 26, 46hr) and quantified by Image J software.

Measurement of oxidative stress markers

To further explore the regulatory effects of candesartan on oxidant/antioxidant balance in tumor tissues, the catalase (CAT), superoxide dismutase (SOD), malondialdehyde (MDA) and total thiol levels were determined in homogenized colon samples as described previously (Asgharzadeh et al., 2021).

Table 1: qPCR primer sequences

Gene	Source	Primer	Sequence
GAPDH	Mouse	Forward	CAACGACCCCTTCATTGACC
		Reverse	CTTCCCATTCTCGGCCTTGA
Cyclin D1	Mouse	Forward	GCGTACCCTGACACCAATCTC
		Reverse	ACTTGAAGTAAGATACGGAGGGC
Survivin	Mouse	Forward	GCATTGTCAGAAGCCCTTGT
		Reverse	TGTCCCCTTCAGCCAATACT
MMP-2	Mouse	Forward	AACTGTTGCTTTTGTATGCCCT
		Reverse	CGATGTCAGACAACCCGAGT
MMP-9	Mouse	Forward	GCGTCGTGATCCCCACTTAC
		Reverse	CAGGCCGAATAGGAGCGTC
E-cad	Mouse	Forward	GTCTACCAAAGTGACGCTGA
		Reverse	GGGAAACATGAGCAGCTCTG

Docking study

The interaction of the anti-cancer drug (Candesartan Cilexetil) with the active site of the target proteins was examined by Molecular Operating Environment docking experiment (MOE, Chemical Computing Group Inc. Montreal, <http://www.chemcomp.com>). No previous report was published associate to interaction and inhibition of mentioned target proteins in this article by Candesartan Cilexetil. The chemical structure of Candesartan was drawn using a software program (ChemDraw Ultra 7.0) and its energy minimized by MOE, respectively. Then we downloaded the crystal structure of proteins from the RCSB Protein Data Bank. PDB ID used in this study were 4APJ (for ACE), 5QIN (for TGF-beta type II), 3TZM (for TGF-beta type I), 4ZUD (for AT1), 4G9L (for MMP3), 3D91 (for Renin), 1GKC (for MMP9), 1XOX (for Survivin), 5cvb (for COL1A1), 2O72 (for E-cadherin), 5CTI (for COL1A2), 3KBH (for ACE2), 1ALU (for IL-6), 7JRA (for TNF- α), and 5VZU (for Cyclin D1). The docking procedure was performed using the default settings of the MOE-DOCK. The final docking scores were evaluated using the GBVI/WSA dG scoring function with the Generalized Born solvation model (GBVI) (Shakour et al., 2020). The GBVI/WSA dG is a force field-based scoring function, which estimates the free energy of binding of the ligand from a given pose. The inhibition constant (K_i) was computed through the binding free energy estimated with the GBVI/WSA dG scoring function according to the following equation: $\Delta G = RT \ln(K_i)$ where R represented the gas constant and T the temperature in kelvin. The K_i was computed from the binding free energy values at a fixed temperature (298 K).

Statistical analysis

All assays were undertaken in triplicate and verified in at least two independent experiments. All results were expressed as mean \pm SD, and analyzed by Student's t-test following by Tukey's multiple comparison tests or Kruskal Wallis tests and Mann Wilcoxon U

tests. The statistical significance was considered lower than 0.05.

RESULTS

Candesartan inhibits cell viability

To investigate the cytotoxic effects of candesartan in CRC, the CT-26 and SW-480 cells were treated with different concentrations (0-1000 μ M) of candesartan for 24 hours. Our data demonstrated that candesartan inhibited the growth of CRC cells in dose-dependent manner (Figure 1A). To further assess the anti-cancer effects of candesartan on CRC, the CT-26 and SW-480 cells were treated with candesartan (20 μ M) for 24 hours and apoptosis was determined by flow cytometry method. As shown in Figure 1B, candesartan significantly induced cell death in CRC cells.

Candesartan inhibits migratory behaviors of CRC cells

To identify the regulatory mechanism of candesartan on migratory behaviors of CRC cells, we evaluated the effect of Candesartan on the expression of E-cadherin and matrix metalloproteases-3 and -9. Administration of candesartan decreased the expression of MMP-3 and -9 while significantly increased expression of E-cadherin as a key regulatory factor involved in maintaining cell-to-cell adhesion (Figure 2A). Consistent with these findings, the migration of candesartan treated CRC cells was significantly inhibited as compared with control group (Figure 2B).

Candesartan suppresses the tumor growth

To further evaluate the regulatory effects of candesartan on colon tumor growth *in vivo*, a mouse model was used in which mice were treated with candesartan either alone or in combination with 5-FU and tumor size and weight were measured and compared between different groups (Figure 3A). In accordance with the *in vitro* data, intra-tumoral injection of candesartan reduced the size and weight of tumors in mice (n=6 in each group) (Figure 3B, C). Additionally, compared with the 5-FU




Figure 1: Growth inhibitory and cytotoxic effects of Candesartan on CRC cells. (A) SW-480 and CT-26 cells were treated with different concentrations of Candesartan for 24 and the cell viability was assessed by MTT assay. (B) SW-480 and CT-26 cells were treated with Candesartan for 24 hours and the early and late apoptosis were detected by flow cytometry method. All data was obtained from three independent experiments.




Figure 2: Candesartan inhibits migratory behaviors of CRC cells. (A) Expression level of E-cadherin and MMP3/9 in SW-480 and CT-26 cells in the presence of candesartan was detected by real time RT-PCR. (B) Inhibitory effect of candesartan on the migration of CT-26 cells was measured. All results were obtained from three independent experiments. * $p < 0.05$.

group, the group receiving adjuvant candesartan demonstrated a more potent anti-cancer effect and the tumor size and weight was lower in the combination group. Interestingly, histological staining of tumor tissues showed higher tissue necrosis in the combination group (Figure 3D) suggesting that candesartan synergistically increases the anti-cancer activity of 5-FU in this CRC mouse model. To

further determine the anti-proliferative activity of Candesartan, the expression of cyclin D1 and survivin mRNA levels were investigated in CRC tissues. As shown in Figure 3E, candesartan significantly decreased the expression of cyclin D1 and survivin suggesting the mechanism by which candesartan can inhibit tumor cell proliferation.

Effects of candesartan on fibrosis

To explore the regulatory effects of candesartan on fibrosis, the tumor tissues were stained with Masson's trichrome. Colonic fibrosis is characterized by excessive collagen deposition and Histopathological staining findings demonstrated that candesartan significantly inhibits tissue fibrosis and decreased collagen deposition in tumor tissues. Moreover, the tissue fibrosis was also shown to be significantly suppressed in the combination group suggesting that candesartan enhances the anti-fibrosis effects of 5-FU (Figure 4A, B).

Effects of candesartan on oxidant/antioxidant markers

To further investigate the role of candesartan on oxidant/antioxidant balance, the levels of SOD, CAT, MDA, and thiol were determined in homogenized colon tissues and results demonstrated in Figure 5A-D. The MDA level as a biomarker of oxidative stress increased in treated groups while the enzymatic activity of SOD and CAT and thiol concentrations were significantly decreased compared with the control group. To further confirm that candesartan exerted regulatory effects on oxidant/antioxidant balance, we evaluated the protein levels of inflammatory cytokines in colon tissues. As shown in Figure 5E, candesartan significantly decreased the expression of TNF- α and Il-6 in comparison with the control group.

Docking on Candesartan with potential markers associated with RAS pathway

While previous studies have reported that this drug is the inhibitor for angiotensin II type 1 receptor (AT1) (Bregonzio et al., 2008;




Figure 3: Candesartan suppresses tumor growth in colorectal cancer cells and mouse model (n=6 in each group). **(A)** Schematic representation of murine tumor model and treatment. **(B, C)** Tumor size and weight in CRC mouse model treated with Candesartan, 5-FU and their combination. **(D)** Histological staining of tumor tissues by H&E for evaluating aggregation of tumor cells (T) and necrotic area (N). **(E)** Expression of cyclin D1 and survivin were measured in CRC tumor tissues using Western blotting. *p < 0.05; **p < 0.01




Figure 4: Candesartan suppresses tumor fibrosis in colorectal cancer mouse model (n=6 in each group). (A, B) Trichrome staining for investigating the deposition of collagen fibers in CRC tissues

Baguet et al., 2009; Mengden et al., 2009), we studied the affinity of Candesartan Cilexetil (ligand) and potential markers associated with RAS pathway (Figure 6). Docking results are shown that Candesartan could directly interact and inhibit eight proteins as well as AT1 according to Figure 6 (Score ≤ -7.4 kcal/mol). In particular, ACE had $k_i = 0.179 \mu\text{M}$ through two H acceptor bonding among oxygen ligand

to oxygen and carbon of THR_302 and THR_301 residues of angiotensin-I-converting enzyme with distance 2.89 and 3.22 respectively and one pi-cation binding among of its 5-ring to nitrogen of LYS_449 with distance 3.89, followed by TGF-beta type II ($k_i = 0.420 \mu\text{M}$; one H acceptor binding of ligand carbonyl group oxygen to carbon of GLY_251 residue with distances 3.52; one H




Figure 5: Candesartan induces oxidative stress in homogenized colon samples. The regulatory effect of Candesartan on the enzymatic activity of SOD (A) and catalase (B) and on the MDA (C), total thiol (D) and inflammatory cytokines (E) were investigated in CRC tumor tissues.




Figure 6: The protein structure in 3D dimensions (left column) and 2D dimensions (right column) related to interaction Candesartan Cilexetil and target proteins

donor binding of ligand carbon to sulfur of CYS_396 residue with distances 3.86; three pi-H binding of ligand (two 5-ring and one 6-ring) to carbon of GLY_253, VAL_258, and SER_383 residues with distances 3.71, 3.96, and 3.65 respectively.

Moreover TGF-beta type I with $k_i = 0.430 \mu\text{M}$; had one H acceptor bonding of ligand carbonyl group oxygen to nitrogen of LYS_232 residue with distances 3.37, and two pi-H binding among its 5-ring and 6-ring to carbon of ILE_211 and LEU_340 residues with distances 3.69 and 3.79. Also, AT1R with $k_i = 0.768 \mu\text{M}$; had three H acceptor bonding of ligand nitrogen to nitrogen of ARG_167 residue with distances 3.42, 3.34, and 3.53. Similar data were also detected for MMP3 ($k_i = 0.796 \mu\text{M}$; two H donor binding between carbon and nitrogen of ligand and both of oxygen of LEU_222 residue (distance: 3.29) and nitrogen of HIS_166 residue (distance: 3.49), two H acceptor binding between ligand nitrogen and both of carbon of VAL_163 residue (distance: 3.69) and nitrogen of LEU_164 residue (distance: 3.43), one pi-H binding of ligand carbon and 5-ring of HIS_205 residue with distance 4.60. Furthermore, Renin with $k_i = 0.815 \mu\text{M}$ had three H acceptor between oxygen of THR_85 residue and nitrogen and oxygen of ligand (distances: 2.99, 2.90 and 3.02), one pi-H binding between of 5-ring of ligand and SER_84 residue with distance 3.85.

The next factor was MMP9 with $k_i = 1.636 \mu\text{M}$ with following information: two H acceptor binding of ligand oxygen to nitrogen of LEU_188 and ALA_189 residues with distance 3.26. As shown in Table 2 and Figure 6, Survivin with $p_k_i = 1.779 \mu\text{M}$ had one H acceptor binding between carbonyl group oxygen and nitrogen of LYS_15 residue with distance 3.02, one H donor binding of nitrogen to oxygen of GLN_92 residue with distance 2.79, one pi-H between 5-ring of ligand and nitrogen of PHE_93 residue with distance 4.03. Also, ACE2 with $k_i = 149.862 \mu\text{M}$ had one H donor binding among nitrogen of ligand to oxygen of ASP_30 residue with distance 3.48.

According to the above-mentioned results, ACE, TGF-beta type I and II, AT1R, MMP3, and Renin, with $k_i \leq 0.81$ were found to have a strong bond to inhibit Candesartan, due to have lowest inhibition constant between the mentioned drug and these receptors.

DISCUSSION

The therapeutic potential of inhibiting the renin-angiotensin system using AT1R blockers appears to be promising for the management of various cancers including breast, bladder, pancreas, prostate, renal, and colon cancer (Kosugi et al., 2009; Chen et al., 2013; Alhusban et al., 2014; Okazaki et al., 2014; Asgharzadeh et al., 2018, 2020a). In this study, we used systems and molecular biology to explore the molecular mechanisms of the anti-tumor effects of candesartan in CRC progression in cellular and animal models. Our results suggest that candesartan elicits potent anti-tumor properties by inhibiting cell proliferation, migration, and inducing tumor cell necrosis in CRC.

The renin-angiotensin system is overexpressed in various human cancers which is associated with poor prognosis and aggressive phenotype (Ishikane and Takahashi-Yanaga, 2018; Hashemzahi et al., 2021b). Moreover, angiotensin II has the potential to promote tumor cell proliferation and invasion through inducing Wnt/ β -catenin signaling pathway (Yu et al., 2016; Li et al., 2018). Cyclin D1 as a key downstream target of Wnt/ β -catenin signaling is overexpressed in several human cancers including CRC (Rahmani et al., 2018; Soleimani et al., 2019). Cyclin D1, as a regulatory subunit for cyclin-dependent kinase enzymes, plays an essential role in cell proliferation and cell cycle progression (Peurala et al., 2013). In the present study, we evaluated the anti-tumor effects of candesartan on CRC cells through investigating expression of cyclin D1 and MMPs. Our findings indicated that candesartan treatment inhibits Wnt/ β -

catenin signaling as visualized by downregulation of cyclin D1, survivin, and MMPs mRNA levels.

Recently, inhibition of Wnt/ β -catenin signaling pathway was shown to be related to suppressing tumor cell migration and metastasis (Tong et al., 2016). The MMP enzyme as downstream effectors of Wnt/ β -catenin signaling were reported to be involved in cell migration and invasion (Jia et al., 2017; Chen et al., 2021). In line with this data, we illustrated that treatment with candesartan inhibits cancer cell migration. To further confirm the suppressive effects of candesartan on cellular migration the expression levels of E-cadherin

and MMP3/9 were investigated and our results indicate that candesartan inhibits migratory behavior of CRC cells by induction of E-cadherin and downregulation of MMP3/9.

Further studies showed that inhibition of Wnt signaling induces tumor cell apoptosis in CRC and breast cancer (Bilir et al., 2013). Wnt/ β -catenin signaling was identified as a pro-survival pathway and its inhibition was shown to be involved in cancer cell death and apoptosis (Jia et al., 2019). In agreement with these results, we demonstrated that candesartan induces apoptosis/necrosis in CRC cell lines.

Table 2: Data for the docking interactions of Candesartan Cilexetil at the active sites of following receptors as the molecular targets.

Target Receptors	Docking score (kcal/mol)	K_i (μ M)	H-bonding Interactions (Distance (Å))	pi-H binding Interactions (Distance (Å))	Pi-cation Interactions (Distance (Å))
ACE	-9.193	0.179	THR_301 (3.22) THR_302 (2.89)	-	LYS_449(3.89)
TGF-beta type II	-8.690	0.420	CYS_396 (3.86) GLY_251 (3.52)	GLY_253 (3.71) VAL_258 (3.96) SER_383 (3.65)	LYS_381 (3.68)
TGF-beta type I	-8.675	0.430	LYS_232 (3.37)	ILE_211 (3.69) LEU_340 (3.79)	-
AT1R	-8.332	0.768	ARG_167 (3.42) ARG_167 (3.34) ARG_167 (3.53)	-	-
MMP3	-8.310	0.796	LEU_222 (3.29) HIS_166 (3.49) VAL_163 (3.69) LEU_164 (3.43)	HIS_205 (4.60)	-
Renin	-8.297	0.815	THR_85 (2.99) THR_85 (2.90) THR_85 (3.02)	SER_84 (3.85)	-
MMP9	-7.884	1.636	LEU_188 (3.26) ALA_189 (3.26)	-	-
Survivin	-7.83486	1.779	GLN_92 (2.79) LYS_15 (3.02)	PHE_93 (4.03)	-
COL1A1	-7.417	3.605	GLY_18 (3.46) ARG_17 (2.98) VAL_22 (3.45) ARG_17 (2.54)	-	-
E-cadherin	-5.532	87.123	LYS_25 (3.94)	-	-
COL1A2	-5.350	118.428	GLU_54 (2.93) GLN_55 (3.17)	-	-
ACE2	-5.211	149.862	ASP_30 (3.48)	-	-
IL-6	-5.033	202.365	PRO_65 (3.33) LYS_86 (3.56)	-	-
TNF- α	-4.681	366.739	TYR_227 (3.36)	-	-
CyclinD1	-4.378	612.073	ASP_294 (3.12)	-	ARG_291 (3.52)

We also investigated the anti-cancer effects of candesartan in a colon cancer mouse model. Our results demonstrated that candesartan inhibits cancer progression and development through inducing tumor necrosis and changing the oxidant/antioxidant balance in tumor tissue. Results of our study indicate that administration of candesartan increased MDA level and reduced total thiol, SOD, and catalase levels. These results suggest that alterations of oxidant/antioxidant status can be one of the underlying mechanisms contributed to candesartan anti-tumor activities against colon cancer cells. Taken together, we have shown that candesartan elicits its anti-cancer effects by suppressing CRC cell growth and migration, impairing oxidant/antioxidant balance, and enhancing tumor necrosis in CRC cells and tumors, suggesting the therapeutic potential of targeting renin-angiotensin system in colon cancer treatment. Finally, our study demonstrated that candesartan synergistically enhanced the anti-tumor effects of 5-FU suggesting that the AT1R blockers may be used as novel adjuvants in combination with a chemotherapeutic agent in further clinical studies.

Conflict of interest

The authors have no conflict of interest to disclose.

Acknowledgments

This research is part of a PhD thesis of Mr. Ehsan Tabatabai, which was supported by the Tabriz University of Medical Sciences and Mashhad University of Medical Sciences.

REFERENCES

Abraham HMA, White CM, White WB. The comparative efficacy and safety of the angiotensin receptor blockers in the management of hypertension and other cardiovascular diseases. *Drug Saf.* 2015;38:33-54.

Afsar B, Afsar R, Ertuglu L, Kuwabara M, Ortiz A, Covic A, et al. Renin-angiotensin system and cancer: epidemiology, cell signaling, genetics and epigenetics. *Clin Transl Oncol.* 2021;(4):1-15.

Afshari AR, Roshan MK, Soukhtanloo M, Ghorbani A, Rahmani F, Jalili-Nik M, et al. Cytotoxic effects of auroaptene against a human malignant glioblastoma cell line. *Avicenna J Phytomed.* 2019;9:334-46.

Alhusban A, Al-Azayzih A, Goc A, Gao F, Fagan SC, Somanath PR. Clinically relevant doses of candesartan inhibit growth of prostate tumor xenografts in vivo through modulation of tumor angiogenesis. *J Pharmacol Exp Ther.* 2014;350:635-45.

Amerizadeh F, Rezaei N, Rahmani F, Hassanian SM, Moradi-Marjaneh R, Fiuji H, et al. Crocin synergistically enhances the antiproliferative activity of 5-fluorouracil through Wnt/PI3K pathway in a mouse model of colitis-associated colorectal cancer. *J Cell Biochem.* 2018;119:10250-61.

Arjmand M-H. Elucidating the association between the upregulation of angiotensin type 1-receptors and the development of gastrointestinal malignancies. *J Gastrointest Cancer.* 2020; epub ahead of print. doi: 10.1007/s12029-020-00547-0.

Arrieta O, Villarreal-Garza C, Vizcaíno G, Pineda B, Hernández-Pedro N, Guevara-Salazar P, et al. Association between AT1 and AT2 angiotensin II receptor expression with cell proliferation and angiogenesis in operable breast cancer. *Tumor Biol.* 2015;36:5627-34.

Asgharzadeh F, Hassanian SM, Ferns GA, Khazaei M, Hasanzadeh M. The therapeutic potential of angiotensin-converting enzyme and angiotensin receptor inhibitors in the treatment of colorectal cancer: Rational strategies and recent progress. *Curr Pharm Des.* 2018; 24:4652-8.

Asgharzadeh F, Hashemzahi M, Moradi-Marjaneh R, Hassanian SM, Ferns GA, Khazaei M, et al. Angiotensin-converting enzyme inhibitors and angiotensin receptor blockers as therapeutic options in the treatment of renal cancer: A meta-analysis. *Life Sci.* 2020a;242: 117181.

Asgharzadeh F, Mostafapour A, Amerizadeh F, Avan A, Rahmani F, Sabbaghzadeh R, et al. Inhibition of angiotensin II type I pathway reduced tumor growth and ameliorates fibrosis/inflammation associated with colorectal cancer. *Preprints.* 2020b;2020:2020070508. doi: 10.20944/preprints202007.0508.v1.

Asgharzadeh F, Hashemzadeh A, Yaghoobi A, Avan A, Nazari SE, Soleimanpour S, et al. Therapeutic effects of silver nanoparticle containing sulfasalazine on DSS-induced colitis model. *J Drug Deliv Sci Technol.* 2021;61:102133.

- Baguet J-P, Barone-Rochette G, Neuder Y. Candesartan cilexetil in the treatment of chronic heart failure. *Vasc Health Risk Manag.* 2009;5:257-64.
- Bilir B, Kucuk O, Moreno CS. Wnt signaling blockage inhibits cell proliferation and migration, and induces apoptosis in triple-negative breast cancer cells. *J Transl Med.* 2013;11(1):1-12.
- Bregonzio C, Seltzer A, Armando I, Pavel J, Saavedra J. Angiotensin II AT1 receptor blockade selectively enhances brain AT2 receptor expression, and abolishes the cold-restraint stress-induced increase in tyrosine hydroxylase mRNA in the locus coeruleus of spontaneously hypertensive rats. *Stress.* 2008;11:457-66.
- Chen T, Chen Z, Lian X, Wu W, Chu L, Zhang S, et al. MUC 15 Promotes osteosarcoma cell proliferation, migration and invasion through livin, MMP-2/MMP-9 and Wnt/ β -catenin signal pathway. *J Cancer.* 2021;12:467-73.
- Chen X, Meng Q, Zhao Y, Liu M, Li D, Yang Y, et al. Angiotensin II type 1 receptor antagonists inhibit cell proliferation and angiogenesis in breast cancer. *Cancer Lett.* 2013;328:318-24.
- Childers WK. Interactions of the renin-angiotensin system in colorectal cancer and metastasis. *Int J Colorect Dis.* 2015;30:749-52.
- Du N, Feng J, Hu L-J, Sun X, Sun H-B, Zhao Y, et al. Angiotensin II receptor type 1 blockers suppress the cell proliferation effects of angiotensin II in breast cancer cells by inhibiting AT1R signaling. *Oncol Rep.* 2012;27:1893-903.
- Enofe N, Morris AD, Liu Y, Liang W, Wu CS, Sullivan PS, et al. Receipt of adjuvant chemotherapy in stage ii colon cancer and overall survival: A national cancer database study. *J Surg Res.* 2020;252:69-79.
- Fan F, Tian C, Tao L, Wu H, Liu Z, Shen C, et al. Candesartan attenuates angiogenesis in hepatocellular carcinoma via downregulating AT1R/VEGF pathway. *Biomed Pharmacother.* 2016;83:704-11.
- Ghanaatgar-Kasbi S, Amerizadeh F, Rahmani F, Hassanian SM, Khazaei M, Ferns GA, et al. AMP-kinase inhibitor dorsomorphin reduces the proliferation and migration behavior of colorectal cancer cells by targeting the AKT/mTOR pathway. *IUBMB Life.* 2019;71:1929-36.
- Hashemzahi M, Behnam-Rassouli R, Hassanian SM, Moradi-Binabaj M, Moradi-Marjaneh R, Rahmani F, et al. Phytosomal-curcumin antagonizes cell growth and migration, induced by thrombin through AMP-Kinase in breast cancer. *J Cell Biochem.* 2018;119:5996-6007.
- Hashemzahi M, Rahmani F, Khoshakhlagh M, Avan A, Asgharzadeh F, Barneh F, et al. Angiotensin receptor blocker Losartan inhibits tumor growth of colorectal cancer. *EXCLI J.* 2021a;20:506-21.
- Hashemzahi M, Yavari N, Rahmani F, Asgharzadeh F, Soleimani A, Shakour N, et al. Inhibition of transforming growth factor-beta by Tranilast reduces tumor growth and ameliorates fibrosis in colorectal cancer. *EXCLI J.* 2021b;20:601-13.
- Ishikane S, Takahashi-Yanaga F. The role of angiotensin II in cancer metastasis: potential of renin-angiotensin system blockade as a treatment for cancer metastasis. *Biochem Pharmacol.* 2018;151:96-103.
- Jia S, Qu T, Wang X, Feng M, Yang Y, Feng X, et al. KIAA1199 promotes migration and invasion by Wnt/ β -catenin pathway and MMPs mediated EMT progression and serves as a poor prognosis marker in gastric cancer. *PLoS One.* 2017;12(4):e0175058.
- Jia Y, Chen L, Guo S, Li Y. Baicalin induced colon cancer cells apoptosis through miR-217/DKK1-mediated inhibition of Wnt signaling pathway. *Mol Biol Rep.* 2019;46:1693-700.
- Kosugi M, Miyajima A, Kikuchi E, Kosaka T, Horiguchi Y, Murai M, et al. Angiotensin II type 1 receptor antagonist enhances cis-dichlorodiammineplatinum-induced cytotoxicity in mouse xenograft model of bladder cancer. *Urology.* 2009;73:655-60.
- Li Z, Yan Z, Xu C, Dong Y, Xiong Y, Dai Y. Acetylshikonin attenuates angiotensin II-induced proliferation and motility of human brain smooth muscle cells by inhibiting Wnt/ β -catenin signaling. *Human Cell.* 2018;31:242-50.
- Mc Menamin UC, Murray LJ, Cantwell MM, Hughes CM. Angiotensin-converting enzyme inhibitors and angiotensin receptor blockers in cancer progression and survival: A systematic review. *Cancer Causes Control.* 2012;23:221-30.
- Mengden T, Uen S, Bramlage P. Management of hypertension with fixed dose combinations of candesartan cilexetil and hydrochlorothiazide: Patient perspectives and clinical utility. *Vasc Health Risk Manag.* 2009;5:1043-58.
- Minatoguchi S, Aoyama T, Kawai N, Iwasa M, Oda M, Kida K, et al. Comparative effect of candesartan and amlodipine, and effect of switching from valsartan, losartan, telmisartan and olmesartan to candesartan, on early morning hypertension and heart rate. *Blood Pressure.* 2013;22(Suppl 1):29-37.

- Miyajima A, Kosaka T, Asano T, Asano T, Seta K, Kawai T, et al. Angiotensin II type I antagonist prevents pulmonary metastasis of murine renal cancer by inhibiting tumor angiogenesis. *Cancer Res.* 2002;62:4176-9.
- Molteni A, Heffelfinger S, Moulder JE, Uhal B, Castellani WJ. Potential deployment of angiotensin I converting enzyme inhibitors and of angiotensin II type 1 and type 2 receptor blockers in cancer chemotherapy. *Anticancer Agents Med Chem.* 2006;6:451-60.
- Nakai Y, Isayama H, Ijichi H, Sasaki T, Takahara N, Ito Y, et al. A multicenter phase II trial of gemcitabine and candesartan combination therapy in patients with advanced pancreatic cancer: GECA2. *Invest New Drugs.* 2013;31:1294-9.
- Okazaki M, Fushida S, Harada S, Tsukada T, Kinoshita J, Oyama K, et al. The angiotensin II type 1 receptor blocker candesartan suppresses proliferation and fibrosis in gastric cancer. *Cancer Lett.* 2014;355:46-53.
- Pei N, Mao Y, Wan P, Chen X, Li A, Chen H, et al. Angiotensin II type 2 receptor promotes apoptosis and inhibits angiogenesis in bladder cancer. *J Exp Clin Cancer Res.* 2017;36:1-12.
- Peurala E, Koivunen P, Haapasaari K-M, Bloigu R, Jukkola-Vuorinen A. The prognostic significance and value of cyclin D1, CDK4 and p16 in human breast cancer. *Breast Cancer Res.* 2013;15(1):R5.
- Piastowska-Ciesielska AW, Kozłowski M, Wagner W, Domińska K, Ochędalski T. Effect of an angiotensin II type 1 receptor blocker on caveolin-1 expression in prostate cancer cells. *Arch Med Sci.* 2013;9(4):739.
- Rahmani F, Avan A, Hashemy SI, Hassanian SM. Role of Wnt/ β -catenin signaling regulatory microRNAs in the pathogenesis of colorectal cancer. *J Cell Physiol.* 2018;233:811-7.
- Rahmani F, Amerizadeh F, Hassanian SM, Hashemzahi M, Nasiri SN, Fiuji H, et al. PNU-74654 enhances the antiproliferative effects of 5-FU in breast cancer and antagonizes thrombin-induced cell growth via the Wnt pathway. *J Cell Physiol.* 2019;234:14123-32.
- Rahmani F, Asgharzadeh F, Avan A, Barneh F, Parizadeh MR, Ferns GA, et al. Rigosertib potently protects against colitis-associated intestinal fibrosis and inflammation by regulating PI3K/AKT and NF- κ B signaling pathways. *Life Sci.* 2020;249:117470.
- Renziehausen AWJ. Angiotensin II type 2 receptor signalling promotes glioblastoma growth: A novel target for therapeutic intervention. Ph.D. thesis. London: Imperial College London, 2017.
- Renziehausen A, Wang H, Rao B, Weir L, Nigro CL, Lattanzio L, et al. The renin angiotensin system (RAS) mediates bifunctional growth regulation in melanoma and is a novel target for therapeutic intervention. *Oncogene.* 2019;38:2320-36.
- Shakour N, Ruscica M, Hadizadeh F, Cirtori C, Banach M, Jamialahmadi T, et al. Statins and C-reactive protein: In silico evidence on direct interaction. *Arch Med Sci.* 2020;16:1432-9.
- Soleimani A, Rahmani F, Saeedi N, Ghaffarian R, Khazaei M, Ferns GA, et al. The potential role of regulatory microRNAs of RAS/MAPK signaling pathway in the pathogenesis of colorectal cancer. *J Cell Biochem.* 2019;120:19245-53.
- Soleimani A, Asgharzadeh F, Rahmani F, Avan A, Mehraban S, Fakhraei M, et al. Novel oral transforming growth factor- β signaling inhibitor potently inhibits postsurgical adhesion band formation. *J Cell Physiol.* 2020;235:1349-57.
- Tong Y, Liu Y, Zheng H, Zheng L, Liu W, Wu J, et al. Artemisinin and its derivatives can significantly inhibit lung tumorigenesis and tumor metastasis through Wnt/ β -catenin signaling. *Oncotarget.* 2016;7:31413-28.
- Wong MC, Huang J, Lok V, Wang J, Fung F, Ding H, et al. Differences in incidence and mortality trends of colorectal cancer, worldwide, based on sex, age, and anatomic location. *Clin Gastroenterol Hepatol.* 2021; 19:955-66.e61.
- Yu L, Meng W, Ding J, Cheng M. Klotho inhibits angiotensin II-induced cardiomyocyte hypertrophy through suppression of the AT1R/ β catenin pathway. *Biochem Biophys Res Commun.* 2016;473:455-61.