

.....  
*The European Journal of Public Health*, Vol. 29, No. 6, 1043–1048

© The Author(s) 2019. Published by Oxford University Press on behalf of the European Public Health Association.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.  
 doi:10.1093/eurpub/ckz086 Advance Access published on 23 May 2019  
 .....

## Secular trends in stillbirth by maternal socioeconomic status in Spain 2007–15: a population-based study of 4 million births

Miguel Angel Luque-Fernandez<sup>1,2,3,4,6</sup>, Aurielle Thomas<sup>5</sup>, Bizu Gelaye<sup>3</sup>, Judith Racape<sup>4</sup>, Maria Jose Sanchez<sup>1,6</sup>, Michelle A. Williams<sup>3</sup>

1 Andalusian School of Public Health, Biomedical Research Institute of Granada, University of Granada, Granada, Spain

2 Department of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, UK

3 Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA

4 Public Health School, Centre de Recherche Épidémiologie, Biostatistique et Recherche Clinique, Université Libre de Bruxelles, Brussels, Belgium

5 Department of Population Medicine, Harvard Medical School, Harvard Pilgrim Health Care Institute, Boston, MA, USA

6 Biomedical Research Networking Centres of Epidemiology and Public Health (CIBERESP), Madrid, Spain

**Correspondence:** Miguel Angel Luque-Fernandez, Andalusian School of Public Health, Biomedical Research Institute of Granada, University of Granada, Cuesta del Observatorio, 4, Campus Universitario de la Cartuja., Granada 18010, Spain, Tel: +34 677 903 936, e-mail: miguel.luque.easp@juntadeandalucia.es

**Introduction:** Stillbirth, one of the urgent concerns of preventable perinatal deaths, has wide-reaching consequences for society. We studied secular stillbirth trends by maternal socioeconomic status (SES) in Spain. **Methods:** We developed a population-based observational study, including 4 083 919 births during 2007–15. We estimate stillbirth rates and secular trends by maternal SES. We also evaluated the joint effect of maternal educational attainment and the Human Development Index (HDI) of women's country of origin on the risk of stillbirth. The data and statistical analysis can be accessed for reproducibility in a GitHub repository: <https://github.com/migariane/Stillbirth> **Results:** We found a consistent pattern of socioeconomic inequalities in the risk of delivering a stillborn, mainly characterized by a persistently higher risk, over time, among women with lower SES. Overall, women from countries with low HDIs and low educational attainments had approximately a four times higher risk of stillbirth (RR: 4.44; 95%CI: 3.71–5.32). Furthermore, we found a paradoxical reduction of the stillbirth gap over time between the highest and the lowest SESs, which is mostly due to the significant and increasing trend of stillbirth risk among highly educated women of advanced maternal age. **Conclusion:** Our findings highlight no improvement in stillbirth rates among women of lower SES and an increasing trend among highly educated women of advanced maternal age over recent years. Public health policies developing preventive programmes to reduce stillbirth rates among women with lower SES are needed as well as the necessity of further study to understand the growing trend of age-related stillbirths among highly educated women in Spain.  
 .....

### Introduction

Stillbirths are an important health outcome and a tragedy that deserves attention.<sup>1</sup> Often, stillbirth rates are used as an indicator of the level of development of a country.<sup>2</sup> Despite the fact that western countries' stillbirth rates are lower than those in developing countries, stillbirth rates have remained relatively stable over the last few decades.<sup>3</sup>

Higher stillbirth rates have been found in Western countries among women of lower socioeconomic status (SES) and specific ethnic groups.<sup>3–6</sup> Maternal educational attainment and country of origin at birth have been used by several authors as a proxy for maternal SES.<sup>3,4,7,8</sup> The use of maternal country of origin at birth, in this way, is supported by studies which have shown evidence of worse perinatal outcomes among immigrant women in high-income countries.<sup>9,10</sup>

Europeristat, a European report focussing on perinatal health, recommended that stillbirth rates should be used as an indicator to allow international comparisons.<sup>11</sup> Stillbirth risk provides information on avoidable mortality and reveals problems in the quality of perinatal care. Secular trends in perinatal mortality have been used as a public health indicator, as it is highly sensitive to social and health inequalities.<sup>12</sup>

We described stillbirth secular trends by maternal country of origin at birth, the 2015 Human Development Index (HDI) of their respective country of origin and maternal educational attainment as proxies of maternal SES during the period 2007–15 in Spain. Secondly, we explored joint effects on stillbirth of (i) low maternal educational attainment and advanced maternal age and (ii) low maternal educational attainment and low HDI.

### Methods

#### *Type of study and data source*

We designed a population-based observational study using vital statistics from Spain's National Institute of Statistics (INE). The final dataset contains basic sociodemographic, obstetric and perinatal information for 11 323 stillbirths and 4 179 402 total births in Spain from 2007–15. Infants born before 28 gestational weeks were excluded (Supplementary figure S1).

#### *Main outcome and exposures*

The main outcome of the study is stillbirth, which is defined as foetal death at  $\geq 28$  gestation weeks at delivery.<sup>13</sup> The main exposure of interest was maternal SES. We used maternal educational attainment

at the country of birth, maternal country of origin at birth and the 2015 HDI from the women's country of origin as proxies of maternal SES. The maternal educational attainment refers to the highest degree obtained by the woman at the time of delivery and is based on the International Standard Classification of Education.<sup>14</sup> Maternal educational attainment was categorized as (i) secondary education or lower (mothers with  $\leq 12$  years of the mandatory education in Spain), (ii) upper secondary and first stage of tertiary education (mothers with  $>12$  but  $\leq 15$  years of education) and (iii) tertiary education (mothers who received  $>15$  years of education). The HDI quantifies the social progress of countries, based on a summary measure of average achievement in key dimensions of human development: a long and healthy life, being knowledgeable, and a decent standard of living. The index is quantified as the geometric mean of normalized indices for each of the three previous dimensions.<sup>15</sup> We categorized the HDI into four groups: low, medium, high and very high HDI.<sup>15</sup> Finally, we categorized maternal country of origin into six groups: Africa, America and the Caribbean, Asia and Oceania, other European Union countries with 15-member states (EU15), non-EU15 European Countries, and Spain.<sup>11</sup>

### Other variables

Parity, calendar year, maternal age (years) at the time of delivery were included in the statistical analysis. Parity was dichotomized as nulliparous (women who have never previously given birth) and multiparous (women who have  $\geq 1$  prior births). Maternal age was categorized consistently based on international recommendations as follows:  $\leq 19$ , 20–24, 25–29, 30–34, 35–39 and  $\geq 40$  years.<sup>16</sup>

### Statistical analysis

In descriptive analyses, we summarized the total number of births and stillbirths by baseline characteristics and calendar year. Next, we computed stillbirth rates and univariate rate ratios with 95% CIs by population characteristics. We also computed linear trend tests for stillbirth rates by maternal country of origin and educational attainment over the time-period in analysis. Furthermore, we computed stillbirth rates of change (percent) for maternal educational attainment by calendar period and categories of maternal age ( $<35$ , 35–39 and  $\geq 40$  years).

In multivariable analyses, we used generalized linear regression models with family Poisson and link log to derive rate ratios. We derived 'sandwich' standard errors for statistical inference.<sup>17</sup> The final multivariable model was adjusted for maternal age, educational attainment, parity, HDI and calendar period. We used this model to estimate the joint effect on stillbirth of maternal educational attainment and maternal age at delivery. Age was introduced in the model with four categories ( $<25$ , 25–29, 30–34 and  $\geq 35$ ) as we combined the categories of age  $\geq 19$  and 20–24 given the reduced number of women with tertiary studies among this group of age. Similarly, we also assessed the joint effect on stillbirth of maternal educational attainment and HDI (very high, high, medium and low). Based on the linear combination of the coefficients we derived rate ratios for each of the categories of the linear combination considering women with tertiary education, very high HDI and 25–29 years the reference group against which, women in the other groups were compared.

Data were analyzed using the statistical software Stata Multi-Processor Parallel Edition v.15 (StataCorp, College Station, TX, USA) and R v.3.3.4 (R Foundation for Statistical Computing, Vienna, Austria). The data and statistical analysis can be accessed for reproducibility in a GitHub repository: <https://github.com/migriane/Stillbirth>

## Results

During the 9-year study period (2007–15), 4 207 372 infants were born in Spain. We excluded 11 450 births ( $<28$  gestation weeks and 16 520 (0.4%) births not linked to the 2015 HDI. The final dataset included 4 179 402 births (11 323 stillbirths and 4 168 09 live births) (Supplementary figure S1).

Overall, stillbirth rates per 1000 births were higher among very young ( $\leq 19$  years) and those of advanced reproductive age ( $\geq 35$  years). Furthermore, stillbirth rates were higher among women with an African country of origin, with low educational attainment, and low HDI, respectively. Stillbirth rates were approximately two times higher risk for African women, low educational attainment, and low HDI. Women from low HDI countries showed approximately three-times higher risk of stillbirth with a univariate rate ratio of 2.8 (95%CI 2.4–3.3). There was strong evidence of an increasing linear trend in stillbirth risk across levels of the variables maternal age, educational attainment and HDI (table 1, Supplementary table S1). However, no evidence of a secular trend by calendar period over the 9-year period was observed (test of linear trend  $P$  values = 0.946) (table 1).

Regarding women's country of origin, there was no evidence of a secular trend in stillbirth risk for any of the other five maternal regions of origin including native women (test of linear trend  $P$  values = 0.342; Supplementary figure S2). However, there was strong evidence of a secular trend by maternal educational attainment (test of linear trend  $P$  values  $< 0.001$ ) (figure 1).

The gap in stillbirth rates between the highest and lowest levels of maternal education attainment decreased over the 9-year period; mostly because the average rate of change of stillbirth rates was approximately two times higher among women with tertiary education than among mothers with secondary or lower educational attainment: 5.2% (95%CI: 2.3–8.1%) vs. 2.5% (95%CI: 1.4–3.5%) (figure 1). Among highly educated women (tertiary education), the rate of change of stillbirth was 1.6 times higher for women aged 35–39 years compared with the rate of change of women  $<35$  years of age (rates of change: 5.2%; 95%CI: 2.3–8.1 and 3.2%; 95%CI:  $-0.1$ –7.6%, respectively). Furthermore, the rate of change was also greater for women aged  $\geq 40$  compared with the rate of women aged  $<35$  years, 4.3 vs. 3.2%, respectively (figure 2).

In the multivariable analysis, low HDI, low educational attainment and advanced maternal age were strongly associated with higher risk of stillbirth, showing approximately two times higher risk than women from countries with very high HDI, tertiary education and aged between 20 and 24 years (table 2, model 5).

Overall, the joint effect of women with low educational attainment and low HDI increased approximately four times the risk of delivering a stillborn compared with women with tertiary education and very-high HDI. Similarly, the joint effect of advanced maternal age with low educational attainment increased four times the risk of delivering a stillborn compared with highly educated women aged between 25 and 29 years (Supplementary table S2).

## Discussion


We have found a consistent pattern of socioeconomic inequalities in the risk of delivering a stillborn in Spain during 2007–15, mainly characterized by a persistently higher stillbirth risk over time among women with lower SES (women from countries with low HDI, particularly Africa, and low educational attainment). Furthermore, we have found a paradoxical reduction of the stillbirth gap over time between the highest and the lowest SES mostly due to an increasing trend on stillbirth risk among highly educated women with advanced maternal age.

The persistent increased risk among women with lower SES is an important public health cause of concern in addition to the

**Table 1** Univariate stillbirth rate among women at least 28 weeks' gestation by maternal age, country of origin, SES, parity and period in Spain, during 2007–15 (11 323 stillbirths and 4 179 402 total births)

| Variables  | Total births (n) | Stillbirths (n) | Rate per 1000 births (95%CI) | Rate ratio (95%CI) | P-value |
|--|------------------|-----------------|------------------------------|--------------------|---------|
| Maternal age in years                                  |                  |                 |                              |                    | <0.001  |
| ≤19  | 100 082          | 321             | 3.2 (2.9, 3.6)               | 1.17 (1.02, 1.35)  |         |
| 20–24  | 355 059          | 1015            | 2.8 (2.7, 3.0)               | 1.0 (Reference)    |         |
| 25–29  | 845 889          | 2218            | 2.6 (2.5, 2.7)               | 0.93 (0.85, 1.00)  |         |
| 30–34  | 1 561 448        | 3781            | 2.4 (2.3, 2.5)               | 0.85 (0.79, 0.92)  |         |
| 35–39  | 1 070 715        | 3093            | 2.9 (2.7, 3.0)               | 1.03 (0.95, 1.11)  |         |
| ≥40  | 234 886          | 895             | 3.8 (3.6, 4.1)               | 1.37 (1.24, 1.52)  |         |
| Parity   |                  |                 |                              |                    | <0.001  |
| Nulliparous (first delivery)                           | 2 106 505        | 6373            | 3.0 (2.9, 3.1)               | 1.18 (1.13, 1.23)  |         |
| Multiparous (≥1 deliveries)                            | 2 061 574        | 4950            | 2.4 (2.3, 2.5)               | 1.00 (Reference)   |         |
| Maternal country of origin                             |                  |                 |                              |                    | <0.001  |
| Spain  | 3 372 357        | 8545            | 2.5 (2.4, 2.6)               | 1.00 (Reference)   |         |
| EU15   | 60 990           | 142             | 2.3 (2.0, 2.7)               | 0.93 (0.77, 1.11)  |         |
| Other European countries                               | 160 896          | 435             | 2.7 (2.5, 3.0)               | 1.10 (0.99, 1.23)  |         |
| Africa   | 264 356          | 1290            | 4.9 (4.6, 5.1)               | 2.31 (2.16, 2.47)  |         |
| America  | 248 056          | 720             | 2.9 (2.7, 3.1)               | 1.20 (1.11, 1.31)  |         |
| Asia and Oceania                                       | 61 424           | 191             | 3.1 (2.7, 3.6)               | 1.84 (1.58, 2.14)  |         |
| HDI <sup>a</sup> for maternal country of origin (2015) |                  |                 |                              |                    | <0.001  |
| Low  | 33 686           | 176             | 5.2 (4.4, 6.0)               | 2.78 (2.36, 3.27)  |         |
| Medium   | 340 616          | 1430            | 4.2 (4.0, 4.4)               | 1.96 (1.84, 2.09)  |         |
| High   | 204 019          | 602             | 2.9 (2.7, 3.2)               | 1.24 (1.14, 1.36)  |         |
| Very high  | 3 589 758        | 9115            | 2.5 (2.4, 2.6)               | 1.00 (Reference)   |         |
| Maternal education attainment <sup>b</sup>             |                  |                 |                              |                    | <0.001  |
| Secondary education or lower                           | 2 396 836        | 7087            | 3.0 (2.9, 3.1)               | 2.36 (2.19, 2.54)  |         |
| Upper secondary or first stage of tertiary             | 965 730          | 1413            | 1.5 (1.4, 1.5)               | 1.12 (1.02, 1.23)  |         |
| Tertiary education                                     | 719 034          | 906             | 1.3 (1.2, 1.3)               | 1.00 (Reference)   |         |
| Period   |                  |                 |                              |                    | 0.946   |
| 2007   | 491 288          | 1099            | 2.7 (2.5, 2.8)               | 1.00 (Reference)   |         |
| 2008   | 516 602          | 1409            | 2.7 (2.6, 2.9)               | 0.98 (0.91, 1.06)  |         |
| 2009   | 491 856          | 1344            | 2.7 (2.6, 2.9)               | 0.96 (0.91, 1.05)  |         |
| 2010   | 483 456          | 1303            | 2.7 (2.5, 2.8)               | 0.95 (0.87, 1.03)  |         |
| 2011   | 468 936          | 1288            | 2.7 (2.6, 2.9)               | 0.99 (0.91, 1.07)  |         |
| 2012   | 451 678          | 1259            | 2.8 (2.6, 2.9)               | 1.01 (0.92, 1.09)  |         |
| 2013   | 422 742          | 1188            | 2.8 (2.6, 3.0)               | 1.00 (0.92, 1.09)  |         |
| 2014   | 424 498          | 1111            | 2.6 (2.5, 2.8)               | 0.95 (0.87, 1.04)  |         |
| 2015   | 417 023          | 1112            | 2.7 (2.5, 2.8)               | 0.98 (0.89, 1.06)  |         |

a: HDI (UNESCO).  
 b: 2.1% (88 396) missing values for maternal education attainment.



**Figure 1** Stillbirth rate of births ≥28 gestational weeks by maternal education in Spain from 2007 to 2015 (11 113 stillbirths and 4 083 919 total births)




Figure 2 Stillbirth rate of births ≥28 gestational weeks by maternal education and age in Spain from 2007 to 2015 (11 113 stillbirths and 4 083 919 total births)

Table 2 Stillbirth rate ratios comparing the HDI for maternal countries of origin adjusted for maternal age, SES, parity and period in Spain, during 2007–15 (11 323 stillbirths and 4 179 402 total births)

| Variables                                  | Model 1           | Model 2           | Model 3           | Model 4           | Model 5           |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|
| <b>HDI</b>                                 |                   |                   |                   |                   |                   |
| Low  | 2.05 (1.76, 2.38) | 2.10 (1.80, 2.43) | 1.67 (1.40, 1.96) | 1.74 (1.47, 2.05) | 1.73 (1.46, 2.04) |
| Medium                                     | 1.65 (1.56, 1.75) | 1.69 (1.60, 1.78) | 1.33 (1.24, 1.42) | 1.38 (1.29, 1.46) | 1.37 (1.28, 1.46) |
| High                                       | 1.16 (1.06, 1.26) | 1.17 (1.08, 1.28) | 0.94 (0.85, 1.03) | 0.96 (0.87, 1.05) | 0.97 (0.87, 1.06) |
| Very high                                  | 1.00 (Reference)  | 1.00 (Reference)  | 1.00 (Reference)  | 1.00 (Reference)  | 1.00 (Reference)  |
| <b>Maternal age in years</b>               |                   |                   |                   |                   |                   |
| ≤19  |                   | 1.18 (1.04, 1.34) | 1.15 (1.01, 1.32) | 1.10 (0.96, 1.26) | 1.10 (0.96, 1.26) |
| 20–24                                      |                   | 1.00 (Reference)  | 1.00 (Reference)  | 1.00 (Reference)  | 1.00 (Reference)  |
| 25–29                                      |                   | 1.00 (0.89, 1.04) | 1.02 (0.93, 1.10) | 1.04 (0.96, 1.13) | 1.04 (0.96, 1.13) |
| 30–34                                      |                   | 0.94 (0.88, 1.01) | 1.14 (1.06, 1.23) | 1.22 (1.12, 1.32) | 1.21 (1.12, 1.31) |
| 35–39                                      |                   | 1.14 (1.06, 1.22) | 1.43 (1.32, 1.55) | 1.60 (1.47, 1.73) | 1.58 (1.45, 1.71) |
| ≥40  |                   | 1.48 (1.35, 1.62) | 1.81 (1.64, 2.00) | 2.03 (1.83, 2.24) | 1.99 (1.79, 2.20) |
| <b>Maternal education attainment</b>       |                   |                   |                   |                   |                   |
| Secondary education or lower               |                   |                   | 2.48 (2.31, 2.66) | 2.56 (2.38, 2.75) | 2.57 (2.39, 2.77) |
| Upper secondary or first stage of tertiary |                   |                   | 1.21 (1.11, 1.31) | 1.21 (1.11, 1.31) | 1.21 (1.11, 1.31) |
| Tertiary education                         |                   |                   | 1.00 (Reference)  | 1.00 (Reference)  | 1.00 (Reference)  |
| <b>Parity</b>                              |                   |                   |                   |                   |                   |
| Nulliparous (first delivery)               |                   |                   |                   | 1.31 (1.26, 1.37) | 1.31 (1.26, 1.37) |
| Multiparous (≥1 deliveries)                |                   |                   |                   | 1.00 (Reference)  | 1.00 (Reference)  |
| <b>Period 2007–15</b>                      |                   |                   |                   |                   |                   |
| Per 1-year increase                        |                   |                   |                   |                   | 1.02 (1.01, 1.03) |

Notes: Model 1 univariate logistic model (outcome: stillbirth; exposure: 2015 HDI for maternal country of origin); Model 2 same as model 1 adjusted for age; Model 3 same as model 1 adjusted for age, maternal and education; Model 4 same as model 1 adjusted for age, maternal education and parity; Model 5 same as model 1 adjusted for age, maternal education, parity and period.

increasing stillbirth rates among highly educated women with advanced maternal age during the last 9-year period in Spain. Similar studies provide evidence of greater risk for stillbirth among women with low educational attainment.<sup>3</sup> Furthermore, women living in regions with higher unemployment rates in Spain were shown to have twice the risk of stillbirth compared with women living in regions with less unemployment.<sup>18</sup> However, the increasing trends in stillbirth among older women of higher education may reflect infertility problems associated with delayed childbearing and the use of assisted reproductive technologies.

Multiple studies have shown that lower SES can impact negatively pregnancy outcomes, whether due to differences in access to prenatal

care or with regard to preferences about how to handle high-risk pregnancies.<sup>19</sup> A systematic review examining pregnancy outcomes from immigrant women in western countries showed that women from Sub-Saharan African countries had higher perinatal mortality.<sup>20</sup> Recently, a study showed similar findings in Brussels, Belgium.<sup>21</sup> It has been described that social barriers from non-native foreign women can limit both access to antenatal screening and the ability to detect foetal growth restriction.<sup>22</sup>

To the best of our knowledge, this is the first study in Spain to demonstrate greater stillbirth risk among mothers of low SES. Furthermore, we improve the characterization of maternal SES adding to the maternal country of origin at birth two additional

determinants of maternal SES: (i) HDI from the maternal country of origin at birth (women from countries, with low HDI,<sup>14</sup> are more likely to have been exposed to adverse childhood environment<sup>23</sup>) (ii) and maternal educational attainment. It represents an improvement compared with other studies that only use maternal country of origin at birth allowing to better identify higher stillbirth risk groups.

We consider the migration phenomenon in Spain as a natural experiment given that immigration in Spain is an extremely recent phenomenon (late in the twentieth century), which means that an important number of foreign non-EU15 women included in the study are the first generations of economic migrants.<sup>24</sup> Thus, the maternal country of origin at birth in Spain reflects, more precisely, on the maternal SES given the lower probability of acculturation. For instance, we argue that women from countries of low HDI and low educational attainment (i.e. women from some sub-Saharan countries) are less likely to have been misclassified as low SES.

Previous evidence in Spain showed that stillbirth rates declined from 1996 to 2006, whereas the risk increased among women of advanced maternal age.<sup>25</sup> During the current period of analysis (2007–15) stillbirth rates remain constant (no secular trend), but the risk among highly educated women of advanced maternal age continued to increase. Furthermore, stillbirth rates among women from countries of low HDI remain constantly high showing no improvement during the last 10 years in Spain. We argue that public health policies banning preventive care for undocumented pregnant immigrant women during the 2009 economic crisis might explain the consistently higher rates of stillbirth among women of low SES in Spain. It is not only in Spain where undocumented immigrants have difficulties accessing healthcare. For instance, in Denmark, undocumented migrants have limited medical rights in addition to other barriers accessing healthcare such as arbitrariness in healthcare professionals' attitudes, fear of being reported to the police, poor language skills, lack of social network and lack of knowledge about the healthcare system.<sup>26</sup> In general, undocumented migrants seem to use different types of healthcare services less often than legal residents in most of the European countries. Even when care is utilized, it often seems to be inadequate or insufficient.<sup>27</sup>

Our article extends the important research, which seeks to measure the association between the HDI from women's countries of origin as a proxy for SES and identifying the joint effect between maternal educational attainment and HDI on stillbirth risk. However, to improve our understanding of the underlying causes of higher vulnerability to stillbirth among women with low educational attainment from countries with low HDI in Spain, more information related to immigrant background and culture, such as communication problems due to language skills, accessibility to the health-care system, acceptance of preventive interventions, use of prenatal services and quality of health care received, is needed.<sup>28</sup> Understanding the mechanisms associated with these risk factors could have an important impact on the reduction of the stillbirth rates.

Stillbirth is particularly subjected to under-reporting at low gestational ages (20–27 weeks; early foetal death).<sup>29</sup> However, the exclusion of infants <28 gestational weeks allowed us to minimize the bias due to under-reporting and misclassification of foetal deaths.<sup>30</sup>

Finally, we did not restrict our analysis to just singleton births, and we were not able to identify women delivering more than once during the period in study. It might have induced some confounding. However, sensitivity analysis restricted to singleton births (removing 168 900 multiple births) showed the same results as the unrestricted analysis.

Measuring social inequalities in stillbirth and monitoring trends over time enables target-setting for health policies. Unfortunately, our findings show no improvement in stillbirth rates among women with lower SES and an increasing trend among highly educated

women with advanced maternal age in recent years in Spain. Therefore, the findings highlight the need for developing preventive programmes to reduce stillbirth rates among women with lower SES as well as the necessity of further study to understand the growing trend of age-related stillbirths among highly educated women in Spain.

## Supplementary data

Supplementary data are available at *EURPUB* online.

## Funding

M.A.L.F. is supported by a Miguel Servet I Investigator Award (grant CP17/00206) from the Carlos III Institute of Health (ISCIII), Madrid, Spain. The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the funding agencies. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

*Conflicts of interest:* None declared.

## Key points

- Despite the fact that in Europe stillbirth rates are lower than in developing countries, stillbirth rates remain relatively stable over the last few decades, which is the cause of public health concern.
- Furthermore, evidence shows a higher risk of stillbirth among women of lower socioeconomic status (SES) in Europe.
- The study shows no improvement in stillbirth rates, over the last decade, among women of lower SES in Spain.
- The findings highlight the need for developing preventive programmes to reduce stillbirth rates among women of lower SES.

## References

- 1 World Health Organization. *Every Newborn: an action plan to end preventable deaths*. Geneva: WHO Press, 2014.
- 2 Garcia-Tizon Larroca S, Arevalo-Serrano J, Duran Vila A, et al. Human Development Index (HDI) of the maternal country of origin as a predictor of perinatal outcomes - a longitudinal study conducted in Spain. *BMC Pregnancy Childbirth* 2017;17:314.
- 3 Zeitlin J, Mortensen L, Prunet C, et al. Socioeconomic inequalities in stillbirth rates in Europe: measuring the gap using routine data from the Euro-Peristat Project. *BMC Pregnancy Childbirth* 2016;16:15.
- 4 Farrant BM, Shepherd CC. Maternal ethnicity, stillbirth and neonatal death risk in Western Australia 1998–2010. *Aust N Z J Obstet Gynaecol* 2016;56:532–6.
- 5 Flenady V, Koopmans L, Middleton P, et al. Major risk factors for stillbirth in high-income countries: a systematic review and meta-analysis. *Lancet* 2011;377:1331–40.
- 6 Ravelli AC, Tromp M, Eskes M, et al. Ethnic differences in stillbirth and early neonatal mortality in the Netherlands. *J Epidemiol Community Health* 2011;65:696–701.
- 7 Villadsen SF, Mortensen LH, Andersen AM. Ethnic disparity in stillbirth and infant mortality in Denmark 1981–2003. *J Epidemiol Community Health* 2009;63:106–12.
- 8 Rom AL, Mortensen LH, Chattingius S, et al. A comparative study of educational inequality in the risk of stillbirth in Denmark, Finland, Norway and Sweden 1981–2000. *J Epidemiol Community Health* 2012;66:240–6.
- 9 Essén B, Hanson BS, ÖSTERGREN PO, et al. Increased perinatal mortality among sub-Saharan immigrants in a city-population in Sweden. *Acta Obstet Gynecol Scand* 2000;79:737–43.
- 10 Cacciani L, Asole S, Polo A, et al. Perinatal outcomes among immigrant mothers over two periods in a region of central Italy. *BMC Public Health* 2011;11:294.

- 11 Zeitlin J, Mohangoo A, Cuttini M, et al. The European Perinatal Health Report: comparing the health and care of pregnant women and newborn babies in Europe. *J Epidemiol Community Health* 2009;63:681–2.
- 12 Richardus JH, Graafmans WC, Verloove-Vanhorick SP, et al. Differences in perinatal mortality and suboptimal care between 10 European regions: results of an international audit. *BJOG* 2003;110:97–105.
- 13 WHO. Expert Committee on Health Statistics: report on the second session, Geneva, 18–21 April 1950, including reports on the first sessions of the Subcommittees on Definition of Stillbirth, Registration of Cases of Cancer, Hospital Statistics, 1950.
- 14 United Nations Educational, Scientific and Cultural Organization. *International Standard Classification of Education-ISCED 1997: November 1997*. Paris: UNESCO, 1997.
- 15 United Nations Development Programme. *Human Development Report*. Oxford: Oxford University Press, 1990.
- 16 Vandenbroucke JP, Von Elm E, Altman DG, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *PLoS Med* 2007;4:e297.
- 17 White H. A heteroskedasticity-consistent covariance matrix estimator and a direct test for heteroskedasticity. *Econometrica* 1980;48:817–38.
- 18 Luque-Fernandez MA, Franco M, Gelaye B, et al. Unemployment and stillbirth risk among foreign-born and Spanish pregnant women in Spain, 2007–2010: a multilevel analysis study. *Eur J Epidemiol* 2013;28:991–9.
- 19 Tararbit K, Bui TT, Lelong N, et al. Clinical and socioeconomic predictors of pregnancy termination for fetuses with congenital heart defects: a population-based evaluation. *Prenat Diagn* 2013;33:179–86.
- 20 Gagnon AJ, Zimbeck M, Zeitlin J, et al. Migration to western industrialised countries and perinatal health: a systematic review. *Soc Sci Med* 2009;69:934–46.
- 21 Racape J, De Spiegelaere M, Alexander S, et al. High perinatal mortality rate among immigrants in Brussels. *Eur J Public Health* 2010;20:536–42.
- 22 Gardosi J, Madurasinghe V, Williams M, et al. Maternal and fetal risk factors for stillbirth: population based study. *BMJ* 2013;346:f108.
- 23 Eurostat. First and second-generation immigrants - statistics on main characteristics. 2016.
- 24 Perez G, Gotsens M, Palencia L, et al. Study protocol on the effect of the economic crisis on mortality and reproductive health and health inequalities in Spain. *Gac Sanit* 2016;30:472–6.
- 25 Luque-Fernandez MA, Lone NI, Gutierrez-Garitano I, Bueno-Cavanillas A. Stillbirth risk by maternal socio-economic status and country of origin: a population-based observational study in Spain, 2007–08. *Eur J Public Health* 2012;22:524–9.
- 26 Biswas D, Kristiansen M, Krasnik A, Norredam M. Access to healthcare and alternative health-seeking strategies among undocumented migrants in Denmark. *BMC Public Health* 2011;11:560.
- 27 Winters M, Rechel B, de Jong L, Pavlova M. A systematic review on the use of healthcare services by undocumented migrants in Europe. *BMC Health Serv Res* 2018;18:30.
- 28 Wolff H, Epiney M, Lourenco AP, et al. Undocumented migrants lack access to pregnancy care and prevention. *BMC Public Health* 2008;8:93.
- 29 Graafmans WC, Richardus JH, Macfarlane A, et al. Comparability of published perinatal mortality rates in Western Europe: the quantitative impact of differences in gestational age and birthweight criteria. *BJOG* 2001;108:1237–45.
- 30 Adams MM. *Perinatal Epidemiology for Public Health Practice*. New York: Springer, 2009.


.....  
*The European Journal of Public Health*, Vol. 29, No. 6, 1048–1055

© The Author(s) 2019. Published by Oxford University Press on behalf of the European Public Health Association.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

doi:10.1093/eurpub/ckz120 Advance Access published on 4 July 2019  
 .....

## Perinatal health of refugee and asylum-seeking women in Sweden 2014–17: a register-based cohort study

Can Liu <sup>1,2</sup>, Mia Ahlberg<sup>1,3</sup>, Anders Hjern<sup>1,2</sup>, Olof Stephansson<sup>1,3</sup>

1 Clinical Epidemiology Division, Department of Medicine, Karolinska Institutet, Stockholm, Sweden

2 Department of Public Health Sciences, Centre for Health Equity Studies, Karolinska Institutet/Stockholm University, Stockholm, Sweden

3 Department of Women's and Children's Health, Division of Obstetrics and Gynecology, Karolinska University Hospital and Institutet, Stockholm, Sweden

**Correspondence:** Can Liu, Department of Medicine Solna, Karolinska Institutet, Clinical Epidemiology Division T2, Karolinska University Hospital, SE-171 76 Stockholm, Sweden, Tel: +46 76 1684 323, e-mail: can.liu@ki.se

**Background:** An increasing number of migrants have fled armed conflict, persecution and deteriorating living conditions, many of whom have also endured risky migration journeys to reach Europe. Despite this, little is known about the perinatal health of migrant women who are particularly vulnerable, such as refugees, asylum-seekers, and undocumented migrants, and their access to perinatal care in the host country. **Methods:** Using the Swedish Pregnancy Register, we analyzed indicators of perinatal health and health care usage in 31 897 migrant women from the top five refugee countries of origin between 2014 and 2017. We also compared them to native-born Swedish women. **Results:** Compared to Swedish-born women, migrant women from Syria, Iraq, Somali, Eritrea and Afghanistan had higher risks of poor self-rated health, gestational diabetes, stillbirth and infants with low birthweight. Within the migrant population, asylum-seekers and undocumented migrants had a higher risk of poor maternal self-rated health than refugee women with residency, with an adjusted risk ratio (RR) of 1.84 and 95% confidence interval (95% CI) of 1.72–1.97. They also had a higher risk of preterm birth (RR 1.47, 95% CI 1.21–1.79), inadequate antenatal care (RR 2.56, 95% CI 2.27–2.89) and missed postpartum care visits (RR 1.15, 95% CI 1.10–1.22). **Conclusion:** Refugee, asylum-seeking and undocumented migrant women were vulnerable during pregnancy and childbirth. Living without residence permits negatively affected self-rated health, pregnancy and birth outcomes in asylum-seekers and undocumented migrants. Pregnant migrant women's special needs should be addressed by those involved in the asylum reception process and by health care providers.  
 .....