

SCIENTIFIC REPORTS



OPEN

Long-term growth of temperate broadleaved forests no longer benefits soil C accumulation

Yu-he Ji^{1,*}, Ke Guo^{2,*}, Shi-bo Fang^{1,*}, Xiao-niu Xu³, Zhi-gao Wang⁴ & Shu-dong Wang⁵

Received: 23 May 2016
Accepted: 09 January 2017
Published: 08 February 2017

It is widely recognized that the long-term growth of forests benefits biomass carbon (C) sequestration, but it is not known whether the long-term growth of forests would also benefit soil C sequestration. We selected 79 representative soil profiles and investigated the influence of the forest stand age on the soil C dynamics of three soil layers (0–10, 10–20 and 20–30 cm) in temperate broadleaved forests in East China. The results suggest that the soil C density in temperate broadleaved forests significantly changes with the stand age, following a convex parabolic curve. At an early stand age, the soil C density usually increases, reaching its peak value at a pre-mature stand age (approximately 50 years old). At later stand ages, the soil C density usually decreases. Therefore, our results reveal a turning point in the soil C density at a pre-mature stand age. The long-term growth of temperate broadleaved forests after pre-mature stand age no longer benefits soil C accumulation, probably promotes topsoil C loss. In addition, we found that the soil C density in the upper soil layer usually changes with the forest stand development more significantly than that in deeper soil layers.

Soil carbon (C) in forests has attracted much attention in recent years because its stability contributes to the mitigation of climate change^{1,2}. It was discovered that the soil C pool in temperate forests appears to be stable under disturbances, such as logging, wind storms, and invasive species^{3,4}. It is widely recognized that the soil C pool in forests varies dynamically with the stand age^{5,6}, but there are disagreements on how this occurs, leaving forest managers with uncertainty on how to best update forests for optimal C sequestration in the soil.

Rothstein *et al.* (2004) observed a weak decline in the surface soil C content with the stand age in Michigan jack pine forests⁷. On the contrary, Fonseca *et al.* (2011) discovered that the soil C increased by 1.1 Mg ha⁻¹ yr⁻¹ (1 Mg = 10⁶g) over the stand age range of 4–20 years in secondary tropical forests in Costa Rica⁸. Chen *et al.* (2013) discovered that the soil C pool in a Chinese fir plantation declined at young stand ages and then re-accumulated C at the stand ages of 16–21 years⁹. Shi & Cui (2010) argued, by summarizing 70 publications, that the highest soil C accumulation rate occurred at stand ages of 10–20 years old¹⁰. By summarizing more than 100 publications, Yang *et al.* (2011) argued that the soil C pool did not undergo significant changes during forest stand development in most studies¹¹. Therefore, it remains uncertain how the soil C changes in the long-term growth process of forests.

In China, secondary forests have expanded due to reforestation over the past half century. Seven national-scale forest investigations have been performed since the 1950s, but focused only on the timber volume and forest C biomass, without considering the soil C¹². National-scale soil investigations have also been performed, but unfortunately, they focused on the soil C in different soil types rather than for vegetation types^{13–14}. Some studies estimated the soil C pool in Chinese forests based on process-based BIOME models¹⁵, but could not resolve the complicated relationship between the soil C and forest stand age. Thus, it is necessary to elucidate how the soil C changes with the forest stand age to provide scientific evidence for forest management.

¹State Key Laboratory of Severe Weather (LASW), Chinese Academy of Meteorological Science, Beijing 100081, China. ²State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China. ³School of Forestry & Landscape Architecture, Anhui Agricultural University, Hefei 230036, China. ⁴Department of Life Science, Anqing Normal University, Anqing 246133, China. ⁵State Key Laboratory of Remote Sensing Science, Institute of Remote Sensing and Digital Earth, Chinese Academy of Science, Beijing 100101, China. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to Y.-h.J. (email: yuheji@163.com) or K.G. (email: guoke@ibcas.ac.cn) or S.-B.F. (email: fangshibo@camsma.cn)




Figure 1. Soil C density change with actual stand age in soil layers (0–30 cm) in temperate broadleaved forests in Anhui Province, East China.




Figure 2. Soil C density change with actual stand age in three soil layers (0–10, 10–20, 20–30 cm) in temperate broadleaved forests in Anhui Province, East China.

In this paper, we investigated 79 representative soil profiles in temperate broadleaved forests in eastern China. The objectives were to uncover the relationship between the soil C sequestration and the stand age of temperate broadleaved forests to improve forest management.

Results

Change trend of soil C density with stand age. Regardless of the broadleaved tree species, we use the actual forest stand age as the independent variable to examine how the soil C density changes in three soil layers (0–10, 10–20, 20–30 cm) in temperate broadleaved forests. When the three soil layers were taken as a unit, the results show that the soil C density is significantly correlated with the forest stand age. The soil C density changes with the stand age following a convex parabolic curve ($R^2 = 0.3273$), not a straight line. The soil C density increases at a young stand age, reaches its maximum carbon storage at an average age of approximately 50 years, and then gradually declines with the increasing stand age (Fig. 1). Therefore, the results indicate that there exists a turning point of soil C density in temperate broadleaved forests during stand age development. The soil acts as a C sink following forest establishment, but switches to a C source at approximately 50 years old, implying that the long-term growth of temperate broadleaved forests after 50 years no longer benefits soil C accumulation, but rather contributes to C loss from the soil.

When comparing the three soil layers, a significant change in the soil C density with the forest stand age following a parabolic curve was observed ($R^2 = 0.4309$) in the upper soil layer (0–10 cm). In the soil layer of 10–20 cm, the soil C density also varied in a parabolic curve with the forest stand age ($R^2 = 0.2346$), but the peak of the parabolic curve became lower. In the deeper soil layer of 20–30 cm, the peak of the parabolic curve disappeared ($R^2 = 0.0193$), as the soil C density changed only slightly compared to in the upper soil layers (Fig. 2). As a result, the soil C in the upper layers is more sensitive to the forest stand age than that in the lower soil layers.

Average change rate of soil C density with stand age class. To quantify the soil C dynamics with the stand age class, we divided the entire growth sequence of temperate broadleaved forests into five stand age classes (young, middle, pre-mature, mature and over-mature) (Table 1). When the three soil layers (0–10, 10–20, 20–30 cm) were taken into account as a whole, the results suggest that the soil C density reaches its peak value (approximately 85.6 Mg C/ha) at the pre-mature stand age. On average, the soil C density increased at a rate of

| Period of stand age | Stand age classes | Average stand age | Number of soil profiles | Soil C density of 0–10 cm (Mg C/ha) | Soil C density of 10–20 cm (Mg C/ha) | Soil C density of 20–30 cm (Mg C/ha) |
|---------------------|-------------------|-------------------|-------------------------|-------------------------------------|--------------------------------------|--------------------------------------|
| Young | (0–20] | 12 | 18 | 21.0 ± 12.5 | 16.3 ± 10.6 | 15.8 ± 11.1 |
| Middle | (20–40] | 33 | 26 | 42.0 ± 11.5 | 24.3 ± 8.2 | 12.7 ± 4.4 |
| Pre-mature | (40–60] | 52 | 20 | 46.7 ± 13.0 | 27.0 ± 11.1 | 13.3 ± 3.7 |
| Mature | (60–80] | 74 | 11 | 37.5 ± 7.7 | 22.9 ± 7.0 | 14.6 ± 7.2 |
| Over mature | (80–100] | 91 | 3 | 30.8 ± 7.4 | 16.9 ± 0.9 | 8.4 ± 1.1 |

Table 1. Soil C density at a depth of 0–30 cm in temperate broadleaved forests. Note: stand ages refer to the standards of the “Forest Resource Statistics of China” (Zhang *et al.*¹³).




Figure 3. Soil C dynamics with stand age class in three soil layers (0–10, 10–20, 20–30 cm) in temperate broadleaved forests.

0.813 Mg C/ha per year prior to the pre-mature stand age. Subsequently, it declined at a rate of 0.74 Mg C/ha per year after the pre-mature stand age and to 56.0 Mg C/ha at the over-mature stand age (average 91 years old). Therefore, the quantitative results indicate that the pre-mature stand age (average 52 years old) is a turning point in the soil C dynamics. The soil C accumulated at a rate of 0.813 Mg C/ha per year before the turning point and then exhibited a loss of 0.74 Mg C/ha per year after the turning point (Fig. 3).

Comparing the three soil layers, the upper soil layer (0–10 cm) showed the most significant change in soil C density, with an increase of 0.643 Mg C/ha per year from young to pre-mature and a decrease of 0.398 Mg C/ha per year from pre-mature to over-mature. The middle soil layer (10–20 cm) showed an increase of 0.268 Mg C/ha per year from young to pre-mature and a decrease of 0.253 Mg C/ha per year from pre-mature to over-mature. The deepest soil layer (20–30 cm) showed a slightly fluctuating soil C density without any significant change over the entire growth sequence.

Discussion

Our results demonstrate that the soil C density in temperate broadleaved forest changes with the stand age following a convex parabolic curve, and there exists a turning point with a single peak in the soil C accumulation at approximately 50 years old (Figs 1, 2 and 3). However, the results only show a general (significant single-peak curve) change trend of soil C density, but omits minor changes along the successional process since the data is not enough to identify the minor changes. The soil C density probably changes following a multi-peak curve with no more than one turning point during the entire forest stand age, as some studies reported that the soil C initially decreased or increased slowly for the first decade after afforestation and then began to accumulate quickly with the stand age. The multi-peak curve of soil C density usually exist under the precondition of afforestation. For example, Paul *et al.* (2002) synthesised available world-wide information on changes in soil C after afforestation, and argued that soil C in surface soil (<10 or <30 cm depth) initially declines during the first 5 years after establishing a plantation but recovers by the age of 30 years¹⁶. The initial decline of soil C in Paul *et al.* (2002) came from the average data in the 43 published or unpublished studies, so the decline is not significant since the data are highly variable, with soil C either increasing or decreasing in young (<10 year) forest stands. Li *et al.* (2011) observed the total mineral soil C initially appeared to decline at the early stand age, but recovered by the stand age of 35 years for coniferous plantation forest with Korean Pine (*Pinus koraiensis*). It is a pity that the natural change of soil C is not credible after 35-year-old stand because the soil C suffered from disturbance greatly from human, such as thinning treatment to the 35- and 51-year-old stands¹⁷. Hiltbrunner *et al.* (2013) examined the effects of afforestation with Norway spruce (*Picea abies* L.) in a grazed subalpine pasture in Switzerland on soil organic carbon (SOC), and discovered that soil C stock decreased after tree establishment, reaching a minimum 40–45 years after afforestation, and increased thereafter¹⁸. Barcena *et al.* (2014) analyzed the changes in SOC stocks




Figure 4. Multi-peak curve of topsoil C density change with forest stand age according to our study and other studies.

at the 0–30 cm soil layer following afforestation in Northern Europe by a meta-analysis, revealed that SOC loss generally for barren, cropland, heathland and grass-land at the initial phase following afforestation (0–30 years). The detectable gains in SOC stocks appear in later stages (>30 years), especially for afforestation of croplands¹⁹. Yu *et al.*²⁰ investigated the soil C in four Chinese fir (*Cunninghamia lanceolata* Hook) plantations (Chinese fir was planted in clear-cut sites in natural broad-leaved forest) in Jiangxi Province, south China, discovered that soil C density at the depth of 0–20 cm declined before 16 years, but increased after 16 years, since soil C density declined from 35.98 Mg·ha⁻¹ in the 7-year plantation to 30.12 Mg·ha⁻¹ in the 16-year plantation, and then increased after 16 years old²⁰. Therefore, we can speculate that soil C density probably changes following a multi-peak curve, and another turning point of the soil C density may exist in the early decades of afforestation, besides the large turning point at approximately 50 years old (Fig. 4).

In the later stage of forest development, many reports indicate that old forests are expected to maintain their biomass accumulation for a long time through the development of a multilayer canopy structure²¹, but this does not mean that the soil C can continue to increase as long as the biomass accumulates. Our results show that old-growth forest could not sustain the soil C increase due to the decreasing soil C density after the pre-mature stand age (average stand age 50 years old). However, there seems to be some controversy on this point. It is conventionally accepted that the soil C levels in old-growth forests are in a steady state^{21–22}. However, Zhou *et al.* (2006) reported that soils in the top 20-cm soil layer in preserved old-growth forests (age > 400 years) in southern China accumulated C significantly at an unexpectedly high rate from 1979 to 2003²³. Li & Liu (2014) argued that an old forest (38–56 y) was able to continuously accumulate C in the soil in China's Loess Plateau, even when the biomass significantly decreased²⁴.

These different opinions could be partly attributed to the different definitions of “old forest” with tree species and environment spatial variability, as there is currently no recognized definition. For example, a stand age of 38–56 years old in the study of Li & Liu (2014) was regarded as old forest²⁴, equivalent to the pre-mature stands age (40–60 years old) in our study. It is likely that tree species and environment spatial variability are the leading causes of the different opinions, as different environments and tree species can affect the carbon accumulation-and-release processes^{11,25–27}. The synergistic effects of many factors should be further explored to uncover the complex mechanism of soil C dynamics^{28–31}.

Conclusions

Our results show that soil C in temperate broadleaved forests significantly changes with stand age. Generally, it exhibits a change trend in the shape of a convex parabolic curve with stand age, regardless of the tree species. At the early stage of forest development, the soil C density usually increases, and it reaches its peak value at the pre-mature stand age (approximately 50 years old). At later stages of forest development, the soil C density usually decreases. This phenomenon provides strong evidence that there is a turning point of the soil C density in temperate broadleaved forests at the pre-mature stand age, when the soil switches from being a net C sink to a net C source. Therefore, we drew the conclusion that the long-term growth of temperate broadleaved forests after pre-mature stand age no longer benefits soil C accumulation. Our study also confirmed that the soil C in the upper layers is more sensitive to forest stand age than that of the lower soil layers, as the soil C density in the upper soil layers usually changes significantly with the forest stand development.

Materials and Methods

Study area. The study area covers approximately 139,000 km² in the Anhui Province (114°51′–119°36′E, 29°26′–34°37′N) of East China (Fig. 5). The Asian monsoon circulation creates a temperate continental monsoon climate with an annual average temperature of 14–17 °C and an annual precipitation of 800–1800 mm. The soil type is yellow brown soil³². Three mountain ranges (Dabie, Jiuhua and Huang Mountains) lie in the southwestern




Figure 5. Location of study area and sampling plots of 79 representative soil profiles in temperate broadleaved forests in Anhui Province, East China (The map was generated using ArcGIS for Desktop 10.2, <http://www.esri.com/software/arcgis>).

and southern regions (Fig. 5) and are covered with temperate and subtropical forests. The dominant forest types are temperate deciduous broadleaved forests, coniferous forests, and mixed forests.

In recent decades, large-scale deforestation has been curbed, and reforestation projects have been carried out, providing an opportunity to resume normal forest development³³. Forest managers also allow people to update some forests to obtain timber for money. Therefore, the study area contains various forests with stand ages ranging from 0 to 100 years. However, it is unclear what updating schedule for the forest is the most favourable for soil C sequestration.

Sampling sites (plots). To avoid the effect of the spatial heterogeneity of sampling sites on SOC, we did our utmost to select coincident sampling sites (plots) in vegetation composition, soil type, and the same development process. All the sampling sites must have typical temperate broadleaved forest which is determined by climatic zones, though there are other forests, such as coniferous forests, conifer and broadleaf mixed forests. To avoid of human disturbance, all the sampling sites were selected in protecting natural forests to ensure a natural growth process. Young broadleaved forests being selected should have similar land use process because SOC in young forests suffer more effect from previous land use. We didn't consider the young forests which land use type had been changed greatly by human. Thus, almost all forest vegetation in sampling sites belongs to secondary successional vegetation under the protection of human beings.

We selected the typical forests for every stand ages (young, middle, pre-mature, mature, over-mature) according to the natural succession of temperate broadleaved forest, so there is a slightly inconsistent in vegetation (species) composition for different stand age due to the natural succession. The vegetation composition for different stand ages is listed as follows (Table 2).

The soil in study area is classified as yellow brown soil zone according to “Map of Soil Regionalization of China”³⁴. The sampling sites in our study ensured a typical yellow brown soil, and other soil types were avoided.

Generally, the sampling sites are coincident approximately in vegetation composition, soil type and development process, in spite of existing spatial heterogeneity more or less.

Soil sampling. To examine the dynamics of the soil C with stand age, 79 soil profiles of sampling sites were investigated in representative temperate broadleaved forests in September of both 2011 and 2012 (Fig. 5). The actual stand age of each forest type was recorded by visiting farmers and forest management staff. The investigation focused on soil carbon density in the 0–30 cm soil layer, since soil carbon in the layer accounts for the majority of the soil profile 0–100 cm, and is sensitive to forest stand ages more than that in deeper soil layers. Vertical soil profiles were dug in the sampling plots, and soil samples were collected from three soil layers (0–10, 10–20, 20–30 cm) using a ring knife (volume of 100 cm³) for measuring the soil bulk density and a spade for measuring the soil C content. Soil samples from the surface soil layer (0–10 cm) included the forest floor (O horizons, i.e., the organic horizon), but surface litter was not included in the calculation of the soil C. In addition, the relevant environmental information for each sampling plot was recorded, such as the geographical location with latitude and longitude, forest type and stand age.

Data analysis. The soil samples collected with the ring knife were used to measure the soil bulk density by the drying method in the laboratory. The soil samples collected with the spade were air-dried, ground using a mortar, and passed through a 2-mm sieve to remove all roots and stones. Finally, chemical analyses were performed to measure the soil C concentration by dry combustion using an elemental analyser (vario MACRO cube, Elementar, Germany).

| | Young forest | Middle age forest | Pre-mature forest | Mature and over mature forest |
|-------------|--|---|--|---|
| Tree layer | <i>Quercus glandulifera</i> var. <i>brevipetiolata</i> , <i>Pistacia chinensis</i> , <i>Broussonetia papyrifera</i> , <i>Sorbushemsleyi</i> | <i>Quercus glandulifera</i> var. <i>brevipetiolata</i> , <i>Cyclobalanopsis glauca</i> , <i>Quercus acutissima</i> , <i>Castanopsis eyrei</i> , <i>Castanopsis sclerophylla</i> , <i>Fraxinus sinularis</i> | <i>Quercus glandulifera</i> var. <i>brevipetiolata</i> , <i>Cyclobalanopsis glauca</i> , <i>Quercus acutissima</i> , <i>Castanopsis eyrei</i> , <i>Castanopsis sclerophylla</i> , <i>Platycarya strobilacea</i> | <i>Quercus glandulifera</i> var. <i>brevipetiolata</i> , <i>Cyclobalanopsis glauca</i> , <i>Castanopsis eyrei</i> , <i>Quercus acutissima</i> |
| Shrub layer | <i>Rhododendron simsii</i> , <i>Lespedeza vitorum</i> , <i>Lindera reflexa</i> , <i>Zanthoxylum armatum</i> , <i>Rhizoma Smilacis</i> | <i>Rhododendron simsii</i> , <i>Lespedeza bicolor</i> , <i>Lorpetalum chinense</i> , <i>Lindera glauca</i> , <i>Glochidion puberum</i> , <i>Camellia fraternal</i> , <i>Rhamnus globosa</i> | <i>Rhododendron simsii</i> , <i>Lespedeza</i> <i>Formosa</i> , <i>Lorpetalum chinense</i> , <i>Camellia fraternal</i> | <i>Pleioblastus amarus</i> , <i>Camellia</i> <i>fraternal</i> , <i>Lindera glauca</i> |
| Herb layer | <i>Dryopteris chinensis</i> , <i>Carex tristachya</i> , <i>Oplismenus undulatifolius</i> , <i>Arthraxon hispidus</i> , <i>Commelinabengalensis</i> , <i>Carpesium abrotanoides</i> , <i>Artemisia</i> <i>lavandulaefolia</i> , <i>Leonurus japonicus</i> , <i>Viola concordifolia</i> | <i>Dryopteris chinensis</i> , <i>Woodwardia japonica</i> , <i>Carex tristachya</i> , <i>Oplismenus undulatifolius</i> , <i>Rhizoma imperata</i> , <i>Aster ageratoides</i> , <i>Antenoron filiforme</i> , <i>Vicia amoenaefisch</i> | <i>Dryopteris chinensis</i> , <i>Woodwardia japonica</i> , <i>Carex tristachya</i> , <i>Oplismenus undulatifolius</i> , <i>Lindera aggregate</i> , <i>Phaenopermaglobosa</i> | <i>Dryopteris chinensis</i> , <i>Woodwardia</i> <i>japonica</i> , <i>Carex breviculmis</i> , <i>Oplismenus undulatifolius</i> , <i>Lophatherum gracile</i> , <i>Lindera aggregata</i> |

Table 2. The vegetation composition in sampling sites for different stand ages of temperate broadleaved forests in Anhui Province.

For these soil profiles, we calculated the soil C density (D) of the three soil layers of 0–10, 10–20 and 20–30 cm using Equation (1),

$$D = \sum_{i=1}^n (D_i \times H_i \times C_i), \quad (1)$$

where D is the soil C density (10^5 g C/ha.), D_i is the bulk density (g/cm^3), H_i is the soil depth (cm), C_i is the soil C concentration (%), and i represents the three soil layers.

References

- Liu, S. R., Wang, H. & Luan, J. W. A review of research progress and future prospective of forest soil carbon stock and soil carbon process in China. *Acta Ecol. Sin.* **31**, 5437–5448 (2011).
- Cook, R. L., Binkley, D., Mendes, J. C. T. & Stape, J. L. Soil carbon stocks and forest biomass following conversion of pasture to broadleaf and conifer plantations in southeastern Brazil. *For Ecol. and Manag.* **324**, 37–45 (2014).
- Davis, M. R., Allen, R. B. & Clinton, P. W. Carbon storage along a stand development sequence in a New Zealand *Nothofagus* forest. *For Ecol. and Manag.* **177**, 313–321 (2003).
- Tyrrell, M. L., Ross, J. & Kelty, M. Carbon Dynamics in the Temperate Forest. In M. S. Ashton *et al.* (eds.), *Managing Forest Carbon in a Changing Climate*, 77–101 (2012).
- Wang, S. Q. & Zhou, C. H. Estimating soil carbon reservoir of terrestrial ecosystem in China. *Geogr. Res.* **18**, 349–357 (1999).
- Wu, H. B., Guo, Z. T., Gao, Q. & Peng, C. H. Distribution of soil inorganic carbon storage and its changes due to agricultural land use activity in China. *Agri. Ecosyst. Environ.* **129**, 413–421 (2009).
- Rothstein, D. E., Yermakov, Z. & Buell, A. L. Loss and recovery of ecosystem carbon pools following stand-replacing wildfire in Michigan jack pine forests. *Can. J. of For Res.* **34**, 1908–1918 (2004).
- Fonseca, W., Benayas, J. M. R. & Alice, F. E. Carbon accumulation in the biomass and soil of different aged secondary forests in the humid tropics Costa Rica. *For Ecol. and Manag.* **262**, 1400–1408 (2011).
- Chen, G. S. *et al.* Carbon storage in a chronosequence of Chinese fir plantations in southern China. *For Ecol. and Manag.* **300**, 68–76 (2013).
- Shi, J. & Cui, L. L. Soil carbon change and its affecting factors following afforestation in China. *Landscape Urban Plan* **98**, 75–85 (2010).
- Yang, Y. H., Lu, Y. Q. & Finzi, A. C. Carbon and nitrogen dynamics during forest stand development: a global synthesis. *New Phytol.* **190**, 977–989 (2011).
- Zhang, C. H. *et al.* China's forest biomass carbon sink based on seven inventories from 1973 to 2008. *Clim. Chang.* **118**, 933–948 (2013).
- Yu, D. S. *et al.* Regional patterns of soil organic carbon stocks in China. *J. Environ. Manag.* **85**, 680–689 (2007).
- Yu, D. S. *et al.* National scale analysis of soil organic carbon storage in China based on Chinese soil taxonomy. *Pedosphere* **17**, 11–18 (2007).
- Ni, J. Net primary production, carbon storage and climate change in Chinese biomes. *Nordic J Bot* **20**(4), 415–426 (2000).
- Paul, K. I., Polglase, P. J., Nyakuengama, J. G. & Khanna, P. K. Change in soil carbon following afforestation. *For Ecol. and Manag.* **168**, 241–257 (2002).
- Li, X. D. *et al.* Biomass and carbon storage in an age-sequence of Korean Pine (*Pinus koraiensis*) plantation forests in central Korea. *J. Plant Biol.* **54**, 33–42 (2011).
- Hiltbrunner, D., Zimmermann, S. & Hagedorn, F. Afforestation with Norway spruce on a subalpine pasture alters carbon dynamics but only moderately affects soil carbon storage. *Biogeochemistry* **115**, 251–266 (2013).
- Barcena, T. G. *et al.* Soil carbon stock change following afforestation in Northern Europe: a meta-analysis. *Global Change Biology* **20**, 2393–2405 (2014).
- Yu, S. B., Wang, D., Dai, W. & Li, P. Soil carbon budget in different-aged Chinese fir plantations in south China. *J. For. Res.* **25**, 621–626 (2014).
- Luyssaert, S. *et al.* Old-growth forests as global carbon sinks. *Nature* **455**, 213–215 (2008).
- Seedre, M., Kopacek, J., Janda, P., Bace, R. & Svoboda, M. Carbon pools in a montane old-growth Norway spruce ecosystem in Bohemian Forest: Effects of stand age and elevation. *For Ecol. and Manag.* **346**, 106–113 (2015).
- Zhou G. Y. *et al.* Old-growth forests can accumulate carbon in soils. *Science* **314**, 1417 (2006).
- Li, T. J. & Liu, G. B. Age-related Changes of Carbon Accumulation and Allocation in Plants and Soil of Black Locust Forest on Loess Plateau in Ansai County, Shaanxi Province of China. *Chin. Geogr. Sci.* **24**, 414–422 (2014).
- Kawasaki, M., Ohte, N. & Katsuyama, M. Biogeochemical and hydrological controls on carbon export from a forested catchment in central Japan. *Ecol Res* **20**, 347–358 (2005).
- Wei, Y. W. *et al.* Variation in carbon storage and its distribution by stand age and forest type in boreal and temperate forests in northeastern China. *PLOSone* **8**, 1–9 (2013).
- Vanguelova, E. I. *et al.* Sources of errors and uncertainties in the assessment of forest soil carbon stocks at different scales—review and recommendations. *Environ Monit Assess.* **188**, 1–24 (2016).

28. Xu, W. H., Li, W., Jiang, P., Wang, H. & Bai, E. Distinct temperature sensitivity of soil carbon decomposition in forest organic layer and mineral soil. *Scientific Reports* **4**, 1–6 (2014).
29. Zhou, Z. Y., Xu, M. L., Kang, F. F. & Sun, O. J. X. Maximum temperature accounts for annual soil CO₂ efflux in temperate forests of Northern China. *Scientific Reports* **5**, 1–10 (2015).
30. Tian, Q. X., He, H. B., Cheng, W. X., Bai, Z., Wang, Y. & Zhang, X. D. Factors controlling soil organic carbon stability along a temperate forest altitudinal gradient. *Scientific Reports* **6**, 1–9 (2016).
31. You, Y. M., Wang, J., Sun, X. L., Tang, Z. X., Zhou, Z. Y. & Sun, O. J. X. Differential controls on soil carbon density and mineralization among contrasting forest types in a temperate forest ecosystem. *Scientific Reports* **6**, 1–15 (2016).
32. Xu, X. W., Pan, G. X., Cao, Z. H. & Wang, Y. L. A study on the influence of soil organic carbon density and its spatial distribution in Anhui Province of China. *Geogr. Res.* **26**, 1077–1086 (2007).
33. Fang, J. Y., Chen, A. P., Peng, C. H., Zhao, S. Q. & Ci, L. J. Changes in forest biomass carbon storage in China between 1949 and 1998. *Science* **292**, 2320–2322 (2001).
34. Xi, C. F. & Zhang J. M. The soil regions of China. *Acta Pedologica Sinica*, **19**, 97–110 (1982).

Acknowledgements

The special project “Forest Carbon Sequestration Rate and Potential Research in Anhui Province” (XDA0505020401), the National Natural Science Foundation of China (Grant No. 41571175, 41330531, 41671362) and the Agricultural Project for Coping with Climate Change (Grant No. 132020010406) financially supported this study.

Author Contributions

Professor Ke Guo led this research project. Dr. Yu-he Ji wrote the manuscript. Professor Shi-bo Fang contributed to the revision and polishing of the manuscript. Professor Xiao-niu Xu, Zhi-gao Wang, and Shu-dong Wang participated in field experiments and data processing.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Ji, Y.- *et al.* Long-term growth of temperate broadleaved forests no longer benefits soil C accumulation. *Sci. Rep.* **7**, 42328; doi: 10.1038/srep42328 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017