




Burden of fungal infections in Kenya

Stanley N. Ratemo ^a and David W Denning ^{b,c}

^aResearch Department, Kisii Teaching and Referral Hospital, Kisii, Kenya; ^bManchester Fungal Infection Group, Faculty of Biology, Medicine and Health, Manchester Academic Health Science Centre, University of Manchester, Manchester, UK; ^cGlobal Action for Fungal Infections (GAFFI), Geneva, Switzerland

ABSTRACT

The burden of fungal infections has been on the rise globally and remains a significant public health concern in Kenya. We estimated the incidence and prevalence of fungal infections using all mycology publications in Kenya up to January 2023, and from neighbouring countries where data lacked. We used deterministic modelling using populations at risk to calculate the disease burden. The total burden of serious fungal infections is estimated to affect 6,328,294 persons which translates to 11.57% of the Kenyan population. Those suffering from chronic infections such as chronic pulmonary aspergillosis are estimated to be 100,570 people (0.2% of the population) and probably nearly 200,000 with fungal asthma, all treatable with oral antifungal therapy. Serious acute fungal infections secondary to HIV (cryptococcal meningitis, disseminated histoplasmosis, pneumocystis pneumonia, and mucosal candidiasis) affect 196,543 adults and children (0.4% of the total population), while cancer-related invasive fungal infection cases probably exceed 2,299 and those in intensive care about 1,230 incident cases, including *Candida auris* bloodstream infection. The burden of fungal infections in Kenya is high; however, limited diagnostic test availability, low clinician awareness and inadequate laboratory capacity constrain the country's health system in responding to the syndemic of fungal disease in Kenya.

ARTICLE HISTORY

Received 1 February 2023
Accepted 13 April 2023

KEYWORDS

Mycetoma; cryptococcal meningitis; pneumocystis; candida; aspergillus; asthma

1. Introduction

The global burden of fungal infections is on the rise due to an increase in the number of people with associated risk factors such as HIV, cancer, and organ transplant as well as extremes of age (Maertens et al. 2001; Vallabhaneni et al. 2016). Fungal infections in Kenya are still a significant public health concern, thought to affect almost one in fifteen of the population at any point in time (Guto et al. 2016; Malwal et al. 2016). In order to prioritise healthcare, including training, provision of diagnostics and antifungal therapies, numerical estimates of disease burden are helpful.

The Kenyan population has also increased by an estimated 11 million people over the past decade (Commons 2022). This coupled with improved reporting rates for key health indicators, in tuberculosis and HIV, has led to a substantial increase in the population at risk of fungal infections (Samaranayake et al. 2002; Guarner 2017; Chepkondol et al. 2020; Ngugi et al. 2020). In 2016, it was estimated that about 7% of the Kenyan population was affected by a serious fungal infection, including

millions of children with tinea capitis (Guto et al. 2016). Since then, better fungal estimates and updated literature have been published.

The use of azole fungicides has in the recent past contributed to the thriving of triazole-resistant species of *Aspergillus fumigatus* species in some regions in Kenya (Kemoi et al. 2018). Hitherto, unreported cases of empyema necessitans due to *Aspergillus* are also being diagnosed in Kenya as well as reports of drug-resistant yeast infections (Brooks et al. 2019; Macharia et al. 2019). On the other hand, there is greater coverage of patients with HIV infection with antiretroviral therapy, although only modest progress on reducing tuberculosis mortality, which impacts the chronic pulmonary aspergillosis (CPA) caseload (Mohamed et al. 2022). Currently, about a dozen hospitals are offering kidney transplant services at a much-reduced cost compared to the cost of seeking such services abroad. The number of kidney transplants has therefore significantly increased with two of the dozen hospitals reporting about 400 transplants in the last decade (Ministry of Health Kenya

CONTACT Stanley N. Ratemo  sta.ratz@yahoo.com

© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

2021; News 2021). These shifts in the demographics of the population and those at risk of fungal infections, ongoing gaps in diagnostic provision (GAFFI.org 2022) and new data propelled us into a thorough re-estimation of the incidence and prevalence of serious fungal infections in Kenya.

2. Materials and methods

Basic demographic data on the Kenya population was derived from Indexmundi.com (Mundi 2022) (Table 1). HIV data was taken from UNAIDS, and the assumption made that for those adults not on antiretroviral therapy (ART) it takes 7 years for the CD4 counts and

immunity to decline into the high-risk group. In children, it was assumed that all children were at risk, unless on ART. Furthermore, it was assumed that ART was failing in 12.3% of those on ART (Tsikhutsu et al. 2022). Pulmonary Tuberculosis (PTB) incidence was abstracted from the WHO 2020 annual report for Kenya (StopTB 2021). Asthma prevalence in adults was derived from a 2016 household survey (Turpin et al. 2019). Chronic obstructive pulmonary disease (COPD) was estimated from data from Rwanda, Tanzania, and Uganda, and admissions to hospital from Algeria (Hammond et al. 2020). Lung cancer and acute leukaemia were taken from the International Agency for Research in Cancer

Table 1. Demographics of Kenya, pertinent to fungal infections.

Variable	Result	Source
Population characteristics	Total population: 54,685,051 Total Adult population: 33,357,881 Population under 15 years: 39% Annual population growth rate: 2.15% Population >60 years: 4% Percentage of females: 50% Number of children aged 5–14 years: 21,327,170 Proportion of population 15–55 years: 0.556 Number of women aged 15–55 years: 15,263,254 Total population of women aged 15–49: 13,355,347	Mundi 2022
Health status	Total number of functional health facilities: 9,064 Maternal mortality rate: 342 per 100,000 live births Infant mortality rate: 41.9 per 1000 live births Life expectancy at birth in 2021: 61.4 years Prevalence of underweight (Wt/Age) among children 6–59 months (2SD): 11%	Kenya Gazette Notice No 786; (Elephant 2022; UNICEF 2022a; UNDP 2022; UNICEF 2022b)
HIV	Total HIV/AIDS: 1,400,000 Children with HIV/AIDS: 86,000 Number of adults with HIV/AIDS: 1,314,000 of diagnosed cases not on ARVs: 14% Number of adults on ARVs: 1,160,570 Number of diagnosed adults not receiving ARVs: 153,430 Number of adult HIV patients with CD4 < 200/ μ l: 21,919 Percentage of adults with ARV failure: 11% Number of adult HIV patients at risk of OIs: 164,670 Number of children receiving ART: 69,390 Number of children at risk of OIs: 24,243 AIDS-related deaths: 19,000 Number of AIDS-related deaths in children: 3,100	UNAIDS 2022
TB	Total TB cases: 139,000 PTB annual incidence HIV+: 35,000 PTB annual incidence HIV-: 65,520 PTB annual incidence Total: 86,447 Pulmonary tuberculosis annual survivors: 73,480	StopTB 2021
Asthma, COPD, and Cystic Fibrosis	Asthma rate in adults: 10.00% Asthma number in adults: 3,335,788 COPD prevalence all GOLD stages: 7.8% COPD admissions to hospital per year: 388,665 (assumes 10.5% admissions per year)	Hammond et al. 2020
Malignancies	Number of lung cancer deaths per 100,000 population: 729 Leukaemia rate per 100,000: 2.5 Leukaemia patients per year: 1,367	IARC Kenya Factsheets (International Agency for Cancer on Research WHO 2022; Control UfIC 2014)
Transplant recipient	Allogenic haematologic stem cell transplant: 0 Renal transplants per year: 470 Liver transplants per year: 12	IRODaT 2023

(International Agency for Cancer on Research WHO 2022) and acute myeloid leukaemia from a general WHO figure (Control UfIC 2014). Annual renal and hepatic transplant numbers were from 2019 (IRODaT 2023).

Searches for the incidence and prevalence of different fungal diseases were done. The following search terms were used, "Aspergillosis Kenya" "Candidiasis Kenya" "Pneumocystis Kenya" "Cryptococcus Kenya" "Histoplasmosis Kenya" "Mycetoma Kenya" "Chromoblastomycosis Kenya" "Sporotrichosis Kenya" "Tinea capitis Kenya". The terms "incidence" and "prevalence" were used alongside the above terms. Where no data was found from Kenya, "Uganda", "Sudan", "Tanzania" and "Africa" were used. From these, and from existing papers in our libraries, we have estimated the proportion of people, with or without an underlying disease, who have a fungal disease. In many instances data was missing from Kenya, so nearby country estimates were used as proxy estimates, and if these were not available, then international data was used. All the assumptions and their sources for each fungal infection or disease are documented in Table 2.

Following this, calculations of annual incidence or prevalence were made using straightforward deterministic modeling using population(s) at risk (Table 2). These are estimated partly by underlying disease, where there well-established risk populations (i.e. advanced HIV disease) and by partly by individual fungal disease (i.e. fungal keratitis) on a population basis (Table 1). To use an example, 164,670 adults are at risk of a severe fungal infection (Table 1) and the point prevalence of cryptococcal antigenaemia is 6% to 7.5% (Table 2), so the annual incidence of cryptococcal disease (most with meningitis as there is very little screening) is 9,880 patients, using the lower percentage. The annual incidence of invasive aspergillosis (IA) comprised these underlying risk groups: acute myeloid leukaemia (AML) (13% risk) (Chen et al. 2020) plus an equal number for other moderate and high-risk haematological conditions (i.e. other leukaemia, multiple myeloma, and lymphoma), lung carcinoma (2.6%) (Yan et al. 2009), HIV deaths (4%) (Khoo SD 1994; Denning and Morgan 2022) and 1.3% of all those hospitalised (or severely ill) with a COPD exacerbation (Hammond et al. 2020) (Table 2). Chronic pulmonary aspergillosis (CPA) incidence and prevalence

Table 2. Key assumptions to allow modelling of fungal disease incidence and prevalence in Kenya.

Fungal Infection	Underlying condition	Assumptions	Reference(s)
Recurrent <i>Candida</i> vaginitis ($\geq 4 \times$ /year)	None	6% females aged 15–49	Denning et al. 2018
Candidaemia	Cancer, surgery critical care	5/100,000 (mean of 2–11/100,000), with 33% occurring in critical care	Bongomin et al. 2017
Intraabdominal candidiasis	Abdominal surgery, pancreatitis	50% of the incidence of candidaemia in critical care	Montravers et al. 2011
Oesophageal candidiasis	HIV/AIDS	20% of new HIV/AIDS adults and 0.5% of adults on ARVs	Smith and Orholm 1990; Buchacz et al. 2010
Cryptococcal meningitis	HIV/AIDS	7.5% positive CrAg positives, 6% in HIV/AIDS adults with CD4 < 200/uL	Kanji et al. 2011; Meyer et al. 2013
<i>Pneumocystis</i> pneumonia	HIV/AIDS	15% of newly diagnosed HIV adults, 16% of children with HIV/AIDS	Chakaya et al. 2003; Bii et al. 2006
Disseminated histoplasmosis	HIV/AIDS	4.7% of advanced HIV	Ocansey et al. 2022
Invasive aspergillosis (IA)	HIV/AIDS	13% of AML develop IA and an equal number of non-AML haematological conditions. 2.6% of lung cancer patients and 1.3% of COPD annual admissions, and 2% of renal transplant recipients.	Chen et al. 2014; Yan et al. 2009; Hammond et al. 2020
Chronic pulmonary aspergillosis (CPA) with a TB-like presentation	Respiratory	19% of HIV- and 10% of HIV+ clinically diagnosed PTB, 7% of HIV- and 3% of HIV+ proven PB and 1.5% annual rate after PTB cure.	Oladele et al. 2017; Page et al. 2019; Denning et al. 2023
Chronic pulmonary aspergillosis – all	Respiratory	Assumes 67% of cases occur during or after pulmonary TB	Smith and Denning 2011
Allergic bronchopulmonary aspergillosis (ABPA)	Respiratory	2.5% of adult asthmatics	Denning et al. 2013; Kwizera et al. 2020
Severe asthma with fungal sensitisation (SAFS)	Respiratory	3.3% adult asthmatics, all with severe asthma	Kwizera et al. 2021
Mucormycosis	Diabetes, leukaemia, trauma	0.2/100,000	Sharma and Goel 2022
Fungal keratitis	Eye injury	13.3/100,000	Brown et al. 2021
Tinea capitis	None	23% of children 0–14 years	Bongomin et al. 2021

were estimated from 2020 pulmonary tuberculosis (PTB) data. We also assume that PTB is the underlying cause of CPA in 80% of all CPA cases (Smith and Orholm 1990). For several fungal infections, notably the skin fungal Neglected Tropical Diseases (NTD), a realistic estimate or burden was not possible.

3. Results

3.1. Kenya demographic profile

The Kenyan population in 2021 was estimated to be approaching 55 million (Table 1). This represents an increase of about 11 million people over the course of a decade. The current annual population growth rate stood at 2.15% in 2021. The proportion of the population aged between 15 and 55 years was 39%. The Gross-Domestic-Product (GDP) per capita is \$4,200 (CIA 2022), making Kenya a Lower Middle-Income Country (LMIC) (World Bank 2023).

Kenya is also categorised as having medium development, ranking 152 out of 191 countries in the 2021–2022 United Nations Human Development Index (UNDP 2022). According to the World Bank, the poverty level in Kenya is currently at 34.4%; poverty indices in Kenya have reduced over time.

The health system in Kenya is composed of public and private health-care service providers whose total currently stands at 9,064 health facilities. The public health system in Kenya was devolved a decade ago and is now run by (regional) county governments. This has led to a more responsive healthcare administration that is closer to the people.

The maternal mortality rate in Kenya is 342 per 100,000 live-births (UNICEF 2022a), and the infant mortality rate is at 41.9 per 1000 live-births. The life expectancy at birth is currently 61.4 years, a ten-year improvement from two decades ago at the height of the HIV pandemic mortality in Kenya (Guto et al. 2016; UNDP 2022).

3.2. HIV-related fungal disease in Kenya

UNAIDS reports that 1,400,000 Kenyan adults and 86,000 children were living with HIV/AIDS in 2021. This number has remained nearly constant over the last 10 years. The number of adults on ARVs has in the same period more than doubled, from 548,588 in 2012 (Guto et al. 2016) to 1,160,570 in 2021. About

150,000 HIV-positive diagnosed adults are at risk of opportunistic infections, as are 24,243 children. AIDS-related deaths have reduced threefold over the last decade, from 57,000 in 2012 to 19,000 in 2021.

3.2.1. Cryptococcal meningitis

A recent global estimate put cryptococcal antigen (CrAg) prevalence for eastern and southern Africa at 3.7% (96% CI 3.6–3.8), with Kenya having a prevalence of about 7% (95% CI 5–10) (Rajasingham et al. 2022). Cryptococcal meningitis is rare in children with HIV, so only adults are included in the risk population. This yielded a total annual incidence of about 9,000 cases (Table 3). This estimate was informed by a 6% proportion of patients with CrAg positive with a CD4 < 200 (Kanji et al. 2011; Meyer et al. 2013).

3.2.2. Disseminated histoplasmosis

Histoplasmosis was recognised early as an AIDS-defining illness (Kalata et al. 2019) and has other underlying causes (Ekeng et al. 2021). We only found one case report of disseminated histoplasmosis (Macharia and Walong 2019). However, we assumed a 4.7% incidence in advanced HIV disease (based on data from Ghana) that yielded an estimation of 2,244 incident cases, possibly a conservative estimate, bearing in mind other underlying causes.

3.2.3. *Pneumocystis pneumonia* (PCP)

The prevalence of PCP in children was estimated to be 17%, from a study conducted on Kenyan children who died following severe respiratory infections (Mirza et al. 2018). The proportion of adults found to present with PCP and a CD4 count of less than 200 cells/mm³ was estimated at 16%. This translated to 17,000 incident cases and a rate of 43 per 100,000 persons (Bii et al. 2006). In the present study, we estimated a PCP annual incidence of 7.1 per 100,000 in children and 46.5 per 100,000 people in adults in Kenya (Table 3).

3.2.4. Mucosal candidiasis in those infected with HIV

We assumed an oral candidiasis attack rate of 90% of at-risk HIV patients over 2 years. This resulted in an incidence of 67,312 cases annually. To estimate oesophageal candidiasis, we assumed that adults affected comprise 20% of new HIV/AIDS patients and 0.5% of those HIV patients on ARVs (Table 2). This yielded an annual incidence of 87,945 cases. There is uncertainty

Table 3. Incidence and prevalence of serious fungal infections in Kenya.

Infection	Incidence or prevalence	Number of infections per underlying disorders per year					Rate/ 100 K	Total burden
		None	HIV/AIDS	Respiratory	Cancer	ICU		
PCP in adults	I	-	25,429	-	-	-	46.5	25,429
PCP in children	I	-	3,879	-	-	-	7.1	3,879
Cryptococcal meningitis	I	-	8,975	-	-	-	16.4	8,975
Histoplasmosis	I	-	2,244	-	-	-	4.1	2,244
Oesophageal candidiasis	I	-	87,945	-	-	-	161	87,945
Chronic pulmonary aspergillosis-Post TB	I	-	-	17,738	-	-	32.4	17,738
Chronic pulmonary aspergillosis-Post TB	P	-	-	80,456	-	-	147	80,456
Chronic pulmonary aspergillosis-all	P	-	-	100,570	-	-	184	100,570
Allergic bronchopulmonary aspergillosis (ABPA)	P	-	-	83,395	-	-	153	83,395
Severe asthma with fungal sensitisation (SAFS)	P	-	-	110,081	-	-	201	110,081
Candidaemia	I	-	-	-	1,914	820	5.0	2,804
Candida peritonitis	I	-	-	-	-	410	0.8	410
Oral candidiasis	I	-	67,312	-	-	-	123	67,312
Recurrent vaginal candidiasis (>4 times/year)	P	915,795	-	-	-	-	3,349*	915,795
Tinea capitis	I	4,905,249	-	-	-	-	8,970	4,905,249
Fungal keratitis	I	7,929	-	-	-	-	14.5	7,929
Mycetoma	P	ND	-	-	-	-	-	-
Invasive aspergillosis	I	-	760	5,053	385	-	11.33	6,198
Mucormycosis	I	109	-	-	-	-	0.20	109
Total burden estimated		5,829,083	196,543	299,099	2,299	1,230		6,328,294

Note: *females only
ND- No Data

about the frequency of oesophageal candidiasis in children.

3.3. Aspergillosis

Aspergillosis is a condition that affects individuals who have an underlying pulmonary disease and is highly prevalent in immunocompromised populations (Fosses Vuong and Waymack 2022). Tuberculosis (TB) prevalence in Kenya was found to be higher than previously estimated (Enos et al. 2018) where approximately half of those who contract the disease annually are missed. The PTB incidence in HIV was estimated to be 35,000. This represents about 40% of the annual PTB incidence in Kenya (Table 1) (UNAIDS 2022).

3.3.1. Chronic pulmonary aspergillosis

We estimated the incidence and prevalence of CPA from 2020 WHO pulmonary TB figures. Misdiagnosis of CPA as TB probably occurs in 5,490 HIV-negative and 1,842 HIV-positive people annually, with an additional 2,396 people presenting with dual infection with TB and CPA. An additional 8,424 patients who successfully complete anti-tuberculous therapy develop CPA during that therapy or in the few months following it. An additional 13,918 people develop CPA in the two to five years following TB

cure, with an overall annual incidence of 32,070. Some of these patients will die and so the five-year period prevalence is estimated at 88,015 patients, with an annual mortality related to CPA of 9,172. We have also assumed that 80% of CPA cases are linked to TB, leading to a total prevalence of >110,000 cases.

3.3.2. Invasive aspergillosis and mucormycosis

There is paucity of data on cancers and transplants in Kenya. Available sources have estimated the number of lung cancer deaths at 729 per 100,000 persons (International Agency for Cancer on Research WHO 2022). Approximately 1,367 new leukaemia patients are seen annually, with an annual incidence rate of 2.5/100,000. A total of 470 renal transplants and 12 liver transplants are done each year in Kenya. COPD prevalence for at GOLD stages 2–4 stood at 7.8%, or 3.7 million, of whom an estimated 388,665 are admitted to hospital or die at home (Hammond et al. 2020).

Mucormycosis is a rarely diagnosed condition in Kenya. We used a conservative estimate of 0.2 per 100,000, which gave an annual burden of 109 cases. One of its biggest risk factors is diabetes mellitus. In Kenya, only 20% of those with diabetes were found to have received treatment, and less than 10% of those who knew their diabetes status had their blood

glucose under control (Mohamed et al. 2018). This means the actual number of cases could be significantly more than our estimates.

For IA annual incidence, we assumed a 13% incidence in AML and an equal number of other patients with haematological malignancy and lymphoma (355 cases annually), 2.6% of lung cancer patients (21 cases), 1.3% of COPD admissions (5,053 cases annually), 2% and 4% of renal and liver transplant recipients respectively (10 cases). An estimated 760 HIV/AIDS patients died of invasive aspergillosis each year and 385 cases from cancer and other immunosuppressive conditions, including transplantation. Our annual incidence estimate for invasive aspergillosis incidence 6,198 cases.

3.3.3. Fungal asthma and allergic aspergillosis

Kenya has an asthma prevalence of 10% in adults according to findings from a national survey. Translating to a total burden of 3,335,788 adult cases. Allergic bronchopulmonary aspergillosis (ABPA) was found to have a prevalence of an estimated 83,395 cases almost all with asthma, usually poorly controlled asthma. Cystic fibrosis is barely recognised in Kenya. We also estimated severe asthma with a fungal sensitisation prevalence of 110,081 cases with severe disease. The underlying assumption here was that 33% of 10% adult asthmatics had severe asthma (Denning et al. 2014), and there is some overlap with ABPA. We have not been able to estimate allergic fungal rhinosinusitis in the absence of any data.

3.4. Candidiasis and candidemia

To estimate the annual incidence of candidemia, we assumed a rate of 5/100,000, this being the mean of 2–11/100,000 persons (Arendrup 2010; Bongomin et al. 2017). This translated to a burden of 1,914 cases of individuals with cancer and other immunocompromising conditions and 820 cases from surgery and critical care, summing up to a total figure of 2,734 cases. One 12-month study of 378 patients admitted to intensive care in Nairobi in 2019–2020 found 31 (8.2%) to have a bloodstream *Candida* infection. The commonest 2 species were *Candida auris* (29%) and *Candida albicans* (26%) (Solomon et al. 2021). These data don't allow a national extrapolation as there are

very few intensive care beds in public hospitals in Kenya.

For *Candida* peritonitis, we estimate an annual incidence of 0.75 per 100,000 persons, yielding a total burden of 410 cases among those who are critically ill and have undergone surgical procedures. This is based on a ratio of 2 cases of candidaemia for every case of intraabdominal candidiasis in intensive care (*Candida* peritonitis) (Montravers et al. 2011).

Recurrent *Candida* vaginitis is characterised by four or more episodes of vulvovaginal candidiasis over a year (Lakoh et al. 2021). To estimate the annual incidence we assumed that 6% of females aged between 15 and 49 years are affected (Denning et al. 2018). This gave a total burden of 915,795 cases almost all in individuals with no underlying disease.

3.5. Fungal keratitis

Fungal keratitis is rarely diagnosed in Kenya, although the Ministry of Health collects annual data on microbial keratitis (Brown et al. 2021). Using data from other African countries for the ratio of bacterial to fungal keratitis, we conservatively estimate an annual incidence of 14.5 per 100,000 persons, translating to 7,929 people whose eyes are affected, most losing their sight.

3.6. Tinea capitis and other serious skin fungal diseases

Tinea capitis is a fairly ubiquitous condition in Kenya that affects a large number of rural and urban dwellers, primarily school-going children. The most prevalent aetiological agents in Kenya are *Trichophyton* spp, *Epidermophyton* spp, and *Microsporum* spp. which occur singly, dually, or as a triple infection (Moto et al. 2015). We estimated the annual prevalence of 4,905,249 cases almost all without any underlying disease.

Mycetoma was reported relatively frequently in the 1960s (Cameron et al. 1973a). This case series of 155 cases provided data on both Actinomycetoma and Eumycetoma. The cultures were often negative, but 34 cases were attributed to *Madurella mycetomatis*. We were unable to find any epidemiological studies on chromoblastomycosis, but there is one published case (Cameron et al. 1973b). No studies on sporotrichosis were found.

The total burden of serious fungal infections is estimated to affect 6,328,294 persons which translates to 11.57% of the Kenyan population.

4. Discussion

The last decade saw Kenya's economic status raised to the lower middle-income level. This had significant implications for health-related parameters including the type and number of fungal infections in the country. It also means more and more people can afford basic amenities and have higher expectations of health services (The Kenya Institute for Public Policy Research and Analysis 2020; Chepkondol et al. 2020). There are now more healthcare facilities, more healthcare workers, and improved diagnostics across board. Research outputs also improved within the same period, allowing for more up-to-date fungal burden estimates.

An increase in the number of people affected with non-communicable diseases in Kenya has significantly increased as well. The number of cancer cases, especially has gone up, as has the number of kidney transplants conducted in Kenya. The number of immunocompromised individuals has also increased, in part, due to an increase in the screening and diagnosis of cancer in Kenya, particularly breast and cervical cancer.

The period also saw a transition of the country's health system, from a centralised national system to a more regional, devolved county system. This in effect meant perennially experienced stock-outs of drugs including antifungals were mitigated more swiftly at the local level.

The number of HIV cases has remained fairly constant over the last five years, owing largely to aggressive campaigns targeting HIV diagnosis and recruitment to and retention in care of many more HIV-positive people. Both new HIV infections and mortality have decreased benefitting from multifaceted biomedical and social HIV prevention strategies and the integration of HIV services into the public health system in Kenya. However, evidence points to an increased HIV mortality risk, owing to HIV-positive individuals starting antiretroviral therapy late with progressive HIV disease (Silverman et al. 2019).

Cryptococcal meningitis diagnosis relied on culture but more sensitive and cost-effective techniques such as the lateral flow antigen immunoassay are

increasingly being used in Kenya (Abassi et al. 2015; Gitonga et al. 2019; Lakoh et al. 2022). The overall burden of cryptococcosis has decreased in Kenya compared to the last estimates from 11,900 to 8,975 (Guto et al. 2016). This could be attributed to more conservative estimates used previously as informed by the existing evidence at the time.

In 2015, histoplasmosis was added to Kenya's list of "zoonotic diseases of public health concern" (Munyua et al. 2016). Chest X-rays of Suk (Chok or Chuk) tribesman in 1949 (northeast of Eldoret) showed some with calcification, characteristic of prior resolved histoplasmosis. In 1951, Dr. Hugh Stott skin-tested 768 gold miners (east of Lake Victoria) and male prisoners in the goal in Nakuru (mostly from the Rift Valley) with histoplasmin (Stott 1954). He found 65 (8.5%) to be positive, with the miners having a 14.6% positivity rate and prisoners 5.3%. Chest X-ray calcification was found in 34.8% of those with a positive histoplasmin reaction compared with those with negative tests (9.8%). *Histoplasma capsulatum* was first isolated from soil in the village of Darajani, East Kenya in 1964 near the Chyulu Hills National Park, home to some huge lava tubes (caves) (al-Doory and Kalter 1967). These caves house huge numbers of bats. This was the second isolation of *H. capsulatum* from east Africa, the other being from Tanzania in 1960. Histoplasmosis clearly is present in Kenya, with very few case reports only. *Histoplasma* antigen testing of urine and serum (the most sensitive assay for disseminated histoplasmosis) is not available in Kenya (Lakoh et al. 2022).

PCP has a higher incidence (25,429 cases in adults and 3,879 cases in children) than previously estimated (17,000) (Guto et al. 2016). This corroborates the evolving epidemiological trends that showed an increase in incidence over a recent five-year period (Kolbrink et al. 2022). PCP fatalities are probably also linked with cancer and autoimmune diseases. Currently, most diagnoses are made clinically, but the broader at-risk population and frequent bacterial co-infection in babies with PCP underlines the need to implement *Pneumocystis* PCR diagnosis across the country (Morrow et al. 2014). *Pneumocystis* PCR (a WHO Essential diagnostic) is not done in any public facility in Kenya (Lakoh et al. 2022).

Oral and oesophageal candidiasis still pose a significant health challenge in Kenya. The burden

of infection reduced significantly from the last estimate and this is largely due to the widespread use of HIV opportunistic infection prevention medications in the country. Recurrent vaginal candidiasis cases with no underlying cause increased by 50% from the last estimation (using the same 6% proportion), making it the second most burdensome fungal infection in Kenya. This increase reflects changing population demographics in Kenya. The risk of infection increases with a history of bacterial vaginosis, diabetes, wearing panty liners, drinking acidic juices such as cranberry juice, antibiotic use, having intra-uterine devices, and a short ano-vaginal distance, among others (Patel et al. 2004; Guzel et al. 2011).

CPA is being increasingly recognised in Kenya. Data from across Africa reveals growing interest in CPA as documented from South Africa, Senegal, and Morocco (Olum et al. 2021). Most recently in Kenya, serological evidence of *Aspergillus* infection from HIV-positive patients with difficult-to-treat tuberculosis found at least 46 cases (Mohamed et al. 2022). Our current estimate is nearly eight-fold higher than previously reflecting a better understanding of CPA internationally, supported by national data. Our estimate of over 100,000 prevalence emphasises the need to incorporate *Aspergillus* IgG antibody testing into TB programmes across Kenya. Likewise, with invasive *aspergillosis* mortality in Africa exceeding 60% (Yerbanga et al. 2023), the inclusion of *Aspergillus* antigen testing into leukaemia and intensive care would be appropriate. Ongoing work on fungal adhesins points to a promising potential for addressing invasive fungal infections in immunosuppression both preventively and therapeutically, this will lead to better estimates in future (Kumari et al. 2021).

A substantial number of Kenyan patients were found to have severe persistent asthma; 10% of adults compared with a 2003 estimate used previously of 3.12%. These patients presented with higher rates of adult-onset asthma and lower rates of rhinosinusitis, among other comorbidities (Kirenga et al. 2020). A significant increase in the number of ABPA and SAFS cases was established in this study, a five-fold increase from the last estimate. Fungal asthma in Africa recently quantified by Kwizera et al. found a relatively high prevalence of fungal sensitisations

in asthmatics (3%-52%) and a substantially high prevalence of ABPA estimated at 1.6%-21.2% (Kwizera et al. 2019). More cases are being diagnosed in Kenya than before. More focus should now be put on antifungal treatment as an alternative to long-term prednisolone or biologics to improve asthma control (Moss 2014).

Fungal rhinosinusitis (FRS) is poorly diagnosed and not documented in Kenya, it is divided into invasive and non-invasive FRS. Most of the former are captured in IA and mucormycosis, these are life-threatening and life-altering for survivors because of (usually) extensive surgery (Searyoh et al. 2020). Non-invasive FRS includes sinus fungal ball and fungus-related eosinophilic FRS (Chakrabarti et al. 2009). Invasive fungal rhinosinusitis. More awareness of fungal rhinosinusitis in Kenya is required.

Candida auris is less virulent but as a cause of invasive *candidiasis* may pose a significant challenge to eliminate from Kenyan patients (Adam et al. 2019). Studies have shown the existence of diversity in both genetic and biological characteristics of different *Candida auris* clades that are resistant to at least one antifungal drug (Bing et al. 2022). Our estimate of *Candida* bloodstream infection has increased in line with an increasing population size (5/100,000).

The estimate for invasive *aspergillosis* is substantially higher than the previous estimate. This reflects the addition of deaths from AIDS being included (4%), COPD as an underlying disease being added and increased numbers of renal transplant procedures and lung cancer cases.

We have included an estimate of fungal keratitis incidence, which was absent previously. We know that microbial keratitis is relatively common in Kenya (based on annual County reports), but distinction between bacterial and fungal keratitis is only occasionally made (Brown et al. 2021).

Azole resistant *Aspergillus* is present in nearly all continents, and some variants have a high fatality rate even in Kenya (Jeanvoine et al. 2020; Burks et al. 2021). The drug used, clinical setting and patient characteristics including gene alterations such as *cyp51* gene, all lead to varied levels of resistance (Chowdhary et al. 2017; Guegan et al. 2020).

Tinea capitis was the most burdensome condition accounting for 77% of all fungal infections in Kenya. Mostly affecting school-going children, this

contagious disease spreads quickly among young children who intimately interact with each other during child play. The pooled prevalence in Africa was found to affect more than one in five school-aged children (Bongomin et al. 2021). In the prior estimate, 1.7 million children were thought to be affected (Guto et al. 2016); our current estimate is 4.9 million. This new estimate reflects the large increase in children in Kenya from 17.1 to 21.3 million over 7 years and the use of a more up to date estimate of the proportion affected – from 10% to 23%. Treatment options available in Kenya include griseofulvin, itraconazole, terbinafine, and fluconazole (Alkeswani et al. 2019), most of which are readily available.

Skin fungal infections were excluded from this study due to data paucity occasioned by human resource capacity challenges. Most regions in sub-Saharan Africa have less than 1 dermatologist per million people. This coupled with limited training in dermatology makes it difficult to accurately diagnose skin fungal NTDs (Mosam and Todd 2021). Promising advances in collaborative skin research shine a ray of hope for future quantification of fungal skin NTDs in Kenya and Africa at large (Verschoore and Dlova 2022).

The national infection prevention and control guidelines in Kenya were last updated more than a decade ago. Lack of interventions and measures specifically targeting fungal conditions in Kenya are obviated by the lack of mycology treatment guidelines. This significantly lowers the suspicion indices by healthcare workers, delaying the diagnosis and treatment of fungal infections (Sangoi et al. 2009; Caudron de Coquereaumont et al. 2021) in Kenya.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This study was not externally or otherwise funded.

ORCID

Stanley N. Ratemo  <http://orcid.org/0000-0003-2632-0208>

Author contributions

SNR: Concept design and development, data acquisition, drafting the article, data analysis and interpretation, final approval of the version to be published.

DWD: Concept design and development, data acquisition, drafting the article, data analysis and interpretation, revising critically for important intellectual content, final approval of the version to be published.

References

- Abassi M, Boulware DR, Rhein J. 2015. Cryptococcal meningitis: diagnosis and management update. *Curr Trop Med Rep.* 2 (2):90–99. doi:10.1007/s40475-015-0046-y.
- Adam RD, Revathi G, Okinda N, Fontaine M, Shah J, Kagoto E, Castanheira M, Pfaller MA, Maina D. 2019. Analysis of *Candida auris* fungemia at a single facility in Kenya. *Int J Infect Dis.* 85:182–187. doi:10.1016/j.ijid.2019.06.001.
- al-Doory Y, Kalter SS. 1967. The isolation of *Histoplasma duboisii* and keratinophilic fungi from soils of East Africa. *Mycopathol Mycol Appl.* 31(3–4):289–295. doi:10.1007/BF02053427.
- Alkeswani A, Cantrell W, Elewski B. 2019. Treatment of tinea capitis. *Skin Appendage Disorders.* 5(4):201–210. doi:10.1159/000495909.
- Arendrup MC. 2010. Epidemiology of invasive candidiasis. *Curr Opin Crit Care.* 16(5):445–452. doi:10.1097/MCC.0b013e32833e84d2.
- Bii CC, Kose J, Taguchi H, Amukoye E, Ouko TT, Muita LC, Mugasia O, Wamae N, Kamiya S. 2006. Pneumocystis jirovecii and microbiological findings in children with severe pneumonia in Nairobi, Kenya. *Int J Tuberc Lung Dis.* 10 (11):1286–1291.
- Bing J, Wang S, Xu H, Fan S, Du H, Nobile CJ, Huang G. 2022. A case of *Candida auris* candidemia in Xiamen, China, and a comparative analysis of clinical isolates in China. *Mycology.* 13(1):68–75. doi:10.1080/21501203.2021.1994479.
- Bongomin F, Gago S, Oladele RO, Denning DW. 2017. Global and multi-national prevalence of fungal diseases—estimate precision. *J Fungi (Basel).* 3(4):57. doi:10.3390/jof3040057.
- Bongomin F, Olum R, Nsenga L, Namusobya M, Russell L, de Sousa E, Osaigbovo KR II, Baluku JB, Baluku JB. 2021. Estimation of the burden of tinea capitis among children in Africa [10.1111/myc.13221]. *Mycoses.* 64(4):349–363. doi:10.1111/myc.13221.
- Brooks RB, Walters M, Forsberg K, Vaeth E, Woodworth K, Vallabhaneni S. 2019. *Candida auris* in a U.S. patient with Carbapenemase-producing organisms and recent hospitalization in Kenya. *MMWR Morb Mortal Wkly Rep.* 68 (30):664–666. doi:10.15585/mmwr.mm6830a3.
- Brown L, Leck AK, Gichangi M, Burton MJ, Denning DW. 2021. The global incidence and diagnosis of fungal keratitis. *Lancet Infect Dis.* 21(3):e49–e57. doi:10.1016/S1473-3099(20)30448-5.

- Buchacz K, Baker RK, Palella FJ Jr, Chmiel JS, Lichtenstein KA, Novak RM, Wood KC, Brooks JT. 2010. AIDS-defining opportunistic illnesses in US patients, 1994-2007: a cohort study. *Aids*. 24(10):1549–1559. doi:10.1097/QAD.0b013e32833a3967.
- Burks C, Darby A, Gómez Londoño L, Momany M, Brewer MT. 2021. Azole-resistant *Aspergillus fumigatus* in the environment: identifying key reservoirs and hotspots of antifungal resistance. *PLoS Pathog*. 17(7):e1009711.
- Cameron HM, Gatei D, Bremner AD. 1973a. The deep mycoses in Kenya: a histopathological study. 1. Mycetoma. *East Afr Med J*. 50(8):382–395.
- Cameron HM, Gatei D, Bremner AD. 1973b. The deep mycoses in Kenya: a histopathological study. 3. Chromomycosis. *East Afr Med J*. 50(8):406–412.
- Caudron de Coquereumont G, Couchepin J, Perentes JY, Krueger T, Lovis A, Rotman S, Lamoth F. 2021. Limited index of clinical suspicion and underdiagnosis of histopathologically documented invasive mold infections. *Open Forum Infect Dis*. 8(7):ofab174.
- Chakaya JM, Bii C, Ng'ang'a L, Amukoye E, Ouko T, Muita L, Gathua S, Gitau J, Odongo I, Kabanga JM, et al. 2003. Pneumocystis carinii pneumonia in HIV/AIDS patients at an urban district hospital in Kenya. *East Afr Med J*. 80(1):30–35.
- Chakrabarti A, Denning DW, Ferguson BJ, Ponikau J, Buzina W, Kita H, Marple B, Panda N, Vlamincik S, Kauffmann-Lacroix C, et al. 2009. Fungal rhinosinusitis: a categorization and definitional schema addressing current controversies. *Laryngoscope*. 119(9):1809–1818.
- Chen PY, Chuang YC, Wang JT, Sheng WH, Yu CJ, Chu CC, Hsueh PR, Chang SC, Chen YC. 2014. Comparison of epidemiology and treatment outcome of patients with candidemia at a teaching hospital in Northern Taiwan, in 2002 and 2010. *J Microbiol Immunol Infect*. 47(2):95–103.
- Chen CY, Sheng WH, Tien FM, Lee PC, Huang SY, Tang JL, Tsay W, Tien HF, Hsueh PR. 2020. Clinical characteristics and treatment outcomes of pulmonary invasive fungal infection among adult patients with hematological malignancy in a medical centre in Taiwan, 2008-2013. *J Microbiol Immunol Infect*. 53(1):106–114.
- Chepkondol GK, Jolly PE, Yatich N, Mbowe O, Jaoko WG. 2020. Types and prevalence of HIV-related opportunistic infections/conditions among HIV-positive patients attending Kenyatta National Hospital in Nairobi, Kenya. *Afr Health Sci*. 20(2):615–624.
- Chowdhary A, Sharma C, Meis JF. 2017. Azole-resistant aspergillosis: epidemiology, molecular mechanisms, and treatment. *J Infect Dis*. 216(suppl_3):S436–S444.
- CIA. 2022. CIA The world fact book: kenya-Country summary; [accessed 2022 Oct 1]. <https://www.cia.gov/the-world-factbook/countries/kenya/summaries>.
- Commons D. 2022. Timelines; [accessed 2022 June 21]. https://datacommons.org/tools/timeline#&place=country/KEN&statsVar=Count_Person.
- Control UfC. 2014. Acute Myelogenous leukemia and acute promyelocytic leukemia. Review of cancer medicines on WHO list of essential medicines. Geneva, Switzerland: Union for International Cancer Control.
- Denning DW, Cole DC, Ray A. 2023. New estimation of the prevalence of chronic pulmonary aspergillosis (CPA) related to pulmonary TB - a revised burden for India. *IJID Reg*. 6:7–14.
- Denning DW, Kneale M, Sobel JD, Rautemaa-Richardson R. 2018. Global burden of recurrent vulvovaginal candidiasis: a systematic review. *Lancet Infect Dis*. 18(11):e339–e347.
- Denning DW, Morgan EF. 2022. Quantifying deaths from aspergillosis in HIV positive people. *J Fungi*. doi:10.3390/jof8111131.
- Denning DW, Pashley C, Hartl D, Wardlaw A, Godet C, Del Giacco S, Delhaes L, Sergejeva S. 2014. Fungal allergy in asthma-state of the art and research needs. *Clin Transl Allergy*. 4:14.
- Denning DW, Pleuvry A, Cole DC. 2013. Global burden of allergic bronchopulmonary aspergillosis with asthma and its complication chronic pulmonary aspergillosis in adults. *Med Mycol*. 51(4):361–370.
- Ekenge BE, Edem K, Amamilo I, Panos Z, Denning D, Oladele RO. 2021. Histoplasmosis in Children; HIV/AIDS Not a major driver. *J Fungi*. 7(7):530.
- Elephant T. 2022. Kenya Gazette Notice No. 786 - Categorization of Health Facilities. [accessed 2022 Oct 30th]. <https://www.theelephant.info/documents/kenya-gazette-notice-no-786-categorization-of-health-facilities/>.
- Enos M, Sitienei J, Ong'ang'o J, Mungai B, Kamene M, Wambugu J, Kipruto H, Manduku V, Mburu J, Nyaboke D, et al. 2018. Kenya tuberculosis prevalence survey 2016: challenges and opportunities of ending TB in Kenya. *PLoS One*. 13(12):e0209098.
- Fosses Vuong M, Waymack JR. 2022. Aspergillosis. StatPearls. Treasure Island (FL): statPearls Publishing Copyright © 2022. Tampa, Florida, United States: StatPearls Publishing LLC.
- GAFI.org. 2022. Diagnostics for fungal diseases in Africa; [accessed 2023 Dec 1]. <https://gaffi.org/africa-diagnostic-reports-2/>.
- Gitonga LK, Boru WG, Kwena A, Maritim M, Wamicwe J, Ransom J. 2019. Point of care testing evaluation of lateral flow immunoassay for diagnosis of cryptococcus meningitis in HIV-positive patients at an urban hospital in Nairobi. *BMC Research Notes*. Vol. 12. Springer Nature Switzerland AG; p. 797.
- Guarner J. 2017. Human immunodeficiency virus and fungal infections. *Semin Diagn Pathol*. 34(4):325–331.
- Guegan H, Prat E, Robert-Gangneux F, Gangneux JP. 2020. Azole Resistance in *Aspergillus fumigatus*: a five-year follow up experience in a tertiary hospital with a special focus on cystic fibrosis. *Front Cell Infect Microbiol*. 10:613774.
- Guto JA, Bii CC, Denning DW. 2016. Estimated burden of fungal infections in Kenya. *J Infect Dev Ctries*. 10(8):777–784.
- Guzel AB, Ilkit M, Akar T, Burgut R, Demir SC. 2011. Evaluation of risk factors in patients with vulvovaginal candidiasis and the value of chromID Candida agar versus CHROMagar Candida for recovery and presumptive identification of vaginal yeast species. *Medical Mycology*. 49(1):16–25.

- Hammond EE, McDonald CS, Vestbo J, Denning DW. 2020. The global impact of *Aspergillus* infection on COPD. *BMC Pulm Med.* 20(1):241.
- International Agency for Cancer on Research WHO. 2022. Kenya fact sheets. [accessed 2022 Oct 20th]. <https://gco.iarc.fr/today/data/factsheets/populations/404-kenya-fact-sheets.pdf>.
- IRODaT. 2023. Kenya national tissues and organ transplant authority. kenya deceased organ donor evolution; [accessed 2023 Jan 4]. <https://www.irodat.org/?p=databse&c=KE#data>.
- Jeanvoine A, Rocchi S, Bellanger AP, Reboux G, Millon L. 2020. Azole-resistant *Aspergillus fumigatus*: a global phenomenon originating in the environment? *Med Mal Infect.* 50(5):389–395.
- Kalata KE, Osborne C, Willis A, Navarro K, Fenton LZ, Smith C. 2019. Disseminated histoplasmosis as an AIDS-defining illness presenting as fever of unknown origin in an 11-Year-Old Female. *Case Rep Pediatr.* 2019:9417102.
- Kanji SS, Kaikai S, Onyango RO. 2011. Cryptococcal meningitis among human immunodeficiency virus patients attending major hospitals in Kisumu, Western Kenya. *Arch Clin Microbiol.* 2:1–6.
- Kemoi EK, Nyerere A, Bii CC. 2018. Triazole-Resistant *Aspergillus fumigatus* from fungicide-experienced soils in Naivasha sub-county and Nairobi County, Kenya. *Int J Microbiol.* 2018:7147938.
- The Kenya Institute for Public Policy Research and Analysis. 2020. Creating an enabling environment for inclusive growth in Kenya. Nairobi, Kenya: The Kenya Institute for Public Policy Research and Analysis (KIPPRA).
- Khoo SD DW. 1994. *Aspergillus* infection in the acquired immune deficiency syndrome. *Clin Infect Dis.* 19:541–548.
- Kirenga B, Chakaya J, Yimer G, Nyale G, Haile T, Muttamba W, Mugenyi L, Katagira W, Worodria W, Aanyu-Tukamuhebwa H, et al. 2020. Phenotypic characteristics and asthma severity in an East African cohort of adults and adolescents with asthma: findings from the African severe asthma project. *BMJ Open Respir Res.* 7(1):e000484.
- Kolbrink B, Scheikholeslami-Sabzewari J, Borzikowsky C, von Samson-Himmelstjerna Fa, Ullmann AJ, Kunzendorf U, Schulte K. 2022. Evolving epidemiology of pneumocystis pneumonia: findings from a longitudinal population-based study and a retrospective multi-center study in Germany. *Lancet Regi Health.* 18:100400.
- Kumari A, Tripathi AH, Gautam P, Gahtori R, Pande A, Singh Y, Madan T, Upadhyay SK. 2021. Adhesins in the virulence of opportunistic fungal pathogens of human. *Mycology.* 12(4):296–324.
- Kwizera R, Bongomin F, Olum R, Meya DB, Worodria W, Bwanga F, Fowler SJ, Gore R, Denning DW, Kirenga BJ. 2021. Fungal asthma among Ugandan adult asthmatics. *Med Mycol.* 59(9):923–933.
- Kwizera R, Musaaazi J, Meya DB, Worodria W, Bwanga F, Kajumbula H, Fowler SJ, Kirenga BJ, Gore R, Denning DW. 2019. Burden of fungal asthma in Africa: a systematic review and meta-analysis. *PLoS One.* 14(5):e0216568.
- Kwizera R, Wadda V, Mugenyi L, Aanyu-Tukamuhebwa H, Nyale G, Yimer G, Chakaya J, De Jong C, Van der Molen T, Denning DW, et al. 2020. Skin prick reactivity among asthmatics in East Africa. *World Allergy Organ J.* 13(6):100130.
- Lakoh S, Kamudumuli PS, Penney ROS, Haumba SM, Jarvis JN, Hassan AJ, Moudoute NLE, Ocansey BK, Izco S, Kipkerich S, et al. 2022. Diagnostic capacity for invasive fungal infections in advanced HIV disease in Africa: a continent-wide survey. *Lancet infect dis.* S1473-3099(22):00682–X.
- Lakoh S, Orefuwa E, Kamara MN, Jiba DF, Kamara JB, Kpaka S, Denning DW. 2021. The burden of serious fungal infections in Sierra Leone: a national estimate. *Ther Adv Infect Dis.* 8:204993612111027996.
- Macharia A, Ombajo L, Farah A, Walong E, Mwangi G, Wamalwa D. 2019. Empyema necessitans due to *Aspergillus* in a 3-year-old paediatric patient at a referral hospital in Nairobi, Kenya. *Pan Afr Med J.* 34:86.
- Macharia AM, Walong E. 2019. Disseminated histoplasmosis in a paediatric patient at a referral hospital in Nairobi, Kenya. *Pan Afr Med J.* 34(86). doi:10.11604/pamj.2019.34.86.19469.
- Maertens J, Vrebos M, Boogaerts M. 2001. Assessing risk factors for systemic fungal infections [10.1046/j.1365-2354.2001.00241.x]. *Eur J Cancer Care (Engl).* 10(1):56–62.
- Malwal FM, Muriuki G, Maru SS, Mwangangi EM. 2016. Prevalence of opportunistic infections in HIV-infected adult patients at Kenyatta National Hospital, Kenya. *African J Pharmacol Ther.* 5:163–168.
- Meyer AC, Kendi CK, Penner JA, Odhiambo N, Otieno B, Omondi E, Opiyo E, Bukusi EA, Cohen CR. 2013. The impact of routine cryptococcal antigen screening on survival among HIV-infected individuals with advanced immunosuppression in Kenya. *Trop Med Int Health.* 18(4):495–503.
- Ministry of Health Kenya. 2021. Ten Years and 200 transplants later. KNH Celebrates a Milestone; [accessed 2022 Oct 19]. <https://www.health.go.ke/10-years-and-200-transplants-later-knh-celebrates-a-milestone/>.
- Mirza IB, Gachii AK, Okiro P, Zaki SR. 2018. The prevalence of pneumocystis in lung tissue of Kenyan children who died following severe respiratory infection. *J Trop Dis & Public Health.* 6(5): 1000288.
- Mohamed SF, Mwangi M, Mutua MK, Kibachio J, Hussein A, Ndegwa Z, Owondo S, Asiki G, Kyobutungi C. 2018. Prevalence and factors associated with pre-diabetes and diabetes mellitus in Kenya: results from a national survey. *BMC Public Health.* 18(3):1215.
- Mohamed A, Obanda BA, Njeri HK, Loroyokie SN, Mashedi OM, Ouko TT, Gatumba EM, Korir RK, Yaguchi T, Bii CC. 2022. Serological evidence of chronic pulmonary Aspergillosis in tuberculosis patients in Kenya. *BMC Infect Dis.* 22(1):798.
- Montravers P, Mira JP, Gangneux JP, Leroy O, Lortholary O. 2011. A multicentre study of antifungal strategies and outcome of *Candida* spp. peritonitis in intensive-care units. *Clin Microbiol Infect.* 17(7):1061–1067.
- Morrow BM, Samuel CM, Zampoli M, Whitelaw A, Zar HJ. 2014. Pneumocystis pneumonia in South African children diagnosed by molecular methods. *BMC Res Notes.* 7(1):26.

- Mosam A, Todd G. 2021. Dermatology training in Africa: successes and challenges. *Dermatol Clin.* 39(1):57–71.
- Moss RB. 2014. Treatment options in severe fungal asthma and allergic bronchopulmonary aspergillosis. *European Resp J.* 43(5):1487.
- Moto JN, Maingi JM, Nyamache AK. 2015. Prevalence of *Tinea capitis* in school going children from Mathare, informal settlement in Nairobi, Kenya. *BMC Res Notes.* 8:274.
- Mundi I. 2022. Kenya demographics profile. [accessed 2022 25th Oct 2022 Oct 25th]. https://www.indexmundi.com/kenya/demographics_profile.html.
- Munyua P, Bitek A, Osoro E, Pieracci EG, Muema J, Mwatondo A, Kungu M, Nanyingi M, Gharpure R, Njenga K, et al. 2016. Prioritization of zoonotic diseases in Kenya, 2015. *PLoS One.* 11(8):e0161576.
- News K. 2021. Hospital Marks 151st Kidney transplant; [accessed 2022 Oct 19]. <https://www.kenyanews.go.ke/med/ih-eal-group-of-hospitals-celebrates-successful-kidney-transplant/>.
- Ngugi PN, Gesicho MB, Babic A, Were MC. 2020. Assessment of HIV data reporting performance by facilities during EMR systems implementations in Kenya. 272. 18th annual International Conference on Informatics, Management, and Technology in Healthcare (ICIMTH 2020), virtually in Athens, Greece, from 2020; 2020 Jul 3 – 3; Athens, Greece.
- Ocansey BK, Otoo B, Asamoah I, Ganu V, Berko KP, Oladele O, Amankwa EA, Opoku-Asare B, Agyei M, George L, et al. 2022. Cryptococcal and histoplasma antigen screening among people with human immunodeficiency virus in Ghana and comparative analysis of oidx histoplasma lateral flow assay and IMMY histoplasma enzyme immunoassay. *Open Forum Infect Dis.* 9(7):ofac277.
- Oladele RO, Irurhe NK, Foden P, Akanmu AS, Gbaja-Biamila T, Nwosu A, Ekundayo HA, Ogunsola FT, Richardson MD, Denning DW. 2017. chronic pulmonary aspergillosis as a cause of smear-negative TB and/or TB treatment failure in Nigerians. *Int J Tuberc Lung Dis.* 21(9):1056–1061.
- Olum R, Osaigbovo BJB II, Stemler J, Kwizera R, Bongomin F. 2021. Mapping of Chronic Pulmonary Aspergillosis in Africa. *J Fungi (Basel).* 7(10):790.
- Page ID, Byanyima R, Hosmane S, Onyachi N, Opira C, Richardson M, Sawyer R, Sharman A, Denning DW. 2019. Chronic pulmonary aspergillosis commonly complicates treated pulmonary tuberculosis with residual cavitation. *Eur Respir J.* 53(3):1801184.
- Patel DA, Gillespie B, Sobel JD, Leaman D, Nyirjesy P, Weitz MV, Foxman B. 2004. Risk factors for recurrent vulvovaginal candidiasis in women receiving maintenance antifungal therapy: results of a prospective cohort study. *Am J Obstet Gynecol.* 190(3):644–653.
- Rajasingham R, Govender NP, Jordan A, Loyse A, Shroufi A, Denning DW, Meza DB, Chiller TM, Boulware DR. 2022. The global burden of HIV-associated cryptococcal infection in adults in 2020: a modelling analysis. *Lancet Infect Dis.* 22(12): 1748–1755.
- Samaranayake L, Fidel P, Sweet S, Teanpaisan R, Coogan M, Bignaut E, Wanzala P. 2002. Fungal infections associated with HIV infection. *Oral Dis.* 8:151–160.
- Sangoi AR, Rogers WM, Longacre TA, Montoya JG, Baron EJ, Banaei N. 2009. Challenges and pitfalls of morphologic identification of fungal infections in histologic and cytologic specimens: a ten-year retrospective review at a single institution. *Am J Clin Pathol.* 131(3):364–375.
- Searyoh K, Bankah P, Acquah S. 2020. Complicated sinusitis in developing countries: an African perspective. *Curr Opin Otolaryngol Head Neck Surg.* 28(3):182–187.
- Sharma A, Goel A. 2022. Mucormycosis: risk factors, diagnosis, treatments, and challenges during COVID-19 pandemic. *Folia Microbiol (Praha).* 67(3):363–387.
- Silverman RA, John-Stewart GC, Beck IA, Milne R, Kiptinness C, McGrath CJ, Richardson BA, Chohan B, Sakr SR, Frenkel LM, et al. 2019. Predictors of mortality within the first year of initiating antiretroviral therapy in urban and rural Kenya: a prospective cohort study. *PLoS One.* 14(10):e0223411.
- Smith NL, Denning DW. 2011. Underlying conditions in chronic pulmonary aspergillosis including simple aspergilloma. *Eur Respir J.* 37(4):865–872.
- Smith E, Orholm M. 1990. Trends and patterns of opportunistic diseases in Danish AIDS patients 1980–1990. *Scand J Infect Dis.* 22(6):665–672.
- Solomon DA, Nyerere AK, Kanyua A, Ngugi CW. 2021. Prevalence, species distribution and antifungal susceptibility profile of *candida* species isolated from bloodstream of critical care unit patients in a tertiary care hospital in Kenya. *Open J Med Microbiol.* 11(1):50.
- StopTB. 2021. Tuberculosis Situation in Kenya 2020; [accessed 2022 Nov 1] . https://www.stoptb.org/static_pages/KEN_Dashboard.html.
- Stott H. 1954. Histoplasmin sensitivity and pulmonary calcification in Kenya. *Br Med J.* 1(4852):22–25.
- Tsikhutsu I, Bii M, Dear N, Ganesan K, Kasembeli A, Sing'oei V, Rombosia K, Ochieng C, Desai P, Wolfman V, et al. 2022. Prevalence and correlates of viral load suppression and human immunodeficiency virus (HIV) drug resistance among children and adolescents in South Rift Valley and Kisumu, Kenya. *Clin Infect Dis.* 75(6):936–944.
- Turpin K, Rockers PC, Vian T, Onyango MA, Laing R, Wirtz VJ. 2019. Prevalence and treatment of hypertension, diabetes and asthma in Kenya: a representative household survey in eight counties in 2016. *East Afr Med J.* 95:1286–1297.
- UNAIDS. 2022. UNAIDS DATA 2020. Unaid.org; [accessed 2022 25th Oct 2022 Oct 25th]. https://www.unaids.org/sites/default/files/media_asset/2020_aids_data_book_en.pdf.
- UNDP. 2022. Human development reports. UNDP; [accessed 2022 Oct 25th]. <https://hdr.undp.org/data-center/human-development-index#/indicies/HDI>.
- UNICEF. 2022a. Kenya country profile. [accessed 2022 Oct 30th]. <https://data.unicef.org/country/ken/>.
- UNICEF. 2022b. Kenya nutrition. UNICEF; [accessed 2022 Oct 21st]. <https://www.unicef.org/kenya/nutrition>.

- Vallabhaneni S, Mody RK, Walker T, Chiller T. 2016. The Global Burden of Fungal Diseases. *Infect Dis Clin North Am.* 30(1):1–11.
- Verschoore M, Dlova N. 2022. Advances in dermatology in sub-Saharan Africa in the past 20 years from workshops to the birth of the African Society of Dermatology and Venereology [10.1111/ijd.16073]. *Int J Dermatol.* 61(7):841–847.
- World Bank. 2023. World bank country and lending groups; [accessed 2022 Oct 20]. <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>.
- Yan X, Li M, Jiang M, Zou LQ, Luo F, Jiang Y. 2009. Clinical characteristics of 45 patients with invasive pulmonary aspergillosis: retrospective analysis of 1711 lung cancer cases. *Cancer.* 115(21):5018–5025.
- Yerbanga IW, Nakanabo Diallo S, Rouamba T, Denis O, Rodriguez-Villalobos H, Montesinos I, Bamba S. 2023. A systematic review of epidemiology, risk factors, diagnosis, antifungal resistance, and management of invasive aspergillosis in Africa. *J Medi Mycology.* 33(1):101328.