

REVIEW

Systematic characterization of non-coding RNAs in triple-negative breast cancer

Jie Mei¹  | Leiyu Hao² | Huiyu Wang¹ | Rui Xu² | Yan Liu² | Yichao Zhu^{2,3} | Chaoying Liu¹ 

¹Department of Oncology, Wuxi People's Hospital Affiliated to Nanjing Medical University, Wuxi, China

²Department of Physiology, Nanjing Medical University, Nanjing, China

³State Key Laboratory of Reproductive Medicine, Nanjing Medical University, Nanjing, China

Correspondence

Chaoying Liu, Department of Oncology, Wuxi People's Hospital Affiliated to Nanjing Medical University, 299 Qingyang Road, Wuxi 214023, China.
Email: liuchaoying666@163.com

Yichao Zhu, Department of Physiology, Nanjing Medical University, 101 Longmian Av., Nanjing 211166, China.
Email: zhuyichao@njmu.edu.cn

Funding information

Foundation of Wuxi Health Commission, Grant/Award Number: QNRC003; Natural Science Foundation of Jiangsu Province of China, Grant/Award Number: BE2017626 and BK20181367

Abstract

Triple-negative breast cancer (TNBC) is one of the most aggressive subtypes of breast cancer with negativity for oestrogen receptor (ER), progesterone receptor (PR) and human epidermal growth factor receptor (HER2). Non-coding RNAs (ncRNAs) make up most of the transcriptome and are widely present in eukaryotic cells. In recent years, emerging evidence suggests that ncRNAs, mainly microRNAs (miRNAs), long ncRNAs (lncRNAs) and circular RNAs (circRNAs), play prominent roles in the tumorigenesis and development of TNBC, but the functions of most ncRNAs have not been fully described. In this review, we systematically elucidate the general characteristics and biogenesis of miRNAs, lncRNAs and circRNAs, discuss the emerging functions of these ncRNAs in TNBC and present future perspectives in clinical practice.

1 | INTRODUCTION

In the past few decades, the morbidity of human breast cancer has increased continuously and has led to a great threat to women's lives. According to the statistics gathered by the American Cancer Society, there will be more than 271 000 new cases of breast cancer and approximately 42 260 deaths in 2019.¹ Being a heterogeneous disease, breast cancer can be classified into several main subclasses based on the expression status of oestrogen receptor (ER), progesterone receptor (PR), human epidermal growth factor receptor (HER2) and antigen ki-67 (Ki-67).² Among known

subclasses, triple-negative breast cancer (TNBC) is the most aggressive subtype, which is characterized by negativity for ER, PR and HER2. Great efforts have been made to understand the mechanisms of TNBC carcinogenesis, especially focus on the role of non-coding RNAs (ncRNAs).

Non-coding RNAs make up most of the transcriptome, while protein-coding RNAs only account for 3% of the genome; the remaining 97% is composed of "dark matter" of transcripts with molecular functions.³ It has been proven that the genome "dark matter" can be transcribed into various RNA species, most of which do not encode proteins, namely, ncRNAs, but exert

Jie Mei and Leiyu Hao equally contributed to this study.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Cell Proliferation* Published by John Wiley & Sons Ltd.

significant functions mainly responsible for phenotypic regulation.⁴ The emerging functions of ncRNAs have been generally determined in the cancer research field. Currently, studies of ncRNA-related cancer are commonly concentrated on miRNAs, lncRNAs and circRNAs. Here, we review the general characteristics and functions of ncRNAs and discuss their underlying mechanisms in the carcinogenesis and development of TNBC.

2 | CATEGORIES AND GENERAL CHARACTERISTICS OF NCRNAs

Depending on the number of nucleotides (nt), ncRNAs can be categorized into two main groups: (a) short ncRNAs, which include microRNA (miRNA), small interfering RNA (siRNA), small nucleolar RNA (snoRNA), small nuclear RNA (snRNA), piwi-interacting RNA (piRNA), tRNA-derived stress-induced RNA (tiRNA) and tRNA-derived small non-coding RNA (tDR); (b) long ncRNAs (lncRNAs), which have transcripts with more than 200 nt in length and include long intergenic non-coding RNA (lincRNA), natural antisense transcript (NAT), circular RNA (circRNA), pseudogene transcript, transcribed ultraconserved region (T-UCR) and telomerase RNA component (TERC).³ Although circRNA belonging to the lncRNA family, researchers tend to discuss them separately distinguishing from lncRNAs due to their unique structure. The general characteristics and functions of common ncRNAs are summarized in Table 1.

3 | MICRORNAs

3.1 | Biogenesis of miRNAs

MicroRNAs are endogenous, non-coding small RNAs with approximately 20-24 nt in length which mainly participate in regulating gene expression. In the canonical biogenesis pathway, genes encoding miRNAs are transcribed into long initial transcript by RNA polymerase II, namely pri-miRNAs, 300-1000 base pairs (bp) in length.⁵ The pri-miRNAs are cleaved to pre-miRNAs (about 70 bp in length) with a stem-loop structure under the action of RNase III Drosha.⁶ Next, the pre-miRNAs are transported from the nucleus to the cytoplasm by the Ran-GTP-dependent transporter Exportin 5.⁷ The pre-miRNAs are further processed by Dicer enzyme, a double-stranded RNA-specific RNA endonuclease, cleaving into 20-24 nt double-stranded miRNAs. To act as gene expression regulators, firstly, the mature miRNA binds to Dicer's complementary sequence to form a double helix, then the double helix unwinds, one of which with the lower stability in the 5' end preferentially binds to the RNA-induced silencing complex (RISC) to form a complex including Argonaute (AGO) proteins, the mRNA strand and several cofactors. At last, the complex would combine its target mRNA, causing the inhibition of gene expression (Figure 1).

3.2 | Functions of miRNAs

3.2.1 | MiRNAs cause mRNA degradation

MicroRNAs play significant regulatory roles in eukaryotes by binding to corresponding mRNA transcripts, leading to their degradation at mRNA level and/or translational repression. There is currently no clear conclusion on how miRNAs induce mRNA degradation. Two different perspectives have been issued currently: (a) ordinary mechanism of mRNA degradation. This view holds that miRNAs merely act as guides, they do not directly degrade mRNAs through AGO proteins, but only label and send mRNAs that need to be degraded to the normal mRNA degradation mechanism. Giraldez et al⁸ found that miR-430 induced the deadenylation of its targets, resulting in the recruitment to processing bodies, where mRNAs were decapped and degraded. (b) MiRISC-mediated mRNA degradation. This view holds that a variety of related enzymes and complexes, such as AGO proteins, decapping enzymes and their cofactors, all of which may be involved in miRNA-mediated degradation of mRNAs. As a recognized mechanism, miRNAs bind to AGO proteins in miRISCs to recognize their mRNA targets. AGO proteins interact with a GW182 protein, in turn, interacts with cytoplasmic poly(A)-binding protein (PABP) and with the cytoplasmic deadenylase complexes CCR4-NOT and PAN2-PAN3, which catalyse the deadenylation of the mRNA targets, and then, deadenylated mRNAs are decapped and rapidly degraded by 5' to 3' exoribonuclease 1 (XRN1).⁹⁻¹¹

3.2.2 | MiRNAs repress protein translation

Several studies have proven that miRNA-mediated gene silencing is first manifested through repressing translation and subsequently consolidated by mRNA deadenylation and decay.^{12,13} Interestingly, Beilharz et al¹⁴ observed that miRNA-mediated mRNA deadenylation contributed to translational repression in mammalian cells as well. Multiple perspectives have been issued to explain this mechanism, such as inhibition of ribosome assembly and degradation of neopeptide,¹⁵ but there are still no unified views up to now. The emerging recognition is that miRNAs inhibited translation initiation by interfering with the activity and/or assembly of the eukaryotic initiation factor 4F (eIF4F) complex, consisting of the cap-binding protein eIF4E, the adaptor protein eIF4G and the box protein 6 (DEAD) box RNA helicase eIF4A.¹⁶ Whatever, it is now widely accepted that both mRNA degradation and translational repression participate in miRNA-mediated gene silencing.

3.2.3 | Unconventional functions of miRNAs

Although negative regulation of target gene expression is the most common mechanism of miRNAs participating in biological functions, other non-canonical mechanisms also play irreplaceable roles. It has been proven that miRNAs can interact with non-AGO

proteins. Eiring et al¹⁷ found that miR-328 interacted with hnRNP E2, leading to release of CEBPA mRNA from hnRNP E2-mediated translational inhibition. To our surprise, miRNAs can directly activate

transcription. Matsui et al¹⁸ found that miR-589, in complex with AGO2 and GW182, bound the promoter RNA of COX2, leading to transcriptional activation of COX2. Besides, other non-canonical

TABLE 1 The main types of non-coding RNAs and their features

Types of ncRNAs	Abbreviation	Length (nt)	Localization	Main functions
Short ncRNAs				
MicroRNA	miRNA	20-24	Nucleus, cytoplasm	Translation suppression
Small interfering RNA	siRNA	20-30	Cytoplasm	Translation suppression
Small nucleolar RNA	snoRNA	60-200	Nucleus, cytoplasm	2'-O-methylation and pseudouridylation of rRNA
Small nuclear/cytoplasmic RNA	snRNA/scRNA	100-200	Nucleus, cytoplasm	Component of spliceosome
piwi-interacting RNA	piRNA	24-31	Cytoplasm	Translation suppression, modulation of transposons
tRNA-derived stress-induced RNA	tiRNA	30-40	Cytoplasm	Translation suppression, signalling molecule
tRNA-derived small non-coding RNA	TDRs	20	nucleus	Translation suppression, target transportable element
Long ncRNAs				
Long intergenic non-coding RNA	lincRNA	>200	Nucleus, cytoplasm	miRNA sponge, regulation of gene transcription
Natural antisense transcript	NAT	>200	Nucleus, cytoplasm	inhibition of the mRNA, epigenetic gene silencing
Circular RNA	circRNA	200-800	Nucleus, cytoplasm	miRNA sponge, regulation of gene transcription
Pseudogene transcript	–	>200	Nucleus, cytoplasm	translation repression, miRNA sponge
Transcribed ultraconserved region	T-UCR	>200	Cytoplasm	miRNA sponge
Telomerase RNA component	TERC	451	Nucleus	Telomere length maintenance




FIGURE 1 Mechanism of miRNA biogenesis. The miRNA gene is preliminarily transcribed into primary miRNA (pri-miRNA, 300-1000 bp) in the nucleus. A processing complex consisting of Drosha and DGCR8 cleaves the pri-miRNA, resulting in the generation of precursor miRNA (pre-miRNA, 70 bp). Exportin 5 exports the pre-miRNA from the nucleus to the cytoplasm. In the cytoplasm, the pre-miRNA is further processed by the Dicer complex into double-stranded miRNA. One strand of the miRNA duplex (mature miRNA, 20-24 nt) is selected to assemble the RNA-induced silencing complex (RISC). The miRISC includes the miRNA, corresponding mRNA and Argonaute (AGO) and causes mRNA destabilization and translational inhibition relying on miRNA binding to the complementary sequence in the 3' UTR of its target mRNA

mechanisms were observed as well, including upregulating protein expression, activating Toll-like receptors, targeting mitochondrial transcripts and targeting nuclear ncRNAs.¹⁹

3.3 | MiRNAs in TNBC

MicroRNAs have been exhibited to act as multiple roles in TNBC, including functioning as diagnostic and prognostic biomarkers and exerting multifaceted effects on tumour progression, such as the modulation of tumour growth, metastasis and manipulation of chemoresistance, and regulation of metabolism (Table 2).

3.3.1 | MiRNAs function as diagnostic and prognostic biomarkers

Most functional miRNAs tend to exhibit various expression patterns between TNBC patients and healthy individuals even other subtype breast cancer samples. Furthermore, miRNAs can indicate cancer progression and monitor the prognosis of TNBC patients, which have potential to act as biomarkers in clinical practice.

Circulating cell-free miRNAs extracted from blood are a hotspot in the field of diagnostic biomarkers.²⁰ According to published literature, circulating cell-free miR-199a-5p could function as a TNBC-specific diagnostic biomarker distinguishing TNBC patients from non-TNBC and healthy controls.²¹ Circulating miR-105 and miR-93-3p also acted as a diagnostic biomarker for TNBC.²² Niedźwiecki et al²³ found that the expression level of miR-200c was lower in TNBC patients' serum when compared with the levels in the ER/PR-positive group, revealing that miR-200c is a diagnostic biomarker to distinguish the subtype of breast cancer.

Since the prognosis of TNBC patients is quite poor, prognostic evaluation is essential for both clinicians and patients. Positive miR-34b, miR-301a and miR-361-5p expression was significantly associated with poor overall survival (OS) in TNBC patients.²⁴⁻²⁶ In addition, high expression of miR-9 and miR-155 exhibited a tight association with poor distant metastasis-free survival (DMFS) in TNBC patients, and the level of miR-9 expression also correlated with worse disease-free survival (DFS).²⁷ On the other hand, several miRNAs have been shown to function as tumour suppressors in TNBC. MiR-148a suppressed metastasis of TNBC and served as a prognostic indicator.²⁸ Moreover, stroma-derived miRNAs exhibited promising value in the prognostic evaluation of TNBC. Stromal expression of miR-21 was remarkably upregulated and associated with recurrence-free survival (RFS).²⁹

Furthermore, prognostic models developed based on combinations of multiple miRNA signatures have also been shown excellent values. Two independent miRNA signatures developed by Cascione et al³⁰ had predictive values for OS and DMFS in TNBC patients, respectively. In addition, a prognostic model developed based on the expression levels of miR-155, miR-493, miR-30e and miR-27a accurately identified high-risk and low-risk groups in TNBC patients' cohort.³¹

Besides, Kleivi et al³² performed genome-wide serum miRNA expression detection and identified a four-miRNA signature that predicted RFS and OS; this prediction model may bring better treatment options for patients with TNBC.

3.3.2 | MiRNAs modulate cancer cell growth

Cell proliferation is crucial for cancer progression and is commonly mediated by the dysregulation of cell cycle-related proteins. Tumour growth tends to contribute to the occurrence of tumour angiogenesis and metastasis. Emerging studies have shown that miRNAs regulate cell proliferation in TNBC.

Loss of p53, a classic tumour suppressor, always occurs in tumours. MiR-205 is directly transactivated by p53 and is downregulated in TNBC. Re-expression of miR-205 significantly reduced cell proliferation in vitro and inhibited tumour growth in vivo. The underlying mechanism by which miR-205 inhibited cancer cell growth was that overexpressed miR-205 directly targeted E2F1 to regulate cell cycle progression and LAMC1, thus modulating cell adhesion, proliferation and migration.³³ More targets regulated by miRNAs in TNBC have recently been revealed. Re-expression of the silenced miRNAs miR-17-5p and miR-3178 in TNBC inhibited cell proliferation and migration by targeting ETV1 and NOTCH1, respectively.^{34,35} In addition, miR-185 suppressed tumour proliferation by directly downregulating E2F6 and DNMT1, thus indirectly upregulating BRCA1 in TNBC.³⁶

On the other hand, oncogenic miRNAs also play crucial roles in TNBC. MiR-498, miR-374a-5p and miR-761 were significantly overexpressed in TNBC and promoted cell proliferation and migration by downregulating PTEN, ARRB1 and TRIM29, respectively.³⁷⁻³⁹ Moreover, the miR-106b-25 cluster mediated oncogenesis in breast cancer by activation of NOTCH1 by targeting NEDD4L in both ER+ and TNBC breast cancer cells, suggesting that the miR-106b-25/NEDD4L/NOTCH1 axis played a crucial role in breast cancer.⁴⁰ Cantini et al described a novel algorithm named ClustMMRA, which was applied to explore clustered miRNAs potentially driving cancer molecular subtypes. Using ClustMMRA to analyse breast cancer patient data, the authors identified miR-199/miR-214 as a novel cluster that promoted TNBC cell proliferation and further validated these findings in vitro.⁴¹

3.3.3 | MiRNAs affect migration and metastasis

Although the aetiology and onco-genetic mechanism of TNBC have been initially investigated, there are no defined conclusions to explain metastasis in clinical practice, which causes thousands of cancer-related deaths. Emerging evidence has demonstrated that miRNAs are associated with the metastatic process in TNBC, suggesting novel strategies to control metastasis and improve prognosis.

Emerging evidence suggests that miRNAs are associated with the metastatic process in TNBC. Dysregulated expression

TABLE 2 Summarization of the cellular functions of miRNAs in tumorigenesis of TNBC

miRNAs	Role in TNBC	Cancer phenotype	Target gene	References
miR-498	Oncogene	Promoted proliferation and migration	PTEN	37
miR-374a-5p	Oncogene	Promoted proliferation and migration	ARRB1	38
miR-761	Oncogene	Promoted proliferation migration and invasion	TRIM29	39
miR-106b-25 cluster	Oncogene	Induced tumour initiating cell phenotypes	NEDD4L	40
miR-199	Oncogene	Promoted proliferation and migration	/	41
miR-214	Oncogene	Promoted proliferation and migration	/	41
miR-200c	Oncogene	Promoted migration and invasion	/	46
miR-423-5p	Oncogene	Promoted migration and invasion, increased resistance to cisplatin	/	58
miR-18a	Oncogene	increased resistance to cisplatin	Dicer	60
miR-181a	Oncogene	Promoted migration and invasion, increased resistance to doxorubicin	BAX	62
miR-221	Oncogene	Promoted proliferation, migration and invasion	PTEN	66
miR-939	Oncogene	Increased tumour cell trans-endothelial migration	VE-cadherin	67
miR-205	Tumour suppressor	Inhibited proliferation, cell cycle progression	E2F1, LAMC1	33
miR-17-5p	Tumour suppressor	Inhibited proliferation and migration	ETV1	34
miR-3178	Tumour suppressor	Inhibited proliferation, invasion and migration	NOTCH1	35
miR-185	Tumour suppressor	Inhibited proliferation	E2F6, DNMT1	36
miR-200a	Tumour suppressor	Inhibited migration	EPHA2	44
miR-200b	Tumour suppressor	Inhibited migration and invasion	PKC α , ARHGAP18	45,55
miR-655	Tumour suppressor	Inhibited migration and invasion	PRRX1	48
miR-125b	Tumour suppressor	Inhibited proliferation, migration and invasion	MAP2K7	49
miR-212-5p	Tumour suppressor	Inhibited migration and invasion	PRRX2	50
miR-613	Tumour suppressor	Inhibited migration and invasion	DAAM1	51

(Continues)

TABLE 2 (Continued)

MiRNAs	Role in TNBC	Cancer phenotype	Target gene	References
miR-638	Tumour suppressor	Inhibited proliferation and migration, increased sensitivity to cisplatin	BRCA1	57
miR-101	Tumour suppressor	Inhibited proliferation, induces apoptosis, increased sensitivity to paclitaxel	MCL1	59
miR-130a-3p	Tumour suppressor	Increased sensitivity to doxorubicin	/	61
miR-451a	Tumour suppressor	Increased sensitivity to doxorubicin	/	61
miR-34a	Tumour suppressor	Inhibited proliferation and invasion, increased sensitivity to dasatinib	c-SRC	64

of miR-200 family members has been observed in multiple cancers.^{42,43} However, the roles of miR-200 family members in the metastasis of breast cancer are still disputable. MiR-200a and miR-200b inhibited the migration of TNBC cells by directly targeting EPHA2 and PKC α , respectively.^{44,45} In contrast, overexpression of miR-200c promoted the migration and invasion capacity of TNBC cells by secreting VEGF-A through activation of the FAK and PI3K/AKT signalling pathways.⁴⁶ Epithelial-mesenchymal transition (EMT) is a critical process in cancer cell invasion, characterized by the downregulation of cell adhesion markers with the concomitant acquisition of mesenchymal molecules.⁴⁷ MiRNAs serve as crucial regulators in the EMT process, and miR-655, miR-125b and miR-212-5p have been reported to suppress the EMT process by targeting PRRX1, MAP2K7 and PRRX2 in TNBC, respectively.⁴⁸⁻⁵⁰

RhoA is a classic small GTPase that is generally thought to be essential for the formation of stress fibres and invasion of cancer cells. Xiong et al⁵¹ reported that miR-613 suppressed TNBC cell migration and invasion by targeting DAAM1, a novel indirect regulator of RhoA activation.⁵²⁻⁵⁴ However, to our surprise, Humphries et al⁵⁵ found that downregulation of ARHGAP18 by miR-200b inhibited metastasis of TNBC by increasing the activation of RhoA, suggesting the tumour suppressive role of RhoA in TNBC.

3.3.4 | MiRNAs regulate the sensitivity to therapeutic drugs

Chemotherapy remains an indispensable therapeutic strategy for TNBC, but the acquisition of chemoresistance is the primary obstacle for successful treatment. Accordingly, tremendous effort has been made in recent years to elucidate the mechanisms of TNBC chemoresistance with the aim of identifying new molecular targets.⁵⁶ It has become evident that ncRNAs act as significant regulators in the development of TNBC chemoresistance.

Cisplatin is a common anti-cancer drug applied in clinical practice. Overexpression of miR-638 increased sensitivity to cisplatin by regulating the expression of BRCA1 in TNBC.⁵⁷ In contrast, miRNAs could also promote resistance to chemotherapy. Cisplatin-resistant MDA-MB-231 cell-derived miR-423-5p increased the resistance of recipient cells in an exosomal-dependent manner.⁵⁸ The bidirectional effects of miRNAs on chemoresistance have also been demonstrated for paclitaxel and doxorubicin. MiR-101 increased paclitaxel sensitivity by downregulating MCL1 expression,⁵⁹ while miR-18a conferred paclitaxel resistance by targeting Dicer in TNBC.⁶⁰ Furthermore, overexpression of miR-130a-3p and miR-451a in TNBC cells significantly enhanced the cell sensitivity to doxorubicin,⁶¹ but high expression levels of miR-181a contributed to chemotherapy resistance and predicted poor DFS and OS in patients receiving these treatments.⁶² MiRNAs also regulate the curative effect of targeted therapies. Decreased miR-206 expression was observed in BRCA1 wild-type TNBC cells after concomitant treatment with gemcitabine and PARP1 inhibitor, suggesting that miR-206 served as a curative regulator of PARP1 inhibitor combination chemotherapy for BRCA1 wild-type TNBC patients.⁶³ Expression of miR-34a was silenced in TNBC, and re-expressed miR-34a in cell lines promoted sensitivity to dasatinib and suppressed proliferation and invasion by downregulating c-SRC.⁶⁴

3.3.5 | Exosomes-mediated miRNAs in TNBC

Exosomes are becoming a novel hotspot in the field of cancer research, which are a subclass of extracellular vesicles involved in intercellular communication.⁶⁵ Not surprisingly, exosomes-mediated miRNAs play significant roles in tumorigenesis of TNBC. As described before, cisplatin-resistant MDA-MB-231 cells transmitted resistance by exosomes containing miR-423-5p.⁵⁸ In addition, cancer cell-derived exosomes have been shown to transfer

miR-221 to recipient cells to promote EMT, thereby promoting metastasis.⁶⁶ Modica et al⁶⁷ observed the extracellular pro-tumorigenic role of tumour-derived, exosome-associated miR-939 in targeting VE-cadherin, explaining its association with poor prognosis in TNBC. Moreover, exosomal miRNAs could serve as monitoring indicators for therapeutic response. Stevic et al⁶⁸ found that 17 miRNAs in 224 TNBC patients who underwent neoadjuvant therapy were significantly deregulated, which were significantly associated with the clinic-pathological and risk factors. To sum up, as a novel hotspot of cancer research, exosome-mediated miRNAs may act as multiple roles in TNBC, which should be further explored in the future.

4 | LONG NON-CODING RNAs

4.1 | Biogenesis of lncRNAs

lncRNA is one of ncRNAs subclass that most researched in the field of cancer research. In mammalian genome, more than 58 000 lncRNAs have been identified, and approximately 30 000 lncRNAs have been curated in GENCODE v29.⁶⁹ As we all know, lncRNAs exist widely in human organism, and it is vital to human gene expression modulation and physiological and pathological processes. Most types of lncRNAs are transcribed by RNA polymerase II, capped at

the 5' end, polyadenylated at the 3' end and edited by a series of splicing processes which occurs in the nucleus. Besides, there are other mechanisms participating in the process of lncRNA maturation. For instance, precursor lncRNAs (pre-lncRNAs) are cleaved by RNaseP to achieve mature ends.⁷⁰ lncRNAs are pervasively interspersed in the genome based on their various transcriptional origins, including whole or partial natural antisense transcripts, coding genes, between coding genes, within introns, promoter and enhancer (Figure 2). Due to the complexity and diversity of lncRNAs, the biogenesis and regulation mechanism of different lncRNAs have not been completely summarized. In the future years, by virtue of the advanced techniques, including chromatin isolation by ChIRP-Seq, CRISPR and CLIP,⁷¹⁻⁷³ it is worthy of believing that the mechanism of biogenesis and synthesis of lncRNAs will be further deeply cognized.

4.2 | Functions of lncRNAs

4.2.1 | lncRNAs act as ceRNA

Long ncRNAs have diversity of regulatory functions, which can modulate chromatin remodelling, transcriptional regulation and post-transcriptional processing, translation.^{74,75} Although multiple functions of lncRNAs have been observed, competing endogenous




FIGURE 2 Mechanism of lncRNA biogenesis. lncRNA is transcribed by RNA polymerase II, capped at the 5' end, polyadenylated at the 3' end and edited by a series of splicing events that occur in the nucleus. This figure preliminarily summarizes the diverse biogenesis of lncRNAs based on their transcriptional origin, namely, promoter-associated lncRNAs, enhancer-associated lncRNAs, exonic lncRNAs, intronic lncRNAs, bidirectional lncRNAs, intergenic lncRNAs, antisense lncRNAs and sense lncRNAs

RNA (ceRNA) or miRNA sponge is the most focused function. For instance, LINC01123 was found to be upregulated in non-small-cell lung cancer and associated with poor prognosis. By a series of functional experiment, researchers revealed that LINC01123 might increase c-MYC mRNA expression by sponging miR-199a-5p, which was a direct transcriptional regulator for c-MYC.⁷⁶ Even if the function of lncRNAs as ceRNA has been widely accepted, the mechanism concerning these remain unknown to some extent and require to further deeper investigation.

4.2.2 | LncRNAs modulate transcriptional process

Long ncRNAs can directly participate in mediating transcription by different mechanisms.⁷⁷ R-loops are defined as a novel DNA-RNA hybrid abundant at CpG islands, and it can be formed when the nascent RNA reanneals into the template DNA.⁷⁸ Arab et al⁷⁹ uncovered that lncRNA TARID could form an R-loop at the TCF21 promoter and regulated its transcriptional level. Causally, there is a CG skew within exon 2 of TARID, rendering TARID overlap with CpG islands around the transcription start site of TCF21. By DRIP-qPCR assay, it revealed that R-loops could coincide with the TSS-proximal GC skew. When silencing TCF21 by promoter hypermethylation or losing the TARID promoter, the R-loop structure was not observed in cells. In addition, upregulation of RNase H1 degrading RNA within DNA-RNA hybrids would reduce the level of R-loops at TCF21 and the level of TARID. Moreover, lncRNAs can regulate transcription through binding to histone-modifying complexes, to transcription factors and even to RNA polymerase II as well.⁸⁰

4.2.3 | LncRNAs modulate translation

Distinctly, lncRNAs are defined as a sort of ncRNA without translation function. However, during the process of translation, lncRNAs play indispensable roles. As recently reviewed, it has been found that lncRNA MEG3 significantly decreased in human invasive bladder cancers, and its exogenous expression can inhibit the invasiveness of human bladder cancer cells. The main reason is that MEG3 negatively regulated the expression of c-MYC by promoting PHLPP2 protein translation.⁸¹ In recent years, ribosome profiling has identified ribosome-associated lncRNAs, which accords with the idea that some annotated lncRNAs may be translated, which remains under investigation. Besides, some lncRNAs can be used as a scaffold for assembly of whole protein complexes, for example NEAT1 for paraspeckle proteins and HOTAIR for the HBXIP/HOTAIR/LSD1 complex.^{82,83}

4.2.4 | LncRNAs are involved in chromatin action

There are lots of lncRNAs accumulating in the nucleus to regulate chromatin architecture, participate in chromatin remodelling and

promote the recruitment of chromatin modifiers. For example, the genome multipotent stem cells are equipped with the form of higher-order chromatin architecture, with a variety of intra- and inter-chromosomal interactions. The peculiarity of stem cell pluripotency is directly determined by the promoter region of stem cell core factor genes around the architecture. In many studies, lncRNAs can take part in the composition of chromatin architecture to establish the stemness state, chiefly bringing distant enhancer elements into proximity of the core promoter.⁸⁴ Besides, lncRNAs also can interact with chromatin-modifying enzymes, catalysing covalent changes of histones or DNA on the chromatin to affect genetic expression information, such as GADD45a, DNMT1 and PRC.⁸⁵

4.2.5 | LncRNAs regulate mRNA stability

Studies show that lncRNAs can regulate mRNA stability via corresponding miRNAs, recruiting proteins to degrade mRNA and acting as molecular decoys for RBPs taking part in mRNA decay. For instance, the binding of hnRNP E1 to a nucleic acid structural element located in exon 12 of PNUTS pre-RNA can regulate its alternative splicing. In breast cancer cells, the expression of lncRNA PNUTS was elevated and associated with levels of ZEB mRNAs. Furthermore, PNUTS also served as a competitive sponge for miR-205 during epithelial-mesenchymal transition.⁸⁶ Besides, PDCD4-AS1 regulated breast cancer progression through stabilizing PDCD4 RNA by forming RNA duplex and controlling the interaction between PDCD4 RNA and RNA decay-promoting factors such as HuR.⁸⁷ Although the role of lncRNAs in regulating mRNA stability has been rarely observed, it is a novel mechanism that lncRNA-related biological process needs to be further explored in the future.

4.3 | LncRNAs in TNBC

Long ncRNAs play crucial roles in human beings, not only do lncRNAs participate in regulating normal mammalian gene expression function or other body biological processes, but also they have notable influences on human diseases, such as various cancers, neuropsychiatric disorders and atherosclerosis.⁸⁸⁻⁹⁰ Apparently, there are gradually increasing explorations in the mechanism of lncRNAs functioning in TNBC in recent years. Accumulated evidence suggested that lncRNAs can modulate proliferation, invasion, differentiation, chemoresistance of TNBC cells in positive or negative manner (Table 3). Even a few studies indicated that lncRNAs can work as a biomarker for diagnosing and evaluating the prognosis in TNBC.

4.3.1 | LncRNAs regulate cancer cell growth

Long ncRNA CCAT1 expression is higher in TNBC cells than in normal breast epithelial cells. Functional analysis indicated that

TABLE 3 Summarization of the cellular functions of lncRNAs in tumorigenesis of TNBC

lncRNAs	Role in TNBC	Cancer phenotype	Molecular mechanism	References
CCAT1	Oncogene	Promoted proliferation and migration	Regulated miR-218/ZFX axis	91
TROJAN	Oncogene	Promoted proliferation	Induced ZMYND8 degradation	92
LINC00339	Oncogene	Promoted proliferation and inhibited apoptosis	Regulated miR-377-3p/HOXC6 axis	93
MIR100HG	Oncogene	Promoted proliferation and induced cell arrest in the G1 phase	Negatively regulated p27 gene expression	94
NRAD1	Oncogene	Promoted proliferation	Positively regulated by ALDH1A3	95
DANCR	Oncogene	Promoted proliferation	Bound to RXRA and enhanced PI3K/AKT signals	96
NAMPT-AS	Oncogene	Promoted migration	Recruited POU2F2 to activate NAMPT Regulated miR-548b-3p/NAMPT axis	10
ROR	Oncogene	Promoted migration	Regulated miR-145/ARF6 axis	11
Linc-ZNF469-3	Oncogene	Promoted migration	Regulated miR-574-5p/ZEB1 axis	12
HULC	Oncogene	Promoted migration	Upregulated MMP2 and MMP9	13
NEAT1	Oncogene	Inhibited apoptosis, regulated cell cycle progression and promoted chemoresistance	Upregulated ATP7A and ATP7B	16
BORG	Oncogene	Increased resistance to doxorubicin	Activated RPA1 and NF- κ B signals	17
HCP5	Oncogene	Increased resistance to cisplatin	Downregulated PTEN	18
SONE	Tumour suppressor	Inhibited proliferation and migration	Positively regulated TP53 and negatively regulated c-MYC	97
RMST	Tumour suppressor	Inhibited proliferation and migration, induced apoptosis	/	98
PTCSC3	Tumour suppressor	Inhibited proliferation	Downregulated lncRNA H19	99
LOC554202	Tumour suppressor	Inhibited migration	Promoter hypermethylated	14
NEF	Tumour suppressor	Inhibited migration	Negatively regulated miR-155	15
GAS5	Tumour suppressor	Increased sensitivity to paclitaxel, induced apoptosis	Regulated miR-378a-5p/SUFU axis	19

silencing CCAT1 inhibited cell proliferation and migration in TNBC by regulating the miR-218/ZFX axis.⁹¹ Similarly, lncRNA TROJAN promoted the proliferation of TNBC cells by inducing ZMYND8 degradation.⁹² LINC00339 regulated TNBC cell growth by promoting proliferation and inhibiting apoptosis *via* regulation of miR-377-3p/HOXC6 expression.⁹³ Moreover, there are also some non-classical lncRNA-involved mechanisms in the regulation of TNBC progression. MIR100HG was identified as a pro-oncogene for TNBC progression with a high expression level in TNBC and reduced MIR100HG significantly inhibited cell proliferation and induced cell cycle arrest in the G1 phase. Furthermore, MIR100HG negatively regulated p27 gene expression to control the cell cycle by forming RNA-DNA triplex structures, impacting the progression of TNBC.⁹⁴ lncRNA NRAD1 was regulated by ALDH1A3 and was a therapeutic target for TNBC for its regulation of gene expression and effect on cancer cell survival.⁹⁵ lncRNA DANCR enhanced PI3K/AKT signals and TNBC proliferation by binding to RXRA and increasing its serine 49/78 phosphorylation to activate PIK3CA transcription.⁹⁶

In contrast, several lncRNAs also play tumour suppressor roles in TNBC progression. Downregulating lncRNA SONE resulted in a

remarkable TP53 decrease and c-MYC increase, which could alter the expression of downstream tumour suppressor miRNAs, leading to increased cell proliferation and migration.⁹⁷ lncRNA RMST functioned as a tumour suppressor in TNBC by decreasing cell proliferation and migration, modulating the cell cycle and enhancing apoptosis.⁹⁸ Moreover, lncRNA PTCSC3 inhibited TNBC cell proliferation by downregulating lncRNA H19, exhibiting a novel RNA-RNA interacting mechanism in TNBC.⁹⁹

4.3.2 | lncRNAs mediate migration and metastasis

As long non-coding antisense transcript of NAMPT, NAMPT-AS was upregulated in TNBC and promoted cell migration. NAMPT-AS epigenetically regulated the expression of NAMPT in two different ways. One was that the transcription of NAMPT was activated by NAMPT-AS, thereby recruiting POU2F2. The other was that NAMPT-AS acted as ceRNA to rescue NAMPT degradation from miR-548b-3p.¹⁰ A previous report indicated that a novel miRNA sponge, lincRNA ROR, was dramatically upregulated in TNBC and interacted with miR-145 to regulate cancer cell invasion by targeting

ARF6.¹¹ Wang et al¹² found that linc-ZNF469-3 interacted with miR-574-5p and overexpression of linc-ZNF469-3 upregulated ZEB1 expression, which stimulated lung metastasis of TNBC. Furthermore, proteins of the MMP family have been shown to be involved in the breakdown of the extracellular matrix, which is essential for cancer metastasis. Shi et al¹³ demonstrated that lincRNA HULC expression was increased in TNBC tissues and silencing HULC expression effectively suppressed cell metastasis through suppressed MMP2 and MMP9 expression.

In addition, it has been proven that lincRNAs also function as cell migration inhibitors in various ways. lincRNA LOC554202 was downregulated in TNBC by promoter hypermethylation, and LOC554202 loss promoted cell migration, accounting for the aggressive phenotype of TNBC to some extent.¹⁴ In addition, lincRNA NEF was found to be downregulated in TNBC, and NEF overexpression inhibited cell migration by negatively regulating miR-155.¹⁵

4.3.3 | LncRNAs dominate the sensitivity to chemotherapy

Studies have shown that lincRNA NEAT1 expression is upregulated in cisplatin- and taxol-resistant cells compared with parental cells.

By qRT-PCR assay, it was demonstrated that when knocking down NEAT1 in sh-NEAT1 cells, drug transporter genes, such as ATP7A and ATP7B, were downregulated, and functional analysis indicated that NEAT1 knockdown sensitized cells to chemotherapy through a synergistic effect.¹⁶ BORG, an oncogenic lincRNA, was greatly responsive to cytotoxic drug treatment, particularly doxorubicin. The mechanism of drug BORG-associated resistance relied on its significant activation of the NF- κ B signalling pathway through the BORG-mediated feed-forward signalling loop and its ability to activate RPA1, making BORG-expressing TNBC sensitive to doxorubicin-induced cytotoxicity.¹⁷ In addition, upregulation of lincRNA HCP5 contributed to cisplatin resistance in TNBC, and inhibition of HCP5 reversed resistance to cisplatin by upregulating PTEN expression.¹⁸ However, tumour suppressive lincRNA GAS5 could enhance the sensitivity of TNBC to paclitaxel and induce apoptosis in TNBC cells by targeting miR-378a-5p/SUFU signalling.¹⁹

Moreover, lincRNAs not only participate in the process of chemoresistance, but can also be used for predicting the response to neoadjuvant chemotherapy in TNBC. Zheng et al¹⁰ systematically compared gene expression between TNBC patients with pathological complete response and those without a complete response to neoadjuvant chemotherapy and ultimately developed a gene signature of 2 coding genes and 3 lincRNAs to predict the response to neoadjuvant chemotherapy of TNBC patients.




FIGURE 3 Mechanism of exonic circRNA and ElciRNA biogenesis. A, Intron-pairing-driven circularization: Complementary sequence motifs in introns flanking the exons can directly pair each other to induce circularization, and as a result, exonic circRNAs or ElciRNAs are produced; B, RBP-dependent circularization: Interaction between RBPs bound to sequence motifs in both introns flanking the exons to be circularized facilitates the head-to-tail end-joining of exon 2 and exon 3; C, Lariat-driven circularization: Folding of a region of pre-RNA can result in exon skipping; furthermore, the splice donor in 3' end of exon 1 and the splice acceptor in 5' end of exon 4 are covalently joined together to form a lariat containing exon 2 and exon 3, ultimately producing a circRNA. SF: splicing factor; RBP: RNA-binding protein




FIGURE 4 Mechanism of intronic circRNA biogenesis. A, Group II intron-mediated circRNA formation: Circle formation requires prior release of the 3' end exon. The terminal 2'-OH group of the intron attacks the 5'-splice site, creating a circular RNA by 2',5'-phosphodiester formation. B, Group I intron-supported regular splicing: An exogenous guanosine (exoG) bound to the intron structure serves as a nucleophile attacking the 5'-splice site. (a) Upon first transesterification, the 5'-exon is cut-off and exoG becomes linked to the intron. (b) The terminal 3'-OH group of the 5'-exon then attacks the 3'-splice site, and the ligated exons and a linear intron are released. Eventually, the linear intron is circularized by nucleophilic attack of the 2'-OH group of the terminal guanosine (ω G) onto a phosphodiester bond close to the 3'-end and release of a short 3'-tail. Note that in this case, a 2'-5'-phosphodiester bridge closes the circle. C, Circular intron RNA: Hydrolysis of the 3'-exon allows circle formation by direct nucleophilic attack of ω G onto the 5'-splice site, producing an RNA lariat circularized with a 2',5'-phosphodiester

4.3.4 | LncRNAs serve as potential biomarkers for diagnosis and prognosis

At present, with the deepening of TNBC research, lncRNAs have generally been revealed as promising diagnostic and prognostic biomarkers.¹¹¹ ANRIL, HIF1A-AS2 and UCA1 have been reported to be markedly upregulated in the plasma of patients with TNBC compared with patients with non-TNBC. In addition, a 3-lncRNA signature obtained using the logistic model exhibited excellent diagnostic values with an AUC of 0.934.¹¹² In addition, an epigenome-wide association study (EWAS) conducted by Bermejo et al¹¹³ revealed that hypermethylation of LINC00299 in the peripheral blood of TNBC patients could function as a useful circulating biomarker for TNBC and exhibited excellent diagnostic value. Jiang et al¹¹⁴ proposed an integrated mRNA-lncRNA signature based on a combination of mRNA and lncRNA species and found that lncRNA HIF1A-AS2 and AK124454 could be involved in mediating TNBC cell proliferation, invasion and paclitaxel resistance and also exhibited good prognostic value.

Furthermore, Lv et al¹¹⁵ compared the expression levels of lncRNAs in TNBC and non-TNBC tissues separately and found that

dysregulated lncRNAs participated in important biological processes. They further validated these lncRNA expression levels and confirmed that four dysregulated lncRNAs were significantly correlated with TNBC occurrence. Other systemic analyses revealed that seven prognosis-related lncRNAs were significantly associated with poor RFS in TNBC patients.¹¹⁶

4.3.5 | LncRNAs mediate immunomodulation

Immunotherapy, including PD-1 and/or PD-L1 blockade, is an important cancer therapeutic method to restrict cancer progression.¹¹⁷ However, during immunotherapy, the loss of antigenicity and evasion of immune checkpoints in malignant tumour cells is a puzzling issue that deserves further exploration. Hu et al¹¹⁸ reported that tissue-specific lncRNA LINK-A expression facilitated the crosstalk between GPCR signalling and upregulated K48-polyubiquitination-mediated degradation of the antigen PLC and intrinsic tumour suppressors Rb and p53. Furthermore, the treatment with LINK-A locked nucleic acids or GPCR antagonists stabilized the PLC components Rb and p53 and sensitized breast cancer

TABLE 4 Summarization of the cellular functions of circRNAs in tumorigenesis of TNBC

CircRNAs	Circbase ID	Role in TNBC	Cancer phenotype	Sponge miRNAs	Target genes	References
circGFRA1	hsa_circ_005239	Oncogene	Promoted proliferation and induced apoptosis	miR-34a	GFRA1	134
circEPST11	hsa_circ_000479	Oncogene	Promoted proliferation and induced apoptosis	miR-4753 miR-6809	BCL11A	135
circUBAP2	hsa_circ_0001846	Oncogene	Promoted proliferation and migration and induced apoptosis	miR-661	MTA1	136
circAGFG1	hsa_circ_0058514	Oncogene	Promoted proliferation, migration and invasion	miR-195-5p	CCNE1	137
circKIF4A	hsa_circ_0007255	Oncogene	Promoted proliferation and migration	miR-375	KIF4A	138
circPLK1	hsa_circ_0038632	Oncogene	Promoted proliferation and migration	miR-296-5p	PLK1	139
circRAD18	hsa_circ_0002453	Oncogene	Promoted proliferation and migration and induced apoptosis	miR-208a/miR-3164	IGF1/FGF2	140
circRNA_069718	hsa_circ_069718	Oncogene	Promoted proliferation and invasion	/	/	141
circANKS1B	hsa_circ_0007294	Oncogene	Promoted migration and invasion	miR-148a-3p miR-152-3p	USF1	142
ciRS-7	/	Oncogene	Promoted migration and invasion	miR-1299	MMPs	143
circITCH	/	Tumour suppressor	Inhibited proliferation, migration and invasion	miR-214/miR-17	ITCH1	144
circTADA2A-E6	hsa_circ_0006220	Tumour suppressor	Inhibited proliferation, migration and invasion	miR-203a-3p	SOC33	145
circFBXW7	hsa_circ_0001451	Tumour suppressor	Inhibited proliferation, migration and invasion	miR-197-3p	FBXW7	146

cells to immune checkpoint blockers. In the clinical practice, TNBC patients with PD-1 blockade resistance exhibited upregulated LINK-A expression and downregulated PLC components. Although no additional literature has focused on this hotspot, lncRNA-dependent immunomodulation is next promising research direction.

5 | CIRCULAR RNAs

5.1 | Biogenesis of circRNAs

Circular RNA was first discovered in the 1970s; the first observation of circRNA in plant-infected *Viroids* by electron microscopy was reported by Sanger et al in 1976.¹¹⁹ So far, scholars have found that more than 10% of protein-encoding genes in a variety of biological cells and tissues can produce circRNAs,¹²⁰ suggesting that circRNAs are ancient molecules that with evolutionary conservation. CircRNAs are derived from precursor mRNAs (pre-mRNAs), which are transcribed by RNA polymerase II. CircRNAs can be divided into three subclasses according to their various position and forming mechanism, exonic circRNA, intronic circRNA and lncRNA.¹²¹ Although increasing studies concentrated on the cycling processes, the exact mechanisms of circRNAs maturation have not been fully elucidated. At present, three hypothetical models explain the formation of exonic circRNA and/or lncRNA: intron-pairing-driven circularization, RNA-binding protein (RBP)-dependent circularization and lariat-driven circularization (Figure 3).¹²² There are also three hypothetical models expounding the formation of intronic circRNA have been issued: Group II intron-mediated circRNA formation, group I intron-supported regular splicing and circular intron RNA (ciRNA; Figure 4).^{123,124}

5.2 | Functions of circRNAs

5.2.1 | CircRNAs act as ceRNA or miRNA sponges

Similar to lncRNA, the most common mechanism of circRNA regulating biological process is the ceRNA model. An increasing number of studies have proven that multiple circRNAs act as miRNA sponges. For example, circTP63 functioned as a ceRNA to upregulate FOXM1, thus promoting lung cancer progression.¹²⁵ CircHIPK3 mediated autophagy via sponging miR-124-3p to regulate STAT3/PRKAA/AMPK α signalling in STK11-mutant lung cancer.¹²⁶ CircMYLK promoted hepatocellular carcinoma progression by upregulating Rab23 via sponging miR-362-3p.¹²⁷ To sum up, these studies above support the idea that circRNAs functioning as miRNA sponges may be a common mechanism in cancerous diseases.

5.2.2 | CircRNAs regulate gene transcription

Although most research focuses on the role of circRNAs as sponges for miRNAs, some scholars have uncovered that some




FIGURE 5 Summary of the function of ncRNAs in TNBC. NcRNAs play multifaceted roles in tumour initiation and development, which can control cell proliferation, apoptosis, migration and therapy resistance by orchestrating their downstream targets. Obviously, one ncRNA tends to affect tumour progression by regulating multiple attributes of TNBC cells

intronic circRNAs and lncRNAs could regulate protein level by regulating gene expression at transcriptional or post-transcriptional level. Logically, these circRNAs always exist in the nucleus. For example, Li et al¹²⁸ found that circEIF3J and circPAIP2 up-regulated the expression of their parental genes in *cis* and raised a novel regulatory strategy for transcriptional control *via* specific RNA-RNA interaction between U1 snRNP and lncRNAs. CiRNAs can also regulate gene transcription. Besides, intronic circRNAs also participate in gene regulation. Scholars uncovered that ci-ANKRD52 and ci-SIRT7 can act as positive regulators to up-regulate their parental gene transcription *via* interacting with RNA polymerase II.¹²³

5.2.3 | CircRNAs interact with functional proteins

Similar to linear RNAs, which have been reported to interact with proteins, several studies have also exhibited that some circRNAs can work as protein partner such as circANRIL and circFOXO3. CircANRIL had been revealed to directly bind to PES1, an important 60S-preribosomal assembly factor and next control maturation of ribosomal RNA and modulating pathways of atherosclerosis.¹²⁹ Besides, circFOXO3 was found to be highly expressed in non-cancer cells and correlated with cell cycle progression. Functional analysis

indicated that overexpression of circFOXO3 can repress cell cycle progression *via* binding to CDK2 and cyclin-dependent kinase inhibitor 1 (or p21), resulting in the formation of a ternary complex.¹³⁰

5.2.4 | CircRNAs can be translated into proteins

Most circRNAs are derived from exons and predominantly present in the cytoplasm, suggesting that they can be loaded into ribosomes for translation into polypeptides. CircZNF609 was derived from the circularization of the second exon of its host gene. A 753 nt ORF was present in circZNF609, which could be translated into a protein in a splicing-dependent and cap-independent manner, providing a novel example of a protein-coding circRNA in eukaryotes.¹³¹ Ivano et al¹³² revealed that circFBXW7 was highly expressed in the normal human brain tissues, which can be translated into novel 21 kDa protein termed as FBXW7-185aa. Besides, upregulation of FBXW7-185aa inhibited proliferation and cell cycle acceleration in tumour cells. In addition, circSHPRH can be translated into 17 kDa named SHPRH-146aa. Both circSHPRH and SHPRH-146aa were abundantly expressed in normal human brains and downregulated in glioblastoma tissues. The re-expression of SHPRH-146aa in glioblastoma cells suppressed their malignant behaviour and tumorigenicity *in vitro* and *in vivo*.¹³³ As can be seen from the above examples, the definition of circRNA as ncRNA is somewhat limited.

5.3 | CircRNAs in TNBC

In recent years, circRNAs have gradually become a novel hotspot in the ncRNAs and cancer research field. However, the functions of circRNAs in cancerous diseases, especially in TNBC, have not been fully understood. The known studies about circRNAs mostly focused on the mechanisms of ceRNA in TNBC progression. Besides, circRNAs act as biomarkers for diagnosis and prognosis in TNBC is also widely observed (Table 4).

5.3.1 | CircRNAs act as tumour promoters

Derived from gene GFRA1, circGFRA1 was upregulated in TNBC and high expression of circGFRA1 was correlated with poor OS. Knockdown of circGFRA1 suppressed proliferation and promoted apoptosis via binding to miR-34a and upregulating GFRA1 expression in TNBC.¹³⁴ Besides, silencing of circEPST11 inhibited cell proliferation and induced apoptosis via sponging miR-4753 and miR-6809 to increase BCL11A in TNBC.¹³⁵ It was also revealed that circUBAP2 was correlated with tumour size, advanced TNM stage and worse prognosis in TNBC and promoted tumour progression by sponging miR-661 to upregulate MTA1.¹³⁶ Furthermore, circAGFG1, circKIF4A, circPLK1 and circRAD18 were revealed to promote malignant progression of TNBC by sponging corresponding tumour suppressive miRNAs as well.¹³⁷⁻¹⁴⁰ In addition, Zhang et al¹⁴¹ found that circRNA_069718 can promote the proliferation and invasion of cancer cells by activating the Wnt/ β -catenin pathway, thereby promoting the malignant progression of TNBC.

On the other hand, there is emerging evidence that circRNAs are associated with the metastasis of TNBC. Zeng et al¹⁴² found that circANKS1B was remarkably upregulated in TNBC tissues and increased circANKS1B expression was correlated with lymph node metastasis. Functional studies uncovered that circANKS1B promoted breast cancer cell migration both in vitro and in vivo, whereas it had no effect on breast cancer growth via sponging miR-148a-3p and miR-152-3p to upregulate USF1, leading to transcriptional activation of TGF- β 1, which could upregulate TGF- β 1/Smad signalling to promote EMT. In addition, Sang et al¹⁴³ found that circRNA ciRS-7 was highly expressed in TNBC tissues and upregulated the expression of multiple MMPs by sponging miR-1299, promoting tumour invasion and metastasis in vitro and in vivo.

5.3.2 | CircRNAs act as tumour suppressors

Several studies have found that circRNAs function as tumour suppressors in TNBC as well. CircITCH was remarkably downregulated in TNBC tissues and predicted poor prognosis. Overexpression of circITCH significantly suppressed cell proliferation and migration via acting as a sponge for miR-214 and miR-17 to upregulate ITCH1, thus inactivating Wnt/ β -catenin signalling.¹⁴⁴ Based on screening circRNA profiles, Xu et al¹⁴⁵ identified circTADA2A-E6

spliced from exon 6 of TADA2A gene as a prognostic biomarker in TNBC and overexpression of circTADA2A-E6 significantly inhibited cell proliferation, migration and clonogenicity via regulating miR-203a-3p/SOCS3 axis. Interestingly, circRNA-encoded proteins were also involved in TNBC progression. CircFBXW7 inhibited the malignant progression of TNBC by sponging miR-197-3p and encoding a tumour suppressor FBXW7-185aa.¹⁴⁶ To sum up, limited numbers of studies on the tumour suppressor roles of circRNAs in TNBC are available currently, so it is still urgent to explore further mechanism of circRNAs participating in the suppression of TNBC.

6 | OTHER NON-CODING RNAs

Although most research related to TNBC are concentrated on the field of miRNAs, lncRNAs and circRNAs, other ncRNAs also have great significant functions in the carcinogenesis of TNBC. tDR is one of the other concentrated ncRNAs in TNBC. tRNA has been for a long time deemed to transcripts with non-coding capacity, but with well-established functions in the translation process. However, due to the discovery of tRNA fragments with the function that can regulate gene expression, their crystallized roles have changed over the last decade. Recently, tDRs have been detected in several human diseases and biological processes, including TNBC.^{147,148} TDR-000620 was found to downregulated in TNBC stem cells by RNA sequencing and validation of qPCR. Besides, low tDR-000620 expression served as an independent predictive factor for RFS of TNBC patients.¹⁴⁹ Besides, the expression of tDRs from specific tRNA loci has been found to be associated with the observed race disparities in TNBC, such as the nuclear tRNA^{Gly} and tRNA^{Leu}, the mitochondrial tRNA^{Val} and tRNA^{Pro}.¹⁵⁰ In addition to serving as biomarkers, tDRs also participate in the process of TNBC. Cui et al¹⁵¹ found that tDR-0009 and tDR-7336 were notably upregulated in the SUM-1315 cell lines stimulated by hypoxia; further, bioinformatic analysis indicated that these two upregulated tDRs might be involved in the chemoresistance to doxorubicin in TNBC via mediating the activation of phosphorylation of STAT3. As for other ncRNAs, such as snoRNA and snRNA, their tumour-associated functions have been observed in multiple cancers,¹⁵²⁻¹⁵⁴ but no exact roles have been defined in TNBC, which need to be further explored in the future.

7 | PERSPECTIVES IN CLINICAL PRACTICE

7.1 | NcRNAs as diagnostic and prognostic biomarkers in cancer assessment

Non-coding RNAs were demonstrated to have strong diagnostic and prognostic values in multiple cancers, which were revealed by pan-cancer analysis.^{155,156} With the development of RNA-sequencing technology and advanced analysis methods, the roles of ncRNAs have been well summarized. Guo et al¹⁵⁷ performed small RNA profiling of

26 TNBC cell lines and compared the abundance of ncRNAs among the transcriptional subtypes of TNBC, which identified a mass of dysregulated small ncRNAs, highlighting potential biomarkers for future studies. Although there are still quite a few challenges that need to be addressed, the roles of ncRNAs in clinical practice are being a novel hotspot and many scholars try to employ ncRNAs as biomarkers and therapeutic targets to diagnose, treat and monitor TNBC. As mentioned before, lots of ncRNAs are dysregulated in TNBC tissues compared with non-TNBC and healthy samples, which might serve as potential biomarkers to diagnose TNBC. However, more value will be aggrandized if dysregulated ncRNAs can be detected in the peripheral blood. Circulating miR-199a-5p, miR-105, miR-93-3p and miR-200c had been reported to serve as TNBC-specific diagnostic biomarkers.²¹⁻²³ Besides, the dysregulated lncRNA ANRIL, HIF1A-AS2 and UCA1 can be detected in plasma of TNBC patients and the 3-lncRNAs combined signature exhibited excellent diagnostic value.¹¹² Moreover, even hypermethylation of LINC00299 in peripheral blood of TNBC patients also served as a useful circulating biomarker for TNBC.¹¹³ However, the circulating circRNAs have not been found to be biomarkers in TNBC. Theoretically, circRNAs are covalently closed and generally resistant to the degradation of ribonucleases, suggesting they might be more stabilized in peripheral blood. Thus, detection of the dysregulated circRNAs in the peripheral blood of TNBC patients should next hotspot in the field of ncRNA research.

7.2 | NcRNAs as therapeutic tool in cancer treatment

It has been proven that abnormal small ncRNA expression levels in tumour cells can affect tumour development and therefore can be used as effective targeted drugs for the treatment of tumours.¹⁵⁸ Although small ncRNAs have been shown to be promising and effective therapeutic drugs in vitro, due to the degradation of nucleases in the body, naked ncRNAs have a short half-life in the blood, and the low bioavailability of these nucleic acid drugs in vivo is a major challenge. Therefore, ncRNAs need to be transported to the target tissue by a suitable carrier to exert their effect. Various small ncRNA carriers or systems have been proposed and widely explored, including nanoparticles, ncRNA modification and oncolytic adenovirus strategies.¹⁵⁹ Nanoparticle-based small ncRNA carriers are the most common strategies. Shu et al¹⁶⁰ developed a 15 nm nanoparticle with a 58 nt phi29 pRNA-three-way junction (3WJ), an 8 nt sequence complementary to the seed region of miR-21 and a 39 nt EGFR targeting aptamer. They successfully applied this novel RNA nanotechnology for efficient delivery of anti-miR-21 to block the growth of TNBC in orthotopic mouse models. Moreover, Yin et al¹⁶¹ utilized the thermodynamically and chemically stable 3WJ motif as a scaffold to carry an RNA aptamer binding to CD133 and a locked nucleic acid sequence for miR-21 inhibition to develop a unique delivery strategy to control TNBC progression. Although ncRNA modification and oncolytic adenovirus strategies are also effective technology to enhance the precision and durability of small ncRNAs

in targeting functional genes in cancer, the relevant research on their utility in TNBC is not currently available. Thus, continued progress in the development of ncRNA carrier strategies might allow these approaches to be important and powerful alternative tools to treat TNBC.

8 | CONCLUSION

To conclude, ncRNAs have notable effects on TNBC progression. We summarize the roles of ncRNAs and their mechanisms in TNBC in Figure 5. Although the functions of miRNAs and lncRNAs on the carcinogenesis and development of TNBC have been widely studied and have well-established roles, many other ncRNAs including circRNAs also have a significant influence on tumour progression but have long been neglected. With more research interests concentrated on the lesser-known ncRNAs and their relationship with TNBC, we believe that the mystery of the ncRNA world will eventually be solved and the clinical practice of TNBC management will be largely improved in the future.

ACKNOWLEDGEMENTS

This work was supported by grants from the Natural Science Foundation of Jiangsu Province of China (BE2017626, BK20181367) and the Foundation of Wuxi Health Commission (QNRC003).

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

AUTHOR CONTRIBUTIONS

YZ, CL and JM conceptualized the review. JM and LH wrote the manuscript. LH, HW, RX and YL prepared the figures and tables. YZ and CL critically reviewed and edited the manuscript. All authors read and approved the final manuscript.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Jie Mei  <https://orcid.org/0000-0001-7799-8129>

Chaoying Liu  <https://orcid.org/0000-0003-0484-9522>

REFERENCES

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin.* 2019;69:7-34.
2. Onitilo AA, Engel JM, Greenlee RT, Mukesh BN. Breast cancer subtypes based on ER/PR and Her2 expression: comparison of clinicopathologic features and survival. *Clin Med Res.* 2009;7:4-13.
3. Braicu C, Zimta AA, Harangus A et al The function of non-coding RNAs in lung cancer tumorigenesis. *Cancers (Basel).* 2019;11:605.
4. Ling H, Girnita L, Buda O, Calin GA. Non-coding RNAs: the cancer genome dark matter that matters!. *Clin Chem Lab Med.* 2017;55:705-714.

5. Lee Y, Kim M, Han J et al MicroRNA genes are transcribed by RNA polymerase II. *EMBO J*. 2004;23:4051-4060.
6. Lee Y, Ahn C, Han J et al The nuclear RNase III Drosha initiates microRNA processing. *Nature*. 2003;425:415-419.
7. Yi R, Qin Y, Macara IG, Cullen BR. Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev*. 2003;17:3011-3016.
8. Giraldez AJ, Mishima Y, Rihel J et al Zebrafish MiR-430 promotes deadenylation and clearance of maternal mRNAs. *Science*. 2006;312:75-79.
9. Wahle E, Winkler GS. RNA decay machines: deadenylation by the Ccr4-not and Pan2-Pan3 complexes. *Biochim Biophys Acta*. 2013;1829:561-570.
10. Huntzinger E, Kuzuoglu-Ozturk D, Braun JE, Eulalio A, Wohlbald L, Izaurralde E. The interactions of GW182 proteins with PABP and deadenylases are required for both translational repression and degradation of miRNA targets. *Nucleic Acids Res*. 2013;41:978-994.
11. Nagarajan VK, Jones CI, Newbury SF, Green PJ. XRN 5'→3' exoribonucleases: structure, mechanisms and functions. *Biochim Biophys Acta*. 2013;1829:590-603.
12. Mathonnet G, Fabian MR, Svitkin YV et al MicroRNA inhibition of translation initiation in vitro by targeting the cap-binding complex eIF4F. *Science*. 2007;317:1764-1767.
13. Djuranovic S, Nahvi A, Green R. miRNA-mediated gene silencing by translational repression followed by mRNA deadenylation and decay. *Science*. 2012;336:237-240.
14. Beilharz TH, Humphreys DT, Clancy JL et al microRNA-mediated messenger RNA deadenylation contributes to translational repression in mammalian cells. *PLoS ONE*. 2009;4:e6783.
15. Chendrimada TP, Finn KJ, Ji X et al MicroRNA silencing through RISC recruitment of eIF6. *Nature*. 2007;447:823-828.
16. Jonas S, Izaurralde E. Towards a molecular understanding of microRNA-mediated gene silencing. *Nat Rev Genet*. 2015;16:421-433.
17. Eiring AM, Harb JG, Neviani P et al miR-328 functions as an RNA decoy to modulate hnRNP E2 regulation of mRNA translation in leukemic blasts. *Cell*. 2010;140:652-665.
18. Matsui M, Chu Y, Zhang H et al Promoter RNA links transcriptional regulation of inflammatory pathway genes. *Nucleic Acids Res*. 2013;41:10086-10109.
19. Dragomir MP, Knutsen E, Calin GA. SnapShot: unconventional miRNA functions. *Cell*. 2018;174:1038-1038 e1031.
20. Armand-Labit V, Pradines A. Circulating cell-free microRNAs as clinical cancer biomarkers. *Biomol Concepts*. 2017;8:61-81.
21. Shin VY, Siu JM, Cheuk I, Ng EK, Kwong A. Circulating cell-free miRNAs as biomarker for triple-negative breast cancer. *Br J Cancer*. 2015;112:1751-1759.
22. Li HY, Liang JL, Kuo YL et al miR-105/93-3p promotes chemoresistance and circulating miR-105/93-3p acts as a diagnostic biomarker for triple negative breast cancer. *Breast Cancer Res*. 2017;19:133.
23. Niedzwiecki S, Piekarski J, Szymanska B, Pawlowska Z, Jeziorski A. Serum levels of circulating miRNA-21, miRNA-10b and miRNA-200c in triple-negative breast cancer patients. *Ginekol Pol*. 2018;89:415-420.
24. Svoboda M, Sana J, Redova M et al MiR-34b is associated with clinical outcome in triple-negative breast cancer patients. *Diagn Pathol*. 2012;7:31.
25. Yu H, Li H, Qian H et al Upregulation of miR-301a correlates with poor prognosis in triple-negative breast cancer. *Med Oncol*. 2014;31:283.
26. Cao ZG, Huang YN, Yao L et al Positive expression of miR-361-5p indicates better prognosis for breast cancer patients. *J Thorac Dis*. 2016;8:1772-1779.
27. Jang MH, Kim HJ, Gwak JM, Chung YR, Park SY. Prognostic value of microRNA-9 and microRNA-155 expression in triple-negative breast cancer. *Hum Pathol*. 2017;68:69-78.
28. Xu X, Zhang Y, Jasper J et al MiR-148a functions to suppress metastasis and serves as a prognostic indicator in triple-negative breast cancer. *Oncotarget*. 2016;7:20381-20394.
29. MacKenzie TA, Schwartz GN, Calderone HM et al Stromal expression of miR-21 identifies high-risk group in triple-negative breast cancer. *Am J Pathol*. 2014;184:3217-3225.
30. Cascione L, Gasparini P, Lovat F et al Integrated microRNA and mRNA signatures associated with survival in triple negative breast cancer. *PLoS ONE*. 2013;8:e55910.
31. Gasparini P, Cascione L, Fassin M et al microRNA expression profiling identifies a four microRNA signature as a novel diagnostic and prognostic biomarker in triple negative breast cancers. *Oncotarget*. 2014;5:1174-1184.
32. Kleivi Sahlberg K, Bottai G, Naume B et al A serum microRNA signature predicts tumor relapse and survival in triple-negative breast cancer patients. *Clin Cancer Res*. 2015;21:1207-1214.
33. Piovan C, Palmieri D, Di Leva G et al Oncosuppressive role of p53-induced miR-205 in triple negative breast cancer. *Mol Oncol*. 2012;6:458-472.
34. Li J, Lai Y, Ma J et al miR-17-5p suppresses cell proliferation and invasion by targeting ETV1 in triple-negative breast cancer. *BMC Cancer*. 2017;17:745.
35. Kong P, Chen L, Yu M et al miR-3178 inhibits cell proliferation and metastasis by targeting Notch1 in triple-negative breast cancer. *Cell Death Dis*. 2018;9:1059.
36. Tang H, Liu P, Yang L et al miR-185 suppresses tumor proliferation by directly targeting E2F6 and DNMT1 and indirectly upregulating BRCA1 in triple-negative breast cancer. *Mol Cancer Ther*. 2014;13:3185-3197.
37. Chai C, Wu H, Wang B, Eisenstat DD, Leng RP. MicroRNA-498 promotes proliferation and migration by targeting the tumor suppressor PTEN in breast cancer cells. *Carcinogenesis*. 2018;39:1185-1196.
38. Son D, Kim Y, Lim S et al miR-374a-5p promotes tumor progression by targeting ARRB1 in triple negative breast cancer. *Cancer Lett*. 2019;454:224-233.
39. Guo GC, Wang JX, Han ML, Zhang LP, Li L. microRNA-761 induces aggressive phenotypes in triple-negative breast cancer cells by repressing TRIM29 expression. *Cell Oncol (Dordt)*. 2017;40:157-166.
40. Guarnieri AL, Towers CG, Drasin DJ et al The miR-106b-25 cluster mediates breast tumor initiation through activation of NOTCH1 via direct repression of NEDD4L. *Oncogene*. 2018;37:3879-3893.
41. Cantini L, Bertoli G, Cava C et al Identification of microRNA clusters cooperatively acting on epithelial to mesenchymal transition in triple negative breast cancer. *Nucleic Acids Res*. 2019;47:2205-2215.
42. Diaz-Riascos ZV, Ginesta MM, Fabregat J et al Expression and role of microRNAs from the miR-200 family in the tumor formation and metastatic propensity of pancreatic cancer. *Mol Ther Nucleic Acids*. 2019;17:491-503.
43. Title AC, Hong SJ, Pires ND et al Genetic dissection of the miR-200-Zeb1 axis reveals its importance in tumor differentiation and invasion. *Nat Commun*. 2018;9:4671.
44. Tsouko E, Wang J, Frigo DE, Aydogdu E, Williams C. miR-200a inhibits migration of triple-negative breast cancer cells through direct repression of the EPHA2 oncogene. *Carcinogenesis*. 2015;36:1051-1060.
45. Humphries B, Wang Z, Oom AL et al MicroRNA-200b targets protein kinase Calpha and suppresses triple-negative breast cancer metastasis. *Carcinogenesis*. 2014;35:2254-2263.
46. Choi SK, Kim HS, Jin T, Hwang EH, Jung M, Moon WK. Overexpression of the miR-141/200c cluster promotes the migratory and invasive ability of triple-negative breast cancer cells

- through the activation of the FAK and PI3K/AKT signaling pathways by secreting VEGF-A. *BMC Cancer*. 2016;16:570.
47. Brabletz T, Kalluri R, Nieto MA, Weinberg RA. EMT in cancer. *Nat Rev Cancer*. 2018;18:128-134.
 48. Lv ZD, Kong B, Liu XP et al miR-655 suppresses epithelial-to-mesenchymal transition by targeting Prrx1 in triple-negative breast cancer. *J Cell Mol Med*. 2016;20:864-873.
 49. Hong L, Pan F, Jiang H et al miR-125b inhibited epithelial-mesenchymal transition of triple-negative breast cancer by targeting MAP2K7. *Onco Targets Ther*. 2016;9:2639-2648.
 50. Lv ZD, Yang DX, Liu XP et al MiR-212-5p suppresses the epithelial-mesenchymal transition in triple-negative breast cancer by targeting Prrx2. *Cell Physiol Biochem*. 2017;44:1785-1795.
 51. Xiong H, Yan T, Zhang W et al miR-613 inhibits cell migration and invasion by downregulating Daam1 in triple-negative breast cancer. *Cell Signal*. 2018;44:33-42.
 52. Habas R, Kato Y, He X. Wnt/Frizzled activation of Rho regulates vertebrate gastrulation and requires a novel Formin homology protein Daam1. *Cell*. 2001;107:843-854.
 53. Zhu Y, Tian Y, Du J et al Dvl2-dependent activation of Daam1 and RhoA regulates Wnt5a-induced breast cancer cell migration. *PLoS ONE*. 2012;7:e37823.
 54. Yan T, Zhang A, Shi F et al Integrin alphavbeta3-associated DAAM1 is essential for collagen-induced invadopodia extension and cell haptotaxis in breast cancer cells. *J Biol Chem*. 2018;293:10172-10185.
 55. Humphries B, Wang Z, Li Y, Jhan JR, Jiang Y, Yang C. ARHGAP18 downregulation by miR-200b suppresses metastasis of triple-negative breast cancer by enhancing activation of RhoA. *Cancer Res*. 2017;77:4051-4064.
 56. Nedeljkovic M, Damjanovic A. Mechanisms of chemotherapy resistance in triple-negative breast cancer-how we can rise to the challenge. *Cells*. 2019;8:957.
 57. Tan X, Peng J, Fu Y et al miR-638 mediated regulation of BRCA1 affects DNA repair and sensitivity to UV and cisplatin in triple-negative breast cancer. *Breast Cancer Res*. 2014;16:435.
 58. Wang B, Zhang Y, Ye M, Wu J, Ma L, Chen H. Cisplatin-resistant MDA-MB-231 cell-derived exosomes increase the resistance of recipient cells in an exosomal miR-423-5p-dependent manner. *Curr Drug Metab*. 2019;20:804-814.
 59. Liu X, Tang H, Chen J et al MicroRNA-101 inhibits cell progression and increases paclitaxel sensitivity by suppressing MCL-1 expression in human triple-negative breast cancer. *Oncotarget*. 2015;6:20070-20083.
 60. Sha LY, Zhang Y, Wang W et al MiR-18a upregulation decreases Dicer expression and confers paclitaxel resistance in triple negative breast cancer. *Eur Rev Med Pharmacol Sci*. 2016;20:2201-2208.
 61. Ouyang M, Li Y, Ye S et al MicroRNA profiling implies new markers of chemoresistance of triple-negative breast cancer. *PLoS ONE*. 2014;9:e96228.
 62. Niu J, Xue A, Chi Y et al Induction of miRNA-181a by genotoxic treatments promotes chemotherapeutic resistance and metastasis in breast cancer. *Oncogene*. 2016;35:1302-1313.
 63. Sasaki A, Tsunoda Y, Tsuji M et al Decreased miR-206 expression in BRCA1 wild-type triple-negative breast cancer cells after concomitant treatment with gemcitabine and a Poly(ADP-ribose) polymerase-1 inhibitor. *Anticancer Res*. 2014;34:4893-4897.
 64. Adams BD, Wali VB, Cheng CJ et al miR-34a Silences c-SRC to attenuate tumor growth in triple-negative breast cancer. *Cancer Res*. 2016;76:927-939.
 65. Ruivo CF, Adem B, Silva M, Melo SA. The biology of cancer exosomes: insights and new perspectives. *Cancer Res*. 2017;77:6480-6488.
 66. Das K, Paul S, Singh A et al Triple negative breast cancer-derived microvesicles transfer microRNA221 to the recipient cells and thereby promote epithelial-to-mesenchymal transition. *J Biol Chem*. 2019;294:13681-13696.
 67. Di Modica M, Regondi V, Sandri M et al Breast cancer-secreted miR-939 downregulates VE-cadherin and destroys the barrier function of endothelial monolayers. *Cancer Lett*. 2017;384:94-100.
 68. Stevic I, Muller V, Weber K et al Specific microRNA signatures in exosomes of triple-negative and HER2-positive breast cancer patients undergoing neoadjuvant therapy within the GeparSixto trial. *BMC Med*. 2018;16:179.
 69. Lim LJ, Wong SYS, Huang F et al Roles and regulation of long noncoding RNAs in hepatocellular carcinoma. *Cancer Res*. 2019;79:5131-5139.
 70. Quinn JJ, Chang HY. Unique features of long non-coding RNA biogenesis and function. *Nat Rev Genet*. 2016;17:47-62.
 71. Boldogkoi Z, Moldovan N, Balazs Z, Snyder M, Tombacz D. Long-read sequencing – a powerful tool in viral transcriptome research. *Trends Microbiol*. 2019;27:578-592.
 72. Engreitz J, Abudayyeh O, Gootenberg J, Zhang F. CRISPR tools for systematic studies of RNA regulation. *Cold Spring Harb Perspect Biol*. 2019;11:a035386.
 73. Ramanathan M, Porter DF, Khavari PA. Methods to study RNA-protein interactions. *Nat Methods*. 2019;16:225-234.
 74. Dahariya S, Paddibhatla I, Kumar S, Raghuvanshi S, Pallepati A, Gutti RK. Long non-coding RNA: classification, biogenesis and functions in blood cells. *Mol Immunol*. 2019;112:82-92.
 75. Fico A, Fiorenzano A, Pascale E, Patriarca EJ, Minchiotti G. Long non-coding RNA in stem cell pluripotency and lineage commitment: functions and evolutionary conservation. *Cell Mol Life Sci*. 2019;76:1459-1471.
 76. Hua Q, Jin M, Mi B et al LINC01123, a c-Myc-activated long non-coding RNA, promotes proliferation and aerobic glycolysis of non-small cell lung cancer through miR-199a-5p/c-Myc axis. *J Hematol Oncol*. 2019;12:91.
 77. Yao RW, Wang Y, Chen LL. Cellular functions of long noncoding RNAs. *Nat Cell Biol*. 2019;21:542-551.
 78. Chedin F. Nascent connections: R-loops and chromatin patterning. *Trends Genet*. 2016;32:828-838.
 79. Arab K, Karaulanov E, Musheev M et al GADD45A binds R-loops and recruits TET1 to CpG island promoters. *Nat Genet*. 2019;51:217-223.
 80. Long Y, Wang X, Youmans DT, Cech TR. How do lncRNAs regulate transcription? *Sci Adv*. 2017;3:eaao2110.
 81. Huang C, Liao X, Jin H et al MEG3, as a competing endogenous RNA, binds with miR-27a to promote PHLPP2 protein translation and impairs bladder cancer invasion. *Mol Ther Nucleic Acids*. 2019;16:51-62.
 82. Clemson CM, Hutchinson JN, Sara SA et al An architectural role for a nuclear noncoding RNA: NEAT1 RNA is essential for the structure of paraspeckles. *Mol Cell*. 2009;33:717-726.
 83. Li Y, Wang Z, Shi H et al HBXIP and LSD1 scaffolded by lncRNA hotair mediate transcriptional activation by c-Myc. *Cancer Res*. 2016;76:293-304.
 84. Zhang S, Wang Y, Jia L et al Profiling the long noncoding RNA interaction network in the regulatory elements of target genes by chromatin in situ reverse transcription sequencing. *Genome Res*. 2019;29:1521-1532.
 85. Han P, Chang CP. Long non-coding RNA and chromatin remodeling. *RNA Biol*. 2015;12:1094-1098.
 86. Grelet S, Link LA, Howley B et al A regulated PNUITS mRNA to lncRNA splice switch mediates EMT and tumour progression. *Nat Cell Biol*. 2017;19:1105-1115.
 87. Jadaliha M, Gholamalamdari O, Tang W et al A natural antisense lncRNA controls breast cancer progression by promoting tumor suppressor gene mRNA stability. *PLoS Genet*. 2018;14:e1007802.

88. Guo FX, Wu Q, Li P et al The role of the lncRNA-FA2H-2-MLKL pathway in atherosclerosis by regulation of autophagy flux and inflammation through mTOR-dependent signaling. *Cell Death Differ.* 2019;26:1670-1687.
89. Hosseini E, Bagheri-Hosseinabadi Z, De Toma I, Jafarisani M, Sadeghi I. The importance of long non-coding RNAs in neuropsychiatric disorders. *Mol Aspects Med.* 2019;70:127-140.
90. Wang CJ, Zhu CC, Xu J et al Correction to: The lncRNA UCA1 promotes proliferation, migration, immune escape and inhibits apoptosis in gastric cancer by sponging anti-tumor miRNAs. *Mol Cancer.* 2019;18:129.
91. Han C, Li X, Fan Q, Liu G, Yin J. CCAT1 promotes triple-negative breast cancer progression by suppressing miR-218/ZFX signaling. *Aging (Albany NY).* 2019;11:4858-4875.
92. Luo L, Tang H, Ling L et al LINC01638 lncRNA activates MTDH-Twist1 signaling by preventing SPOP-mediated c-Myc degradation in triple-negative breast cancer. *Oncogene.* 2018;37:6166-6179.
93. Wang X, Chen T, Zhang Y et al Long noncoding RNA linc00339 promotes triple-negative breast cancer progression through miR-377-3p/HOXC6 signaling pathway. *J Cell Physiol.* 2019;234:13303-13317.
94. Wang S, Ke H, Zhang H et al lncRNA MIR100HG promotes cell proliferation in triple-negative breast cancer through triplex formation with p27 loci. *Cell Death Dis.* 2018;9:805.
95. Vidovic D, Huynh TT, Konda P et al ALDH1A3-regulated long non-coding RNA NRAD1 is a potential novel target for triple-negative breast tumors and cancer stem cells. *Cell Death Differ.* 2019;27:363-378.
96. Tang J, Zhong G, Zhang H et al lncRNA DANCR upregulates PI3K/AKT signaling through activating serine phosphorylation of RXRA. *Cell Death Dis.* 2018;9:1167.
97. Youness RA, Hafez HM, Khallaf E, Assal RA, Abdel Motaal A, Gad MZ. The long noncoding RNA sONE represses triple-negative breast cancer aggressiveness through inducing the expression of miR-34a, miR-15a, miR-16, and let-7a. *J Cell Physiol.* 2019;234:20286-20297.
98. Wang L, Liu D, Wu X et al Long non-coding RNA (lncRNA) RMST in triple-negative breast cancer (TNBC): expression analysis and biological roles research. *J Cell Physiol.* 2018;233:6603-6612.
99. Wang N, Hou M, Zhan Y, Sheng X. lncRNA PTCSC3 inhibits triple-negative breast cancer cell proliferation by downregulating lncRNA H19. *J Cell Biochem.* 2019;120:15083-15088.
100. Zhang H, Zhang N, Liu Y et al Epigenetic regulation of NAMPT by NAMPT-AS drives metastatic progression in triple-negative breast cancer. *Can Res.* 2019;79:3347-3359.
101. Eades G, Wolfson B, Zhang Y, Li Q, Yao Y, Zhou Q. lincRNA-RoR and miR-145 regulate invasion in triple-negative breast cancer via targeting ARF6. *Mol Cancer Res.* 2015;13:330-338.
102. Wang PS, Chou CH, Lin CH et al A novel long non-coding RNA linc-ZNF469-3 promotes lung metastasis through miR-574-5p-ZEB1 axis in triple negative breast cancer. *Oncogene.* 2018;37:4662-4678.
103. Shi F, Xiao F, Ding P, Qin H, Huang R. Long noncoding RNA highly up-regulated in liver cancer predicts unfavorable outcome and regulates metastasis by MMPs in triple-negative breast cancer. *Arch Med Res.* 2016;47:446-453.
104. Augoff K, McCue B, Plow EF, Sossey-Alaoui K. miR-31 and its host gene lncRNA LOC554202 are regulated by promoter hypermethylation in triple-negative breast cancer. *Mol Cancer.* 2012;11:5.
105. Song X, Liu Z, Yu Z. lncRNA NEF is downregulated in triple negative breast cancer and correlated with poor prognosis. *Acta Biochim Biophys Sin (Shanghai).* 2019;51:386-392.
106. Shin VY, Chen J, Cheuk IW et al Long non-coding RNA NEAT1 confers oncogenic role in triple-negative breast cancer through modulating chemoresistance and cancer stemness. *Cell Death Dis.* 2019;10:270.
107. Gooding AJ, Zhang B, Gunawardane L, Beard A, Valadkhan S, Schiemann WP. The lncRNA BORG facilitates the survival and chemoresistance of triple-negative breast cancers. *Oncogene.* 2019;38:2020-2041.
108. Wu J, Chen H, Ye M et al Long noncoding RNA HCP5 contributes to cisplatin resistance in human triple-negative breast cancer via regulation of PTEN expression. *Biomed Pharmacother.* 2019;115:108869.
109. Zheng S, Li M, Miao K, Xu H. lncRNA GAS5-promoted apoptosis in triple-negative breast cancer by targeting miR-378a-5p/SUFU signaling. *J Cell Biochem.* 2020;121:2225-2235.
110. Zheng T, Pang Z, Zhao Z. A gene signature predicts response to neoadjuvant chemotherapy in triple-negative breast cancer patients. *Biosci Rep.* 2019;39:BSR20190414.
111. Liu YR, Jiang YZ, Xu XE, Hu X, Yu KD, Shao ZM. Comprehensive transcriptome profiling reveals multigene signatures in triple-negative breast cancer. *Clin Cancer Res.* 2016;22:1653-1662.
112. Liu M, Xing LQ, Liu YJ. A three-long noncoding RNA signature as a diagnostic biomarker for differentiating between triple-negative and non-triple-negative breast cancers. *Medicine (Baltimore).* 2017;96:e6222.
113. Bermejo JL, Huang G, Manoochehri M et al Long intergenic non-coding RNA 299 methylation in peripheral blood is a biomarker for triple-negative breast cancer. *Epigenomics.* 2019;11:81-93.
114. Jiang Y-Z, Liu Y-R, Xu X-E et al Transcriptome analysis of triple-negative breast cancer reveals an integrated mRNA-lncRNA signature with predictive and prognostic value. *Can Res.* 2016;76:2105-2114.
115. Lv M, Xu P, Wu Y et al lncRNAs as new biomarkers to differentiate triple negative breast cancer from non-triple negative breast cancer. *Oncotarget.* 2016;7:13047-13059.
116. Tian T, Gong Z, Wang M et al Identification of long non-coding RNA signatures in triple-negative breast cancer. *Cancer Cell Int.* 2018;18:103.
117. Topalian SL, Drake CG, Pardoll DM. Immune checkpoint blockade: a common denominator approach to cancer therapy. *Cancer Cell.* 2015;27:450-461.
118. Hu Q, Ye Y, Chan LC et al Oncogenic lncRNA downregulates cancer cell antigen presentation and intrinsic tumor suppression. *Nat Immunol.* 2019;20:835-851.
119. Sanger HL, Klotz G, Riesner D, Gross HJ, Kleinschmidt AK. Viroids are single-stranded covalently closed circular RNA molecules existing as highly base-paired rod-like structures. *Proc Natl Acad Sci USA.* 1976;73:3852-3856.
120. Kelly S, Greenman C, Cook PR, Papantonis A. Exon skipping is correlated with exon circularization. *J Mol Biol.* 2015;427:2414-2417.
121. Shang Q, Yang Z, Jia R, Ge S. The novel roles of circRNAs in human cancer. *Mol Cancer.* 2019;18:6.
122. Jeck WR, Sorrentino JA, Wang K et al Circular RNAs are abundant, conserved, and associated with ALU repeats. *RNA.* 2013;19:141-157.
123. Zhang Y, Zhang XO, Chen T et al Circular intronic long noncoding RNAs. *Mol Cell.* 2013;51:792-806.
124. Petkovic S, Muller S. RNA circularization strategies in vivo and in vitro. *Nucleic Acids Res.* 2015;43:2454-2465.
125. Cheng Z, Yu C, Cui S et al circTP63 functions as a ceRNA to promote lung squamous cell carcinoma progression by upregulating FOXM1. *Nat Commun.* 2019;10:3200.
126. Chen X, Mao R, Su W et al Circular RNA circHIPK3 modulates autophagy via MIR124-3p-STAT3-PRKAA/AMPKalpha signaling in STK11 mutant lung cancer. *Autophagy.* 2020;16:659-671.

127. Li Z, Hu Y, Zeng Q et al Circular RNA MYLK promotes hepatocellular carcinoma progression by increasing Rab23 expression by sponging miR-362-3p. *Cancer Cell Int.* 2019;19:211.
128. Li Z, Huang C, Bao C et al Exon-intron circular RNAs regulate transcription in the nucleus. *Nat Struct Mol Biol.* 2015;22:256-264.
129. Holdt LM, Stahringer A, Sass K et al Circular non-coding RNA ANRIL modulates ribosomal RNA maturation and atherosclerosis in humans. *Nat Commun.* 2016;7:12429.
130. Du WW, Yang W, Liu E, Yang Z, Dhaliwal P, Yang BB. Foxo3 circular RNA retards cell cycle progression via forming ternary complexes with p21 and CDK2. *Nucleic Acids Res.* 2016;44:2846-2858.
131. Legnini I, Di Timoteo G, Rossi F et al Circ-ZNF609 is a circular RNA that can be translated and functions in myogenesis. *Mol Cell.* 2017;66:22-37 e29.
132. Yang Y, Gao X, Zhang M et al Novel role of FBXW7 circular RNA in repressing glioma tumorigenesis. *J Natl Cancer Inst.* 2018;110:304-315.
133. Zhang M, Huang N, Yang X et al A novel protein encoded by the circular form of the SHPRH gene suppresses glioma tumorigenesis. *Oncogene.* 2018;37:1805-1814.
134. He R, Liu P, Xie X et al circGFRA1 and GFRA1 act as ceRNAs in triple negative breast cancer by regulating miR-34a. *J Exp Clin Cancer Res.* 2017;36:145.
135. Chen B, Wei W, Huang X et al circEPST11 as a prognostic marker and mediator of triple-negative breast cancer progression. *Theranostics.* 2018;8:4003-4015.
136. Wang S, Li Q, Wang Y et al Upregulation of circ-UBAP2 predicts poor prognosis and promotes triple-negative breast cancer progression through the miR-661/MTA1 pathway. *Biochem Biophys Res Commun.* 2018;505:996-1002.
137. Yang R, Xing L, Zheng X, Sun Y, Wang X, Chen J. The circRNA circAGFG1 acts as a sponge of miR-195-5p to promote triple-negative breast cancer progression through regulating CCNE1 expression. *Mol Cancer.* 2019;18:4.
138. Tang H, Huang X, Wang J et al circKIF4A acts as a prognostic factor and mediator to regulate the progression of triple-negative breast cancer. *Mol Cancer.* 2019;18:23.
139. Kong Y, Yang L, Wei W et al CircPLK1 sponges miR-296-5p to facilitate triple-negative breast cancer progression. *Epigenomics.* 2019;11:1163-1176.
140. Zou Y, Zheng S, Xiao W et al circRAD18 sponges miR-208a/3164 to promote triple-negative breast cancer progression through regulating IGF1 and FGF2 expression. *Carcinogenesis.* 2019;40:1469-1479.
141. Zhang J, Xu HD, Xing XJ, Liang ZT, Xia ZH, Zhao Y. CircRNA_069718 promotes cell proliferation and invasion in triple-negative breast cancer by activating Wnt/beta-catenin pathway. *Eur Rev Med Pharmacol Sci.* 2019;23:5315-5322.
142. Zeng K, He B, Yang BB et al The pro-metastasis effect of circANKS1B in breast cancer. *Mol Cancer.* 2018;17:160.
143. Sang M, Meng L, Liu S et al Circular RNA ciRS-7 maintains metastatic phenotypes as a ceRNA of miR-1299 to target MMPs. *Mol Cancer Res.* 2018;16:1665-1675.
144. Wang ST, Liu LB, Li XM et al Circ-ITCH regulates triple-negative breast cancer progression through the Wnt/beta-catenin pathway. *Neoplasma.* 2019;66:232-239.
145. Xu JZ, Shao CC, Wang XJ et al circTADA2As suppress breast cancer progression and metastasis via targeting miR-203a-3p/SOCS3 axis. *Cell Death Dis.* 2019;10:175.
146. Ye F, Gao G, Zou Y et al circFBXW7 inhibits malignant progression by sponging miR-197-3p and encoding a 185-aa protein in triple-negative breast cancer. *Mol Ther Nucleic Acids.* 2019;18:88-98.
147. Cole C, Sobala A, Lu C et al Filtering of deep sequencing data reveals the existence of abundant Dicer-dependent small RNAs derived from tRNAs. *RNA.* 2009;15:2147-2160.
148. Anderson P, Ivanov P. tRNA fragments in human health and disease. *FEBS Lett.* 2014;588:4297-4304.
149. Feng W, Li Y, Chu J et al Identification of tRNA-derived small non-coding RNAs as potential biomarkers for prediction of recurrence in triple-negative breast cancer. *Cancer Med.* 2018;7:5130-5144.
150. Telonis AG, Rigoutsos I. Race disparities in the contribution of miRNA isoforms and tRNA-derived fragments to triple-negative breast cancer. *Cancer Res.* 2018;78:1140-1154.
151. Cui Y, Huang Y, Wu X et al Hypoxia-induced tRNA-derived fragments, novel regulatory factor for doxorubicin resistance in triple-negative breast cancer. *J Cell Physiol.* 2019;234:8740-8751.
152. Gong J, Li Y, Liu CJ et al A pan-cancer analysis of the expression and clinical relevance of small nucleolar RNAs in human cancer. *Cell Rep.* 2017;21:1968-1981.
153. McMahon M, Contreras A, Holm M et al A single H/ACA small nucleolar RNA mediates tumor suppression downstream of oncogenic RAS. *eLife.* 2019;8:e48847.
154. Liang J, Wen J, Huang Z, Chen XP, Zhang BX, Chu L. Small nucleolar RNAs: insight into their function in cancer. *Front Oncol.* 2019;9:587.
155. Cheerla N, Gevaert O. MicroRNA based pan-cancer diagnosis and treatment recommendation. *BMC Bioinformatics.* 2017;18:32.
156. Ye B, Shi J, Kang H et al Advancing pan-cancer gene expression survival analysis by inclusion of non-coding RNA. *RNA Biol.* 2019;1-8.
157. Guo Y, Yu H, Wang J et al The landscape of small non-coding RNAs in triple-negative breast cancer. *Genes (Basel).* 2018;9:29.
158. Iman M, Mostafavi SS, Arab SS, Azimzadeh S, Poorebrahim M. HOXB7 and Hsa-miR-222 as the potential therapeutic candidates for metastatic colorectal cancer. *Recent Pat Anticancer Drug Discov.* 2016;11:434-443.
159. Wang WT, Han C, Sun YM, Chen TQ, Chen YQ. Noncoding RNAs in cancer therapy resistance and targeted drug development. *J Hematol Oncol.* 2019;12:55.
160. Shu D, Li H, Shu Y et al Systemic delivery of anti-miRNA for suppression of triple negative breast cancer utilizing RNA nanotechnology. *ACS Nano.* 2015;9:9731-9740.
161. Yin H, Xiong G, Guo S et al Delivery of anti-miRNA for triple-negative breast cancer therapy using RNA nanoparticles targeting stem cell marker CD133. *Mol Ther.* 2019;27:1252-1261.

How to cite this article: Mei J, Hao L, Wang H, et al. Systematic characterization of non-coding RNAs in triple-negative breast cancer. *Cell Prolif.* 2020;53:e12801. <https://doi.org/10.1111/cpr.12801>