

# Wrist Fracture in a Child Irreducible Due to Soft Tissue Interposition

Shinsuke Takeda, MD\*†  
 Katsuyuki Iwatsuki, MD, PhD†  
 Akihiko Tabuchi, MD\*  
 Sadahiro Kubo, MD\*  
 Satoshi Teranishi, MD\*  
 Hitoshi Hirata, MD, PhD†

\*Anjo Kosei Hospital, Emergency and Critical Care Center, Anjo, Japan  
 †Nagoya University Graduate School of Medicine, Department of Hand Surgery, Nagoya, Japan

Section Editor: Manish Amin, DO

Submission history: Submitted May 9, 2018; Revision received June 24, 2018; Accepted July 5, 2018

Electronically published August 16, 2018

Full text available through open access at [http://escholarship.org/uc/uciem\\_cpccm](http://escholarship.org/uc/uciem_cpccm)

DOI: 10.5811/cpcem.2018.7.38952

[Clin Pract Cases Emerg Med. 2018;2(4):363-364.]

## CASE PRESENTATION

An 11-year-old boy fell onto his outstretched arm. He presented to the emergency department with a deformity of his left wrist. Radiograph revealed a greenstick fracture with volar angulation of the distal radius. The distal ulnar physis was disrupted (Salter-Harris type II) and the proximal metaphyseal fragment was displaced dorsally; however, the distal radioulnar joint was intact (Image 1). Closed reduction of the distal ulna under axillary block failed. Three-dimensional computed tomography (3DCT) was performed before open reduction.

## DIAGNOSIS

This type of fracture is known as a Galeazzi-equivalent fracture. Galeazzi-fracture dislocation is a well-known injury,

consisting of a distal radial shaft fracture and dislocation of the distal radioulnar joint (DRUJ). It is rare in adults, and even more uncommon in children. In contrast, Galeazzi-equivalent fractures consist of a fracture at the distal radial metadiaphyseal area with complete distal ulnar epiphyseal separation instead of the more common pattern of DRUJ dislocation.<sup>1</sup> The ulnar physeal fracture in a Galeazzi-equivalent injury can be irreducible due to soft tissue interposition (periosteum, extensor tendons, or joint capsule). It is important to identify and analyze these fractures precisely, as growth arrest has been reported after such injuries.<sup>2</sup> 3DCT revealed the interposition of the extensor carpi ulnaris between the fragments, which hindered the reduction; this was confirmed intraoperatively (Image 2). The patient required



**Image 1.** Anteroposterior (left) and lateral (right) radiographs of the distal forearm and wrist joint. Radiographs show the radius and ulnar fractures (arrows).



**Image 2.** Three-dimensional computed tomography shows the interposition of the extensor carpi ulnaris between the fragments (arrows).

open reduction and fixation of the ulnar physeal fracture with two Kirschner wires. He has regained wrist range of motion, with no complications at two-year follow-up.

Documented patient informed consent and/or Institutional Review Board approval has been obtained and filed for publication of this case report.

---

*Address for Correspondence:* Katsuyuki Iwatsuki, MD, PhD, Nagoya University Graduate School of Medicine, Department of Hand Surgery, 65 Tsurumai-cho, Shouwa-ku, Nagoya, Aichi 466-0065, Japan. Email: [kiwatsuki@med.nagoya-u.ac.jp](mailto:kiwatsuki@med.nagoya-u.ac.jp).

*Conflicts of Interest:* By the CPC-EM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

*Copyright:* © 2018 Takeda et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

---

### *CPC-EM Capsule*

What do we already know about this clinical entity?

*The ulnar physeal fracture in a Galeazzi-equivalent fracture can be irreducible due to soft tissue interposition, such as periosteum, extensor tendons, or joint capsule.*

What is the major impact of the image(s)?

*The interposition of the extensor carpi ulnaris (ECU) between the fragments, which hindered the reduction was revealed by three-dimensional computed tomography (3DCT).*

How might this improve emergency medicine practice?

*This case report reveals the ECU interposition of this Galeazzi-equivalent fracture by 3DCT and shows the difficulty of closed reduction in the emergency department.*

## REFERENCES

1. Galanopoulos I, Fogg Q, Ashwood N, et al. A widely displaced Galeazzi-equivalent lesion with median nerve compromise. *BMJ Case Rep.* 2012;pii:bcr2012006395.
2. Suthar A and Kothari A. Galeazzi-equivalent pronation type Injury with splitting of ulnar epiphyseal plate into two fragments – A rare case report and review of literature. *J Orthop Case Rep.* 2014;4(4):25-28.