

METHODOLOGY ARTICLE

Open Access



An Eigenvalue test for spatial principal component analysis

V. Montano^{1*}  and T. Jombart²

Abstract

Background: The spatial Principal Component Analysis (sPCA, Jombart (Heredity 101:92-103, 2008) is designed to investigate non-random spatial distributions of genetic variation. Unfortunately, the associated tests used for assessing the existence of spatial patterns (*global and local test*; (Heredity 101:92-103, 2008) lack statistical power and may fail to reveal existing spatial patterns. Here, we present a non-parametric test for the significance of specific patterns recovered by sPCA.

Results: We compared the performance of this new test to the original *global and local* tests using datasets simulated under classical population genetic models. Results show that our test outperforms the original *global and local* tests, exhibiting improved statistical power while retaining similar, and reliable type I errors. Moreover, by allowing to test various sets of axes, it can be used to guide the selection of retained sPCA components.

Conclusions: As such, our test represents a valuable complement to the original analysis, and should prove useful for the investigation of spatial genetic patterns.

Keywords: Eigenvalues, sPCA, Spatial genetic patterns, Monte-Carlo

Background

The principal component analysis (PCA; [1, 2]) is one of the most common multivariate approaches in population genetics [3]. Although PCA is not explicitly accounting for spatial information, it has often been used for investigating spatial genetic patterns [4]. As a complement to PCA, the spatial principal component analysis [5] has been introduced to explicitly include spatial information in the analysis of genetic variation, and gain more power for investigating spatial genetic structures.

sPCA finds synthetic variables, the principal components (PCs), which maximise both the genetic variance and the spatial autocorrelation as measured by Moran's *I* [6]. As such, PCs can reveal two types of patterns: '*global*' structures, which correspond to positive autocorrelation typically observed in the presence of patches or clines, and '*local*' structures, which correspond to negative autocorrelation, whereby neighboring individuals are more genetically distinct than expected at random (for a more detailed explanation on the meaning of *global* and *local*

structures see [5]). The *global* and *local* tests have been developed for detecting the presence of global and local patterns, respectively [5]. Unfortunately, while these tests have robust type I error, they also typically lack power, and can therefore fail to identify existing spatial genetic patterns [5]. Moreover, they can only be used to diagnose the presence or absence of spatial patterns, and are unable to test the significance of specific structures revealed by sPCA axes.

In this paper, we introduce an alternative statistical test which addresses these issues. This approach relies on computing the cumulative sum of a defined set of sPCA eigenvalues as a test statistic, and uses a Monte-Carlo procedure to generate null distributions of the test statistics and approximate *p*-values. After describing our approach, we compare its performances to the global and local tests using simulated datasets, investigating several standard spatial population genetics models. Our approach is implemented as the function *sPCA_randtest* in the package *adeigenet* [7, 8] for the R software [9].

* Correspondence: mirainoshjo@gmail.com

¹School of Biology, University of St Andrews, Bute Building, St Andrews KY16 9TS, UK

Full list of author information is available at the end of the article



Methods

Test statistic

As in most multivariate analyses of genetic markers, our approach analyses a table of centred allele frequencies (i.e. set to a mean frequency of zero), in which rows represent individuals or populations, and columns correspond to alleles of various loci [3, 5, 10]. We note X the resulting matrix, and n the number of individuals analysed. In addition, the sPCA introduces spatial data in the form of a n by n matrix of spatial weights L , in which the i^{th} row contains weights reflecting the spatial proximity of all individuals to individual i . The PCs of sPCA are then found by the eigen-analysis of the symmetric matrix (Jombart et al. [5]):

$$1/(2n)X^T(L^T + L)X$$

We note λ the corresponding non-zero eigenvalues. We differentiate the r positive eigenvalues λ^+ , corresponding to global structures, and the s negative eigenvalues λ^- , corresponding to local structures, so that $\lambda = \{\lambda^+, \lambda^-\}$. Without loss of generality, we assume both sets of eigenvalues are ordered by decreasing absolute value, so that $\lambda_1^+ > \lambda_2^+ > \dots > \lambda_r^+$ and $|\lambda_1^-| > |\lambda_2^-| > \dots > |\lambda_s^-|$. Simply put, each eigenvalue quantifies the magnitude of the spatial genetic patterns in the corresponding PC: larger absolute values indicate stronger global (respectively local)

structures. We note $V^+ = \{v_1^+, \dots, v_r^+\}$ and $V^- = \{v_1^-, \dots, v_s^-\}$ the sets of corresponding PCs. The most natural choice of test statistic to assess whether a given PC contains significant structure would seem to be the corresponding eigenvalue. This would, however, not account for the dependence on previous PCs: v_j^+ (respectively v_j^-) can only be significant if all previous PCs $\{v_1^+, \dots, v_{j-1}^+\}$ are also significant. To account for this, we define the test statistic for v_j^+ as:

$$f_i^+ = \sum_{i=1, \dots, j} \lambda_i^+$$

and as:

$$f_i^- = \sum_{i=1, \dots, j} |\lambda_i^-|$$

for v_j^- .

f_i^+ and f_i^- become larger in the presence of strong global or local structures in the first i^{th} global/local PCs. Therefore, they can be used as test statistics against the null hypotheses of absence of global or local structures in these PCs. The expected distribution of f_i^+ and f_i^- in the absence of spatial structure is not known analytically. Fortunately, it can be approximated using a Monte-Carlo procedure, in which at each permutation individual genotypes are shuffled to be assigned to a different pair of coordinates than in the observed original dataset and f_i^+ and f_i^- are computed. Note that the original values of the




Fig. 1 Flow chart illustrating the steps of the *sPCA_randtest*. The first step on the top panel shows one permutation that is used to obtain one value of f_i^+ and f_i^- . To assess the statistical significance of global either local patterns permutations are repeated x times to obtain empirical distributions of f_i^+ and f_i^- that are thus compared to the observed values of f_i^+ and f_i^- . If at least one of the two is significant, the second step of the test exploits the eigenvalue distribution recorded over the permutations to obtain an empirical p -value for each eigenvalue, starting from the most positive (or most negative). As the first eigenvalue is significant in comparison with a chosen threshold, the following is tested and compared to a more stringent threshold (Bonferroni correction) until a non-significant eigenvalue is found and the routine stops

test statistic are also included in these distributions, as the initial spatial configuration is by definition a possible random outcome. The p -values are then computed as the relative frequencies of permuted statistics equal to or greater than the initial value of f_i^* or f_i^- .


To guide the selection of global and local PCs to retain, the simulated values of each eigenvalue (from most positive to most negative), which make up the f_i^* and f_i^- statistics, are also recorded during the permutation procedure. In this way, if global or local structures are detected to be significant, an observed p -value for each observed eigenvalue can be estimated by comparison with its simulated eigenvalue distribution. Note that the number of eigenvalues produced by an sPCA does not change between the observed and permuted datasets, so each observed eigenvalue can be compared with the distribution of the corresponding simulated one. This testing procedure can be used with increasing numbers of retained axes, testing the significance of a new axis as long as previous axes showed significant structure. As one test is performed per axis, we use Bonferroni correction to avoid the inflation of type I error, so that the significance level for the i^{th} PC will be α/i , where α is the target type I error. Hence, the correction implies that if the most positive (or negative) eigenvalue is significant in regards with the chosen p -value threshold, the second

eigenvalue is tested for a p -value threshold that is the half of the previous and so on. The entire testing procedure is implemented in the function *spca_randtest* in the package *adeget* [7, 8] for R [9]. A flow chart of the test procedure is shown in Fig. 1.

Simulation study

To assess the performance of our test, we simulated genetic data under three migration models: island (IS) and stepping stone (SS), using the software *GenomePop* 2.7 [11], and isolation by distance (IBD), using *IBDSimV2.0* [12]. We simulated the IS and SS models with 4 populations, each with 25 individuals, and a single population under IBD with 100 individuals. 200 unlinked biallelic diploid loci (or single nucleotide polymorphisms; SNPs) were simulated. Populations evolved under constant effective population size $\theta = 20$, and interchanged migrants at three different symmetric and homogeneous rates (0.005, 0.01, and 0.1). We performed 100 independent runs for each of the three migration rates, for a total of 300 simulated dataset per migration model. An example of input file for *GenomePop* 2.7 and *IBDSimV2.0* are included as Additional files 1 and 2.

To quantify type I error rates for the *spca_randtest*, *global* and *local* tests, we extracted 100 random coordinates from 10 square 2D grids, using the function



spsample from the *spdep* package [13]. In order to evaluate the rate of false negatives for global patterns, we manually generated 10 sets of 100 pairs of coordinates simulating gradients and/or patches from 2D grids. An example of simulated global patterns is presented in Fig. 2. To test for the rate of false negatives for local patterns, we perform a principal component analysis on 10 random datasets simulated under the SS model with 0.005 migration rate. We used the coordinates of the individuals on the first principal component and set the second coordinate to zero for all individuals (1D). With the coordinates so produced, we used the function *chooseCN* in *adegenet* to obtain 10 neighbouring graphs where the most genetically distinct individuals (falling in the upper quartile of the pairwise genetic distances) are considered as neighbors, while the others are non-neighbors.

We tested 100 simulations each for all the 30 sets of geographic coordinates (random, positive and negative), for each of the three migration rates (0.005, 0.01 and 0.1), for each of the three migration models (IS, SS, IBD; total of 9000 tests per migration model). We repeated all tests using a subset of 40 SNPs per individual, for a total of 18,000 tests in the absence of spatial structures, and 36,000 tests in the presence of global or local structures.

Results

Statistical power of the *spca_randtest*

We compared the performances of the *spca_randtest* with the *global* and *local* tests in three settings: in the absence of spatial structure, and in the presence of global, and local structures. The results obtained in the absence of spatial structure show that all tests have reliable type I errors (Table 1 and 2). The *spca_randtest* exhibited consistently better performances for detecting existing structures in the data than both *global* and *local tests* (Table 1 and 2). Although our simulated local spatial patterns turned out more difficult to detect than global patterns, the *spca_randtest* is twice to five times more effective than the *local test* (Table 1 and 2). Generally, the underlying migration model, the migration rate and the number of loci affect the ability of all tests to detect non-random spatial patterns. Both *spca_randtest* and *global* and *local tests* have in fact a lower sensitivity in presence of island migratory schemes, while results for stepping stone and isolation by distance models are more satisfying (Table 1 and 2). Increasing migration rates lead to a higher rates of false negatives for all tests, which can be overcome using more loci (Table 1 and 2).

Significant eigenvalues are assessed using a hierarchical Bonferroni correction which accounts for non-independence of eigenvalues and multiple testing (Fig. 2). Strong patterns

Table 1 Significant results for *global test* (g test), *local tests* (l test), and *spca_randtest* (r test +/-) for random, global and local patterns using 200 loci per individual. IS, SS, IBD indicate the migration models (see Methods); different migration rates are coded by number: 1 = 0.005, 2 = 0.01 and 3 = 0.1

200 SNPs		Random Patterns				Global Patterns				Local Patterns			
Models	Significance level	g test	r test (+)	l test	r test (-)	g test	r test (+)	l test	rt est. (-)	g test	r test (+)	l test	r test (-)
IS-1	.05	0.054	0.059	<i>0.041</i>	<i>0.047</i>	0.947	0.985	<i>0.029</i>	<i>0.001</i>	<i>0.047</i>	0.071	0.061	0.284
	.01	0.011	<i>0.007</i>	<i>0.009</i>	0.010	0.822	0.948	<i>0.005</i>	<i>0.001</i>	<i>0.008</i>	0.010	0.015	0.113
IS-2	.05	<i>0.040</i>	<i>0.041</i>	0.058	0.056	0.227	0.564	<i>0.044</i>	<i>0.018</i>	0.056	0.059	0.050	0.123
	.01	<i>0.007</i>	<i>0.009</i>	<i>0.009</i>	0.013	0.067	0.302	<i>0.005</i>	<i>0.002</i>	0.011	<i>0.007</i>	0.012	0.026
IS-3	.05	0.051	<i>0.040</i>	0.053	<i>0.041</i>	0.055	<i>0.049</i>	<i>0.045</i>	<i>0.047</i>	<i>0.049</i>	<i>0.047</i>	<i>0.044</i>	0.059
	.01	0.010	0.014	0.013	<i>0.008</i>	0.010	0.013	<i>0.007</i>	0.013	<i>0.002</i>	<i>0.014</i>	<i>0.008</i>	0.019
SS-1	.05	0.053	0.058	0.053	0.050	0.986	0.996	<i>0.022</i>	<i>0.000</i>	0.063	0.064	0.124	0.582
	.01	<i>0.007</i>	0.011	0.010	0.010	0.960	0.988	<i>0.002</i>	<i>0.000</i>	0.017	0.010	<i>0.041</i>	0.398
SS-2	.05	<i>0.044</i>	0.058	0.058	0.063	0.798	0.909	<i>0.047</i>	<i>0.004</i>	<i>0.034</i>	<i>0.044</i>	0.059	0.316
	.01	0.011	0.011	0.013	0.016	0.676	0.771	<i>0.010</i>	<i>0.000</i>	<i>0.004</i>	<i>0.005</i>	0.014	0.147
SS-3	.05	<i>0.047</i>	<i>0.046</i>	0.057	<i>0.049</i>	0.054	0.128	<i>0.040</i>	<i>0.042</i>	<i>0.044</i>	0.054	<i>0.049</i>	0.071
	.01	0.014	<i>0.007</i>	0.011	0.013	0.014	0.036	<i>0.006</i>	0.010	<i>0.003</i>	<i>0.009</i>	<i>0.006</i>	<i>0.009</i>
IBD-1	.05	<i>0.044</i>	0.050	0.053	<i>0.048</i>	0.962	0.999	<i>0.021</i>	<i>0.000</i>	<i>0.025</i>	0.087	0.438	0.809
	.01	<i>0.008</i>	0.012	<i>0.009</i>	0.010	0.926	0.997	<i>0.003</i>	<i>0.000</i>	<i>0.009</i>	0.023	0.192	0.694
IBD-2	.05	0.052	<i>0.045</i>	0.061	<i>0.038</i>	0.967	0.998	<i>0.023</i>	<i>0.000</i>	<i>0.046</i>	0.076	0.451	0.794
	.01	<i>0.009</i>	<i>0.008</i>	0.011	<i>0.009</i>	0.932	0.997	<i>0.004</i>	<i>0.000</i>	<i>0.009</i>	0.018	0.208	0.672
IBD-3	.05	0.052	<i>0.046</i>	0.053	0.050	0.977	0.999	<i>0.015</i>	<i>0.000</i>	0.050	0.083	0.441	0.824
	.01	0.013	<i>0.009</i>	0.011	0.012	0.939	0.999	<i>0.005</i>	<i>0.000</i>	<i>0.009</i>	0.023	0.225	0.684

**p-values* are in italic when non significant and in bold when the fraction of true positive is above 20%
 Results show the proportion of significant tests over 1000 replicates, based on 1000 permutations with thresholds .05 and .01

Table 2 Results for the same simulations reported in Table 1 using a subset of 40 loci per individual

40 SNPs		Random Patterns				Global Patterns				Local Patterns			
Models	Significance level	g test	r test (+)	l test	r test (-)	g test	r test (+)	l test	r test (-)	g test	r test (+)	l test	r test (-)
IS-1	.05	0.052	0.061	0.046	0.050	0.591	0.807	0.033	0.004	0.036	0.000	0.055	0.077
	.01	0.016	0.013	0.010	0.007	0.393	0.592	0.005	0.000	0.004	0.000	0.015	0.022
IS-2	.05	0.053	0.047	0.038	0.042	0.103	0.226	0.046	0.020	0.073	0.000	0.057	0.038
	.01	0.011	0.009	0.006	0.006	0.022	0.072	0.011	0.005	0.012	0.000	0.010	0.006
IS-3	.05	0.047	0.050	0.050	0.045	0.048	0.060	0.044	0.042	0.036	0.000	0.053	0.026
	.01	0.009	0.011	0.008	0.007	0.009	0.011	0.011	0.011	0.002	0.000	0.013	0.001
SS-1	.05	0.052	0.054	0.039	0.049	0.898	0.949	0.017	0.000	0.050	0.001	0.067	0.169
	.01	0.009	0.012	0.005	0.011	0.826	0.865	0.006	0.000	0.007	0.000	0.021	0.052
SS-2	.05	0.046	0.045	0.050	0.046	0.528	0.588	0.044	0.009	0.052	0.000	0.048	0.081
	.01	0.013	0.010	0.010	0.015	0.377	0.370	0.016	0.000	0.005	0.000	0.011	0.014
SS-3	.05	0.068	0.040	0.050	0.048	0.066	0.055	0.053	0.033	0.026	0.000	0.047	0.023
	.01	0.014	0.005	0.013	0.012	0.012	0.009	0.005	0.006	0.006	0.000	0.008	0.000
IBD-1	.05	0.049	0.053	0.052	0.057	0.822	0.883	0.027	0.002	0.034	0.055	0.124	0.480
	.01	0.005	0.008	0.013	0.013	0.755	0.742	0.004	0.000	0.005	0.008	0.032	0.278
IBD-2	.05	0.043	0.054	0.060	0.049	0.835	0.880	0.028	0.001	0.043	0.051	0.111	0.458
	.01	0.011	0.007	0.015	0.009	0.755	0.732	0.005	0.000	0.008	0.015	0.026	0.259
IBD-3	.05	0.043	0.042	0.051	0.050	0.844	0.899	0.026	0.002	0.048	0.058	0.115	0.465
	.01	0.012	0.013	0.012	0.010	0.763	0.756	0.007	0.000	0.009	0.010	0.023	0.263

**p-values* are in italic when non significant and in bold when the fraction of true positive is above 20%

(e.g. IBD) tend to produce a higher number of significant components than weak patterns (e.g. island models with high migration rates), which are otherwise captured by fewer to no components.

Application to real data

We have run the sPCA to compare the new *spca_randtest* and previous tests to a real dataset of human mitochondrial DNA (mtDNA). We used a dataset of 85 populations from Central-Western Africa that spans a big portion of the African continent (from Gabon to Senegal; [14]). Previous analysis on these data detected a clear genetic structure from West to Central Africa with ongoing stepping stone migration movements. We therefore expected that this spatial distribution of genetic variation would be detected as significant. In the sPCA, populations were treated as units of the analysis, for which allele frequencies of mtDNA

polymorphisms are calculated per population. The same approach was used in [14] to run a discriminant analysis of principal components (DAPC; [10]) and detect population genetic structure. The sPCA analysis is found non significant by *global* and *local* tests after 10,000 permutations (*p-value* >0.5), while the *spca_randtest* detects a significant global pattern already with 500 permutations, and with 10,000 permutations the *p-value* for global patterns is 0.005. The second step of the test on single eigenvalues finds the three most positive components to be significant after Bonferroni correction (Table 3). Significant axes can thus be plotted against the spatial network to give a biological interpretation to the results (Fig. 3).


Discussion

We introduced a new statistical test associated to the sPCA to evaluate the statistical significance of global and

Table 3 Results of the *spca_randtest* with 10,000 permutations on the human mtDNA dataset (Montano et al., [14])

Spatial patterns	Observed <i>p-value</i>	Decreasing Positive Eigenvalues	Observed <i>p-value</i>	Bonferroni corrected significant level
Global pattern	0.0058	3.4e-2	0.0105	0.05
Local pattern	0.8826	8.5e-3	0.0137	0.025
		4.1e3	0.0136	0.016
		1.6e-3	0.506	0.0125

The simulated distribution of the f_i^* and f_i^{\sim} statistics are compared to the f_i^* and f_i^{\sim} statistics observed for the original dataset. A significant global pattern (or significant f_i^* observed statistics) is found with the *spca_randtest* (*p-value* <0.01). Thus, each positive eigenvalue is compared with its simulated distribution and assigned to be significant if its observed *p-value* is lower than the corrected Bonferroni *p-value*, with starting threshold of 0.05. Significant observed *p-values* as compared with Bonferroni corrected *p-values* are highlighted in bold



local spatial patterns. Using simulated data, we show that this new approach outperforms previously implemented tests, having greater statistical power (lower type II errors) whilst retaining consistent type I errors. Our simulations also suggest that demographic settings and migratory models can substantially impact the ability to detect spatial patterns. Indeed, high migration rates, non-hierarchical migration models, such as island model, and low amount of loci can hamper or worsen the performance of the test, preventing the detection of actual spatial patterns. In lack of previous information on the demographic history and/or the movement ecology of the population under study, it is certainly useful to exploit all the available genetic information. In this regards, our simulations show how an increased number of loci does improve the ability of the test to provide meaningful results.

The impact of specific factors such as the effective population size or the number of individuals sampled per population remain to be investigated. A more extensive simulation study, possibly comparing different non-model based methods such as sPCA, would clarify the extent of the spatial information that can be obtained with such methods without comparing explicit evolutionary hypotheses. In fact, the sPCA and the associated *spca_randtest* cannot distinguish between explicit migration models. However, the possibility to detect which eigenvalues contain the spatial information provides the user with further information to interpret the biological meaning of the spatial structure, by focusing on few meaningful dimensions.

Our data application seems to confirm that the *spca_randtest* is more effective than *global* or *local* tests. We chose indeed a previously published dataset of human populations which span a subcontinental area of Africa and had been originally detected to be a highly structured dataset with a geographic cline of population differentiation (Montano et al. [14]). On the basis of the original results, we would have expected a spatial global structure to be present in the data and thus detected with

an sPCA. While the *global* test failed to provide statistical significance, the *spca_randtest* did obtain significant results and pointed to the three first most positive components to be also significant after Bonferroni correction. In agreement with the original interpretation of the genetic structure within the samples, spatial component 1 (SP1) shows a clear differentiation of populations in the Gabon-Congo region, while SP2 detects differentiation of Central Nigerian and North Cameroonian populations, on one hand, and extreme Western populations of Senegal, on the other hand, and extreme Western populations of Senegal, on the other hand (Fig. 3). The colored combination of the first and second most positive component (Fig. 3) also correctly detects a more fragmented differentiation across Central forested areas (Cameroon, Gabon and Congo) compared to more homogeneous Central-Western populations, which was the main result of the original publication based on very different approaches [14]. We limited the analysis to these two component as the third did not add much information to the previous.

Conclusions

Our simulation approach coupled with a real data application well illustrates the informativeness of our new test to retrieve significant spatial patterns, being these global or local structures and highlights the usefulness of selecting a specific number of significant components to interpret the biological meaning of the results.

Additional files

Additional file 1: Example of GenomePop Input with a parameter setting to simulate an island model of migration. (TXT 363 kb)

Additional file 2: Example of IbdSettings for IBD_Sim with parameter setting to simulate an isolation by distance evolutionary model. (TXT 635 kb)

Additional file 3: Figure S1. Distributions of significant eigenvalues detected in the presence of global (blue bars) and local (green bars) spatial patterns after hierarchical Bonferroni correction, for 100 significantly positive and 100 significantly negative patterns. Black bars

correspond to eigenvalues which are significant without Bonferroni correction. Bars' height indicates the frequency of observing a significant eigenvalue in a certain position (from most positive to most negative) over the 100 tested patterns. (PDF 1150 kb)

Acknowledgements

We thank two anonymous reviewers for insightful comments and feedback which greatly improved the original version of the manuscript.

Funding

No funds were used for this study.

Availability of data and materials

https://github.com/thibautjombart/adegenet/blob/master/R/spca_randtest.R

Author contributions

Test development: VM and TJ. Data analysis: VM. Wrote the manuscript: VM and TJ.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Competing interests

The authors declare no conflict of interest.

Author details

¹School of Biology, University of St Andrews, Bute Building, St Andrews KY16 9TS, UK. ²Department of Infectious Disease Epidemiology, MRC Centre for Outbreak Analysis and Modelling, Imperial College, St Mary's Campus, Norfolk Place, London W2 1PG, UK.

Received: 1 July 2017 Accepted: 5 December 2017

Published online: 16 December 2017

References

- Pearson K. On lines and planes of closest fit to systems of points in space. *Philosophical Magazine Series*. 1901;6:559–572.
- Hotelling H. Analysis of a complex of statistical variables into principal components. *J Ed Psychol*. 1933;24:417.
- Jombart T, Pontier D, Dufour AB. Genetic markers in the playground of multivariate analysis. *Heredity*. 2009;102:330–41.
- Novembre J, Stephens M. Interpreting principal component analyses of spatial population genetic variation. *Nat Genet*. 2008;40:646–9.
- Jombart T, Devillard S, Dufour AB, Pontier D. Revealing cryptic spatial patterns in genetic variability by a new multivariate method. *Heredity*. 2008;101:92–103.
- Moran PAP. Notes on continuous stochastic phenomena. *Biometrika*. 1950; 37:17–23.
- Jombart T. Adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics*. 2008;24:1403–5.
- Jombart T, Ahmed I. Adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. *Bioinformatics*. 2011;27:3070–1.
- R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2017. URL <https://www.R-project.org/>.
- Jombart T, Devillard S, Balloux F. Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genet*. 2010;11:94.
- Carvajal-Rodríguez A. GENOMEPOP: a program to simulate genomes in populations. *BMC Bioinformatics*. 2008;9:223.
- Leblois R, Estoup A, Rousset F. IBDSim: a computer program to simulate genotypic data under isolation by distance. *Mol Ecol Resour*. 2009;9:107–9.
- Bivand RS, Pebesma E, Gómez-Rubio V. *Applied spatial data analysis with R*. New York: Springer; 2013. p. 378pp.
- Montano V, Marcari V, Pavanello M, Anyaele O, Comas D, Destro-Bisol G, Batini C. The influence of habitats on female mobility in Central and Western Africa inferred from human mitochondrial variation. *BMC Evol Biol*. 2013;13:1:24.
- South A. Rworldmap: a new R package for mapping global data. *R J*. 2011;3:35–43.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

