

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Tris(5-amino-1*H*-1,2,4-triazol-4-ium) dihydrogenphosphate hydrogenphosphate trihydrate

 Mohamed Lahbib Mrad,^a Matthias Zeller,^b Kristen J. Hernandez,^b Mohamed Rzaigui^a and Cherif Ben Nasr^{a*}

^aLaboratoire de Chimie des Matériaux, Faculté des sciences de Bizerte, 7021 Zarzouna, Tunisia, and ^bYoungstown State University, Department of Chemistry, One University Plaza, Youngstown, Ohio 44555-3663, USA
Correspondence e-mail: cherif_bennasr@yahoo.fr


Received 7 October 2012; accepted 26 October 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{N}-\text{C}) = 0.002$ Å; R factor = 0.023; wR factor = 0.059; data-to-parameter ratio = 19.3.

In the crystal structure of the title molecular salt, $3\text{C}_2\text{H}_5\text{N}_4^{+\cdot}\cdot\text{HPO}_4^{2-}\cdot\text{H}_2\text{PO}_4^{-}\cdot 3\text{H}_2\text{O}$, the phosphate-based framework is built upon layers parallel to (010) made up from the $\text{H}_2\text{PO}_4^{-}$ and HPO_4^{2-} anions and water molecules, which are interconnected through $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds. The organic cations are located between the phosphate-water layers and are connected to them *via* $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds. The bond-length features are consistent with an imino resonance form for the exocyclic amino group, as is commonly found for a C–N single bond involving sp^2 -hybridized C and N atoms.

Related literature

For applications of organic phosphate complexes, see: Bringley & Rajeswaran (2006); Dai *et al.* (2002); Masse *et al.* (1993). For graph-set motifs and theory, see: Bernstein *et al.* (1995). For reference structural data, see: Kaabi *et al.* (2004); Shanmuga Sundara Raj *et al.* (2000). For P–OH bond lengths, see: Chtioui & Jouini (2005).



Experimental

Crystal data

$3\text{C}_2\text{H}_5\text{N}_4^{+\cdot}\cdot\text{HO}_4\text{P}^{2-}\cdot\text{H}_2\text{O}_4\text{P}^{-}\cdot 3\text{H}_2\text{O}$
 $M_r = 502.31$
 Monoclinic, Pc
 $a = 10.4793$ (13) Å
 $b = 8.7655$ (11) Å
 $c = 11.4536$ (14) Å
 $\beta = 107.489$ (2)°

$V = 1003.5$ (2) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.30$ mm⁻¹
 $T = 100$ K
 $0.60 \times 0.35 \times 0.18$ mm

Data collection

Bruker SMART APEX CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2011)
 $T_{\min} = 0.693$, $T_{\max} = 0.746$

13833 measured reflections
 6229 independent reflections
 6132 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.059$
 $S = 1.04$
 6229 reflections
 322 parameters
 32 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.33$ e Å⁻³
 $\Delta\rho_{\min} = -0.19$ e Å⁻³
 Absolute structure: Flack (1983), 2950 Friedel pairs
 Flack parameter: -0.02 (4)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1A}-\text{H1A1}\cdots\text{O3B}$	0.85 (1)	2.31 (1)	3.1356 (13)	162 (2)
$\text{N1A}-\text{H1A2}\cdots\text{N3A}^i$	0.85 (1)	2.19 (1)	3.0305 (15)	170 (2)
$\text{N2A}-\text{H2A}\cdots\text{O4B}$	0.88	1.77	2.6130 (13)	161
$\text{N4A}-\text{H4A}\cdots\text{O4B}^i$	0.88	1.76	2.6314 (13)	171
$\text{N1B}-\text{H1B1}\cdots\text{O2}$	0.86 (1)	1.96 (1)	2.8214 (13)	178 (2)
$\text{N1B}-\text{H1B2}\cdots\text{N3B}^i$	0.82 (1)	2.28 (1)	3.0639 (15)	160 (2)
$\text{N2B}-\text{H2B1}\cdots\text{O3}^{ii}$	0.88	1.84	2.6824 (13)	159
$\text{N4B}-\text{H4B}\cdots\text{O1A}^{iii}$	0.88	1.87	2.7376 (12)	167
$\text{N1C}-\text{H1C1}\cdots\text{N3C}^i$	0.83 (1)	2.18 (1)	3.0028 (14)	172 (2)
$\text{N1C}-\text{H1C2}\cdots\text{O3A}$	0.86 (1)	2.24 (1)	3.0589 (13)	160 (2)
$\text{N2C}-\text{H2C}\cdots\text{O4A}$	0.88	1.78	2.6278 (12)	161
$\text{N4C}-\text{H4C}\cdots\text{O4A}^i$	0.88	1.79	2.6645 (12)	170
$\text{O2A}-\text{H2AB}\cdots\text{O3B}^{iv}$	0.76	1.95	2.6593 (11)	155
$\text{O1B}-\text{H1B}\cdots\text{O3A}$	0.77	1.80	2.5495 (12)	161
$\text{O2B}-\text{H2BA}\cdots\text{O1}^{iv}$	0.83	1.73	2.5552 (12)	176
$\text{O1}-\text{H1D}\cdots\text{O2}$	0.84 (1)	1.93 (1)	2.7439 (12)	166 (2)
$\text{O1}-\text{H1E}\cdots\text{O3B}$	0.80 (1)	1.91 (1)	2.6968 (12)	168 (2)
$\text{O2}-\text{H2D}\cdots\text{O1A}^v$	0.82 (1)	1.90 (1)	2.7024 (11)	168 (2)
$\text{O2}-\text{H2E}\cdots\text{O3A}^{iii}$	0.81 (1)	1.95 (1)	2.7566 (12)	178 (2)
$\text{O3}-\text{H3D}\cdots\text{O2B}$	0.79 (1)	2.14 (2)	2.8515 (12)	149 (2)
$\text{O3}-\text{H3E}\cdots\text{O1A}^{iii}$	0.80 (1)	1.92 (1)	2.7085 (11)	176 (2)

Symmetry codes: (i) $x, -y, z + \frac{1}{2}$; (ii) $x, y - 1, z$; (iii) $x + 1, -y + 1, z + \frac{1}{2}$; (iv) $x, -y + 1, z - \frac{1}{2}$; (v) $x + 1, y, z + 1$.

Data collection: *APEX2* (Bruker, 2011); cell refinement: *SAINT* (Bruker, 2011); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL* (Hübschle *et al.*, 2011); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* and *pubCIF* (Westrip, 2010).

We would like to acknowledge the support provided by the Secretary of State for Scientific Research and Technology of Tunisia. The diffractometer was funded by NSF grant 0087210, by Ohio Board of Regents grant CAP-491, and by YSU.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RU2044).

References

- Bernstein, J., Davids, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Bringley, J. F. & Rajeswaran, M. (2006). *Acta Cryst.* **E62**, m1304–m1305.
 Bruker (2011). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Chtioui, A. & Jouini, A. (2005). *Mater. Res. Bull.* **41**, 569–575.

- Dai, J.-C., Wu, X.-T., Fu, Z.-Y., Cui, C.-P., Wu, S.-M., Du, W.-X., Wu, L.-M., Zhang, H.-H. & Sun, Q.-Q. (2002). *Inorg. Chem.* **41**, 1391–1396.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Hübschle, C. B., Sheldrick, G. M. & Dittrich, B. (2011). *J. Appl. Cryst.* **44**, 1281–1284.
- Kaabi, K., Ben Nasr, C. & Lefebvre, F. (2004). *Mater. Res. Bull.* **39**, 205–215.
- Masse, R., Bagieu-Beucher, M., Pecaut, J., Levy, J. P. & Zyss, J. (1993). *J. Nonlinear Opt.* **5**, 413–423.
- Shanmuga Sundara Raj, S., Fun, H.-K., Zhao, P.-S., Jian, F.-F., Lu, L.-D., Yang, X.-J. & Wang, X. (2000). *Acta Cryst.* **C56**, 742–743.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, o3257–o3258 [doi:10.1107/S1600536812044492]

Tris(5-amino-1*H*-1,2,4-triazol-4-ium) dihydrogenphosphate hydrogenphosphate trihydrate

Mohamed Lahbib Mrad, Matthias Zeller, Kristen J. Hernandez, Mohamed Rzaigui and Cherif Ben Nasr

Comment

Inorganic–organic hybrid compounds provide a class of materials with interesting technological applications (Bringley & Rajeswaran, 2006; Dai *et al.*, 2002). Among these materials, compounds with noncentrosymmetric crystallographic structures are interesting for their applications in quadratic non-linear optical materials research (Masse *et al.*, 1993). Their abilities to combine the rigidity and high cohesion of inorganic host matrices with the enhanced polarizability of organic guest chromophores within one molecular scale assists in better performance of optical signal-processing devices. The use of organic-inorganic polar crystalline materials for quadratic nonlinear optical applications is supported by two observations:

- (i) the organic molecules, especially if they contain a delocalized π -system with asymmetric substitution by electron donor-acceptor groups, are highly polarizable entities ideally suited for NLO applications. Being organic materials, the nature of the substituents can be tailored so as to not affect optical transparency;
- (ii) the ionic inorganic host matrices are able to increase the packing cohesion, can induce noncentrosymmetry, and also shift the transparency of crystal towards blue wavelengths.

Within a systematic investigation of new materials resulting from the association of organic chromophores with inorganic species, we report here the synthesis and the characterization of a new hybrid phosphate-amine material, $(\text{C}_2\text{H}_5\text{N}_4)_3(\text{HPO}_4)(\text{H}_2\text{PO}_4)\cdot 3\text{H}_2\text{O}$, which includes the 3-amino-1*H*-1,2,4-triazolium cations, a chromophore which could be efficient in the blue-U.V. wavelength region. The title compound could exhibit a richness of interesting physical properties such as ferroelectricity and nonlinear optic phenomena like second harmonic generation. It crystallizes in a non-centrosymmetric setting in the space group *Pc*. The structure of this organic-inorganic hybrid material consists of one dihydrogenmonophosphate anion, one monohydrogenmonophosphate dianion, three crystallographically independent 3-amino-1*H*-1,2,4-triazolium cations and three water molecules (Fig. 1). The atomic arrangement is a typical layered organization as it is very often encountered in this kind of inorganic-organic hybrid compounds (Kaabi *et al.*, 2004). The H_2PO_4^- anions are hydrogen bonded with the HPO_4^{2-} groups and one of the water molecules (that of O3) to form corrugated chains running parallel to the *a*-axis at (0, 0, 0) and (0, 0, 1/2). These chains are interconnected, *via* O(water)—H \cdots O and O—H \cdots O(water) hydrogen bonds, with the two remaining water molecules $\text{H}_2\text{O}(1)$ and $\text{H}_2\text{O}(2)$, associated through O1—H \cdots O2 hydrogen bonds, on one hand, and with the HPO_4^{2-} anions of the adjacent chain, through O—H \cdots O hydrogen bonds, on the other hand. These hydrogen bonds link the different inorganic units into infinite planar layers parallel to the (0 1 0) plane (Fig. 2) crossing the unit cell at $y = (2n + 1)/2$ (Fig. 3). Within the layers, various graph-set motifs (Bernstein *et al.*, 1995) are apparent, including $R_5^5(10)$ and $R_4^4(12)$ loops. The 3-amino-1*H*-1,2,4-triazolium cations are interconnected *via* weak N—H \cdots N hydrogen bonds, with D—H \cdots A distances between 3.003 (1) and 3.064 (1)

Å, to form organic chains spreading along the *c*-axis at $x \sim (n + 1)/3$ (Fig. 4). The chains are built from the three crystallographically independent organic cations, labelled A, B and C, in such a way that each N—H···N connected chain incorporates only one type of cation: Molecules of type A are located at $x \sim 1/3$, chains at $x \sim 0$ consist of molecules of type B, and the chains at $x \sim 2/3$ are made up of molecules C. Alternating molecules in each of these chains are created by the *c*-glide plane. In two of the chains, that of molecules A and C, alternating molecules are roughly coplanar. In the third, molecules are twisted against each other by an angle of 34.37°. The chains are roughly parallel to each other and weakly π -stacked, with interplanar distances between the mean planes of chains between 3.21 Å (between A and C), and up to 3.52 Å (for A and B). Despite of the quite close interplanar distances, π - π stacking interactions are limited due to molecule offsets in parallel layers, and the non-coplanarity of neighboring molecules in the chains of molecules B. The organic chains are anchored to the inorganic layers through N—H···O hydrogen bonds whose geometrical characteristics are given in Table 2. The projection of the whole arrangement along the *a*-axis (Fig. 4) shows how the organic chains alternate as to fill the space separating parallel inorganic layers. In this structure, three 3-amino-1*H*-1,2,4-triazolium cationic groups compensate the negative charges of the dihydrogenmonophosphate and the mono-hydrogenmonophosphate anions, leading to charge neutrality for the structure as a whole.

The sum of the angles around the N1A, N1B and N1C nitrogen atoms are 360° and the C—N bond distances of the NH₂ groups are 1.332 (1) Å for N1A—C1A, 1.327 (1) Å for N1B—C1B and 1.330 (1) Å for N1C—C1C, which are short for C—N single bonds, but still not quite as contracted as one would expect for a fully established C=N double bond. These bond length features are consistent with an imino resonance form as it is commonly found for a C—N single bond involving *sp*² hybridized C and N atoms (Shanmuga Sundara Raj *et al.*, 2000). In agreement with this, the amino groups are not pyramidal but the electron densities of the hydrogen atoms of the amino groups were found to be in plane with the 3-amino-1*H*-1,2,4-triazolium skeleton. The detailed geometry of the HP(1A)O₄²⁻ and H₂P(1B)O₄⁻ anions shows two kinds of P—O distances. The shortest ones, 1.5243 (8), 1.5294 (8) and 1.5364 Å for the first anion (labelled A) and 1.5132 (8) and 1.5163 (8) Å for the second one (labelled B), correspond to the phosphorous atom doubly bonded to the oxygen atom, while the largest ones 1.5845 (8) Å and (1.5612 (8), 1.5741 (8) Å, respectively, can be attributed to the P—OH bond length. This is in agreement with the literature data (Chtioui & Jouini, 2005). Refining the structure in the asymmetric space group gives a value of -0.02 (4) for the Flack parameter (Flack, 1983), confirming the absolute structure and absence of twinning.

Experimental

Crystals of the title compound were prepared at room temperature by slow addition of a solution of orthophosphoric acid (8 mmol in 30 ml of water) to an alcoholic solution of 3-amino-1*H*-1,2,4-triazole (12 mmol in 30 ml of ethanol). The acid was added until the alcoholic solution became turbid. After filtration, the solution was allowed to slowly evaporate at room temperature over several days leading to formation of transparent prismatic crystals with suitable dimensions for single-crystal structural analysis (1.2 mg, 2.4 mmol, yield 60%). The crystals are stable for months under normal conditions of temperature and humidity.

Refinement

H atoms were placed in calculated positions with the exception of water and NH₂ H atoms, which were located in difference density maps and were refined. C—H distances were set to 0.95 Å, N_{ring}—H distances to 0.88 Å. H atoms of P-bound hydroxy groups were placed geometrically with fixed P—O—H angles, but with variable torional angles and O—H distances to best fit the experimental electron density (AFIX 148 in *SHELXTL*, Sheldrick 2008). All H₂O O—H distances were restrained to be similar within a standard deviation of 0.02 Å. All amino N—H distances were also

restrained to be similar within the same standard deviation. U_{iso} values of H atoms were set to 1.2 or 1.5 times U_{eq} of their respective carrier atom for amino and O-bound H atoms respectively.

Computing details

Data collection: *APEX2* (Bruker, 2011); cell refinement: *S SAINT* (Bruker, 2011); data reduction: *S SAINT* (Bruker, 2011); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL* (Hübschle *et al.*, 2011); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *publCIF* (Westrip, 2010).






Figure 1


A view of the title compound, showing 40% probability displacement ellipsoids and arbitrary spheres for the H atoms.

**Figure 2**

Projection along the *b*-axis of the inorganic layers in the structure of the title compound. PO₄ is given in the tetrahedral representation. Hydrogen bonds are shown as broken lines.

**Figure 3**

The packing diagram of the compound viewed down the *a*-axis. PO₄ is given in the tetrahedral representation. Hydrogen bonds are shown as broken lines.

**Figure 4**

Projection along the *b* axis of the organic chains in the structure of the title compound. Hydrogen bonds are shown as broken lines. Numbers are interplanar spacings between layers of organic molecules of type A, B and C.

Tris(5-amino-1*H*-1,2,4-triazol-4-ium) dihydrogenphosphate hydrogenphosphate trihydrate

Crystal data

$3\text{C}_2\text{H}_5\text{N}_4^+\cdot\text{HO}_4\text{P}^{2-}\cdot\text{H}_2\text{O}_4\text{P}^- \cdot 3\text{H}_2\text{O}$

$M_r = 502.31$

Monoclinic, *Pc*

Hall symbol: P -2yc

$a = 10.4793$ (13) Å

$b = 8.7655$ (11) Å

$c = 11.4536$ (14) Å

$\beta = 107.489$ (2)°

$V = 1003.5$ (2) Å³

$Z = 2$

$F(000) = 524$

$D_x = 1.662$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5573 reflections

$\theta = 3.0\text{--}31.8^\circ$

$\mu = 0.30$ mm⁻¹

$T = 100$ K

Block, colourless

$0.60 \times 0.35 \times 0.18$ mm

Data collection

Bruker SMART APEX CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2011)

$T_{\min} = 0.693$, $T_{\max} = 0.746$

13833 measured reflections

6229 independent reflections

6132 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.016$

$\theta_{\max} = 32.0^\circ$, $\theta_{\min} = 2.3^\circ$

$h = -14 \rightarrow 15$

$k = -12 \rightarrow 13$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.023$

$wR(F^2) = 0.059$

$S = 1.04$

6229 reflections

322 parameters

32 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0385P)^2 + 0.0562P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$

Absolute structure: Flack (1983), 2950 Friedel
pairs

Flack parameter: -0.02 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1A	0.62032 (11)	0.15272 (11)	0.77449 (9)	0.01772 (18)
H1A1	0.6049 (17)	0.2392 (16)	0.7386 (16)	0.021*
H1A2	0.6042 (17)	0.1391 (19)	0.8421 (13)	0.021*
N2A	0.60906 (10)	0.04376 (11)	0.58172 (9)	0.01488 (17)
H2A	0.6141	0.1293	0.5432	0.018*
N3A	0.59896 (10)	-0.09961 (11)	0.52940 (9)	0.01524 (17)
N4A	0.60216 (9)	-0.11435 (11)	0.72352 (9)	0.01336 (16)
H4A	0.6017	-0.1543	0.7939	0.016*
C1A	0.61011 (10)	0.03524 (12)	0.69840 (10)	0.01299 (18)
C2A	0.59494 (11)	-0.19168 (13)	0.61746 (10)	0.01451 (18)
H2AA	0.5879	-0.2995	0.6093	0.017*
N1B	0.95278 (11)	0.16763 (12)	0.67577 (9)	0.01933 (19)
H1B1	0.9835 (17)	0.2502 (16)	0.7144 (15)	0.023*
H1B2	0.9369 (18)	0.0936 (17)	0.7129 (16)	0.023*
N2B	0.92010 (10)	0.02876 (11)	0.48939 (9)	0.01598 (17)
H2B1	0.8958	-0.0591	0.5132	0.019*
N3B	0.93026 (11)	0.05748 (11)	0.37327 (10)	0.01706 (18)
N4B	0.98316 (9)	0.26338 (10)	0.49075 (8)	0.01343 (16)
H4B	1.0082	0.3574	0.5133	0.016*
C1B	0.95240 (10)	0.15303 (12)	0.56033 (10)	0.01362 (18)
C2B	0.96750 (11)	0.19962 (13)	0.37803 (10)	0.01543 (19)
H2B2	0.9821	0.2531	0.3110	0.019*
N1C	0.28072 (10)	0.15370 (11)	0.42295 (9)	0.01621 (17)
H1C1	0.2900 (16)	0.137 (2)	0.4966 (12)	0.019*

H1C2	0.2796 (17)	0.2424 (15)	0.3916 (15)	0.019*
N2C	0.28411 (10)	0.04801 (11)	0.23314 (8)	0.01377 (16)
H2C	0.2843	0.1345	0.1942	0.017*
N3C	0.28565 (10)	-0.09510 (11)	0.18257 (9)	0.01465 (17)
N4C	0.28158 (9)	-0.11349 (11)	0.37482 (8)	0.01238 (16)
H4C	0.2795	-0.1547	0.4443	0.015*
C1C	0.28234 (10)	0.03715 (12)	0.34900 (9)	0.01169 (17)
C2C	0.28461 (10)	-0.18892 (12)	0.27069 (10)	0.01394 (18)
H2CA	0.2858	-0.2969	0.2636	0.017*
P1A	0.23842 (2)	0.41894 (3)	0.11253 (2)	0.00876 (5)
O1A	0.09137 (7)	0.45148 (9)	0.04480 (7)	0.01176 (13)
O2A	0.31520 (8)	0.55501 (9)	0.07195 (7)	0.01273 (14)
H2AB	0.3902 (19)	0.5380 (10)	0.0918 (15)	0.019*
O3A	0.26391 (8)	0.42577 (8)	0.25172 (7)	0.01225 (14)
O4A	0.28676 (8)	0.26682 (9)	0.07706 (7)	0.01314 (14)
P1B	0.59788 (2)	0.42266 (3)	0.46934 (2)	0.00962 (5)
O1B	0.49521 (8)	0.52280 (9)	0.37270 (7)	0.01584 (15)
H1B	0.4300 (18)	0.4769 (15)	0.3450 (15)	0.024*
O2B	0.73507 (8)	0.50517 (10)	0.48323 (7)	0.01466 (14)
H2BA	0.7526 (10)	0.5008 (18)	0.4177 (17)	0.022*
O3B	0.57479 (8)	0.43008 (9)	0.59371 (7)	0.01269 (14)
O4B	0.59985 (9)	0.26114 (9)	0.42328 (7)	0.01614 (15)
O1	0.78679 (8)	0.52112 (11)	0.77991 (8)	0.01834 (16)
H1D	0.8624 (15)	0.502 (2)	0.7733 (18)	0.028*
H1E	0.7308 (17)	0.484 (2)	0.7240 (15)	0.028*
O2	1.04601 (8)	0.44288 (10)	0.79982 (7)	0.01506 (15)
H2D	1.0714 (18)	0.4441 (19)	0.8747 (12)	0.023*
H2E	1.1091 (15)	0.481 (2)	0.7840 (17)	0.023*
O3	0.91292 (8)	0.75025 (9)	0.58426 (8)	0.01633 (15)
H3D	0.8480 (15)	0.699 (2)	0.5712 (17)	0.025*
H3E	0.9680 (16)	0.6944 (19)	0.5732 (17)	0.025*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1A	0.0286 (5)	0.0117 (4)	0.0144 (4)	-0.0010 (4)	0.0086 (4)	-0.0005 (3)
N2A	0.0231 (4)	0.0110 (4)	0.0115 (4)	-0.0008 (3)	0.0067 (3)	0.0012 (3)
N3A	0.0212 (4)	0.0128 (4)	0.0127 (4)	-0.0002 (3)	0.0067 (3)	-0.0007 (3)
N4A	0.0176 (4)	0.0118 (4)	0.0114 (4)	0.0000 (3)	0.0054 (3)	0.0021 (3)
C1A	0.0142 (4)	0.0125 (5)	0.0123 (5)	-0.0003 (3)	0.0041 (3)	0.0017 (3)
C2A	0.0177 (4)	0.0134 (5)	0.0134 (5)	-0.0004 (4)	0.0061 (4)	0.0002 (3)
N1B	0.0291 (5)	0.0157 (4)	0.0145 (4)	-0.0063 (4)	0.0085 (4)	-0.0010 (3)
N2B	0.0207 (4)	0.0123 (4)	0.0155 (4)	-0.0025 (3)	0.0063 (3)	-0.0011 (3)
N3B	0.0207 (4)	0.0154 (4)	0.0154 (4)	-0.0004 (3)	0.0058 (3)	-0.0014 (3)
N4B	0.0169 (4)	0.0105 (4)	0.0129 (4)	-0.0015 (3)	0.0045 (3)	-0.0001 (3)
C1B	0.0139 (4)	0.0119 (4)	0.0146 (5)	-0.0004 (3)	0.0037 (3)	0.0000 (3)
C2B	0.0172 (4)	0.0154 (5)	0.0139 (5)	-0.0002 (4)	0.0048 (4)	-0.0006 (4)
N1C	0.0248 (4)	0.0124 (4)	0.0129 (4)	-0.0016 (3)	0.0079 (3)	-0.0013 (3)
N2C	0.0222 (4)	0.0099 (4)	0.0106 (4)	0.0010 (3)	0.0070 (3)	0.0005 (3)
N3C	0.0225 (4)	0.0101 (4)	0.0123 (4)	0.0017 (3)	0.0066 (3)	0.0002 (3)

N4C	0.0167 (4)	0.0104 (4)	0.0109 (4)	-0.0008 (3)	0.0054 (3)	0.0008 (3)
C1C	0.0135 (4)	0.0112 (5)	0.0107 (4)	-0.0004 (3)	0.0040 (3)	0.0008 (3)
C2C	0.0186 (5)	0.0113 (4)	0.0126 (4)	0.0005 (4)	0.0056 (4)	0.0006 (3)
P1A	0.01123 (10)	0.00815 (11)	0.00727 (11)	0.00019 (8)	0.00335 (8)	0.00021 (8)
O1A	0.0118 (3)	0.0126 (3)	0.0107 (3)	0.0004 (3)	0.0031 (2)	0.0007 (3)
O2A	0.0126 (3)	0.0103 (3)	0.0161 (4)	0.0001 (3)	0.0057 (3)	0.0030 (3)
O3A	0.0152 (3)	0.0137 (3)	0.0078 (3)	0.0001 (3)	0.0034 (3)	-0.0008 (2)
O4A	0.0200 (3)	0.0094 (3)	0.0113 (3)	0.0015 (3)	0.0067 (3)	-0.0003 (3)
P1B	0.01187 (10)	0.00900 (11)	0.00812 (11)	-0.00035 (8)	0.00319 (8)	-0.00054 (8)
O1B	0.0154 (3)	0.0138 (4)	0.0150 (4)	-0.0008 (3)	-0.0006 (3)	0.0036 (3)
O2B	0.0131 (3)	0.0193 (4)	0.0127 (3)	-0.0036 (3)	0.0057 (3)	-0.0023 (3)
O3B	0.0136 (3)	0.0160 (4)	0.0095 (3)	-0.0003 (3)	0.0051 (3)	-0.0022 (3)
O4B	0.0291 (4)	0.0099 (3)	0.0115 (3)	-0.0003 (3)	0.0093 (3)	-0.0011 (3)
O1	0.0131 (3)	0.0292 (5)	0.0127 (4)	-0.0018 (3)	0.0039 (3)	-0.0045 (3)
O2	0.0152 (3)	0.0208 (4)	0.0099 (3)	-0.0028 (3)	0.0047 (3)	-0.0007 (3)
O3	0.0162 (3)	0.0120 (4)	0.0225 (4)	-0.0001 (3)	0.0083 (3)	-0.0010 (3)

Geometric parameters (Å, °)

N1A—C1A	1.3325 (14)	N2C—C1C	1.3361 (13)
N1A—H1A1	0.854 (13)	N2C—N3C	1.3838 (13)
N1A—H1A2	0.849 (13)	N2C—H2C	0.8800
N2A—C1A	1.3353 (14)	N3C—C2C	1.3044 (14)
N2A—N3A	1.3826 (13)	N4C—C1C	1.3537 (14)
N2A—H2A	0.8800	N4C—C2C	1.3724 (14)
N3A—C2A	1.3022 (14)	N4C—H4C	0.8800
N4A—C1A	1.3504 (15)	C2C—H2CA	0.9500
N4A—C2A	1.3732 (14)	P1A—O4A	1.5243 (8)
N4A—H4A	0.8800	P1A—O1A	1.5294 (8)
C2A—H2AA	0.9500	P1A—O3A	1.5364 (8)
N1B—C1B	1.3272 (15)	P1A—O2A	1.5845 (8)
N1B—H1B1	0.859 (13)	O2A—H2AB	0.7639
N1B—H1B2	0.820 (13)	P1B—O4B	1.5132 (8)
N2B—C1B	1.3401 (14)	P1B—O3B	1.5163 (8)
N2B—N3B	1.3888 (14)	P1B—O1B	1.5612 (8)
N2B—H2B1	0.8800	P1B—O2B	1.5741 (8)
N3B—C2B	1.3018 (15)	O1B—H1B	0.7742
N4B—C1B	1.3524 (14)	O2B—H2BA	0.8262
N4B—C2B	1.3706 (14)	O1—H1D	0.835 (14)
N4B—H4B	0.8800	O1—H1E	0.796 (14)
C2B—H2B2	0.9500	O2—H2D	0.819 (13)
N1C—C1C	1.3304 (14)	O2—H2E	0.808 (13)
N1C—H1C1	0.833 (13)	O3—H3D	0.790 (13)
N1C—H1C2	0.855 (13)	O3—H3E	0.795 (13)
C1A—N1A—H1A1	113.9 (12)	C1C—N1C—H1C2	115.6 (11)
C1A—N1A—H1A2	119.3 (12)	H1C1—N1C—H1C2	124.9 (16)
H1A1—N1A—H1A2	120.2 (17)	C1C—N2C—N3C	110.88 (9)
C1A—N2A—N3A	111.07 (9)	C1C—N2C—H2C	124.6
C1A—N2A—H2A	124.5	N3C—N2C—H2C	124.6

N3A—N2A—H2A	124.5	C2C—N3C—N2C	104.11 (9)
C2A—N3A—N2A	104.11 (9)	C1C—N4C—C2C	106.08 (9)
C1A—N4A—C2A	106.34 (9)	C1C—N4C—H4C	127.0
C1A—N4A—H4A	126.8	C2C—N4C—H4C	127.0
C2A—N4A—H4A	126.8	N1C—C1C—N2C	125.74 (10)
N1A—C1A—N2A	125.90 (10)	N1C—C1C—N4C	127.44 (10)
N1A—C1A—N4A	127.52 (10)	N2C—C1C—N4C	106.81 (9)
N2A—C1A—N4A	106.56 (9)	N3C—C2C—N4C	112.11 (10)
N3A—C2A—N4A	111.92 (10)	N3C—C2C—H2CA	123.9
N3A—C2A—H2AA	124.0	N4C—C2C—H2CA	123.9
N4A—C2A—H2AA	124.0	O4A—P1A—O1A	113.13 (4)
C1B—N1B—H1B1	118.9 (12)	O4A—P1A—O3A	110.03 (4)
C1B—N1B—H1B2	120.0 (13)	O1A—P1A—O3A	110.70 (4)
H1B1—N1B—H1B2	120.2 (18)	O4A—P1A—O2A	109.97 (5)
C1B—N2B—N3B	110.80 (9)	O1A—P1A—O2A	103.52 (4)
C1B—N2B—H2B1	124.6	O3A—P1A—O2A	109.26 (4)
N3B—N2B—H2B1	124.6	P1A—O2A—H2AB	109.5
C2B—N3B—N2B	103.98 (9)	O4B—P1B—O3B	112.98 (5)
C1B—N4B—C2B	106.33 (9)	O4B—P1B—O1B	110.95 (5)
C1B—N4B—H4B	126.8	O3B—P1B—O1B	111.80 (5)
C2B—N4B—H4B	126.8	O4B—P1B—O2B	110.94 (5)
N1B—C1B—N2B	127.37 (10)	O3B—P1B—O2B	106.50 (4)
N1B—C1B—N4B	126.03 (10)	O1B—P1B—O2B	103.13 (5)
N2B—C1B—N4B	106.60 (10)	P1B—O1B—H1B	109.5
N3B—C2B—N4B	112.29 (10)	P1B—O2B—H2BA	109.5
N3B—C2B—H2B2	123.9	H1D—O1—H1E	109.7 (19)
N4B—C2B—H2B2	123.9	H2D—O2—H2E	101.5 (17)
C1C—N1C—H1C1	119.1 (12)	H3D—O3—H3E	104.3 (18)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1A—H1A1 \cdots O3B	0.85 (1)	2.31 (1)	3.1356 (13)	162 (2)
N1A—H1A2 \cdots N3A ⁱ	0.85 (1)	2.19 (1)	3.0305 (15)	170 (2)
N2A—H2A \cdots O4B	0.88	1.77	2.6130 (13)	161
N4A—H4A \cdots O4B ⁱ	0.88	1.76	2.6314 (13)	171
N1B—H1B1 \cdots O2	0.86 (1)	1.96 (1)	2.8214 (13)	178 (2)
N1B—H1B2 \cdots N3B ⁱ	0.82 (1)	2.28 (1)	3.0639 (15)	160 (2)
N2B—H2B1 \cdots O3 ⁱⁱ	0.88	1.84	2.6824 (13)	159
N4B—H4B \cdots O1A ⁱⁱⁱ	0.88	1.87	2.7376 (12)	167
N1C—H1C1 \cdots N3C ⁱ	0.83 (1)	2.18 (1)	3.0028 (14)	172 (2)
N1C—H1C2 \cdots O3A	0.86 (1)	2.24 (1)	3.0589 (13)	160 (2)
N2C—H2C \cdots O4A	0.88	1.78	2.6278 (12)	161
N4C—H4C \cdots O4A ⁱ	0.88	1.79	2.6645 (12)	170
O2A—H2AB \cdots O3B ^{iv}	0.76	1.95	2.6593 (11)	155
O1B—H1B \cdots O3A	0.77	1.80	2.5495 (12)	161
O2B—H2BA \cdots O1 ^{iv}	0.83	1.73	2.5552 (12)	176
O1—H1D \cdots O2	0.84 (1)	1.93 (1)	2.7439 (12)	166 (2)
O1—H1E \cdots O3B	0.80 (1)	1.91 (1)	2.6968 (12)	168 (2)

O2—H2D···O1A ^v	0.82 (1)	1.90 (1)	2.7024 (11)	168 (2)
O2—H2E···O3A ⁱⁱⁱ	0.81 (1)	1.95 (1)	2.7566 (12)	178 (2)
O3—H3D···O2B	0.79 (1)	2.14 (2)	2.8515 (12)	149 (2)
O3—H3E···O1A ⁱⁱⁱ	0.80 (1)	1.92 (1)	2.7085 (11)	176 (2)

Symmetry codes: (i) $x, -y, z+1/2$; (ii) $x, y-1, z$; (iii) $x+1, -y+1, z+1/2$; (iv) $x, -y+1, z-1/2$; (v) $x+1, y, z+1$.