

**Introduction:** Bipolar female patients have clinical and evolutionary features which involve different factors related to the women specificities.

**Objectives:** Establish clinical and evolutionary features in a population of bipolar female patients attending to Gabes psychiatry department.

**Methods:** A retrospective descriptive and analytical study was undertaken including female patients referred to psychiatry department of Gabes regional hospital, for the first time in a 6-year period (January 1<sup>st</sup>, 2010 to December 31, 2016) and who were already diagnosed with bipolar disorder (BD). Sociodemographic, clinical and evolutionary data were assessed. Patients were divided into two groups according to gender. The collected data were compared between the two groups. The statistical analysis was executed on the software SPSS (20<sup>th</sup> edition).

**Results:** From the 193 bipolar patients, 103 were women. The mean age of the disorder's onset amongst Female patients was 32.4 years old [14 - 63]. The mean duration of the disorder was 7.6 years [2 - 30]. The polarity of the first episode was a depressive one in 74.7% of cases. It was associated to psychotic features in 43.7% of cases. Seasonal pattern was noted in 10.6% amongst female patients and rapid cycling bipolar disorder in 6.2% of cases. Analytical study showed that women began the BD more often with a depressive episode ( $p=0.004$ ) and were more frequently diagnosed with BD type 2 ( $p<0.001$ ). Men had significantly more auditory ( $p=0.002$ ) and visual hallucinations ( $p=0.019$ ).

**Conclusions:** There were clinical specificities of women with BD from which important to be considered.

**Keywords:** bipolar disorder; clinical features; Gender differences; evolutionary features

## EPP0067

### Music composers and bipolar disorders: Where do we stand?

V. Giannouli\*

Institute Of Neurobiology, Bulgarian Academy of Sciences, Sofia, Bulgaria

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.537

**Introduction:** The intersection between bipolar disorders and creativity has been investigated in western literature. Although psychopathology has been proposed for famous artists in painting, writing, music and other forms of art, there is no systematic study examining bipolar disorders and music production in composers.

**Objectives:** The aim of this review is to investigate this relationship by providing an overview of published studies.

**Methods:** The search included papers published in English as abstracts as well as in full length until October 2020. The literature search was conducted using the MEDLINE, EMBASE, PUBMED and GOOGLE SCHOLAR databases. For all the searches, the terms/key words that were used were "bipolar disorder", "music", "creativity", and "composers".

**Results:** Search results for composers from different music genres and musical periods indicated that the proposed origin of the overall bipolar pattern is attributed to several stressful environmental factors which are taking the form of interpersonal problems regarding the expressed emotion, life events, and paucity of stress-management skills. In addition to that, bipolar psychopathological

patterns seem to influence the quantity of music composing activity indirectly due to changes in everyday functional abilities.

**Conclusions:** Published reports, although based on biographical research, do provide evidence in support of a strong bipolarity-music creativity/production link for famous composers. Further well-designed studies in living music professionals engaged in music composition are needed.

**Keywords:** Creativity; Bipolar Disorders; Music

## EPP0070

### Pediatric bipolar disorder: Preliminary results of a retrospective study using a nationwide administrative database

A. Campos<sup>1\*</sup>, M. Gonçalves-Pinho<sup>2,3,4</sup>, A.R. Ferreira<sup>2</sup>, A. Freitas<sup>2,4</sup> and L. Fernandes<sup>1,2,5</sup>

<sup>1</sup>Faculty Of Medicine, University of Porto, Porto, Portugal; <sup>2</sup>Cintesis – Center For Health Technology And Services Research, Faculty of Medicine, University of Porto, Porto, Portugal; <sup>3</sup>Department Of Psychiatry And Mental Health, Centro Hospitalar do Tâmega e Sousa, Penafiel, Portugal, Penafiel, Portugal; <sup>4</sup>Department Of Community, Information And Health Decision Sciences (medcids), Faculty of Medicine, University of Porto, Porto, Portugal, Porto, Portugal and <sup>5</sup>Psychiatry Service, Centro Hospitalar Universitário de São João, Porto, Portugal, Porto, Portugal

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.538

**Introduction:** Bipolar disorder (BD) is a severe and chronic illness characterized by episodic changes in mood. The average onset of BD symptoms is estimated between 18 and 22 years. However, many adults retrospectively report symptoms onset in childhood or adolescence. Over the last decades, pediatric bipolar disease (PBD) has been the focus of increased attention mainly due to controversies surrounding its prevalence, diagnosis and treatment in the pediatric population.

**Objectives:** To analyze pediatric hospitalizations related to BD held in mainland Portuguese public hospitals between 2000 and 2015.

**Methods:** This retrospective observational study analyzed all pediatric (<18 years old) inpatient episodes from 2000 to 2015 with a primary BD diagnosis, using an anonymized administrative database including all hospitalization from mainland Portuguese public hospitals. ICD-9-CM codes 296.x were used (excluding codes 296.2x; .3x and .9x). Age at admission, admission type and date, sex, charges and length of stay (LoS) were analyzed.

**Results:** A total of 348 hospitalizations were analyzed from 258 patients. Patients were mainly young girls (60.6%), with a mean age of  $15.24 \pm 1.87$  years. The majority of the admissions were urgent (81.0%), and the median LoS was 14 days (IQR: 7; 24). Mean hospitalization charges were 3503.1€ with a total sum of 1.2M€ for all the episodes.

**Conclusions:** PBD hospitalizations occur predominantly in female patients during adolescence. The majority of them are urgent admissions. Descriptive studies will help to describe and characterize socio-demographic and clinical trends in PBD in order to better prevent acute hospitalizations with inevitable social and economic implications.

**Keywords:** bipolar disorder; Hospital admissions; adolescents; epidemiology