

# Integrative analysis of ferroptosis-related genes in ulcerative colitis

De-jun Cui\* , Chen Chen\*, Wen-qiang Yuan, Yun-han Yang and Lu Han

## Abstract

**Objective:** The aim of this study was to identify and validate ferroptosis-related markers in ulcerative colitis (UC) to explore new directions for UC diagnosis and treatment.

**Methods:** We screened UC chips and ferroptosis-related genes from the Gene Expression Omnibus (GEO), FerrDb, and GeneCards databases. The differentially expressed genes (DEGs) and ferroptosis-related DEGs between the UC group and normal controls were analyzed using bioinformatics methods. Enrichment analysis, protein–protein interaction analysis, and hub genes were screened. Peripheral blood chip and animal experiments were used to validate the ferroptosis-related hub genes. Finally, hub gene–transcription factor, hub gene–microRNA (miRNA), and hub gene–drug interaction networks were constructed.

**Results:** Overall, 26 ferroptosis-related DEGs were identified that were significantly enriched in energy pathways and metabolism. We identified ten ferroptosis-related hub genes from the protein–protein interaction network: *IL6*, *PTGS2*, *HIF1A*, *CD44*, *MUC1*, *CAVI*, *NOS2*, *CXCL2*, *SCD*, and *ACSL4*. In the peripheral blood chip GSE94648, *CD44* and *MUC1* were upregulated, which was consistent with the expression trend in GSE75214. Animal experiments showed that *CD44* expression was significantly increased in the colon.

**Conclusions:** Our findings indicate that *CD44* and *MUC1* may be ferroptosis-related markers in UC.

## Keywords

Gene expression omnibus database, ulcerative colitis, ferroptosis, biomarker, bioinformatics, hub genes

Date received: 11 May 2021; accepted: 21 July 2021

\*These authors contributed equally to this work.

### Corresponding author:

De-jun Cui, Department of Gastroenterology, Guizhou Provincial People's Hospital, No. 83, East Zhongshan Road, Guiyang, Guizhou Province 550002, PR China.  
Email: [hxcuidj@163.com](mailto:hxcuidj@163.com)

Department of Gastroenterology, National Institution of Drug Clinical Trial, Guizhou Provincial People's Hospital, Medical College of Guizhou University, Guiyang, China



## Introduction

Ulcerative colitis (UC) is a chronic relapsing colorectal inflammatory disease with unknown etiology. The incidence of UC is increasing globally, but it is difficult to cure. Approximately 15% of UC patients will experience an aggressive course, and some will even develop dysplasia and colorectal cancer (CRC). The current research on the pathogenesis of UC has mainly focused on genetics, microbiota, immunity, environment, and the intestinal mucosal barrier.<sup>1-4</sup> However, the exact pathogenesis of UC is still not fully understood. Therefore, exploring the etiology and pathogenesis of UC is of great significance to the diagnosis and treatment of this disease.

Recent evidence indicates that ferroptosis is closely related to the pathogenesis of UC.<sup>5,6</sup> Ferroptosis is a recently discovered type of regulated cell death, and its occurrence is associated with the excessive accumulation of iron-dependent lipid reactive oxygen species (ROS) and consumption of polyunsaturated fatty acids in the plasma membrane.<sup>6,7</sup> Excessive iron aggravates dextran sulfate sodium (DSS) colitis in rats, and a human diet high in iron also increases the risk of UC.<sup>8,9</sup> Iron chelators reduce ROS and improve UC.<sup>10</sup> However, the anti-inflammatory drug sulfasalazine, used for the treatment of UC, can also induce ferroptosis in cells by inhibiting the function of the antiporter system Xc<sup>-</sup>.<sup>7</sup> This may be one cause of adverse reactions to sulfasalazine. Regulating ferroptosis can ameliorate DSS-induced UC.<sup>11,12</sup> However, the current research on ferroptosis in UC is still at an early stage, and the role of ferroptosis in the pathogenesis of UC is not fully understood.<sup>6</sup> Some studies have reported contrasting results. Uncovering and studying UC ferroptosis-related markers may provide new directions for UC diagnosis and treatment.

Here, we searched for ferroptosis-related genes in UC to address the lack of new ferroptosis-specific pharmacodynamic markers or biomarkers.<sup>13</sup> The differentially expressed genes (DEGs) related to ferroptosis in UC were enriched and analyzed, and a protein-protein interaction (PPI) network was constructed. Then, we screened for ferroptosis-related hub genes. Peripheral blood chip and animal experiments were used to validate these ferroptosis-related hub genes, and then microRNAs (miRNAs), transcription factors (TFs), or drugs targeting the hub genes were predicted.

## Materials and methods

### Screening of microarray data and ferroptosis-related genes

We searched Gene Expression Omnibus (GEO) (<https://www.ncbi.nlm.nih.gov/geo/>), the ferroptosis-related database FerrDb (<http://www.zhounan.org/ferrdb/index.html>), and the GeneCards database (<https://www.genecards.org/>) to find microarray data and ferroptosis-related genes.<sup>14</sup> GEO search formula: (((inflammatory bowel diseases) OR Colitis, Ulcerative) OR Crohn Disease) OR colitis) AND "Homo sapiens". The driver, suppressor, and marker genes related to ferroptosis were downloaded from FerrDb. The GeneCards database was searched using "ferroptosis". The GEO inclusion criteria were human UC intestinal biopsy specimens or peripheral blood specimens, and more than eight cases. The exclusion criterion was colitis-associated CRC.

### Identification of DEGs related to ferroptosis

GEO2R was used for GEO chips. The parameter options used were: 1) apply adjustment to the *P*-values: Benjamini and Hochberg (false discovery rate); 2) apply

log transformation to the data: auto-detect; 3) apply limma precision weights (vooma): yes; 4) force normalization or quantile normalization: yes; 5) category of platform annotation to display on results: NCBI generated; 6) significance level cut-off: 0.05; 7) DEGs filter criteria:  $|\log$  fold change (FC)  $\geq 1$  and adj.  $P$ -value  $< 0.05$ . SangerBox software (<http://soft.sangerbox.com/>) was used to generate volcano maps. The abscissa was  $-\log_{10}$  ( $P$ -value), the ordinate was  $\log_2$ (FC), cut FC ( $-1.0$  to  $1.0$ ), and cut  $P$ -value: 0.05. The red nodes represent upregulated DEGs, the green nodes are downregulated DEGs, and the black nodes are genes with no difference. Then, the DEGs and the ferroptosis-related genes from FerrDb and GeneCards were intersected using online Bioinformatics and Evolutionary Genomics to generate a Venn diagram (<http://bioinformatics.psb.ugent.be/webtools/Venn/>).

### Gene enrichment analysis

FunRich-3.1.3 tool (<http://www.funrich.org/>) was used for gene enrichment analysis, including cell components, molecular functions, and biological processes. Based on Kyoto Encyclopedia of Genes and Genomes (KEGG), Enrichr (<https://maayanlab.cloud/Enrichr/>) was used for pathway analysis. A  $P$ -value less than 0.05 was defined as statistically significant.<sup>15,16</sup>

### Construction of PPI network and identification of hub genes

A PPI network was constructed by GeneMANIA software (<http://genemania.org/>) using the automatically selected weighting method. Different colored edges represent different bioinformatics methods, including co-expression, physical interactions, colocalization, predicted pathways, shared protein domains, and genetic interactions.

The ferroptosis-related PPI network was constructed with STRING (version 11.0, <https://string-db.org/>). The organism was limited to *Homo sapiens*. The minimum effective binding score was set to 0.4. Then, the PPI results were imported into Cytoscape (version 3.6.0) for software analysis.<sup>17</sup> CytoHubba in Cytoscape was used to screen the top ten hub genes, and the algorithm used was the degree of connectivity.<sup>18</sup>

### Validation of hub genes

The peripheral blood chip GSE94648 was used to validate the hub genes. GraphPad Prism v 5.0 statistical software was used to analyze the expression of the hub genes in peripheral blood. The statistical method used was an unpaired t test, and differences were considered statistically significant with  $P$ -values  $< 0.05$ . The hub gene verification method used was the same as that in our previous study.<sup>19</sup> Then, the statistically significant hub genes were further verified with animal experiments. The experimental animals were male BALB/c mice purchased from Tengxin Biotechnology (Chongqing, China), aged 6 to 8 weeks and weighing 18 to 22 g. Sixteen mice were randomly divided into two groups of eight animals each. The animal experiments were approved by the Ethics Committee of Guizhou Provincial People's Hospital on 8 April 2020 (No. 2020-256). We used 3% DSS (molecular weight 36–50 kDa; MP Biomedicals, Aurora, CO, USA) to induce a model of UC. Immunohistochemistry (IHC) was used to detect the expression of the hub genes in colon tissues. A rabbit anti-MUC1 antibody (1:200, ABclonal, Wuhan, China) and a rabbit anti-CD44 antibody (1:200, Affinity Biosciences, Cincinnati, OH, USA) were used as primary antibodies for IHC. A goat anti-rabbit horseradish peroxidase-conjugated secondary antibody (Beijing Zhongshan Jinqiao

Biotechnology, Beijing, China) was used for detection. Analysis of variance (ANOVA) was used for statistical analysis. The experimental method was the same as that in our previous study.<sup>20</sup>

### Construction of miRNA (or TF)–hub gene network

The target genes and miRNA or TF interaction networks were generated by NetworkAnalyst (<https://www.networkanalyst.ca/>).<sup>21</sup> The parameters used were as follows: specify organism: *H. sapiens* (human); set ID type: Official Gene Symbol; Gene–miRNA interaction database: miRTarBase v8.0, TarBase v8.0 and miRecords databases; and TF–gene interaction database: ENCODE, JASPAR, and ChEA databases. The analysis results of NetworkAnalyst were then imported into Cytoscape software for further visualization.

### Potential drug prediction


Potential drugs targeting the hub genes were predicted by The Drug Gene Interaction Database (DGIdb, <https://dgidb.genome.wustl.edu/>). The parameters

used were: search for drug–gene interactions by gene; preset filters: none; advanced filters: none. Then the drug–gene interactions found in DGIdb were imported into Cytoscape software for further visualization.

## Results

### Ferroptosis-related DEGs in UC

We searched the GEO database to obtain 9984 data points from inflammatory bowel disease-related chips. Further screening resulted in two chips: GSE75214 and GSE94648. GSE75214, containing mucosal gene expression profiling of the colon, included 97 UC patients and 11 controls. GSE94648, used for validating the hub genes found in colon tissue, contained peripheral whole blood expression data of 22 healthy controls and 25 UC patients. We identified 545 upregulated DEGs and 300 downregulated DEGs from GSE75214, as shown in Figure 1a. There were 259 ferroptosis-related genes in FerrDb and 103 ferroptosis-related genes in GeneCards. After finding overlapping genes between



**Figure 1.** The differentially expressed genes (DEGs). (a) Volcano plot of GSE75214. The cutoff logFC (fold change) was  $-1/1$ , and  $P$ -value is 0.05. Red dots indicate upregulated genes; green dots indicate downregulated genes; black dots indicate non-statistically significant genes. (b) Venn diagram of the ferroptosis-related DEGs.

the GSE75214 DEGs and the FerrDb and GeneCards genes, 26 ferroptosis-related DEGs were obtained, as shown in Figure 1b.


### Results of enrichment analysis

The results of the GO enrichment analysis indicated that for cell components, the ferroptosis-related DEGs were enriched for the terms integral to membrane, integral to plasma membrane, extracellular, mitochondrion, lysosome, endoplasmic reticulum, nucleus, exosomes, cytoplasm, and plasma membrane. In terms of the molecular functions, the ferroptosis-related DEGs were mainly enriched in chemokine activity, ligase activity, cytokine activity, oxidoreductase activity, auxiliary transport protein activity, peroxidase activity, transcription factor activity, receptor activity, catalytic activity, and molecular function unknown. For biological processes, the ferroptosis-related DEGs were mainly enriched in cell communication, apoptosis, regulation of nucleobase, nucleoside, nucleotide and nucleic acid metabolism, cell growth and/or maintenance, transport, immune response, signal transduction, biological process unknown, energy pathways, and metabolism.

nucleobase and its analogue, cell growth and/or maintenance, transport, immune response, signal transduction, and biological process unknown. They were particularly significantly enriched in energy pathways and metabolism ( $P < 0.05$ ). The KEGG enrichment results indicated that the ferroptosis-related DEGs were mainly enriched in biosynthesis of unsaturated fatty acids, legionellosis, leishmaniasis, the PPAR signaling pathway, ferroptosis, fatty acid biosynthesis, salmonella infection, the interleukin-17 (IL-17) signaling pathway, the hypoxia inducible factor-1 (HIF-1) signaling pathway, and the tumor necrosis factor (TNF) signaling pathway ( $P < 0.05$ ) (Figure 2).


### PPI and hub genes

The PPI network constructed with GeneMANIA is shown in Figure 3a. In the PPI network, co-expression accounts for 68.60%, physical interactions account



**Figure 2.** Enrichment analysis of the ferroptosis-related differentially expressed genes (DEGs).

(a) Cell component analysis for the ferroptosis-related DEGs. (b) Molecular functions analysis for the ferroptosis-related DEGs. (c) Biological processes analysis for the ferroptosis-related DEGs. (d) Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis for the ferroptosis-related DEGs.



**Figure 3.** The protein–protein interaction (PPI) network and the ferroptosis-related hub genes. (a) The PPI network constructed with GeneMANIA. (b) The top ten hub genes. The red nodes indicate higher degrees of connectivity and the yellow nodes indicate lower degrees.

for 14.07%, colocalization accounts for 10.50%, predicted accounts for 2.65%, pathway accounts for 2.31%, shared protein domains account for 1.62%, and genetic interactions account for 0.24%. The functions of the ferroptosis-related DEGs are shown in Figure 3a.

Using the STRING tool, we obtained 26 nodes, 42 edges, and an average node degree of 3.23 in the PPI network. The top ten hub genes were obtained using the cytoHubba plugin in Cytoscape to calculate the degree of connectivity, as shown in Figure 3b and Table 1.

### Validation of hub genes in peripheral blood and animal experiments

The peripheral blood chip GSE94648 was used to validate hub gene expression in peripheral blood. The results showed that the expression levels of *CD44* and cell surface associated mucin 1 (*MUC1*) in peripheral blood were significantly altered ( $P < 0.05$ , Figure 4a and b). In addition, *CD44* and *MUC1* were both upregulated in GSE94648 and GSE75214. The animal

experiments suggested that the expression of *CD44* in the DSS-induced colitis model was significantly different from the control ( $P < 0.05$ ), while the expression of *MUC1* showed no significant difference (Figure 4c).


### Predictive results of miRNA, TF, or drugs targeting the hub genes

The target gene–miRNA network of the hub genes *CD44* and *MUC1* was constructed, which showed that hsa-miR-125b-5p and hsa-miR-145-5p could regulate *CD44* and *MUC1* expression (Figure 5a). The target gene–TF network of the hub genes *CD44* and *MUC1* was constructed, which identified Sin3A associated protein 30 (*SAP30*), tripartite motif containing 22 (*TRIM22*), zinc finger protein 24 (*ZNF24*), lysine demethylase 5B (*KDM5B*), PHD finger protein 8 (*PHF8*), and SIN3 transcription regulator family member A (*SIN3A*) as modulators of *CD44* and *MUC1* (Figure 5b).

Using DGIdb to predict drugs that can target the hub genes, five potential drugs or compounds for *CD44* were found, including

**Table 1.** Top ten hub genes with higher scores.

| Rank | Gene symbol | Score | Regulation in GSE75214 | Description                                    |
|------|-------------|-------|------------------------|--|
| 1    | IL6         | 13    | up                     | interleukin 6                                  |
| 2    | PTGS2       | 9     | up                     | prostaglandin-endoperoxide synthase 2          |
| 3    | HIF1A       | 8     | up                     | hypoxia inducible factor 1 alpha               |
| 4    | CD44        | 6     | up                     | CD44 molecule (Indian blood group)             |
| 5    | MUC1        | 5     | up                     | mucin 1, cell surface associated               |
| 5    | CAVI        | 5     | up                     | caveolin 1                                     |
| 5    | NOS2        | 5     | up                     | nitric oxide synthase 2                        |
| 8    | CXCL2       | 4     | up                     | C-X-C motif chemokine ligand 2                 |
| 8    | SCD         | 4     | up                     | stearoyl-CoA desaturase                        |
| 10   | ACSL4       | 3     | up                     | acyl-CoA synthetase long chain family member 4 |




**Figure 4.** Validation of hub genes in peripheral blood and animal experiments. (a) Expression of CD44 in peripheral blood ( $P < 0.05$ ). (b) Expression of MUC1 in peripheral blood ( $P < 0.05$ ). (c) Expression of CD44 in colon tissue of BALB/c mice (immunohistochemistry,  $\times 400$ ). Cytoplasmic brown is positive expression UC, ulcerative colitis; DSS, dextran sulfate sodium.

gentamicin, progesterone, hyaluronan, hyaluronate sodium, and bivatuzumab (Figure 5c). In addition, four potential drugs or compounds for *MUC1* were predicted, including HuHMGF1, SAR566658, emepipimut-S, and GO-203-2c (Figure 5d).

## Discussion

Intestinal mucosal barrier damage can occur before a clinical diagnosis of UC. This is a variable factor in the pathogenesis of UC, and may also provide a future strategy for preventing this disease.<sup>4</sup> Recent



**Figure 5.** Predictive networks of microRNAs (miRNAs), transcription factors (TFs), or drugs targeting the hub genes. (a) Target gene-miRNA network. (b) Target gene-TF network. The red circle nodes represent hub genes and the yellow squares represent miRNAs or TFs. (c) Drug-CD44 interaction network. (d) Drug-MUC1 interaction network. The red circle nodes are the hub genes and the yellow diamond nodes are the predicted drugs or molecular compounds.

studies have found that ferroptosis is closely related to intestinal mucosal barrier dysfunction, p53, and the immune system.<sup>5,6,22</sup> Thus, it is important to explore the ferroptosis-related markers of intestinal mucosa injury and immunotherapy in UC. Using bioinformatics analysis, peripheral blood chip, and animal experiments, we found *MUC1* and *CD44* to be ferroptosis-related hub genes in UC. This will provide a basis for studying UC pathogenesis and identifying potential novel treatments.

*MUC1*, secreted by intestinal goblet and absorptive cells, is a membrane-bound glycoprotein in the mucus layer. *MUC1* protein expression increases during intestinal inflammation to protect the intestinal

mucosal barrier, and plays an important role in UC pathogenesis.<sup>23</sup> This gene can regulate cell signal transduction and participate in immune regulation. *MUC1* expression was shown to significantly increase during the transition from colitis to tumorigenesis, but the specific mechanism of action was unclear.<sup>24</sup> A recent study reported that *MUC1* is related to ferroptosis.<sup>25</sup> Our research also found that the expression of *MUC1* was upregulated in UC, which is possibly related to ferroptosis. However, further research is necessary because an increase in *MUC1* expression was not observed in animal experiments.

The protein encoded by the *CD44* gene is a cell surface glycoprotein involved in



cell–cell interactions, cell adhesion, and migration. A recent study found that CD44 expression inhibited ferroptosis.<sup>26</sup> Interestingly, multi-bioresponsive silk fibroin-based nanoparticles with the capacity for on-demand cytoplasmic drug release targeted to CD44 could alleviate UC.<sup>27</sup> In our study, the *CD44* expression levels were significantly increased in human colon tissue and peripheral blood. In our animal experiments, *CD44* expression also increased significantly, which is consistent with previous research reports. However, the relationship between *CD44* and ferroptosis proteins requires further study.

Our data suggest that the ferroptosis-related markers *MUC1* and *CD44* are related to the pathogenesis of UC. However, this study has some limitations, including a lack of unique morphological characteristics of ferroptosis and lipid ROS, a lack of research on the effect of ferroptosis-related markers on immunity, and the possibility of selection bias in the inclusion of the chips. To improve the prevention, diagnosis, treatment, and prognosis of UC, future research is necessary to identify ferroptosis-specific pharmacodynamic markers or small-molecule ferroptosis modulators, as well as to study the role of ferroptosis in intestinal mucosal barrier damage, the immune response, an iron-containing diet, and carcinogenesis.

### Declaration of conflicting interest


The authors declare that there is no conflict of interest.

### Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was financially supported by Cultivation Fund of National Natural Science Foundation (Grant No. qiankehe2018-5764-11), Doctor Foundation of Guizhou Provincial People's Hospital [No. GZSYBS(2017)09], and Beijing Medical and

Health Foundation [No. YWJKJJKYJJ-B184053].

### ORCID iD

De-jun Cui  <https://orcid.org/0000-0002-3653-3058>

### References

1. Kobayashi T, Siegmund B, Le Berre C, et al. Ulcerative colitis. *Nat Rev Dis Primers*. 2020; 6: 74. DOI: 10.1038/s41572-020-0205-x.
2. Feuerstein JD, Isaacs KL, Schneider Y, et al. AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. *Gastroenterology*. 2020; 158:1450–1461. DOI: 10.1053/j.gastro.2020.01.006.
3. Rubin DT, Ananthakrishnan AN, Siegel CA, et al. ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol*. 2019; 114: 384–413. DOI: 10.14309/ajg.0000000000000152.
4. Mehandru S and Colombel JF. The intestinal barrier, an arbitrator turned provocateur in IBD. *Nat Rev Gastroenterol Hepatol*. 2021; 18: 83–84. DOI: 10.1038/s41575-020-00399-w.
5. Xu M, Tao J, Yang Y, et al. Ferroptosis involves in intestinal epithelial cell death in ulcerative colitis. *Cell Death Dis*. 2020; 11: 86. DOI: 10.1038/s41419-020-2299-1.
6. Xu S, He Y, Lin L, et al. The emerging role of ferroptosis in intestinal disease. *Cell Death Dis*. 2021; 12: 289. DOI: 10.1038/s41419-021-03559-1.
7. Dixon SJ, Lemberg KM, Lamprecht MR, et al. Ferroptosis: an iron-dependent form of nonapoptotic cell death. *Cell*. 2012; 149: 1060–1072. DOI: 10.1016/j.cell.2012.03.042.
8. Carrier JC, Aghdassi E, Jeejeebhoy K, et al. Exacerbation of dextran sulfate sodium-induced colitis by dietary iron supplementation: role of NF-kappaB. *Int J Colorectal Dis*. 2006; 21: 381–387. DOI: 10.1007/s00384-005-0011-7.
9. Kobayashi Y, Ohfuji S, Kondo K, et al. Association between dietary iron and zinc intake and development of ulcerative colitis: A case-control study in Japan.

- J Gastroenterol Hepatol.* 2019; 34: 1703–1710. DOI: 10.1111/jgh.14642.
10. Millar AD, Rampton DS and Blake DR. Effects of iron and iron chelation in vitro on mucosal oxidant activity in ulcerative colitis. *Aliment Pharmacol Ther.* 2000; 14: 1163–1168. DOI: 10.1046/j.1365-2036.2000.00828.x.
  11. Chen Y, Zhang P, Chen W, et al. Ferroptosis mediated DSS-induced ulcerative colitis associated with Nrf2/HO-1 signaling pathway. *Immunol Lett.* 2020; 225: 9–15. DOI: 10.1016/j.imlet.2020.06.005.
  12. Wang S, Liu W, Wang J, et al. Curculigoside inhibits ferroptosis in ulcerative colitis through the induction of GPX4. *Life Sci.* 2020; 259: 118356. doi: 10.1016/j.lfs.2020.118356.
  13. Conrad M, Lorenz SM, Proneth B. Targeting Ferroptosis: New Hope for As-Yet-Incurable Diseases. *Trends Mol Med.* 2021; 27: 113–122. DOI:10.1016/j.molmed.2020.08.010.
  14. Zhou N and Bao J. FerrDb: a manually curated resource for regulators and markers of ferroptosis and ferroptosis-disease associations. *Database (Oxford).* 2020; 2020: baaa021. DOI: 10.1093/database/baaa021.
  15. Chen EY, Tan CM, Kou Y, et al. Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics.* 2013; 14: 128. DOI: 10.1186/1471-2105-14-128.
  16. Kuleshov MV, Jones MR, Rouillard AD, et al. Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res.* 2016; 44: W90–W97. DOI: 10.1093/nar/gkw377.
  17. Shannon P, Markiel A, Ozier O, et al. Cytoscape: a software environment for integrated models of Biomolecular interaction networks. *Genome Res.* 2003; 13: 2498–2504. DOI:10.1101/gr.1239303.
  18. Chin CH, Chen SH, Wu HH, et al. cytoHubba: identifying hub objects and sub-networks from complex interactome. *BMC Syst Biol.* 2014; 8: S11. DOI: 10.1186/1752-0509-8-S4-S11.
  19. Cui D, Yuan W, Chen C, et al. Identification of colorectal cancer-associated macrophage biomarkers by integrated bioinformatic analysis. *Int J Clin Exp Pathol.* 2021; 14: 1–8.
  20. Cui DJ, Yang XL, Okuda S, et al. Gallincin ameliorates colitis-associated inflammation and barrier function in mice based on network pharmacology prediction. *J Int Med Res.* 2020; 48: 300060520951023. DOI: 10.1177/0300060520951023.
  21. Zhou G, Soufan O, Ewald J, et al. NetworkAnalyst 3.0: a visual analytics platform for comprehensive gene expression profiling and meta-analysis. *Nucleic Acids Res.* 2019; 47: W234–W241. DOI: 10.1093/nar/gkz240.
  22. Xu S, Min JX, Wang FD. Ferroptosis: an emerging player in immune cells. *Science Bulletin.* 2021. DOI: 10.1016/j.scib.2021.02.026.
  23. Vancamelbeke M, Vanuytsel T, Farré R, et al. Genetic and Transcriptomic Bases of Intestinal Epithelial Barrier Dysfunction in Inflammatory Bowel Disease. *Inflamm Bowel Dis.* 2017; 23: 1718–1729. DOI: 10.1097/MIB.0000000000001246.
  24. Sun Y, Fan L, Mian W, et al. Modified apple polysaccharide influences MUC-1 expression to prevent ICR mice from colitis-associated carcinogenesis. *Int J Biol Macromol.* 2018; 120(Pt B): 1387–1395. DOI: 10.1016/j.ijbiomac.2018.09.142.
  25. Hasegawa M, Takahashi H, Rajabi H, et al. Functional interactions of the cystine/glutamate antiporter, CD44v and MUC1-C oncoprotein in triple-negative breast cancer cells. *Oncotarget.* 2016; 7: 11756–11769. DOI: 10.18632/oncotarget.7598.
  26. Liu T, Jiang L, Tavana O, et al. The Deubiquitylase OTUB1 Mediates Ferroptosis via Stabilization of SLC7A11. *Cancer Res.* 2019; 79: 1913–1924. DOI: 10.1158/0008-5472.CAN-18-3037.
  27. Gou S, Huang Y, Wan Y, et al. Multi-bioresponsive silk fibroin-based nanoparticles with on-demand cytoplasmic drug release capacity for CD44-targeted alleviation of ulcerative colitis. *Biomaterials.* 2019; 212: 39–54. DOI: 10.1016/j.biomaterials.2019.05.012.