





Review Article

Effects of Polyphenols on Oxidative Stress, Inflammation, and Interconnected Pathways during Spinal Cord Injury

Sajad Fakhri ¹, Fatemeh Abbaszadeh,^{2,3} Seyed Zachariah Moradi ^{1,4}, Hui Cao,⁵
Haroon Khan ⁶, and Jianbo Xiao ^{5,7}

¹Pharmaceutical Sciences Research Center, Health Institute, Kermanshah University of Medical Sciences, Kermanshah 6734667149, Iran

²Department of Neuroscience, Faculty of Advanced Technologies in Medical Sciences, Iran University of Medical Sciences, Tehran, Iran

³Neurobiology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴Medical Biology Research Center, Health Technology Institute, Kermanshah University of Medical Sciences, Kermanshah 6734667149, Iran

⁵Nutrition and Bromatology Group, Department of Analytical Chemistry and Food Science, Faculty of Food Science and Technology, University of Vigo-Ourense Campus, E-32004 Ourense, Spain

⁶Department of Pharmacy, Abdul Wali Khan University Mardan, 23200, Pakistan

⁷Institute of Food Safety and Nutrition, Jinan University, Guangzhou 510632, China

Correspondence should be addressed to Sajad Fakhri; sajad.fakhri@kums.ac.ir, Haroon Khan; haroonkhan@awkum.edu.pk, and Jianbo Xiao; jianboxiao@uvigo.es

Received 15 October 2021; Accepted 11 December 2021; Published 7 January 2022

Academic Editor: Dragan Hrnčić

Copyright © 2022 Sajad Fakhri et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Despite the progression in targeting the complex pathophysiological mechanisms of neurodegenerative diseases (NDDs) and spinal cord injury (SCI), there is a lack of effective treatments. Moreover, conventional therapies suffer from associated side effects and low efficacy, raising the need for finding potential alternative therapies. In this regard, a comprehensive review was done regarding revealing the main neurological dysregulated pathways and providing alternative therapeutic agents following SCI. From the mechanistic point, oxidative stress and inflammatory pathways are major upstream orchestras of cross-linked dysregulated pathways (e.g., apoptosis, autophagy, and extrinsic mechanisms) following SCI. It urges the need for developing multitarget therapies against SCI complications. Polyphenols, as plant-derived secondary metabolites, have the potential of being introduced as alternative therapeutic agents to pave the way for treating SCI. Such secondary metabolites presented modulatory effects on neuronal oxidative stress, neuroinflammatory, and extrinsic axonal dysregulated pathways in the onset and progression of SCI. In the present review, the potential role of phenolic compounds as critical phytochemicals has also been revealed in regulating upstream dysregulated oxidative stress/inflammatory signaling mediators and extrinsic mechanisms of axonal regeneration after SCI in preclinical and clinical studies. Additionally, the coadministration of polyphenols and stem cells has shown a promising strategy for improving post-SCI complications.

1. Introduction

As pivotal causes of disability and death, neurological diseases (NDDs) increasingly affect millions of people worldwide. Despite the advances in revealing multiple dysregulated pathways in the pathogenesis of NDDs, their main

pathophysiological mechanisms have remained to be elucidated [1, 2]. Spinal cord injury (SCI) is a NDD with sensory-motor dysfunctionalities, which extremely reduces the quality of life. The worldwide prevalence of SCI is on the rise, more in 14.6-67.6-year-old individuals and four times more prevalent in men [3, 4]. In the United States,

the annual incidence of traumatic SCI is estimated at about 12,000 new cases per year, equates to 40 cases per million of population, and bears high costs [5].

From the pathophysiological point of view, SCI involves primary and secondary phases. While the primary injury occurs after the spinal mechanical trauma, the secondary phase consists of intrinsic oxidative stress, inflammatory, apoptotic, and autophagic cascades [6]. On the other hand, extrinsic pathways also play critical roles in the progression of SCI, including glial scar formation and axonal degeneration [7]. Of intrinsic pathways, oxidative stress, neuroinflammation, and neuroapoptosis are interconnected with extrinsic pathways (e.g., axonal signaling). Thus, the aforementioned pathological pathways play destructive roles in neuronal cell death and neurodegenerative processes, leading to cell death. The modulatory effects of antioxidant defenses on neuroinflammatory and neuroapoptotic responses play key roles in the onset and progression of SCI via affecting microglia, astrocytes, and associated mediators [8, 9].

Despite progressions in clinical healthcare, prevention/treatment of SCI has remained a clinical challenge. This challenge could be due to the complex pathophysiological mechanisms of SCI, which raises the need for developing novel multitarget therapeutics. Such treatments could target both the intrinsic and extrinsic pathways [3].

It is worth mentioning that providing new drugs from plants has a long and successful history in complementary and alternative medicine. In this line, the plant kingdom has shown promising results against SCI. Among natural entities, polyphenols/phenolic compounds are multitarget and accessible phytochemicals with high specificity and low toxicity levels. They are currently used in modern medicine to design and develop new therapeutic agents because of their extensive biological activities and health benefits. These compounds have been considered safe dietary agents with potential inhibitory effects against oxidative stress and inflammation in various NDDs [3, 10]. In recent years, growing evidence has focused on the administration of natural neuroprotective polyphenols with potential antioxidant activities to manage SCI and other NDDs [11].

Previous reviews indicated the crucial role of oxidative stress and axonal dysregulations in the pathogenesis of Parkinson's disease [11], epilepsy [12], and some other NDDs [13]. Accordingly, cytokines [14], nitric oxide (NO), antioxidant response element (ARE) [15], and other oxidative stress/inflammatory mediators, as well as extrinsic dysregulated pathways, seem to be potential regulators of such pathological conditions. We have also previously provided the role of apoptosis and autophagy following SCI [3]; however, no focus has been yet done on the pivotal role of oxidative stress after SCI. Therefore, the present study comprehensively reviews oxidative stress pathways and their effects on the inflammatory and apoptotic pathways during SCI. The modulatory effects of polyphenols/phenolic compounds on the intrinsic upstream oxidative stress pathways and extrinsic axonal pathways have also been revealed following SCI. Besides, the coadministration of polyphenols and stem cells has been shown as a new gate in combating post-SCI complications.

2. The Role of Oxidative Stress, Inflammation, and Apoptosis after SCI: Intrinsic Pathway

Oxidative stress has been described as a disturbance and perturbation in the balance among the production and manifestation of reactive oxygen species (ROS) and the biological antioxidant defense ability. It causes toxic effects, tissue injury, inflammation, apoptosis, autophagy, and necrosis via the generation of free radicals and peroxides, which damages DNA, proteins, lipids, and carbohydrates [16]. ROS presuppose both the free radical and nonfree radical oxygen moderators like superoxide, hydrogen peroxide (H_2O_2), hydroxyl radical ($\cdot OH$), and singlet oxygen. Even though ROS could operate as redox signaling messengers, oxidative stress can also cause perturbation in normal cellular signaling mechanisms. From the latter point, oxidative stress is considered one of the most important factors in variant abnormality conditions such as cancer, chronic obstructive pulmonary disease, cardiovascular disease, chronic kidney disease, and NDDs [17, 18].

Nowadays, SCI has been a critical health problem among NDDs. The pathophysiology of SCI is defined through a primary injury/initial trauma, followed by the second cascades of injuries that are consequences of the indirect achievement of the primary injury. The secondary injury continues at different time points after the primary injury and is not restricted to the same primary injury site. Oxidative stress is one of the critical components of the later phase, which is receiving increasing attention. Many studies have been performed to investigate the destructive activity of oxidative stress after SCI and its confounding effects on improving the functional recovery following SCI [19].

Various factors predispose the neurons of the central nervous system to electrophilic/oxidative stress. Among those factors, polyunsaturated fatty acids (e.g., arachidonic acid and linoleic acid), low antioxidant capacity, reactive oxygen metabolites, and neural mitochondrial changes are the critical ones [9, 20]. Decreasing glutathione (GSH) and the levels of copper/zinc superoxide dismutase (SOD), as well as increasing malondialdehyde (MDA), and acrolein are considered critical hallmarks of oxidative stress after SCI [9]. In parallel, the mechanical injury detected in the plasma membrane resulting from SCI leads to the disruption of normal ion balance and obviously to the enhancement of the intracellular concentration of sodium and calcium. Besides oxidative stress and increasing glutamate (Glu), there are other aggravating factors for increasing intracellular calcium. Consequently, Na^+/K^+ ATPase works to improve the electrochemical gradient of the neuronal cell membrane via decreasing the intracellular sodium that leads to reversing the activity of the Na^+/Ca^{2+} exchanger, thereby removing sodium and increasing the intracellular concentration of calcium and related oxidative toxicity. Increased intracellular calcium causes mitochondrial damage and ROS production and activates phospholipase, caspases, and proteases, followed by an increase in the release of membrane fatty acids [20]. Also, SCI impairs ATP production, recruits neutrophils to the injury site, stimulates xanthine oxidase (XO), and produces more ROS [21]. In the same

way, lipid peroxidation (LPO) can increase oxidative stress via the generation of more free radicals and reactive aldehydes [20]. Since oxidative stress and ROS are key elements in the pathophysiology of SCI, reduction of oxidative stress is considered one of the possible effective mechanisms in the treatment/improvement of SCI patients. In recent years, several studies have investigated the antioxidative effects of various bioactive compounds in animal models of SCI.

In addition to oxidative stress, inflammatory responses are essential components of SCI secondary injury that play significant roles in regulating the pathogenesis of acute and chronic phases of SCI and play a special role in nerve injury and regenerative responses [22]. After SCI, ruptured blood vessels disturb the blood-brain barrier and infiltrate the neutrophils to the injury site. The proliferation or migration of activated glial and peripherally derived immune cells initiates the inflammatory responses in the central nervous system after SCI. T cells are one of the main pillars of the inflammatory and immune response and necessary factors for mounting macrophages in these processes [22]. Also, the activated myelin basic protein- (MBP-) reactive T cells lead to transient paralysis and neuroinflammation in SCI. Microglia and macrophages are involved in the inflammatory response and secondary pathology via facilitating the release of cytokines, interferon, interleukin- (IL-) 1, IL-6, IL-10, and tumor necrosis factor (TNF), and the activation of interleukin receptors (IL-2R and IL-4R). Furthermore, cytokines produce inflammatory responses in the central nervous system by enhancing the expression of additional reactive oxygen, cytokines, NO, and chemokines [22]. After SCI, IL-6 and TNF- α are upregulated around the damaged area. It was documented that TNF- α could increase the neuronal cell death mediated by Glu in the rat spinal cord and TNF- α antagonists decrease the development and extension of inflammation as well as tissue injury events dependent on SCI. Suspiciously similar treatment with the IL-6 receptor monoclonal antibody diminishes the differentiation of astrocytes, reduces the number of inflammatory cells, and detracts the formation of connective tissue scars [22]. Cytokines showed an important role in SCI-induced inflammatory processes, and enhancing the production of IL-1 family cytokines (like IL-1 α) is considered a vital trigger for this process. IL-1 α and IL-18 are known as intense mediators of inflammation that can initiate and/or augment an extensive type of effects associated with host responses to tissue injury and innate immunity. SCI induces the activation of caspase-1, enhances the operation of IL-1 α and IL-18, and cleaves the X-linked inhibitors of apoptosis protein (XIAP) [22]. Accordingly, decreasing inflammation could modulate the functional deficit and secondary degeneration after SCI.

Apoptosis is another important physiological process for removing cells, and its implementation is necessary for the regular function and development of multicellular organisms. Apoptosis signaling occurs via several pathways inchoate from triggering events within or outside the cell [23]. Two main pathways are known for the mechanism of cellular apoptosis. During the first pathway, mitochondrial dysfunction begins with different triggers such as ATP

depletion or hypoxia, leading to the release of cytochrome from mitochondria and caspase-9 activation. On the other hand, the extrinsic apoptosis pathway is associated with a group of cell surface receptors known as cell death receptors (DRs) and belongs to the TNF receptor (TNFR) family. TNFR consists of Fas, Fas ligand, TNFR1, DR3, and p75, leading to the activation of caspase-8 and caspase-10 via their intracellular death domains [8, 9]. Caspase-9 and caspase-8 are the major caspases in mitochondrial-dependent pathways and the DR, respectively, which regulate the process; downstream effector caspases (as caspase-3) are activated through proteolytic cleavage and initiator caspases. The Bcl-2 family of proteins, p21-activated kinase, caspase-activated DNase inhibitor, cytoskeletal proteins (such as gelsolin), proteins involved in mRNA splicing, DNA repair, DNA replication, and focal adhesion kinase are some essential proteins among the identified target substrates for caspase-3 [22, 24]. Following SCI, the Fas and p75 receptors are expressed on the spinal cord's microglia, astrocytes, and oligodendrocytes. Some of the extracellular excitatory amino acids such as Glu can increase glial and neuronal cell death's delayed apoptosis via the activation of the N-methyl-D-aspartate (NMDA) receptor. Although SCI's exact mechanism remains unclear, calcium-dependent and TNF- α -mediated activation, inducible nitric oxide synthase (iNOS), and NO production are possible factors involved in this process, and it is proven that Glu-induced cell death was amplified by TNF- α . Also, TNF- α activates the nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) and leads to the enhancement of the transcriptional activation of iNOS and more cell damage. These mechanisms are partially inhibited by the NMDA receptor antagonist and may be used as an appropriate therapeutic target [8, 9].

Altogether, it seems that oxidative stress, inflammation, and apoptosis play key destructive roles in the pathophysiology of SCI. Figure 1 shows the critical role of inflammation, oxidative stress, and interconnected pathways during SCI (Figure 1).

3. The Pivotal Role of Oxidative Stress after SCI

Oxidative stress seems to play a more important role than other factors and mechanisms involved in SCI's pathophysiological processes. Several studies have investigated oxidative stress parameters and antioxidant markers in SCI models. Compression, contusion, transection, hemisection, and ischemia-reperfusion are among the most common models of SCI *in vivo* [25]. The results have shown dysregulated levels of advanced oxidation protein products (AOPP), MDA, GSH peroxidase (GPx), SOD, and catalase (CAT) following SCI [26, 27]. Furthermore, the production of superoxide anion O_2^- , H_2O_2 , and other oxidative derivatives such as nitrogen dioxide, peroxyxynitrite, carbonate, and $\cdot\text{OH}$ is increased [28]. Oxidative stress increases the level of NO, and the production of oxidized protein plays an important role in the pathogenesis of SCI. The produced NO leads to cytotoxic effects, vessel dilation, neuronal apoptosis, neurodysfunction, and neurodegeneration.




FIGURE 1: The pathophysiology and proposed targets of SCI. CAT: catalase; CSPGs: chondroitin sulfate proteoglycans; GSH: glutathione; GPx: glutathione peroxidase; IL: interleukin; MIF: macrophage migration inhibitory factor; NF- κ B: nuclear factor- κ B; NR2B: N-methyl-D-aspartate receptor subtype 2B; ROS: reactive oxygen species; SCI: spinal cord injury; SOD: superoxide dismutase; TNF- α : tumor necrosis factor-alpha.

Enhanced intracellular calcium, increasing Glu, and mitochondrial ROS production can activate the posttraumatic cell death cascade and other harmful mechanisms in the SCI [26, 27]. Besides, oxidative stress exacerbates or initiates various inflammatory and cellular apoptotic pathways, so inhibition of oxidative pathways can significantly reduce the severity of post-SCI. It could be considered one of the most appropriate strategies to treat or prevent the progression of SCI [26]. One of the most substantial cellular defense mechanisms versus xenobiotic damage and oxidative stress is the Keap1-nuclear factor erythroid 2-related factor 2 (Nrf2)/ARE pathway. Attenuating this signaling pathway is one of the best ways to control and completely enclose oxidative stress as a suitable target for preventing and treating related disorders and conditions such as neurodegeneration, malignancy, and cardiovascular and inflammatory diseases [29]. ARE is a *cis*-regulatory element or enhancer sequence that can be found in the promoter region of variant genes encoding cytoprotective proteins and detoxification enzymes. In this line, the transcriptional regulation of ARE-dependent expression proteins and enzymes can be induced and controlled by activating Nrf2 signaling. The Keap1 is an inhibitor of Nrf2, and oxidative stress dissociates Nrf2 from Keap1 and activates the antioxidant genes via binding to the ARE [29].

Many of the studies showed appropriate and acceptable effects in controlling injuries after SCI and exerted their effects through the aforementioned pathway, which shows

the great importance of oxidative stress in injuries caused following SCI [30]. Also, activating the Keap1-Nrf2/ARE pathway mentioned as the main mechanism of the neuroprotective effect of sulforaphane could increase the levels of Glu-cysteine ligase and Nrf2 and GSH and diminish the levels of inflammatory cytokines, TNF- α and IL-1 β [31]. In addition, other molecular mechanisms involved in the oxidative stress process have been targeted by active compounds and agents capable of suppressing oxidative stress. Enhancing the activity and the levels of CAT, SOD, glutathione reductase (GR), GPx, MDA, phosphoinositol, GSH, and nicotinamide adenine dinucleotide (NADH), as well as suppressing the expression of ROS, iNOS, phospholipase C gamma (PLC γ), nicotinamide adenine dinucleotide phosphate- (NADPH-) dependent LPO, and XO, could be targeted by phytochemicals. Besides, decreasing the level of myeloperoxidase, restoring the membrane potential of mitochondria, enhancing the activity of the phosphoinositide 3-kinase (PI3K)/Akt signaling pathway, and increasing the levels of thioredoxin- (TRX-) 1/TRX-2 mRNA are some of the mechanisms that have been targeted by other bioactive compounds in modulating oxidative stress [9, 32–37].

The involvement of oxidative stress and ROS in the pathogenesis of SCI leads to extensive neuroprotective effects of variant compounds with antioxidative activities in different SCI models. Some of these antioxidants may slow the progression and development of spinal cord damage, and antioxidant-based strategies are considered a suitable approach for SCI treatment.

4. Related Interconnection between Oxidative Stress and Inflammatory/Apoptotic Pathways after SCI

As mentioned earlier, oxidative stress, inflammatory reactions, and apoptosis are the leading causes of neuronal damage after SCI, which induce their harmful effects through different mediators. Although the signaling pathways of these factors are somewhat different, they are interconnected to each other, like interconnected links in a chain, and exacerbate each other's effects. Accurate identification of the connections and mechanisms between these factors after SCI can be helpful in the treatment or prevention of SCI.

ROS are critical mediators of oxidative stress pathways and play a significant role in the onset and progression of inflammatory responses. Accordingly, increased ROS generation in the injury site leads to endothelial dysfunction, inflammation, and tissue injury. The oxidative stress-mediated inflammatory response leads to the opening of interendothelial junctions and increases the migration and penetration of inflammatory cells across the endothelial barrier. Besides, overexpression of the antioxidant enzyme such as SOD diminishes adhesion molecules' expression and reduces the leukocyte endothelium binding at the site of inflammation [38, 39].

On the other hand, the progression of oxidative stress leads to DNA breaks and mitochondrial damage, followed by a reduction in the alteration of membrane permeability and transmembrane potential, facilitating the release and production of apoptotic factors. It is documented that ROS could play both the inhibitory and activating effects in NF- κ B signaling [38, 40]. The lower level of ROS leads to the induction of cell survival responses, while the apoptosis and death processes activate in higher doses of ROS [41]. Inactivating the phosphorylation of I κ B α , as an inhibitory protein, is considered the main transcription factor of ROS, while H₂O₂ has an impact on the phosphorylation of I κ B α , which leads to I κ B α degradation and NF- κ B activation. Accordingly, the NF- κ B signaling pathway affects variant antioxidant proteins such as CAT, SOD, TRX-1, and TRX-2. In this line, activating the transcription of main enzymes involved in ROS production, such as inducible NO synthase, NADPH oxidase, cyclooxygenase (COX), and lipoxygenases, affects the intracellular levels of ROS [38, 40].

Furthermore, oxidation protein products in the Sprague Dawley rat model of SCI could affect the microglia-mediated neuroinflammation through the mitogen-activated protein kinase (MAPK)/NF- κ B signaling pathway, activating NADPH oxidase and causing pyroptosis. The process could pass through expressing proinflammatory cytokines and cleavage of Gasdermin-D (GSDMD) during the secondary phase of SCI [42]. Besides, the role of NADPH oxidase and myeloperoxidase generated by immune cells in damaged areas contributes to oxidative damage and inflammation.

As mentioned earlier, mechanical injury to the cord tissue leads to damage to blood vessels. It causes sequestration and penetration of inflammatory molecules such as TNF- α and other inflammatory molecules to the affected area.

Besides, neuronal damage causes the release of Glu into the extracellular space, which leads to the activation of cytotoxic cascades and an increase in glial and neuronal calcium, ultimately increasing neuronal death. Glu activates microglia and enhances the release of inflammatory molecules, like CD95 ligand and TNF- α that, combined with Glu, facilitate necrotic cell death and activate apoptotic pathways [22, 43]. As studies have shown, inflammatory responses, like oxidative stress responses, lead to neuronal apoptosis and neuronal damage. It seems that anti-inflammatory drugs may be a viable option to prevent the progression of nerve cell damage after SCI [43].

From the apoptotic point of view, ROS activate the tumor suppressor protein p53 in severe stress. p53, in turn, downregulates the prosurvival proteins like survivin, B cell lymphoma 2 (Bcl-2), Bcl-XL, and IAPs and upregulates the proapoptotic proteins that actuate the regulation of the apoptotic transcriptional profile. Also, the transcription of proapoptotic genes such as Bid, Bax, p53 upregulated modulator of apoptosis (PUMA), apoptotic protease activating factor 1 (Apaf-1), and Noxa is activated by p53. Cytosolic p53 penetrates the mitochondria and directly interacts with antiapoptotic and proapoptotic proteins (e.g., Bcl-2, Bcl-XL, Mcl-1, Bax, and Bak, respectively), facilitating apoptosis and releasing proapoptotic factors. Direct activation of the Bax through its structural rearrangement and increasing the mitochondrial membrane permeabilization are other mechanisms of p53. In addition, DR-4, DR-5, Fas, and FasL are some of the other extrinsic proapoptotic factors that are altered by p53 [41, 44] (Figure 1).

5. Secondary Metabolites in Combating Oxidative Stress after SCI

Plant secondary metabolites are constructed by plants and considered important classes of natural products that can show significant direct or indirect roles in the normal growth, reproduction, and development of the organism. Furthermore, plant secondary metabolites have undeniable and vital effects on the plant defense system against environmental stresses and pathogenic attacks [45]. It was documented that these compounds have shown remarkable biological activities such as antioxidant, anticancer, anti-inflammatory, antimutagenic, immunostimulatory, antibacterial, antiapoptosis, and neuroprotective effects [46–48]. Phenolic compounds, alkaloids, terpenoids, and sulfur-containing compounds are the most known classes of secondary metabolites that express their neuroprotective effects through antioxidant and anti-inflammatory effects [47–49].

As mentioned earlier, one of the most influential factors after SCI is oxidative stress, inflammation, or apoptosis, and secondary metabolites affect and inhibit these pathways through various mechanisms. Suppressing the expression, activity, and levels of MDA, matrix metalloproteinase (MMP-) 9, TNF- α , COX-2, IL-1 β , IL-6, NADPH oxidase, thiobarbituric acid reactive substance (TBARS), myeloperoxidase (MPO), eNOS, nitrotyrosine, and poly(ADP-ribose) polymerase (PARP) formation and enhancing the levels of SOD, CAT, GPx, and GSH are the main antioxidant, anti-

inflammatory, and antiapoptotic mechanisms mentioned for secondary metabolites after SCI. Also, it has been proved that these compounds apply the aforementioned alterations by interfering with the PI3K/Akt/mammalian target of rapamycin (mTOR), Akt/Nrf2, extracellular signal-regulated kinase (ERK) 1/2, Janus kinase- (JAK-) signal transducer and activator of transcription (STAT), and Keap1-Nrf2/heme oxygenase-1 (HO-1). Additionally, catalytic/modifying subunit of glutamate-cysteine ligase (GCLC/GCLC), p38MAPK, transforming growth factor-beta (TGF- β)/Smad, MAPK/JNK/NF- κ B, sirtuin 1 (SIRT1)/peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1 α), ERK/NADPH oxidase, and toll-like receptor-4 (TLR4) signaling pathways are modulated by phytochemicals [9, 50–54].

6. Polyphenols Target Upstream Mediators of Oxidative Stress after SCI

Different natural polyphenolic/phenolic compounds have been used to alleviate oxidative stress after SCI [55]. These compounds modulate several dysregulated pathways/mediators, including preventing the formation of \cdot OH and inhibiting the repair of molecules after a free radical attack. Such phytochemicals have previously been shown as potential neuroprotective agents in other neurodegenerative conditions associated with oxidative stress [56, 57].

6.1. Quercetin. Quercetin (C₁₅H₁₀O₇) (Figure 2) is a flavonoid extracted from various vegetables, fruits, and grains. It has shown beneficial biological properties, including anti-inflammatory, anticarcinogenic, antioxidant, and antiviral effects. Quercetin has also shown the ability to inhibit lipid peroxidation and capillary permeability to stimulate mitochondrial biogenesis in ameliorating neuronal dysregulation and mental/physical dysfunction [58, 59]. Quercetin can directly scavenge \cdot OH, superoxide anions, and LPO via its phenolic hydroxyl groups [60]. In addition, quercetin can bind transition metals and attenuate oxidation and reduction to form metal chelates to inactivate transition metals, such as copper and iron [61]. The neuroprotective effects of quercetin have been extensively demonstrated in several animal models. It significantly suppressed GSH levels and MPO activity following brain trauma [62]. Quercetin also increased the activity of GPx, SOD, and CAT in traumatic brain injury [63], decreased the elevated MMP-9 level [64], and also activated the brain-derived neurotrophic factor (BDNF), tropomyosin receptor kinase B (TrkB), and PI3K/Akt signaling pathway in cerebral focal ischemia [65]. Additionally, quercetin (30 mg/kg, i.p.) inhibited oxidative stress, spinal production of cytokine, and glial cell activation of glial fibrillary acidic protein (GFAP) and ionized calcium-binding adaptor protein 1 (Iba1) mRNA expression [66]. In another study, i.p. administration of the quercetin in the dose of 20 mg/kg/day for 10 days reduced monosodium Glu-induced neurotoxicity in spinal motor neurons by inhibiting p38MAPK, exerting anti-inflammatory effects, antagonizing oxidative stress, and enhancing GFAP expression [67]. Azevedo et al. showed that quercetin (25, 50, and 100 mg/kg, i.p.) inhibited oxidative stress-induced damage

in dorsal horn neurons by suppressing LPO [68], which was in line with the results of Liu et al. [69]. Sustained-release poly(lactic-co-glycolic acid)- (PLGA-) polyethylene glycol- (PEG-) PLGA hydrogel formulation of quercetin (50 and 100 mg/mL) reduced oxidative damage and inflammation in the spinal cord and also promoted nerve cell survival, nerve regeneration, and motor function recovery in rats with brachial plexus avulsion [70]. Following SCI, i.p. injection of 20 mg/kg of quercetin for 7 days targets BDNF and JAK2/STAT3 signaling pathways [71] and regulated the secondary oxidative stress by inhibiting the p38MAPK/iNOS signaling pathway [61]. Schültke et al. reported that administration of quercetin in a dose of 0.25 μ mol/kg led to a lower MPO expression in damaged spinal cord tissue [72]. In addition, single-dose quercetin within the first 72 hours after SCI enhanced total antioxidant levels and decreased the levels of NO and MDA [73]. In this regard, quercetin also elevated total antioxidant capacity and paraoxonase activity after SCI in rats [74]. In another study, the administration of 20 mg/kg quercetin showed advantageous effects against SCI-induced oxidative damage via its antioxidant and anti-inflammatory effects [75]. In a rat model of SCI, Wang et al. reported that quercetin (50 μ mol/kg) combined with human umbilical cord mesenchymal stromal cells reduced proinflammatory cytokines (e.g., IL-1 β and IL-6) while increased anti-inflammatory cytokines (e.g., IL-4 and IL-10), which are in a near interconnection with oxidative pathways. The treatment also improved macrophage polarization, neurological functional recovery, and axonal preservation and decreased the cystic cavity size [76]. An *in vivo* and *in vitro* study revealed that quercetin (7.5 mg/kg, i.p.) inhibited necroptosis of oligodendrocytes via suppression of the STAT1 and NF- κ B pathway after SCI [77]. Jiang et al. reported that 100 mg/kg of quercetin remarkably decreased ROS production, TNF- α , IL-1 β , and IL-18 after SCI in female rats [78].

Overall, quercetin seems to be a hopeful agent in modulating oxidative stress following neuronal damage and SCI.

6.2. Epigallocatechin Gallate. Epigallocatechin gallate (C₂₂H₁₈O₁) (EGCG) or epicatechin is the main compound of tea catechins. Most of the biological activities of green tea extracts are linked to this composition [79]. Several experimental studies have reported that EGCG causes neuroprotection against neurodegenerative diseases [80], brain injury [81], SCI [82], and peripheral nerve damage [83] through its antiapoptotic, anti-inflammatory, and antioxidant properties [84]. The presence of hydroxyl groups in rings B and D of the catechins makes them bind free radicals [85]. Different concentrations (50–200 μ g/mL) of green tea polyphenols for 24 hours protected spinal neurons against oxidative stress caused by H₂O₂ [86]. An *in vitro* study displayed that treating PC12 cells with 0, 125, 250, 500, 1000, and 2000 μ mol/L concentrations of EGCG reduced ROS production through the SIRT1/PGC-1 α signaling pathway [87]. Different doses (10, 25, or 50 mg/kg, i.p.) of EGCG also attenuated NADPH/neuronal nitric oxide synthase (nNOS) expression after peripheral nerve injury [83] and inhibited neurotoxicity through activation of the cAMP response




FIGURE 2: Selected chemical structures of polyphenols.

element-binding protein (CREB)/BDNF/TrkB and PI3K/Akt signaling pathway in doses of 25, 50, or 75 mg/kg for 18 days in mice [88]. A 50 mg/kg dose of EGCG protected the rat spinal cord from secondary damage by decreasing MPO activity, iNOS, TNF- α , IL-1 β , COX-2, and PARP expression [89]. Khalatbary et al. also found that i.p. injection of EGCG in the dose of 50 mg/kg, immediately and 1 h after SCI, reduced MDA [90]. Consequently, an *in vitro* study indicated that EGCG at a concentration of 5 μ M for 48 h reduced oxidative stress levels and protected motor neurons in the organic culture of the rat spinal cord [91]. Coadministration of 17 mg/kg of EGCG and 60 mg/kg of curcumin for 28 days modulated macrophage inflammatory protein 1- α (MIP-1 α), IL-1 β , IL-4, and IL-6 as cross-talk inflammatory mediators with oxidative stress [92]. Administration of 30 mg/kg of EGCG (i.p.) for a week after SCI reduced thermal hyperalgesia via downregulation of RhoA and TNF- α in mice [93].

In this line, 50 mg/kg of protocatechuic acid (i.p.), an essential metabolite of antioxidant polyphenols in green tea, significantly decreased the expression of inflammatory mediators such as TNF- α , IL-1 β , COX-2, iNOS, and MMP-9 in rats after SCI [94]. So, EGCG seems to have a bright future in combating oxidative stress post-SCI.

6.3. Caffeic Acid Phenethyl Ester. Caffeic acid phenethyl ester (C₁₇H₁₆O₄) or phenylethyl caffeate is a component of honeybee propolis [95]. Its biological activities include antioxidant [96], anti-inflammatory [97], antibacterial [98], anticancer, and cytotoxic properties [99], which is due to related hydroxyl groups in the catechol ring [100]. Caffeic acid phenethyl ester is a potent inhibitor of NF- κ B [101] and protein tyrosine kinase [102]. It suppresses lipoxygenase activities [103] and blocks calcium-induced cytochrome c release in the hypoxic-ischemic brain injury models [104]. From another view, caffeic acid phenethyl ester repressed

the formation of superoxide anion and XO activity in eukaryotic cells [105] and reduced MPO and Na^+/K^+ ATPase activity during ischemia-reperfusion injury [106]. Caffeic acid phenethyl ester activated the expression of HO-1 through Nrf2 activation associated with the ERK signaling pathway [107]. It binds to Keap1 and disrupts the Nrf2/Keap1 complex, thereby enhancing Nrf2 binding to ARE [108]. Intrathecal injection of 1 $\mu\text{g}/\text{kg}$ caffeic acid phenethyl ester after SCI led to a decrease in MDA, LPO, and total oxidant activity. It increased antioxidative agents [109], while decreasing tissue and serum levels of IL-6 post-SCI [110]. In this line, Ak et al. confirmed the effects of i.p. injection of caffeic acid phenethyl ester (10 $\mu\text{g}/\text{kg}$) in reducing the levels of TNF- α and IL-1 β post-SCI [111]. Administration of 10 $\mu\text{mol}/\text{kg}$ (i.p.) of this phytochemical improved motor function and reduced lesion size by decreasing the expression of IL-1 β , NOS, and COX-2 following SCI [112]. Administering 10 $\mu\text{mol}/\text{kg}$ (i.p.) of caffeic acid phenethyl ester before surgery diminished ischemic injury in the spinal cord through scavenging free radicals and providing a better microcirculatory environment by preventing endothelial cell lysis through proteases from activated leukocytes [113]. It also exerted neuroinflammatory effects via inhibiting ROS and catalytic activity of iNOS in the concentration of 50 $\mu\text{mol}/\text{mL}$ [114].

6.4. Honokiol. Honokiol ($\text{C}_{18}\text{H}_{18}\text{O}_2$) is a pleiotropic lignan presented in *Magnolia grandiflora* [115]. It is known for its therapeutic properties such as antioxidant [116], anti-inflammatory [117], analgesic [118], antidepressant [119], antitumorogenic [120], and neuroprotective [121] effects. Several studies showed that honokiol could decrease oxidative stress factors in various organs such as the heart [122], liver [123], kidney [124], and brain [125]. Honokiol inhibited ROS production via the ERK/NADPH oxidase pathway in microglial cells [126]. It also activated Nrf2 [127], inhibited XO [128], and modulated the PI3K/Akt pathway [117] to exert neuroprotective responses. Besides, honokiol preserved mitochondrial respiratory chain enzyme, inhibited ERK/protein kinase C (PKC) pathways [129], and modulated lipopolysaccharide- (LPS-) induced NO expression by targeting PKC, MAPKs, and NF- κB [130]. In a rat model of SCI, 20 mg/kg of honokiol (i.p.) reduced the production of proinflammatory cytokines and inhibited neutrophil infiltration and microglial activation, which are all in a near link with oxidative mediators [131]. In the ischemic brains, 10 $\mu\text{g}/\text{kg}$ of honokiol (i.p., twice) maintained Na^+/K^+ -ATPase activity and mitochondrial function against oxidative stress [121] and inhibited neutrophil infiltration and ROS production [132].

6.5. Rosmarinic Acid. Rosemary ($\text{C}_{18}\text{H}_{16}\text{O}_8$), with a molecular formula of $\text{C}_{18}\text{H}_{16}\text{O}_8$, contains several bioactive polyphenol components, including carnosic acid, carnosol, genkwanin, rosmarinic acid, and rosmanol [133]. Rosmarinic acid is an ester produced of caffeic acid that was first found in *Rosmarinus officinalis*. Rosmarinic acid has remarkable biological effects, including anticancer, antiviral, antibacterial, antiaging, antioxidant, antidiabetic, hepatopro-

TECTIVE, nephroprotective, cardioprotective, antiallergic, antidepressant, and anti-inflammatory activities [134]. The antioxidant activity of rosmarinic acid, along with its effects on signaling pathways and gene expression, contributes to most of its biological properties [135]. Rosmarinic acid decreased LPO and elevated GSH levels in HepG2 hepatoma cells [136]. Besides, it reduced p38MAPK, p-JNK, NF- κB , IL-1 β , IL-6, and TNF- α but increased activity of CAT, SOD, glutathione S-transferase (GST), and GPx in diabetic rats [137]. Moreover, it inhibited ROS-dependent MMP-2 activity by targeting the Nrf2 antioxidant system in hepatic cells [138], increased SOD, CAT, HO-1, and Nrf2 activity, and was significantly decreased by UVB radiation [134]. Rosmarinic acid (0.5, 2.5, 5, and 10 $\mu\text{g}/\text{mL}$) repressed oxidative stress in C6 glial cells by inhibiting LPO and reducing COX-2 expression and iNOS [139]. Rosmarinic acid significantly attenuated the levels of TNF- α , iNOS, apoptotic factors, Iba1, TLR4, and GFAP after chronic contraction injury in rats [140]. Treatment with rosmarinic acid at the doses of 10, 20, and 40 mg/kg (i.p.) for 14 days suppressed the spinal glial cell activation. It inhibited the expression of inflammatory markers, as well as activating AMPK in peripheral nerves and dorsal root ganglia, which might also have contributed to its neuroprotective actions [141]. An *in vitro* study in PC12 cells showed that the antioxidant effects of 10 mM rosmarinic acid are mediated by the Akt/glycogen synthase kinase-3 β (GSK-3 β) pathway and increasing Nrf2 activity [142].

6.6. Carvacrol. Carvacrol ($\text{C}_{10}\text{H}_{14}\text{O}$) or cymophenol is a phenolic monoterpene derivative from cymene. Carvacrol has shown various pharmacological activities such as anxiolytic [143], antidepressant [144], antibacterial, antioxidant [145], antitumor, antigenotoxic, antimutagenic [146], analgesic, anti-inflammatory [147], hepatoprotective, and anti-hepatotoxic activities [148]. Carvacrol protects the brain, liver, and kidney against oxidative stress and significantly increases GSH, SOD, CAT, and GPx levels [149]. Additionally, carvacrol enhanced the expressions of Nrf2 and ERK1 inhibited by cadmium in PC12 cells [150]. It also made cytoprotective effects through HO-1 in cells exposed to H_2O_2 [151] and inhibited the MAPK/JNK-NF- κB signaling pathway in cells exposed to iron ions [152]. The overload of iron ions leads to a reaction with oxygen and the production of oxidative damage, including mitochondrial dysfunction and LPO via Fenton reaction [153]. Carvacrol also exhibited modulatory effects on the expression of pro- and anti-inflammatory cytokines [154]. It has also suppressed transient receptor potential ankyrin 1 (TRPA1) receptors and decreased the production/release of proinflammatory cytokines and markers like TNF- α , IL-1 β , MPO, NF- κB , and COX-2, as well as oxidative stress factors, including MDA, GSH, and NO levels following i.p. administration (25, 75, and 150 mg/kg) [155]. However, more experiments are needed toward a better future of the neuroprotective effects of carvacrol after SCI via suppressing oxidative pathways.

6.7. Rutin. Rutin ($\text{C}_{27}\text{H}_{30}\text{O}_{16}$), also known as rutoside or vitamin P ($\text{C}_{27}\text{H}_{30}\text{O}_{16}$), is a flavonol glycoside obtained from buckwheat [156]. Rutin is known for various

pharmacological activities, including cytoprotective, antioxidant [157], anticancer [158], vasoprotection [159], neuroprotection [160], and post-SCI neuroinflammation [161] activities. Rutin modulated the MAPK [162] and iNOS/Nrf2 signaling pathway [163] and reduced oxidative stress by an enhancement in CAT activity along with a decline in LPO and protein carbonyl content [164]. Rutin also attenuated ischemic neural apoptosis by inhibiting LPO and p53 expression, an increment in antioxidant defense enzymes [161], and a decrement in ROS production [165]. It improved diabetic neuropathy by reducing oxidative stress through HO-1 and Nrf2 in rats [166]. Employing 100 mg/kg of rutin also upregulated BDNF, CREB, and ERK1 gene expression in the hippocampus [167] and protected PC12 cells exposed with sodium nitroprusside via modulating the PI3K/Akt/mTOR and ERK1/2 pathway [168]. Oral administration of 10 mg/kg of rutin for 3 weeks inhibited oxidative stress and inflammation by targeting the NOS-mediated NF- κ B signaling pathway [169]. Besides, Zhang and Ma showed that 10 μ mol/kg of rutin (i.p.) reduced the expression levels of MIP-2 and p-Akt in a rat model of SCI [170]. Also, rutin (30 mg/kg) decreased oxidative stress-related markers and inflammatory cytokines (e.g., IL-1 β , IL-6, and TNF- α) by suppressing the p38MAPK pathway after SCI [171]. Another study showed that i.p. injection of rutin (50 and 100 mg/kg) for 3 days significantly attenuated the levels of ROS, MDA, IL-1 β , IL-18, and TNF- α [161]. In an *in vitro* study, rutin protected cells from H₂O₂-induced oxidative stress and apoptosis through regulation of ROS, suppression of LPO, protection of the intracellular antioxidant system, and regulation of the Bax/Bcl-2 ratio and NF- κ B/p65 signaling pathway [172]. Rutin also prevented oxidative DNA damage and neuronal death induced by nutrient deprivation conditions [173]. Also, it was reported that i.p. injection of 30 mg/kg of rutin for 3 days after SCI plus mild hypothermia reduced inflammatory factors through repressing the TGF- β /Smad pathway in the SCI model [174].

6.8. Hesperidin. Hesperidin (C₂₈H₃₄O₁₅) is a flavanoglycone isolated from citrus fruits that possesses anti-inflammatory, antioxidant [175], anticancer [176, 177], and antiapoptotic [178] activities. Hesperidin attenuated oxidative stress and inflammation via modulation of TGF- β 1/Smad3, Nrf2/ARE/HO-1, and peroxisome proliferator-activated receptor γ (PPAR γ) signaling [179]. Additionally, it inhibited LPO in Parkinson's disease [180], increased GSH and total antioxidant capacity, and reduced H₂O₂ and MDA levels. Hesperidin inhibited GSK-3 β activity in Alzheimer's disease [181] and also activated antioxidant enzymes (XO, GST, and GR) [182]. An *in vitro* study based on oxidative stress showed that activation of the ERK/MAPK signaling by hesperidin is involved in the expression of HO-1 and Nrf2 [183]. Hesperidin also upregulated the Keap1-Nrf2/HO-1 pathway *in vitro* [184] and improved the antioxidant defense system via activation of Nrf2/ARE/HO-1 in kidney tissue [185]. Consequently, it suppressed oxidative stress and inflammation through activating Nrf2/HO-1/ARE and PPAR γ pathways [179]. In a recent study by Heo et al., hesperidin (100 mg/kg, i.p.) for 14 days (from 7 days prior to

SCI to day 7 post-SCI) decreased oxidative stress and inflammation by upregulation of the HO-1/Nrf2 pathway after SCI [186]. At the same dose, hesperidin led to a decrease in TNF- α , IL-1 β , and IL-6 in the spinal cord tissue [187].

6.9. Resveratrol. Resveratrol (C₁₄H₁₂O₃) is a polyphenol belonging to stilbenoids, possessing two phenol rings and an ethylene group. Resveratrol is present in various plants such as plums, peanuts, mulberries, and grapes [188]. In addition to its antiapoptotic [189], anti-inflammatory, antinociceptive [190], and anticancer [191] properties, the antioxidant potential of resveratrol is of the most important biological effects of resveratrol [192]. Resveratrol scavenges free radicals through antioxidant effects and suppresses the iNOS/p38MAPK pathway [193]. Treatment with 50 μ M of resveratrol maintained the spinal cord dorsal column against hypoxic injury through activating Nrf2 [194] and decreasing pancreatic oxidative damage via downregulating NF- κ B and PI3K pathways [195]. Besides, resveratrol diminished inflammatory hyperalgesia by targeting antioxidant enzymes, ROS, and ERK activation [196]. It also exerted antioxidant effects following myocardial ischemia through the Nrf2/ARE signaling pathway [197]. In another study by Tamaki et al., oral administration of resveratrol in the dose of 10 mg/kg enhanced antioxidant gene expression by activating SIRT1/AMPK and Nrf2 pathways [198]. Resveratrol (5 mg/kg orally for 30 days) protected against oxidative stress through Nrf2/Keap1 signaling [199]. An *in vitro* study exhibited that resveratrol protected astrocytes from oxidative damage by inhibiting ROS and NF- κ B, while upregulating Nrf2 expression. They reported that resveratrol was more effective in late hours after oxidative trauma [200]. Resveratrol in the dose of 10 mg/kg protected the spinal cord from spinal ischemia-reperfusion injury via increasing NO release and decreasing oxidative stress [74]. Ates et al. showed that administration of resveratrol (100 mg/kg, i.p.) and methylprednisolone (30 mg/kg, i.p.) immediately after trauma led to a decrease in MDA, XO, and NO activity and an increase in the GSH level following SCI [201]. Also, in another study, the same dose of resveratrol was associated with the regulation of SIRT1 expression and the suppression of NF- κ B activity in lung tissue after SCI [202]. A meta-analysis reported that resveratrol significantly improved motor function after SCI and also elevated SOD levels while diminished MDA levels following SCI [203]. Resveratrol administration (7 and 14 mg/kg) from 4 days before the experiment, until the end of the experiment, resulted in the suppression of proinflammatory cytokines (NF- κ B and TNF- α) and upregulation of antioxidant mediators, including Nrf2, NAD(P)H:quinone oxidoreductase (NQO1), and HO-1 [204]. As one of the resveratrol derivatives, oxyresveratrol showed twice more antioxidant capacity [205]. Treatment of SCI with oxyresveratrol (10 and 20 mg/kg, for 4 weeks) reduced inflammation and oxidative stress by enhancement of Nrf2, SOD, GPx, and GSH and decrement of MDA, IL-1 β , IL-6, TNF- α , NF- κ B, COX-2, and iNOS [206]. Administration of 40 mg/kg (i.p.) polydatin, as a glycoside and precursor form of resveratrol [207], improved

SCI by inhibiting oxidative stress via Nrf2/HO-1 signaling [208]. Polydatin in 20 and 40 mg/kg doses (i.p., 30 min after surgery) attenuated SCI complications by reducing microglial inflammation via suppressing the iNOS and NLR family pyrin domain containing 3 (NLRP3) inflammasome pathway [209]. Polydiatin, in combination with bone marrow stem cells (BMSCs), ameliorated SCI by increasing cell survival and reducing oxidative stress through the Nrf2/ARE pathway [210].

In another study, 100 mg/kg of resveratrol (i.p.) exosomes improved motor function by inhibiting apoptosis and activating autophagy through the PI3K signaling pathway [211]. Resveratrol also blocks the Notch signaling pathway to limit SCI [212]. Liu et al. reported that resveratrol was an effective treatment for SCI by reducing neuronal cell apoptosis and inflammation [213]. In a recent study by Song et al., 10 mg/kg of resveratrol and puerarin-loaded nanoparticles attenuated CAT, SOD, and GSH antioxidant levels [214]. Resveratrol protected the spinal cord against ischemia-reperfusion injury via decreasing oxidative stress and NO release [215]. Treatment with 400 mg/kg of resveratrol for 10 days attenuated inflammation and oxidative stress after a rat model of SCI [216]. The i.p. injection of resveratrol (50 and 100 mg/kg) immediately after SCI markedly improved the activity of Na⁺/K⁺-ATPase and reduced the production of MDA in the injured spinal cord tissue [217].

6.10. Curcumin. Curcumin (C₂₁H₂₀O₆) is a natural phenol extracted from *Curcuma longa*. It has shown several pharmacological effects, such as anticancer [218], antiapoptotic, antioxidant, and anti-inflammatory properties [219]. Unlike other natural antioxidants containing either diketonic or phenolic groups, curcumin has both groups that help scavenge free radicals [220]. Curcumin decreased oxidative stress by activating the Keap1/Nrf2/ARE signaling in diabetic rats [221]. Two other studies confirmed that curcumin decreased MDA levels and increased SOD and GPx levels after spinal cord ischemia-reperfusion injury [222, 223]. Curcumin (300 mg/kg, i.p.) also improved ischemic brain injury by upregulating the Akt/Nrf2 pathway in a rat model of neurotoxicity [224]. It also activated the Nrf2/ARE signaling pathway after traumatic brain injury [225]. It upregulated HO-1 through PI3K/Akt/Nrf2 and reduced HO-2 in SH-SY5Y cells exposed to oxidative stress [226]. In a recent study, Daverey and Agrawal showed that 10 μM of curcumin and riluzole protected astrocytes derived from the human spinal cord against oxidative stress via Nrf2/HO-1 signaling [227]. Previously, Jiang et al. showed that curcumin in different concentrations (5, 10, 15, 20, and 25 mM) for 2 h activated Nrf2 target genes in primary spinal cord astrocytes, attenuated the level of intracellular ROS, and decreased oxidative damage and mitochondrial dysfunction [228]. Additionally, i.v. injection of curcumin (50, 100, and 200 mg/kg) for 7 days decreased p38MAPKs and MAPK kinase phosphorylation levels as key mediators in the microglia-mediated inflammatory response. They also showed that curcumin downregulated the expression levels of NF-κB post-SCI [229]. Curcumin (100 mg/kg) induced the Nrf2 signaling pathway in the SCI model [230] and decreased the levels of NO,

OH, and MDA following SCI [231]. In another study, curcumin (200 mg/kg, i.p.) immediately after the injury increased GPx, SOD, and CAT post-SCI [232]. Curcumin also decreased MDA concentration at the dose of 300 mg/kg [233].

Curcumin protected astrocytes from oxidative stress through the Nrf2/HO-1 signaling pathway [227]. In a dose of 200 mg/kg, for 8 weeks (i.p.), curcumin also decreased the levels of IL-6, IL-8, and TNF-α, as well as reduced astrogliosis and ameliorated motor function following SCI [234]. Also, curcumin in the dosage of 30 mg/kg (i.p.) for 7 consecutive days reduced the expression of NOS and NMDA following ischemic SCI [235] and in the dosage of 50 mg/kg (i.p.) significantly reduced the expression of IL-1β and NO in SCI mice [236]. Lin et al. showed the neuroprotective effects of curcumin by decreasing the expression of oxidative mediators (e.g., MDA) and iNOS in the SCI model study both *in vitro* (1 μM) and *in vivo* (40 mg/kg, i.p.) [237]. In another study, i.p. injection of curcumin was associated with decreasing the level of proinflammatory cytokines, such as IL-1β, TNF-α, and NF-κB, as well as inhibition of the generation of TGF-β1 and TGF-β2 [238]. Treatment with 40 mg/kg of curcumin caused a decrease in inflammatory factors and apoptosis following SCI and LPS-treated astrocytes [239].

Several other studies have shown that curcumin has anti-inflammatory, antiapoptotic, and antioxidant effects after SCI [229, 240–244] (Table 1).

6.11. Mangiferin. Mangiferin (C₁₉H₁₈O₁₁) is a bioactive xanthonoid isolated from different parts of the mango. It is a strong antioxidant [245] with numerous health-related properties, including immunomodulatory [246], antiviral [247], anti-inflammatory [248], antidiabetic [249], anticancer [250], and analgesic effects [251]. Mangiferin neutralizes free radicals by forming mangiferin-iron complexes and efficiently suppresses LPO and DNA damage [245]. In an *in vivo* study, 50 μmol/L of mangiferin for 4 h protected mice against cadmium chloride by reducing LPO content and incensement of GSH, CAT, GST, and SOD activity [252]. In an *in vitro* study, mangiferin enhanced the Nrf2 level, modulated NQO1 expression, and induced ROS levels [253]. Also, treatment with 20 μM of mangiferin for 2 h maintained renal cells against oxidative stress by targeting PI3K/Akt and Nrf2 signaling pathways [254]. In addition, in a neuronal injury model, 20 and 100 mg/kg mangiferin (i.p.) activated the Nrf2/HO-1 cascade in a dose-dependent manner [255]. Mangiferin (20 and 100 mg/kg, i.p.) for 30 days effectively reduced the level of MDA while increased SOD and CAT and GPx post-SCI [19]. The neuroprotective effects of mangiferin in doses of 10, 25, and 50 mg/kg for 30 days after SCI were associated with reduced spinal cord edema, inhibition of oxidative stress, inflammatory response, and regulation of Bcl-2 and Bax pathways [256].

6.12. Schisandrin B. Schisandrin B (C₂₃H₂₈O₆) is a dibenzocyclooctadiene derivative extracted from *Schisandra chinensis* [257]. It has shown promising antioxidant [258], anti-inflammatory [259], hepatoprotection [260], and

TABLE 1: Some preclinical studies on the effects of polyphenols in targeting oxidative stress and close interconnected mediators following SCI.

Phytochemicals	Dose	Pharmacological mechanisms	Animal models	References
Mangiferin	20, 40 mg/kg (i.p.); until 30 days post-SCI 10, 25, 50 mg/kg (i.p.); until 30 days post-SCI	↓ MDA; TNF- α ; IL-1 β ; NF- κ B ↑ SOD; GPx; CAT ↓ MDA; NF- κ B; TNF- α ; IL-1 β ; IL-6; caspase-3; caspase-9 ↑ CAT; SOD; GSH	SD male rats; contusion-SCI; T10; 10 g \times 12.5 mm SD rats; contusion-SCI; T8-T10; 6 g \times 100 mm	[19] [256]
Quercetin	0.2 mg/kg (i.p.); 1 h post-SCI 0.25 μ mol/kg (i.p.); 1 h post-SCI 10, 100 mg/kg (i.p.); first 72 h post-SCI 20 mg/kg (i.p.); twice daily until 7 days post-SCI 7.5 mg/kg (i.p.); twice daily until 10 days post-SCI 100 mg/kg (i.p.); until 3 days post-SCI 50 mg/kg (i.p.); immediately and 1 h post-SCI 30 mg/kg (i.p.); until 7 days post-SCI	↓ iNOS; p38MAPK; MDA; SOD ↓ MPO ↓ MDA; NO ↓ MDA; MPO; NO; IL-1 β ; IL-6; TNF- α ; caspase-3 ↑ SOD; GSH ↓ TNF- α ; IL-1 β ; p-STAT1; iNOS; NF- κ B; IL-12 ↑ IL-4; IL-10; TGF- β ↓ ROS; IL-1 β ; IL-18; TNF- α ↓ iNOS; MPO; COX-2; TNF- α ; IL-1 β ; PARP ↓ TNF- α ; RhoA	SD male rats; contusion-SCI; T10; 5 g \times 80 mm Wistar male rats; compression-SCI; T6-T7; 70 g force; 5 s Wistar male rats; compression-SCI; T7-T9; 70 g force 1 min Wistar albino rats; contusion-SCI; T7-T10; 10 g \times 100 mm SD male rats; compression-SCI; T8 SD female rats; compression-SCI; T10 SD male rats Female BALB/c mice; contusion-SCI; T8-T9; 2 g \times 25 mm SD male rats; contusion-SCI; T9-T10; 10 g \times 25 mm Wistar female mice; compression-SCI; T5-T12; 63 g force; 1 min Wistar albino male rats; compression-SCI; T8-T12; 1.37-1.72 N force; 1 min Wistar female rat; hemitranssection-SCI; T10 SD female rats; contusion-SCI; T10; 25 g/cm SD female mice; compression-SCI; T9-T10 SD male rats; contusion-SCI; T8-T9; 10 g \times 50 mm Rats; contusion-SCI T10-T11; 12 g \times 7 mm SD female rats; compression-SCI; T9-T10; 20 g force; 1 min	[61] [72] [73] [75] [77] [78] [89] [93] [94] [109] [111] [112] [131] [161] [171] [174] [186]
EGCG				
Protocatechuic acid	50 mg/kg (i.p.) 10 μ L; 1 μ g/kg (i.t.); 1 h post-SCI	↓ TNF- α ; IL-1 β ; COX-2; iNOS; MMP-9 ↓ MDA; SOD; TOA ↑ TAC		
Caffeic acid phenethyl ester	10 μ g/kg (i.p.); 30 min post-SCI 10 μ mol/kg (i.p.); until 4 weeks post-SCI 20 mg/kg (i.p.) 50, 100 mg/kg (i.p.); until 3 days post-SCI	↓ IL-1 β ; TNF- α ↓ COX-2; NOS; IL-1 β ↓ MDA; ROS; IL-1 β ; IL-18; TNF- α ↓ MDA; ROS; IL-1 β ; IL-18; TNF- α ↓ MDA; p38MAPK; IL-1 β ; IL-6; TNF- α ; NF- κ B; ↑ SOD; GSH; CAT ↓ TNF- α ; MDA; ROS; TGF- β 1; Smad2 ↑ MPO ↓ IL-1 β ; TNF- α ; NF- κ B; PARP; ↑ SOD; CAT; Nr2f1; HO-1; p-p38		
Honokiol				
Rutin				
Hesperidin				

TABLE 1: Continued.

Phytochemicals	Dose	Pharmacological mechanisms	Animal models	References
Oxyresveratrol	10, 20 mg/kg (i.p.); until 4 weeks post-SCI	↓ MDA; IL-1 β ; IL-6; TNF- α ; NF- κ B; COX-2; iNOS; ↑ Nrf2; SOD; GPx; GSH	SD female rats; contusion-SCI; T10; 10 g \times 25 mm	[206]
Polydatin	20, 40 mg/kg (i.p.); 30 min post-SCI 100 mg/kg (i.p.); immediately post-SCI 100 mg/kg (i.p.); until 2 weeks post-SCI 10 mg/kg (i.p.)	↓ MDA; NO; iNOS; IL-1 β ; IL-6; TNF- α ; NF- κ B; ↑ SOD; Nrf2; HO-1 ↓ MDA; NO; XO; ↑ GSH ↓ Caspase-3; ↑ LC3; Beclin1; PI3K ↓ MDA; p38MAPK; ↑ SOD; CAT; GSH	SD male rats; contusion-SCI; T8; 10 g \times 50 mm Wistar male rats; contusion-SCI; T7-T10; 5 g \times 100 mm SD rats; contusion-SCI; T9-T10 Wistar male rats; Spinal cord ischemia-reperfusion injury	[208, 209] [201] [211] [214]
Resveratrol	1 and 10 mg/kg; 30 min pre-SCI 400 mg/kg (p.o.); until 10 days post-SCI 50, 100 mg/kg (i.p.); immediately post-SCI 200 mg/kg (i.p.); until 3 days post-SCI 50, 100, 200 mg/kg (i.v.); until 7 days post-SCI 100 mg/kg (i.p.) 200 mg/kg (i.p.); immediately post-SCI 300 mg/kg (i.p.)	↓ NO; MDA ↓ MDA; IL-6 ↑ TCA ↓ MDA; ↑ Na ⁺ , K ⁺ -ATPase activities ↓ MDA; MPO; IL-1 β ; IL-10; TNF- α ; ↑ SOD ↓ p38MAPK; NF- κ B ↓ MDA; NO; TBARS ↑ SOD; GPx; CAT ↓ MDA	SD male and female rats; contusion-SCI; T8; 10 g \times 25 mm SD rats; contusion-SCI; T10; 10 g \times 25 mm Female mice; contusion-SCI; T9-T10; 3 g \times 30 mm Long Evans female rats; contusion-SCI; 10 g \times 25 mm Wistar male rats; compression-SCI; T7; 70 g force; 1 min Wistar albino female rats; contusion-SCI; T7-T9; 5 g \times 100 mm	[215] [216] [217] [285] [229] [231] [232] [233]
Curcumin	200 mg/kg (i.p.); until 8 weeks post-SCI 30 mg/kg (i.p.); until 7 days post-SCI 50 mg/kg (i.p.); until 7 days post-SCI 40 mg/kg (i.p.); 30 min post-SCI 300, 100, 30 mg/kg (i.p.); until 7 days post-SCI 40 mg/kg (i.p.); 30 min post-SCI	↓ MDA; IL-6; IL-8; TNF- α ; p-ERK; p-p38; p-JNK; p-STAT; GFAP; ↑ SOD ↓ NMDA; iNOS ↓ IL-1 β ; NO; NF- κ B; STAT3 ↓ MDA; iNOS ↓ TNF- α ; IL-1 β ; NF- κ B; TGF- β 1; TGF- β 2 ↓ iNOS; ↑ Bcl-2, Cisd2	SD female rats; contusion-SCI; T9-T10; 30 g force; 2 min SD rats; ischemic SCI BALB/c female mice; compression-SCI; T8-T9; 30 g force, 1 min SD male rats; contusion-SCI; T9-T10; 10 g \times 25 mm SD male rats; compression-SCI; T8-T10; 50 g force; 1 min C57BL/6JNarl mice; hemisection and contusion-SCI; T9-T10; 10 g \times 25 mm	[234] [235] [236] [237] [238] [239]

TABLE 1: Continued.

Phytochemicals	Dose	Pharmacological mechanisms	Animal models	References
	100 mg/kg (i.p.); until 7 days post-SCI	↓ SOX9; NF- κ B; GFAP; CD45; CD11b	SD female rats; compression-SCI; T8-T10; 50 g force; 1 min	[240]
	60, 6 mg/kg (i.p.); until 4 weeks post-SCI	↓ IL-4; IL-1 β ; IL-2; IL-6; IL-12; TNF- α ; MIP-1 α ; NF- κ B; GFAP	Wistar male rats; balloon compression-SCI; T8-T10	[241]
	40 mg/kg (i.p.); 30 min post-SCI	↓ GFAP; ↑ p-JAK2; p-STAT3	SD male rats; contusion-SCI; T8	[242]
	75, 150, 300 mg/kg (i.p.); 20 min and until 3 days post-SCI	↓ TNF- α ; NF- κ B	SD male rats; compression-SCI; T8-T10	[243]
	100 mg/kg (i.p.); 15 min post-SCI	↓ TNF- α ; IL-1 β ; NF- κ B; TLR4	SD male rats; compression-SCI T8-T9; 30 g force	[244]
	200 mg/kg (p.o.)	↓ MDA; ↑ SOD	Wistar albino male rats; contusion-SCI T7-T10; 5 g \times 100 mm	[286]
	200 mg/kg (i.p.); until 7 days post-SCI	↓ MDA; ↑ SOD	SD male rats; contusion-SCI T8	[287]
	60 mg/kg (i.p.); 30 min and weekly for 3 weeks post-SCI	↓ Caspase-3; IL-6; IL-1 β ; TNF- α ; p-mTOR; p-Akt; Iba1; GFAP ↑ IC3	SD male rats; contusion-SCI; T9-T11; 10 g \times 20 mm	[288]
Apocynin	200 mg/kg (i.p.); until 8 weeks post-SCI	↓ Caspase-3; Bax; GFAP; ↑ Bcl-2	SD male rats; contusion-SCI; T9-T10; 10 g \times 12.5 mm	[289]
	50 mg/kg (i.p.); twice daily until 3 days post-SCI	↓ MPO; MDA; TNF- α ; IL-1 β ; IL-6; ↑ GSH; SOD	SD male rats; compression-SCI; T6-T7; 24 g force; 1 min	[268]
Baicalin	50, 100 mg/kg (p.o.)	↓ MDA; IL-1 β ; IL-6; TNF- α ; NF- κ B; ↑ SOD; GPx; CAT	SD male mice; compression-SCI; T7; 23.8 g force; 2 min	[270]
	10, 30, 100 mg/kg (i.p.); immediately and 24 h post-SCI	↓ MDA; NF- κ B; TNF- α ; Bax; ↑ GSH; Bcl-2	SD male rats; contusion-SCI; T12; 50 g	[271]
Gastrodin	100, 200 mg/kg (i.p.); until 5 days post-SCI	↑ SOD; GSH; Nrf2; GCLc; GCLm	SD male rats; contusion-SCI; T10; 8 g \times 30 mm	[272]
Mulberrin	15, 30 mg/kg (p.o.); until 4 weeks post-SCI	↓ TBARS; ROS; IL-1 β ; IL-6; TNF- α ; NF- κ B; PARP; ↑ SOD; GSH; Nrf2; HO-1	SD male rats; compression-SCI; T10; 30 g force; 1 min	[273]
Salvianolic acid A	2.5, 5, 10 mg/kg (i.p.); until 7 days post-SCI	↑ Nrf2; HO-1	SD male rats; compression-SCI; T12; 50 g force; 5 min	[274]
Salvianolic acid B	1, 10, 50 mg/kg (i.p.); until 3 days post-SCI	↓ TNF- α ; NF- κ B; ↑ HO-1	SD male rats; compression-SCI; T12; 50 g force; 5 min	[275]
	50 mg/kg (i.p.); immediately, 6 h and 12 h, and then one day until 7 days post-SCI	↓ MMP-9; TNF- α ; IL-1 β ; IL-6; COX-2; iNOS	SD rats; contusion-SCI; T9-T10; 10 g \times 25 mm	[276]
Gallic acid	10 mg/kg (i.p.); until 10 days post-SCI	↓ COX-2; NF- κ B	Wistar male rats; contusion-SCI; T9; 15 g \times 25 mm	[277]

TABLE 1: Continued.

Phytochemicals	Dose	Pharmacological mechanisms	Animal models	References
Kaempferol	1 mg/kg (i.p.); 1 and 6 hours post-SCI 27 and 2.7 μ M	↓ IL-1 β ; IL-1 β ; ROS; p-p38MAPK; p-JNK; NF- κ B	SD rat; hemicontusion-SCI	[278]
Luteolin and palmitoylethanolamide	50, 100 mg/kg (p.o.); 3 days pre-SCI until 7 days post-SCI 20 mg/kg (i.p.); immediately and 1 h post-SCI	↓ nNOS; iNOS; COX-2	CD1 mice; compression-SCI; T5-T8; 24 g force; <i>in vitro</i> : spinal cord slices	[279]
Naringenin	50, 100 mg/kg (p.o.); 3 days pre-SCI until 7 days post-SCI 20 mg/kg (i.p.); immediately and 1 h post-SCI	↓ TNF- α ; IL8; IL-1 β ; IL-6	SD male rats; compression-SCI T9-T11	[280]
Oleuropein	250 μ L at the dose of 20/40/100 μ g/kg (i.p.); 1 h and 6 h post-SCI	↓ MDA; Bax; ↑ Bcl-2; GSH	SD female rats; contusion-SCI; T9; 10 g \times 25 mm	[281]
Pycnogenol	100 mg/kg (i.p.); 15 min post-SCI	↓ MPO; IL-1 β ; TNF- α ; NF- κ B; LPO; TBARS; ↑ GDNF	CD1 male mice; compression-SCI; T5-T8; 24 g; 1 min	[282]
Silymarin	Injected 1–2 mm caudal and rostral to the epicenter; 5 min post-SCI	↓ Bax; caspase-3; MDA; ↑ Bcl-2; SOD; MMP	Wistar male rats; contusion-SCI; T10; 10 g \times 50 mm	[283]
Carnosol	5 mg/kg (i.p.); 1 h post-SCI	↓ ROS; NF- κ B; COX-2; iNOS; IL-1 β	SD rats; contusion-SCI; T9-T10; 10 g \times 50 mm	[284]
Tetrahydrocurcumin	80 mg/kg (i.p.); until 2 weeks post-SCI	↓ COX-2; NF- κ B; IL-1 β ; IL-6; TNF- α ; ↑ TAC; GSH; GPx; GST; SOD; CAT; p-Akt; Nrf2 ↓ MDA; TNF- α ; IL-6; IL-1 β ; NF- κ B; ↑ SOD; GSH; GPx; p-Akt; FoxO-4	SD female rats; contusion-SCI; T7-T10; 10 g \times 100 mm SD male rats; compression-SCI; T8; 30 g force; 2 min	[290] [291]

Akt: tyrosine kinase B; Bax: Bcl-2-associated X protein; Bcl: B cell lymphoma; caspases: cysteine-dependent aspartate-directed proteases; CAT: catalase; COX: cyclooxygenase; eNOS: endothelial nitric oxide synthase; ERK: extracellular signal-regulated kinases; FoxO-4: forkhead box O4; GCLC: catalytic subunit of glutamate-cysteine ligase; GCLm: modifying subunit of glutamate-cysteine ligase; GFAP: glial fibrillary acidic protein; GSH: glutathione; GST: glutathione S-transferase; GPx: glutathione peroxidase; HO-1: heme oxygenase-1; Iba1: ionized calcium-binding adaptor molecule 1; IL: interleukin; iNOS: inducible nitric oxide synthase; i.p.: intraperitoneal; i.t.: intrathecal; i.v.: intravenous; JAKs: Janus kinases; JNK: c-Jun N-terminal kinase; LC3: light chain 3; LPO: lipid peroxidation; MAPK: mitogen-activated protein kinase; MDA: malondialdehyde; MIP-1 α : macrophage inflammatory protein 1-alpha; MPO: myeloperoxidase; mTOR: mammalian target of rapamycin; NADPH oxidase: nicotinamide adenine dinucleotide phosphate; NF- κ B: nuclear factor kappa B; NO: nitric oxide; Nrf2: nuclear factor erythroid 2-related factor 2; PARP: poly(ADP-ribose) polymerase; p.o.: oral administration; ROS: reactive oxygen species; SCI: spinal cord injury; SD: Sprague Dawley; SOD: superoxide dismutase; SOX9: SRY-box transcription factor 9; STATs: signal transducer and activator of transcription proteins; T: thoracic; TAC: total antioxidant capacity; TBARS: thiobarbituric acid reactive substance; TLR4: transmembrane lipopolysaccharide receptor; TNF- α : tumor necrosis factor-alpha; XO: xanthine oxidase.

neuroprotection activities [261]. Schisandrin B prevented cardiotoxicity by reducing oxidative stress, DNA damage, and inflammation by inhibiting the MAPK/p53 pathway [262]. Schisandrin B (50 mg/kg) prevented the progression of liver fibrosis by modulation of Nrf2/ARE and TGF- β /Smad signaling pathways [263]. Orally administered Schisandrin B at the dose of 30 or 60 mg/kg alleviated oxidative stress through modulation of the Nrf2/Keap1 antioxidant pathway [264]. The ability of Schisandrin B in activating Nrf2 is related to its biotransformation through cytochrome P450-catalyzed reactions [265]. In another study, a 50 mg/kg dose of Schisandrin B (p.o., for 5 days) diminished oxidative stress and inflammatory responses induced by SCI via inhibition of p53 signaling [266]. Oral administration doses of 20, 40, and 80 mg/kg of Schisandrin B for 5 days before SCI attenuated oxidative stress and inflammation through p38MAPK, ERK1/2, and NF- κ B pathways after an ischemia/reperfusion injury model [267].

The antioxidant effects of some other polyphenols/phenolic compounds, including apocynin [268, 269], baicalin [270, 271], gastrodin [272], mangiferin [19], mulberrin [273], salvianolic acid [274, 275], gallic acid [276, 277], kaempferol [278], luteolin [279], naringenin [280], oleuropein [281, 282], pycnogenol [283], silymarin [284], and protocatechuic acid [94], have been shown in Table 1. From the chemical point of view, Figure 2 presents the candidate chemical structures of neuroprotective polyphenols against SCI.

7. Extrinsic Molecular Mechanisms of Regeneration Post-SCI: Therapeutic Approaches to Polyphenols

After SCI, axons and dendrites lose their connection to the main neural pathways, and the process is termed Wallerian degeneration and axon searing. In this line, anterograde and retrograde degeneration of neural fibers could be the main factor in the declining tissue mass at the injured spinal cord [292]. One of the goals of SCI-related research is to repair disrupted neural networks and regenerate axons, which is hoped to reconnect descending neural fibers with their original targets [293]. There are several theories of post-SCI regeneration, predominantly the Nogo signaling pathway, the glial scar mechanism, and the chondroitin sulfate proteoglycan-related pathway.

7.1. The Nogo Signaling Pathway. In 2000, Nogo-A (reticulin 4) was introduced as the myelin-associated neurite outgrowth inhibitor [294]. It is expressed by oligodendroglia and neurons but not astrocytes [295]. Three Nogo receptors have been reported, NgR1, NgR2, and NgR3 [296]. Many studies reported that blocking Nogo receptors promotes axonal sprouting [297–299]. In an *in vitro* study by Gundimeda et al., polyphenols extracted from green tea prevented antineuritogenic properties of Nogo-A through H₂O₂ and laminin receptors [300].

Nogo, myelin-associated glycoprotein (MAG), and oligodendrocyte myelin glycoprotein (OMgp) trigger interconnected receptors that trigger the Ras homolog gene family

member A (RhoA) and its effector, Rho-kinase (ROCK), which lead to the inhibition of axonal growth [301]. RhoA controls cellular motility via the regulation of cytoskeletal rearrangements. It also activates phospholipases A2, C, and D and serine/threonine kinases [302], promoting the expression of COX-2 at the transcription level depending on NF- κ B [303]. Protein kinase A (PKA) via RhoA phosphorylation regulates the morphological and functional subsequences of cAMP. Besides, RhoA modulates several other cellular functions, such as assembly of extracellular fibronectin, potassium channels, neurotransmitter exocytosis, thrombin-induced cell death, gap junctions, cellular uptake of low-density lipoprotein, expression of cyclin D1, basal muscle tone, IL-8, and neurite growth cones [302]. Reports have shown that SCI upregulates Rho proteins. The Y27632 and fasudil (ROCK-specific inhibitors) significantly reduced IL-6 synthesis, contributing to inflammation-induced CNS regeneration [304, 305].

Phosphatase and tensin homolog (PTEN) catalyzes the conversion of PIP3 to PIP2, followed by Akt activation via phosphoinositide-dependent protein kinase 1 (PDK1). Akt, in turn, stimulates inflammatory messengers Raf/Rac/NF- κ B and thereby inhibits tuberous sclerosis complexes 1 and 2 (TSC1/2) to suppress Ras homolog enriched in brain 1 (Rheb1). Rheb1 activates mTOR to stimulate protein synthesis and cell growth. Activation of Akt also inhibited GSK-3 β , which disinhibits Wnt/ β -catenin to stimulate mTOR and cell proliferation. The cAMP turns on phosphokinase A, which inhibits Rho and Rho-kinase. Rho and Rho-kinase inhibit cell growth when activated by axonal growth inhibitors Nogo and CSPG [302]. In line with this, ROS, which increased axonal dieback and degeneration post-SCI, are also needed for axonal regeneration and functional recovery after SCI. ROS via the release of exosomal NADPH oxidase 2 complexes and oxidized PTEN lead to its inactivation, thus stimulating the PI3K/Akt pathway toward axonal regeneration [306, 307].

Some natural substances, such as curcumin, affected Rho expression and reduced the protein levels of phosphokinase C, NF- κ B, and RhoA [308]. In another study, curcumin was highlighted as a novel inhibitor of the Rho/ROCK pathway to develop axonal growth following SCI *in vivo/in vitro* [309]. As reported on other polyphenols, Álvarez-Pérez et al. showed that EGCG meaningfully downregulated RhoA and astroglial and microglial activity in the spinal cord *in vivo* [93].

7.2. The Glial Scar Mechanism. After SCI, astrocytes play a key role in SCI pathology through a phenotypic change called reactive gliosis. Immediately after injury, astrocytes, as a type of glial cells, proliferate and accumulate around the lesion, separating healthy tissue from the surrounding damaged tissue. Reactive astrocytes migrated to the lesion epicenter, and secluded inflammatory cells resulted in tissue repair and functional improvement [310]. Consequently, reactive astrocytes made the astrocytic scar as the main obstacle to axonal regeneration [293]. A number of processes and proteins such as microgliosis [311], nestin [312], neurotrophins [313], albumin [314], oxygen, and glucose

deprivation contribute to gliosis [315]. In addition, after SCI, intensive inflammatory response resulted in the activation of resident microglia and facilitated the infiltration of macrophages into the lesion post-SCI. Macrophages play main roles in the inflammation process and are known as proinflammatory (M1) and anti-inflammatory (M2) mediators [316, 317]. Interventions that relieve inflammation, reduce secondary damage after SCI, and modulate the phenotypes of infiltrating macrophages can be a therapeutic strategy to improve functional recovery post-SCI [318]. Microglia also secrete a set of signaling molecules, such as chemokines, cytokines, and growth factors. Many of these factors act via receptors on oligodendrocyte progenitor cells. For example, TNF- α , secreted by microglia/macrophages as well as astrocytes in mouse cuprizone-induced lesions, induced the proliferation and remyelination of oligodendrocyte precursor cells (OPCs) via TNFR2 on polydendrocytes [319]. Type I interferon, bone morphogenic proteins [302], and STAT3 knockout inhibited reactive astrocytosis [320]. Blocking STAT3 was associated with preventing astrogliosis, leading to a higher level of inflammation, an increase in lesion volume, and a reduction in motor recovery after SCI. Besides, TGF- β 1 plays a critical role in regulating glial activation. Proinflammatory cytokines trigger the astrocytes to secrete chemokines that are antagonized by TGF- β 1 [321].

These findings revealed that curcumin developed axon/neuron protection of the spinal cord through suppressing the formation of glial scars [241] and the inflammatory astrocyte marker (Iba1) [236]. Nanoformulation of curcumin also mitigated glial scar formation and protected the white matter following SCI in rats [322]. By decreasing the formation of glial scars, curcumin and EGCG increased axonal sprouting and modulated MIP-1- α , IL-1 β , IL-4, and IL-6 in rats post-SCI [92]. The synergistic effects of curcumin with mesenchymal stem cells prevented axonal sprouting, glial scar formation, and inflammatory responses in rats following SCI [323]. As glial cells closely connect with NF- κ B, curcumin also regulated glial scar formation by inhibiting NF- κ B and TGF- β -SOX9 [324].

Resveratrol has also shown promising inhibitory effects on NF- κ B following SCI [325]. Another study by Zhang et al. showed that resveratrol improves neuronal regeneration of the spinal cord passing through the attenuation of Notch signaling [212]. As another phenolic compound, quercetin provided functional recovery, axonal regeneration, and astrocyte activation in rats post-SCI through BDNF and JAK2/STAT3 signaling pathways [71]. Besides, treatment with quercetin regulated oligodendrocyte necroptosis by inhibiting M1 macrophage/microglial polarization and reducing myelin/axon loss following SCI in rats [77].

Naringenin promoted the remyelination of the spinal cord by regulating the differentiation of oligodendrocyte precursor cells passing through the GSK-3 β / β -catenin pathway [326]. Consequently, triggering axonal sprouting and positive modulation of glial scars, as well as inflammation (e.g., NF- κ B) suppression, also resulted from treating with EGCG in rats following SCI [82].

In a previous study reported by Subbarayan et al., caffeic acid hydrogel, another phenolic compound, could improve

functional recovery and remyelination after hemitranssection SCI. They found that such hydrogel decreased IL-6, TNF- α , and neutrophil release. Caffeic acid hydrogel also modulated gamma-aminobutyric acid and glutamine levels, as well as regulation of genes/proteins involved in locomotor activity after SCI [327]. Figure 1 presents the cross-talk between extrinsic dysregulated pathways and intrinsic ones in the pathogenesis of SCI.

7.3. Chondroitin Sulfate Proteoglycan-Related Pathways. Studies show that chondroitin sulfate proteoglycans (CSPGs) repel regenerative axons and inhibit oligodendrocyte remyelination/maturation [328]. Destruction of CSPGs by chondroitinase ABC (ChABC) is a potential therapeutic strategy for breaking the inhibitory barrier, synaptic reorganization, and functional recovery of SCI [329]. ChABC may act through multiple mechanisms. Several studies have shown that ChABC degrades chondroitin sulfate proteoglycans and promotes morphological plasticity. It also reduces glial scars and axon dieback, regenerates spinal respiratory pathways, promotes axonal conduction and functional recovery, stimulates remyelination, and enhances migration of oligodendroglial and neural progenitor cell migration [302]. ChABC in combination with neural stem/progenitor cells (NSPCs) improved functional recovery after SCI [330].

Many studies have reported that injured spinal axons will grow across SCI sites transplanted with neural stem cells [331], Schwann cells [332], mesenchymal stem cells [333], olfactory ensheathing glia [334], or BMSCs [335]. The spinal cord has been shown to be regenerated after treatment with Schwann cells [336] or PTEN deletion [337]. Figure 1 also reveals the interconnections between degenerative/regenerative extrinsic mechanisms and oxidative stress post-SCI.

Table 2 indicates the critical role of polyphenols in modulating extrinsic regenerative/degenerative mechanisms following SCI.

8. Combination Therapy of Polyphenols and Stem Cells Post-SCI

Stem cells are undifferentiated cells with the potential to differentiate into different cells. In recent years, stem cell therapy for neurodegenerative diseases has shown a promising prospect. Many studies have reported that polyphenols can target and promote the potential of stem cells against neurodegeneration [338]. For example, curcumin induced neuronal differentiation in neural stem cells by activating the Wnt pathway and upregulating the genes required for cell differentiation [339]. Ruzicka et al. reported that combination therapy of curcumin in a high dose (60 mg/kg) and mesenchymal stem cells synergistically improved recovery from the balloon compression-SCI model and modulated expression of ILs, TNF- α , and MIP-1 α [323]. Previously, these results were shown by Ormond et al. [340]. Wanjiang et al. suggested that curcumin in a dose of 100 mg/kg suppressed the apoptosis of human umbilical cord-derived mesenchymal stem cells (hUC-MSCs) through the ERK1/2 pathway. Additionally, a combined administration of curcumin with hUC-MSCs improved motor function in female

TABLE 2: Targeting extrinsic regenerative/degenerative mechanisms by polyphenols post-SCI.

Phytochemicals	Dose	Pharmacological effects	Animal models	References
Quercetin	20 mg/kg (i.p.); until 7 days post-SCI	↓ Cavity formation; glial scar (GFAP); ↑ axonal regeneration (5-HT; NF200); locomotor recovery	SD male rats; contusion-SCI T8; 50 g × cm	[71]
	7.5 mg/kg (i.p.); twice daily until 10 days post-SCI	↑ Locomotor recovery; preventing necroptosis; preservation of myelin and axonal	SD male rats; compression-SCI T8	[77]
EGCG	50 mg/kg (directly to the spinal cord surface) until 4 weeks post-SCI	↓ Glial scar (GFAP); ↑ growth factors (FGF2; VEGF); axonal sprouting (GAP43); locomotor recovery; preservation of white and grey matter; modulation macrophage markers (M1 and M2)	Wistar rats; balloon compression-SCI T8	[82]
Curcumin and EGCG	60 mg/kg 17 mg/kg (i.p.); until 4 weeks post-SCI	↓ Glial scar (GFAP); ↑ axonal sprouting (GAP43; Olig2); locomotor recovery; tissue regeneration; neuroprotective effects	Wistar male rats; balloon compression- SCI T10	[92]
EGCG	30 mg/kg (i.p.); until 7 days post-SCI	↓ TNF- α ; RhoA; glial scar (GFAP) ↑ Sensory recovery	Female BALB/c mice; contusion-SCI T8-T9; 2 g × 25 mm	[93]
Curcumin	60, 6 mg/kg (i.p.); until 4 weeks post-SCI	↓ Glial scar (GFAP); ↑ axonal sprouting (GAP43; Olig2); locomotor recovery Anti-inflammatory effects	Wistar male rats; balloon compression- SCI T8-T10	[241]
Polymer-curcumin and ependymal progenitor/stem cell	405 mg; 1.1 mmol	↓ ROCK1; RhoA; GAP43; p-Limk1; glial scar (GFAP); microglial activation (Iba1); macrophage infiltration (ED1); ↑ locomotor recovery	SD rats; Horizon Impactor-SCI T8; 250 kdyn	[309]
Nanoformulated curcumin	0.01 mL	↓ GAP43; IL-1 β ; TNF- α ; IFN- γ ; IL-6; glial scar; ↑ functional recovery; preservation of white matter	Wistar male rats; balloon compression- SCI T10	[322]
Curcumin and mesenchymal stem cells	60, 6 mg/kg (i.p.); until 4 weeks post-SCI	↓ Glial scar (GFAP); ↑ axonal sprouting (GAP43); locomotor and sensory recovery; preservation of white and grey matter	Wistar rats; balloon compression-SCI T8	[323]
Naringin	20, 40 mg/kg (p.o.); until 4 weeks post-SCI	↑ Remyelination (GSK-3 β ; β -catenin); axonal sprouting (NG2; Olig2); locomotor recovery	SD female rats; contusion-SCI T10	[326]
Caffeic acid hydrogel and human gingival-derived neural stem cell		↓ Glial scar (GFAP); ↑ axonal growth; recovery of damaged spinal tissue; locomotor recovery Maturation of neuronal cells	Wistar female rats; hemitranssection-SCI T9-T11	[327]

EGCG: epigallocatechin gallate; GAP43: Growth Associated Protein 43; GFAP: glial fibrillary acidic protein; GNSCs: human gingival-derived neural stem cell; GSK3B: glycogen synthase kinase-3 beta; Iba1: ionized calcium-binding adaptor molecule 1; i.p.: intraperitoneal; Olig2: oligodendrocytes; p.o.: oral administration; ROCK1: Rho-associated coiled-coil containing protein kinase 1; SCI: spinal cord injury; SD: Sprague Dawley; T: thoracic; TGF- β : transforming growth factor-beta; TNF- α : tumor necrosis factor-alpha; VEGF: vascular endothelial growth factor.

rats after compression-SCI in a dose-dependent manner [341]. It has also been shown that intrathecal administration of curcumin can improve the functional recovery post-SCI through the spinal cord-NSPCs (SC-NSPCs) by increasing the expression of SC-NSPCs and decreasing the activity of reactive astrogliosis (GFAP) and lesion cavities [342]. In another study, combined therapy of polymer-curcumin with ependymal progenitor/stem cells (epSPCs) by intrathecal injection (10 μ M) enhanced functional recovery and axonal growth post-SCI in rats [309]. In addition, concomitant

use of human-induced neural and mesenchymal stem cells (iPSC-NSC) with a polyacetal-curcumin nanoconjugate provided neuroprotective and immunomodulatory effects that led to the prevention of axonal degeneration and neuronal death, reduced the injury area, and inhibited the formation of glial scars in chronic stages in a rat model of contusive SCI [343].

In an *in vitro* study, to mimic the microenvironment around transplanted cells in the damaged spinal cord, as a glucoside of resveratrol, polydatin protected BMSCs against




FIGURE 3: Oxidative stress and cross-talk signaling mediators post-SCI. The modulatory role of polyphenols. Akt: tyrosine kinase B; ARE: antioxidant response element; Ca^{2+} : calcium; CAT: catalase; COX: cyclooxygenase; GABAR: gamma-aminobutyric acid receptor; GSH: glutathione; GPx: glutathione peroxidase; IKKB: I κ B kinase; IL: interleukin; Keap1: Kelch-like ECH-associated protein 1; LPO: lipid peroxidation; MAPK: mitogen-activated protein kinase; MDA: malondialdehyde; MOR: morphine receptor; NADPHO: nicotinamide adenine dinucleotide phosphate oxidase; Na^+/K^+ : sodium/potassium; NF- κ B: nuclear factor kappa-light-chain-enhancer of activated B; NOS: nitric oxide synthase; NR2B: N-methyl-D-aspartate receptor subtype 2B; Nrf2: nuclear factor erythroid 2-related factor 2; PARP: poly(ADP-ribose) polymerase; PI3K: phosphoinositide 3-kinase; SOD: superoxide dismutase; SCI: spinal cord injury; TNF- α : tumor necrosis factor-alpha.

oxidative injury caused by H_2O_2 through the Nrf2/ARE pathway [210]. Combining BMSCs and a 400 mg/kg dose of green tea polyphenols, for 30 days before SCI, improved motor function by improving blood-spinal cord barrier integrity and NF- κ B decrement [344]. In a recent *in vivo* study on SCI by Subbarayan et al., treatment with human gingival-derived neural stem cells transplanted in the injectable caffeic acid bioconjugated hydrogel was associated with decreased IL-6 and TNF- α and increased IL-10; it was able to regenerate damaged spinal cord [327].

Figure 3 demonstrates the major targets of polyphenols in combating oxidative stress, inflammation, and cross-linked pathways after SCI.

9. Clinical Potentials of Polyphenols against Oxidative/Inflammatory Disorders

Polyphenols are promising secondary metabolites with various health benefits. Curcuminoids have been approved by the US Food and Drug Administration (FDA) as generally

safe and well-tolerated active metabolites in clinical trials, even in doses up to 12,000 mg/day [345].

Medicinal uses of curcumin have been reported with significant effects in protecting the liver and improving gastrointestinal function, respiration, cardiovascular, anti-inflammatory, anticancer, antidiabetic, antiallergic, neuroprotective, and chemoprotective [346] activities. A randomized, double-blind, placebo-controlled (RDBPC) study with 108 male adults aged between 31 and 59 years reported that curcumin in a dose of 1000 mg/day for 6 weeks (p.o.) decreased TNF- α and IL-1 β levels and increased plasma BDNF [347].

In a controlled clinical trial study, the effects of curcumin on inflammatory and stress markers were evaluated in 100 patients of both sexes with osteoarthritis. Those studies showed that administration of curcumin in the dose of 200 mg/day for 8 months (p.o.) reduced inflammatory markers IL-1 β , IL-6, and soluble ligand CD40 (sCD40L) [348]. In another study, prospective randomized open-ended blinded evaluation (PROBE) on 80 patients with knee osteoarthritis, 30 mg/3 times in a day of curcumin (p.o.) for

TABLE 3: Clinical potentials of polyphenols against oxidative/inflammatory disorders.

Polyphenols	Clinical study	Population	Protocol	Therapeutic findings	References
Curcumin	Major depressive disorder DBRPC	$N = 108$; M ; age = 31-59	1000 mg/day of curcumin/soybean powder; 6 weeks	↓ TNF- α and IL-1 β ; ↑ BDNF	[347]
	Osteoarthritis Controlled clinical trial	$N = 100$ Treatment (50): $M = 23$ $W = 27$ Age = 43.6 Control (50): $M = 28$ $W = 22$ Age = 44.2	1000 mg/day of Meriva (containing 200 mg curcumin); 32 weeks	↓ IL-1 β and IL-6	[348]
	Osteoarthritis Randomized controlled trial	$N = 80$ Curcuminoid (39): $M = 15$ $W = 24$ Diclofenac (41): $M = 12$ $W = 29$	30 mg of curcuminoid; 25 mg of diclofenac sodium; 3 times daily, 4 weeks	↓ COX-2	[349]
	Osteoarthritis RBDCT	$N = 53$ Curcuminoid (19): $M = 5$ $W = 14$ Age = 57.32 Placebo (21): $M = 4$ $W = 17$ Age = 57.57	1500 mg/day of curcuminoid/placebo; 6 weeks	↓ TNF- α , TGF- β , IL-6, substance P, and CGRP	[350]
	Type 2 diabetes Clinical trial		400 mg/3 times a day	↓ MDA, IL-6, and TNF- α	[351]
	SCI Randomized clinical trial	$N = 20$; age = 30-67 Treatment (12): $M = 6$ $W = 6$ Control (8): $M = 4$ $W = 4$	Omega-3 250-500 mg/3 times a day; Chlorella 1000 mg/6 times a day; antioxidants (100 mg coenzyme Q10, 200 mg N-acetylcysteine, 150 mg mixed tocopherols, 100 mg DL-alpha-lipoic acid, 60 mg green tea extract, 5.5 mg zinc, and 100 μ g selenium, 2 times a day); curcumin 400 mg/3 times a day; 12 weeks	↓ IL-1 β	[353]
	Multiple sclerosis Randomized controlled trial	$N = 50$ Curcuminoid (25): $M = 9$ $W = 16$ Age = 35.2 Placebo (25): $M = 10$ $W = 15$ Age = 34.6	80 mg/day of nanocurcumin; 24 weeks	↓ TGF- β and IL-10	[354]
	Metabolic syndrome Randomized controlled trial	$N = 117$ Curcuminoid (58): $M = 35$ $W = 23$ Age = 44.80 Placebo (59): $M = 32$ $W = 27$	1000 mg/day of curcuminoid/placebo; 8 weeks	↓ MDA, TNF- α , TGF- β , and IL-6; ↑ SOD	[370, 371]

TABLE 3: Continued.

Polyphenols	Clinical study	Population	Protocol	Therapeutic findings	References
		Age = 43.46			
	Alzheimer's disease RBDCT	N = 119 Resveratrol (56) Placebo (48)	500 mg/day of placebo/resveratrol (with a dose escalation by 500 mg increments every 13 weeks, ending with 1000 mg twice daily) 52 weeks	↓ MMP-9; ↑ MMP-10; FGF-2 and IL-4	[356]
Resveratrol	Type 2 diabetes RBDCT	N = 41 Resveratrol (23): M = 11 W = 12 Age = 54.96 Placebo (18): M = 8 W = 10 Age = 58.72 N = 42; M = 21; W = 21	800 mg/day of resveratrol/placebo; 8 weeks	↑ TAC; Nrf2 and SOD	[357]
	Obesity RBDCT	Treatment: Age = 36.1 Placebo: Age = 38.7	282 mg/day of EGCG and 80 mg/day of resveratrol/placebo; 12 weeks	↑ Mitochondrial capacity and fat oxidation	[358]

BDNF: brain-derived neurotrophic factor; CGRP: calcitonin gene peptide; COX: cyclooxygenase; DBRPC: double-blind, randomized, placebo-controlled; IL: interleukin; M: male; MDA: malondialdehyde; MMP: matrix metalloproteinase; N: number; Nrf2: nuclear factor erythroid 2-related factor 2; PROBE: prospective randomized open-end blinded evaluation, RBDCT: randomized double-blinded controlled trial; SCI: spinal cord injury; SOD: superoxide dismutase; TAC: total antioxidant capacity; TGF- β : transforming growth factor-beta; TNF- α : tumor necrosis factor- α ; W: woman.

four weeks, reduced COX-2 [349]. Panahi et al. also reported that curcumin (1500 mg/day for six weeks) decreased levels of systemic inflammatory mediators, such as TNF- α , TGF- β , IL-6, substance P, and calcitonin gene peptide (CGRP) [350]. Another RDBPC study confirms the anti-inflammatory potential of oral curcumin (400 mg/3 times a day, p.o.) in patients with type 2 diabetes, along with a significant reduction in the levels of MDA, IL-6, and TNF- α [351].

In a controlled clinical trial study on 100 patients with SCI, curcumin (110 mg/day for six months) was notably associated with reducing osteoporosis progression and bone turnover markers after six months [352]. Results of a randomized, parallel-group, controlled, clinical trial on 20 subjects showed that administration of the InflanNox capsule (curcumin 1200 mg/day), with other anti-inflammatory and antioxidant effects, targeted inflammatory pathways, reduced IL-1 β , and improved mood disorders in SCI patients [353]. Administration of nanocurcumin (80 mg/day) in 50 patients with multiple sclerosis was associated with a significant increase in the levels of TGF- β and IL-10 expression [354]. In another study, nanocurcumin was also employed in a randomized, double-blind, placebo-controlled clinical trial of 40 diabetic patients. In their study, nanocurcumin could suppress oxidative stress and free radicals as an antioxidant agent [355].

During a 52-week study, resveratrol in the dose of 500-1000 mg/day markedly modulated neuroinflammation, decreased MMP-9, and induced adaptive immunity [356]. Administration of 800 mg/day resveratrol for eight weeks showed an antioxidant effect in the serum of patients with type 2 diabetes. Expression of SOD, Nrf2, and plasma total

antioxidant capacity was significantly increased [357]. Combined supplementation with resveratrol (80 mg/day) and EGCG (282 mg/day) for 12 weeks increased mitochondrial capacity and fat oxidation in obese humans [358]. As provided by Hendouei et al., resveratrol plus risperidone reduced irritability of patients with autism-associated oxidative stress in a RDBPC trial [359]. An open-label trial also confirmed the health benefits of high-dose resveratrol (1 g and 5 g/day) in 24 patients [360].

In an 8-month multicenter, RDBPC trial, polyphenol supplementation (200 mL/day) potentially modulated plasma concentration of homocysteine in 48 patients with Alzheimer's disease [361]. In a multicenter, double-blind clinical study, 34 diabetic patients with neuropathy (21-72 years old) were subjected to receive a topical formulation containing quercetin to decrease oxidative stress [362]. Regarding evaluating the therapeutic effects of a herbal, polyphenol-rich extract in a randomized controlled trial, Verlaet et al. reported antioxidant potentials for attention-deficit hyperactivity disorder [363]. Besides, foods high in polyphenols showed the potential to speed up the cognitive reserve randomized factorial design with 180 participants with Alzheimer's disease [364]. Additionally, another polyphenol-riched extract showed potential antioxidative effects in healthy individuals and those subjected to neuronal-associated diseases [365-369].

We have also previously reported the clinical potential of the whole phytochemicals against the apoptotic/autophagy aspects of SCI and related neuronal dysfunctions [3]. Table 3 shows the clinical potentials of some polyphenols against oxidative/inflammatory-based disorders with the base of oxidative damage.

10. Conclusion

Oxidative stress seems to orchestrate the complex pathophysiological mechanisms in SCI regarding modulating other interconnected pathways, as neuroinflammation and neuroapoptosis. Consequently, there is a complicated cross-talk between oxidative stress and neuroinflammatory/neuroapoptotic pathways. In this line, Nrf2/Keap1/ARE, SOD, CAT, GSH, MDA, HO-1, and XO have dramatically attenuated the interconnected pathways/mediators, including Bax, Bcl-2, caspases, p53, and TNF- α , IL-1 β , IL-6, and NF- κ B, toward neuroprotection in NDDs and SCI. The aforementioned signaling mediators are critically interconnected with extrinsic molecular mechanisms of regeneration post-SCI, including Nogo signaling pathways, glial scar formation, and chondroitin sulfate proteoglycan toward axonal activity. We also previously reported the complex dysregulated apoptotic and autophagic pathways following SCI (3). The complexity of those destructive signaling pathways of SCI and their interconnections urges the need to find multitarget agents with more tolerability and safety. In this regard, the plant kingdom has shown to be a promising source of neuroprotective agents. Growing studies are focusing on the potential of polyphenols in targeting oxidative stress and the cross-talk mediators (Figure 3). Our previous study also revealed the potential of polyphenols in targeting apoptosis and autophagy post-SCI (3). Additionally, it is worth noting that there are several common pathophysiological mechanisms behind the SCI and other NDDs. So, providing alternative therapeutic candidates for SCI could pave the way to treat other NDDs [372–374].

However, the polyphenol's pharmacokinetic limitations force the researchers to use novel drug delivery systems such as nanoformulations to delete such limitations in the clinical trial (3). So, the aforementioned limitation, along with the beneficial role of nanoparticles in improving the spinal cord drug delivery, proposes using nanoformulations of polyphenols to drawback such limitations. It will improve the beneficial effects of these compounds in SCI and other NDDs. Accordingly, metal nanoparticles (iron oxide, gold, silver, etc.), PLGA, polycaprolactone (PCL), poly-L-lactic acid (PLLA), liposomes, and inorganics are used toward the development of various nanoparticles in combating SCI complications [375]. Using novel drug delivery systems for phenolic compounds in SCI therapy could improve their bioavailability and facilitate the passages through the blood-spinal cord barrier and also modulate the water solubility and their long-lasting half-life following the spinal cord lesion [376].

Similar reviews will help the scientific community, neurosurgeons, neurooncologists, and patients to become aware of the complexity of dysregulated signaling pathways following SCI and the importance of finding novel multitarget alternative natural agents with more safety and efficacy. Further reports are needed to reveal the precise pathophysiological mechanisms and signaling pathways involved in NDDs, as well as the secondary phase of SCI. We propose targeting the upstream destructive mediators to prevent other interconnected pathways. The Nrf2, Keap1, and ARE

seem to be hopeful candidates in the oxidative pathway to stop related pathogenicity. In this line, natural phytochemicals are proposed to be focused on as alternative treatments with lower side effects and higher efficacy. Among natural entities, polyphenols/phenolic compounds are considered secondary metabolites with extensive biological activities and health benefits, which are currently used in modern medicine toward designing and developing new therapeutic agents. Ongoing clinical trials are assessing the beneficial role of polyphenols on NDDs; however, there is a lack of clinical studies to evaluate the possible potential of polyphenols in combating post-SCI complications. So, well-controlled clinical trials will help reveal polyphenols' clinical potential in combating sensory-motor dysfunction following SCI and paving the road to overcome the possible limitations in their administration. In the present review, we highlighted the modulatory role of oxidative stress on the inflammatory and apoptotic cascades of NDDs, while tackling particular attention to SCI. We also focused our attention on the need to develop polyphenols/phenolic compounds as novel multitarget neuroprotective treatment through targeting the intrinsic oxidative mediators and inflammatory/apoptotic pathways, as well as extrinsic axonal associated pathways. The coadministration of polyphenols/phenolic compounds and stem cells could also pave the road in treating SCI complications. Such studies will provide novel therapeutic targets in the prevention, management, and treatment of NDDs and SCI and investigate polyphenols/phenolic compounds as promising phytochemicals. Future studies should focus on additional large-scale, well-controlled clinical trials on the administration of polyphenols/phenolic compounds alone or in combination with other phytochemicals, or stem cells/other drugs, concentrating on the intrinsic and extrinsic pathways of axonal regeneration.

Conflicts of Interest

The authors declare that they have no conflicts of interest

Authors' Contributions

The contributions of the authors involved in this study are as follows: conceptualizing and designing the structure of the manuscript: S.F. and H.K.; drafting the manuscript: S.F., F.A., and S.Z.M.; software: S.F.; reviewing and editing the manuscript: S.F., J.X., and H.K.; and revising: S.F., F.A., J.X., and H.K.

References

- [1] S. Fakhri, I. Y. Aneva, M. H. Farzaei, and E. Sobarzo-Sánchez, "The neuroprotective effects of astaxanthin: therapeutic targets and clinical perspective," *Molecules*, vol. 24, no. 14, p. 2640, 2019.
- [2] K. J. Barnham, C. L. Masters, and A. I. Bush, "Neurodegenerative diseases and oxidative stress," *Nature reviews Drug discovery*, vol. 3, no. 3, pp. 205–214, 2004.
- [3] F. Abbaszadeh, S. Fakhri, and H. Khan, "Targeting apoptosis and autophagy following spinal cord injury: therapeutic

- approaches to polyphenols and candidate phytochemicals," *Pharmacological Research*, vol. 160, article 105069, 2020.
- [4] N. Sezer, S. Akkuş, and F. G. Uğurlu, "Chronic complications of spinal cord injury," *World journal of orthopedics*, vol. 6, no. 1, pp. 24–33, 2015.
 - [5] D. M. Stein and W. A. Knight IV, "Emergency neurological life support: traumatic spine injury," *Neurocritical care*, vol. 27, no. S1, Supplement 1, pp. 170–180, 2017.
 - [6] C. S. Ahuja, J. R. Wilson, S. Nori et al., "Traumatic spinal cord injury," *Nature reviews Disease primers*, vol. 3, no. 1, pp. 1–21, 2017.
 - [7] A. P. Tran, P. M. Warren, and J. Silver, "The biology of regeneration failure and success after spinal cord injury," *Physiological reviews*, vol. 98, no. 2, pp. 881–917, 2018.
 - [8] H. Ob, "Post-traumatic inflammation following spinal cord injury," *Spinal Cord*, vol. 41, no. 7, pp. 369–378, 2003.
 - [9] Z. Jia, H. Zhu, J. Li, X. Wang, H. Misra, and Y. Li, "Oxidative stress in spinal cord injury and antioxidant-based intervention," *Spinal Cord*, vol. 50, no. 4, pp. 264–274, 2012.
 - [10] R. Naseri, F. Farzaei, S. Fakhri et al., "Polyphenols for diabetes associated neuropathy: pharmacological targets and clinical perspective," *Journal of Pharmaceutical Sciences*, vol. 27, no. 2, pp. 781–798, 2019.
 - [11] K. Hassanzadeh and A. Rahimmi, "Oxidative stress and neuroinflammation in the story of Parkinson's disease: could targeting these pathways write a good ending?," *Journal of cellular physiology*, vol. 234, no. 1, pp. 23–32, 2019.
 - [12] X.-Y. Mao, H.-H. Zhou, and W.-L. Jin, "Redox-related neuronal death and crosstalk as drug targets: focus on epilepsy," *Frontiers in neuroscience*, vol. 13, p. 512, 2019.
 - [13] H. Solleiro-Villavicencio and S. Rivas-Arancibia, "Effect of chronic oxidative stress on neuroinflammatory response mediated by CD4⁺ T cells in neurodegenerative diseases," *Frontiers in cellular neuroscience*, vol. 12, p. 114, 2018.
 - [14] N. Šutulović, Ž. Grubač, S. Šuvakov et al., "Chronic prostatitis/chronic pelvic pain syndrome increases susceptibility to seizures in rats and alters brain levels of IL-1 β and IL-6," *Epilepsy research*, vol. 153, pp. 19–27, 2019.
 - [15] N. Šutulović, Ž. Grubač, S. Šuvakov et al., "Experimental chronic prostatitis/chronic pelvic pain syndrome increases anxiety-like behavior: the role of brain oxidative stress, serum corticosterone, and hippocampal parvalbumin-positive interneurons," *Oxidative Medicine and Cellular Longevity*, vol. 2021, Article ID 6687493, 17 pages, 2021.
 - [16] Z. Liu, Z. Ren, J. Zhang et al., "Role of ROS and nutritional antioxidants in human diseases," *Frontiers in physiology*, vol. 9, p. 477, 2018.
 - [17] I. Liguori, G. Russo, F. Curcio et al., "Oxidative stress, aging, and diseases," *Clinical Interventions in Aging*, vol. 13, pp. 757–772, 2018.
 - [18] G. J. Burton and E. Jauniaux, "Oxidative stress," *Best Practice & Research Clinical Obstetrics & Gynaecology*, vol. 25, no. 3, pp. 287–299, 2011.
 - [19] Y. Luo, C. Fu, Z. Wang, Z. Zhang, H. Wang, and Y. Liu, "Mangiferin attenuates contusive spinal cord injury in rats through the regulation of oxidative stress, inflammation and the Bcl-2 and Bax pathway," *Molecular medicine reports*, vol. 12, no. 5, pp. 7132–7138, 2015.
 - [20] K. Hamann and R. Shi, "Acrolein scavenging: a potential novel mechanism of attenuating oxidative stress following spinal cord injury," *Journal of neurochemistry*, vol. 111, no. 6, pp. 1348–1356, 2009.
 - [21] C. Busquets-Cortés, X. Capó, E. Argelich et al., "Effects of millimolar steady-state hydrogen peroxide exposure on inflammatory and redox gene expression in immune cells from humans with metabolic syndrome," *Nutrients*, vol. 10, no. 12, p. 1920, 2018.
 - [22] N. Zhang, Y. Yin, S.-J. Xu, Y.-P. Wu, and W.-S. Chen, "Inflammation & apoptosis in spinal cord injury," *The Indian journal of medical research*, vol. 135, no. 3, p. 287, 2012.
 - [23] A. Strasser, L. O'Connor, and V. M. Dixit, "Apoptosis signaling," *Annual review of biochemistry*, vol. 69, no. 1, pp. 217–245, 2000.
 - [24] V. V. Adjan, K. F. Hauser, G. Bakalkin et al., "Caspase-3 activity is reduced after spinal cord injury in mice lacking dynorphin: differential effects on glia and neurons," *Neuroscience*, vol. 148, no. 3, pp. 724–736, 2007.
 - [25] M. Sharif-Alhoseini, M. Khormali, M. Rezaei et al., "Animal models of spinal cord injury: a systematic review," *Spinal cord*, vol. 55, no. 8, pp. 714–721, 2017.
 - [26] G. Fatima, V. P. Sharma, S. K. Das, and A. A. Mahdi, "Oxidative stress and antioxidative parameters in patients with spinal cord injury: implications in the pathogenesis of disease," *Spinal Cord*, vol. 53, no. 1, pp. 3–6, 2015.
 - [27] D. Varija, K. Kumar, K. Reddy, and V. Reddy, "Prolonged constriction of sciatic nerve affecting oxidative stressors & antioxidant enzymes in rat," *Indian Journal of Medical Research*, vol. 129, no. 5, p. 587, 2009.
 - [28] N. P. Visavadiya, S. P. Patel, J. L. VanRooyen, P. G. Sullivan, and A. G. Rabchevsky, "Cellular and subcellular oxidative stress parameters following severe spinal cord injury," *Redox Biology*, vol. 8, pp. 59–67, 2016.
 - [29] S. Magesh, Y. Chen, and L. Hu, "Small molecule modulators of Keap1-Nrf2-ARE pathway as potential preventive and therapeutic agents," *Medicinal research reviews*, vol. 32, no. 4, pp. 687–726, 2012.
 - [30] Y. Lu, Y.-Y. Yang, M.-W. Zhou et al., "Ketogenic diet attenuates oxidative stress and inflammation after spinal cord injury by activating Nrf2 and suppressing the NF- κ B signaling pathways," *Neuroscience Letters*, vol. 683, pp. 13–18, 2018.
 - [31] M. Jang and I.-H. Cho, "Sulforaphane ameliorates 3-nitropropionic acid-induced striatal toxicity by activating the Keap1-Nrf2-ARE pathway and inhibiting the MAPKs and NF- κ B pathways," *Molecular neurobiology*, vol. 53, no. 4, pp. 2619–2635, 2016.
 - [32] A. Coyoy-Salgado, J. J. Segura-Urbe, C. Guerra-Araiza et al., "The importance of natural antioxidants in the treatment of spinal cord injury in animal models: an overview," *Oxidative Medicine and Cellular Longevity*, vol. 2019, 22 pages, 2019.
 - [33] W. Jiang, M. Li, F. He et al., "Neuroprotective effect of asiatic acid against spinal cord injury in rats," *Life sciences*, vol. 157, pp. 45–51, 2016.
 - [34] P. O. Donkor, Y. Chen, L. Ding, and F. Qiu, "Locally and traditionally used *Ligusticum* species - A review of their phytochemistry, pharmacology and pharmacokinetics," *Journal of ethnopharmacology*, vol. 194, pp. 530–548, 2016.
 - [35] C. Wang, P. Wang, W. Zeng, and W. Li, "Tetramethylpyrazine improves the recovery of spinal cord injury via Akt/Nrf2/HO-1 pathway," *Bioorganic & Medicinal Chemistry Letters*, vol. 26, no. 4, pp. 1287–1291, 2016.

- [36] M. Nakayama, M. Aihara, Y.-N. Chen, M. Araie, K. Tomita-Yokotani, and T. Iwashina, "Neuroprotective effects of flavonoids on hypoxia-, glutamate-, and oxidative stress-induced retinal ganglion cell death," *Molecular vision*, vol. 17, pp. 1784–1793, 2011.
- [37] F. Tian, L.-H. Xu, W. Zhao, L.-J. Tian, and X.-L. Ji, "The optimal therapeutic timing and mechanism of puerarin treatment of spinal cord ischemia-reperfusion injury in rats," *Journal of ethnopharmacology*, vol. 134, no. 3, pp. 892–896, 2011.
- [38] M. Chelombitko, "Role of reactive oxygen species in inflammation: a minireview," *Moscow University Biological Sciences Bulletin*, vol. 73, no. 4, pp. 199–202, 2018.
- [39] J. P. Laurila, L. E. Laatikainen, M. D. Castellone, and M. O. Laukkanen, "SOD3 reduces inflammatory cell migration by regulating adhesion molecule and cytokine expression," *PloS one*, vol. 4, no. 6, article e5786, 2009.
- [40] M. J. Morgan and Z.-g. Liu, "Crosstalk of reactive oxygen species and NF- κ B signaling," *Cell research*, vol. 21, no. 1, pp. 103–115, 2011.
- [41] M. Redza-Dutordoir and D. A. Averill-Bates, "Activation of apoptosis signalling pathways by reactive oxygen species," *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, vol. 1863, no. 12, pp. 2977–2992, 2016.
- [42] Z. Liu, X. Yao, W. Jiang et al., "Advanced oxidation protein products induce microglia-mediated neuroinflammation via MAPKs-NF- κ B signaling pathway and pyroptosis after secondary spinal cord injury," *Journal of Neuroinflammation*, vol. 17, no. 1, p. 90, 2020.
- [43] M. S. Beattie, "Inflammation and apoptosis: linked therapeutic targets in spinal cord injury," *Trends in Molecular Medicine*, vol. 10, no. 12, pp. 580–583, 2004.
- [44] K. Yoshida and Y. Miki, "The cell death machinery governed by the p53 tumor suppressor in response to DNA damage," *Cancer science*, vol. 101, no. 4, pp. 831–835, 2010.
- [45] S. Fakhri, M. Pesce, A. Patruno et al., "Attenuation of Nrf2/Keap1/ARE in Alzheimer's disease by plant secondary metabolites: a mechanistic review," *Molecules*, vol. 25, no. 21, p. 4926, 2020.
- [46] L. Yang, K.-S. Wen, X. Ruan, Y.-X. Zhao, F. Wei, and Q. Wang, "Response of plant secondary metabolites to environmental factors," *Molecules*, vol. 23, no. 4, p. 762, 2018.
- [47] S. Z. Moradi, S. Momtaz, Z. Bayrami, M. H. Farzaei, and M. Abdollahi, "Nanoformulations of herbal extracts in treatment of neurodegenerative disorders," *Frontiers in bioengineering and biotechnology*, vol. 8, p. 238, 2020.
- [48] S. Fakhri, S. Z. Moradi, M. H. Farzaei, and A. Bishayee, "Modulation of dysregulated cancer metabolism by plant secondary metabolites: a mechanistic review," *Seminars in Cancer Biology*, 2020.
- [49] W. Rungsung, K. K. Ratha, S. Dutta, A. K. Dixit, and J. Hazra, "Secondary metabolites of plants in drugs discovery," *World Journal of Pharmaceutical Research*, vol. 4, no. 7, pp. 604–613, 2015.
- [50] A. M. Shohael, M. B. Ali, K. W. Yu, E. J. Hahn, R. Islam, and K. Y. Paek, "Effect of light on oxidative stress, secondary metabolites and induction of antioxidant enzymes in *Eleutherococcus senticosus* somatic embryos in bioreactor," *Process Biochemistry*, vol. 41, no. 5, pp. 1179–1185, 2006.
- [51] B. Uttara, A. V. Singh, P. Zamboni, and R. Mahajan, "Oxidative stress and neurodegenerative diseases: a review of upstream and downstream antioxidant therapeutic options," *Current neuropharmacology*, vol. 7, no. 1, pp. 65–74, 2009.
- [52] G. Ungerer, J. Cui, T. Ndam et al., "Harpagophytum procumbens extract ameliorates allodynia and modulates oxidative and antioxidant stress pathways in a rat model of spinal cord injury," *Neuromolecular Medicine*, vol. 22, no. 2, pp. 278–292, 2020.
- [53] A. Venditti and A. Bianco, "Sulfur-containing secondary metabolites as neuroprotective agents," *Current medicinal chemistry*, vol. 27, no. 26, pp. 4421–4436, 2020.
- [54] F. Lauro, L. A. Giancotti, S. Ilari et al., "Inhibition of spinal oxidative stress by bergamot polyphenolic fraction attenuates the development of morphine induced tolerance and hyperalgesia in mice," *PLoS One*, vol. 11, no. 5, article e0156039, 2016.
- [55] A. R. Khalatbary, "Natural polyphenols and spinal cord injury," *Iranian biomedical journal*, vol. 18, no. 3, pp. 120–129, 2014.
- [56] E. A. Decker, "Phenolics: prooxidants or antioxidants?," *Nutrition reviews*, vol. 55, no. 11, pp. 396–398, 1997.
- [57] D. M. Kasote, S. S. Katyare, M. V. Hegde, and H. Bae, "Significance of antioxidant potential of plants and its relevance to therapeutic applications," *International journal of biological sciences*, vol. 11, no. 8, pp. 982–991, 2015.
- [58] J. M. Davis, E. A. Murphy, and M. D. Carmichael, "Effects of the dietary flavonoid quercetin upon performance and health," *Current sports medicine reports*, vol. 8, no. 4, pp. 206–213, 2009.
- [59] L. Aguirre, N. Arias, M. Teresa Macarulla, A. Gracia, P. Portillo, and M., "Beneficial effects of quercetin on obesity and diabetes," *The Open Nutraceuticals Journal*, vol. 4, no. 1, pp. 189–198, 2011.
- [60] J. P. Spencer, G. G. Kuhnle, R. J. Williams, and C. Rice-Evans, "Intracellular metabolism and bioactivity of quercetin and its in vivo metabolites," *Biochemical Journal*, vol. 372, no. 1, pp. 173–181, 2003.
- [61] Y. Song, J. Liu, F. Zhang, J. Zhang, T. Shi, and Z. Zeng, "Antioxidant effect of quercetin against acute spinal cord injury in rats and its correlation with the p38MAPK/iNOS signaling pathway," *Life sciences*, vol. 92, no. 24–26, pp. 1215–1221, 2013.
- [62] E. Schültke, H. Kamencic, M. Zhao et al., "Neuroprotection following fluid percussion brain trauma: a pilot study using quercetin," *Journal of neurotrauma*, vol. 22, no. 12, pp. 1475–1484, 2005.
- [63] T. Yang, B. Kong, J.-W. Gu et al., "Anti-apoptotic and antioxidant roles of quercetin after traumatic brain injury," *Cellular and molecular neurobiology*, vol. 34, no. 6, pp. 797–804, 2014.
- [64] J.-K. Lee, H.-J. Kwak, M.-S. Piao, J.-W. Jang, S.-H. Kim, and H.-S. Kim, "Quercetin reduces the elevated matrix metalloproteinases-9 level and improves functional outcome after cerebral focal ischemia in rats," *Acta neurochirurgica*, vol. 153, no. 6, pp. 1321–1329, 2011.
- [65] R.-Q. Yao, D.-S. Qi, H.-L. Yu, J. Liu, L.-H. Yang, and X.-X. Wu, "Quercetin attenuates cell apoptosis in focal cerebral ischemia rat brain via activation of BDNF-TrkB-PI3K/Akt signaling pathway," *Neurochemical research*, vol. 37, no. 12, pp. 2777–2786, 2012.
- [66] S. M. Borghi, F. A. Pinho-Ribeiro, V. Fattori et al., "Quercetin inhibits peripheral and spinal cord nociceptive mechanisms to reduce intense acute swimming-induced muscle pain in mice," *PLoS One*, vol. 11, no. 9, article e0162267, 2016.

- [67] A. E.-D. L. Firgany and N. R. Sarhan, "Quercetin mitigates monosodium glutamate-induced excitotoxicity of the spinal cord motoneurons in aged rats via p38 MAPK inhibition," *Acta Histochemica*, vol. 122, no. 5, article 151554, 2020.
- [68] M. I. Azevedo, A. F. Pereira, R. B. Nogueira et al., "The antioxidant effects of the flavonoids rutin and quercetin inhibit oxaliplatin-induced chronic painful peripheral neuropathy," *Molecular pain*, vol. 9, pp. 1744-8069-9-53-1744-8069-8069, 2013.
- [69] J. Liu, T. Tang, and H. Yang, "Antioxidation of quercetin against spinal cord injury in rats," *Chinese journal of traumatology*, vol. 9, no. 5, pp. 303-307, 2006.
- [70] C. Huang, C. Fu, Z.-P. Qi et al., "Localised delivery of quercetin by thermo-sensitive PLGA-PEG-PLGA hydrogels for the treatment of brachial plexus avulsion," *Artificial Cells, Nanomedicine, and Biotechnology*, vol. 48, no. 1, pp. 1010-1021, 2020.
- [71] Y. Wang, W. Li, M. Wang et al., "Quercetin reduces neural tissue damage and promotes astrocyte activation after spinal cord injury in rats," *Journal of cellular biochemistry*, vol. 119, no. 2, pp. 2298-2306, 2018.
- [72] E. Schültke, R. Griebel, and B. Juurlink, "Quercetin attenuates inflammatory processes after spinal cord injury in an animal model," *Spinal cord*, vol. 48, no. 12, pp. 857-861, 2010.
- [73] O. Ocal, A. O. Borcek, O. Pasaoglu, A. C. Gundogdu, G. T. Kaplanoglu, and M. K. Baykaner, "Can quercetin be an option for treatment of spinal cord injury? An experimental study," *Turkish Neurosurgery*, vol. 29, no. 2, pp. 247-253, 2019.
- [74] U. Çiftçi, E. Delen, M. Vural et al., "Efficacy of resveratrol and quercetin after experimental spinal cord injury," *Turkish Journal of Trauma and Emergency Surgery*, vol. 22, no. 5, pp. 423-431, 2016.
- [75] Ö. Çevik, M. Erşahin, T. E. Şener et al., "Beneficial effects of quercetin on rat urinary bladder after spinal cord injury," *Journal of surgical research*, vol. 183, no. 2, pp. 695-703, 2013.
- [76] X. Wang, Y. Wang, L. Zhang, G. Li, and H. Zhang, "Combinatory effect of mesenchymal stromal cells transplantation and quercetin after spinal cord injury in rat," *European Review for Medical and Pharmacological Sciences*, vol. 22, no. 9, pp. 2876-2887, 2018.
- [77] H. Fan, H.-B. Tang, L.-Q. Shan et al., "Quercetin prevents necroptosis of oligodendrocytes by inhibiting macrophages/microglia polarization to M1 phenotype after spinal cord injury in rats," *Journal of neuroinflammation*, vol. 16, no. 1, pp. 1-15, 2019.
- [78] W. Jiang, Y. Huang, N. Han et al., "Quercetin suppresses NLRP3 inflammasome activation and attenuates histopathology in a rat model of spinal cord injury," *Spinal Cord*, vol. 54, no. 8, pp. 592-596, 2016.
- [79] N. T. Zaveri, "Green tea and its polyphenolic catechins: medicinal uses in cancer and noncancer applications," *Life sciences*, vol. 78, no. 18, pp. 2073-2080, 2006.
- [80] M. Pervin, K. Unno, T. Ohishi, H. Tanabe, N. Miyoshi, and Y. Nakamura, "Beneficial effects of green tea catechins on neurodegenerative diseases," *Molecules*, vol. 23, no. 6, p. 1297, 2018.
- [81] Y. Wu and J. Cui, "(-)-Epigallocatechin-3-gallate provides neuroprotection via AMPK activation against traumatic brain injury in a mouse model," *Naunyn-Schmiedeberg's Archives of Pharmacology*, vol. 393, no. 11, pp. 2209-2220, 2020.
- [82] L. Machova Urdzikova, J. Ruzicka, K. Karova et al., "A green tea polyphenol epigallocatechin-3-gallate enhances neuroregeneration after spinal cord injury by altering levels of inflammatory cytokines," *Neuropharmacology*, vol. 126, pp. 213-223, 2017.
- [83] I.-H. Wei, H.-C. Tu, C.-C. Huang, M.-H. Tsai, C.-Y. Tseng, and J.-Y. Shieh, "(-)-Epigallocatechin gallate attenuates NADPH-d/nNOS expression in motor neurons of rats following peripheral nerve injury," *BMC neuroscience*, vol. 12, no. 1, p. 52, 2011.
- [84] A. R. Khalatbary and E. Khademi, "The green tea polyphenolic catechin epigallocatechin gallate and neuroprotection," *Nutritional neuroscience*, vol. 23, no. 4, pp. 281-294, 2020.
- [85] N. Salah, N. J. Miller, G. Paganga, L. Tijburg, G. P. Bolwell, and C. Riceevans, "Polyphenolic Flavonols as Scavengers of Aqueous Phase Radicals and as Chain- Breaking Antioxidants," *Archives of biochemistry and biophysics*, vol. 322, no. 2, pp. 339-346, 1995.
- [86] J. Zhao, S. Fang, Y. Yuan et al., "Green tea polyphenols protect spinal cord neurons against hydrogen peroxide-induced oxidative stress," *Neural regeneration research*, vol. 9, no. 14, pp. 1379-1385, 2014.
- [87] Q. Ye, L. Ye, X. Xu et al., "Epigallocatechin-3-gallate suppresses 1-methyl-4-phenyl-pyridine-induced oxidative stress in PC12 cells via the SIRT1/PGC-1 α signaling pathway," *BMC complementary and alternative medicine*, vol. 12, no. 1, p. 82, 2012.
- [88] M.-I. Ding, H. Ma, Y.-g. Man, and H. Y. LV, "Protective effects of a green tea polyphenol, epigallocatechin-3-gallate, against sevoflurane-induced neuronal apoptosis involve regulation of CREB/BDNF/TrkB and PI3K/Akt/mTOR signaling pathways in neonatal mice," *Canadian journal of physiology and pharmacology*, vol. 95, no. 12, pp. 1396-1405, 2017.
- [89] A. R. Khalatbary and H. Ahmadvand, "Anti-inflammatory effect of the epigallocatechin gallate following spinal cord trauma in rat," *Iranian biomedical journal*, vol. 15, no. 1-2, pp. 31-37, 2011.
- [90] A. R. Khalatbary, T. Tiraihi, M. B. Boroujeni, H. Ahmadvand, M. Tavafi, and A. Tamjidipoor, "Effects of epigallocatechin gallate on tissue protection and functional recovery after contusive spinal cord injury in rats," *Brain research*, vol. 1306, pp. 168-175, 2010.
- [91] F. Che, G. Wang, J. Yu et al., "Effects of epigallocatechin-3-gallate on iron metabolism in spinal cord motor neurons," *Molecular Medicine Reports*, vol. 16, no. 3, pp. 3010-3014, 2017.
- [92] J. Ruzicka, L. M. Urdzikova, B. Svobodova et al., "Does combined therapy of curcumin and epigallocatechin gallate have a synergistic neuroprotective effect against spinal cord injury?," *Neural Regeneration Research*, vol. 13, no. 1, pp. 119-127, 2018.
- [93] B. Álvarez-Pérez, J. Homs, M. Bosch-Mola et al., "Epigallocatechin-3-gallate treatment reduces thermal hyperalgesia after spinal cord injury by down-regulating RhoA expression in mice," *European Journal of Pain*, vol. 20, no. 3, pp. 341-352, 2016.
- [94] C. S. Park, J. Y. Lee, H. Y. Choi, B. G. Ju, I. Youn, and T. Y. Yune, "Protocatechuic acid improves functional recovery after spinal cord injury by attenuating blood-spinal cord barrier disruption and hemorrhage in rats," *Neurochemistry international*, vol. 124, pp. 181-192, 2019.

- [95] G. Murtaza, S. Karim, M. R. Akram et al., "Caffeic acid phenethyl ester and therapeutic potentials," *BioMed Research International*, vol. 2014, 9 pages, 2014.
- [96] G. F. Sud'ina, O. Mirzoeva, M. Pushkareva, G. A. Korshunova, N. Sumbatyan, and S. Varfolomeev, "Caffeic acid phenethyl ester as a lipoxygenase inhibitor with antioxidant properties," *FEBS letters*, vol. 329, no. 1-2, pp. 21–24, 1993.
- [97] F. Armutcu, S. Akyol, S. Ustunsoy, and F. F. Turan, "Therapeutic potential of caffeic acid phenethyl ester and its anti-inflammatory and immunomodulatory effects (Review)," *Experimental and therapeutic medicine*, vol. 9, no. 5, pp. 1582–1588, 2015.
- [98] S. Meyuhas, M. Assali, M. Huleihil, and M. Huleihel, "Antimicrobial activities of caffeic acid phenethyl ester," *Journal of Molecular Biochemistry*, vol. 4, no. 2, 2015.
- [99] G. Ozturk, Z. Ginis, S. Akyol, G. Erden, A. Gurel, and O. Akyol, "The anticancer mechanism of caffeic acid phenethyl ester (CAPE): review of melanomas, lung and prostate cancers," *European Review for Medical and Pharmacological Sciences*, vol. 16, no. 15, pp. 2064–2068, 2012.
- [100] X. Wang, S. Stavchansky, P. D. Bowman, and S. M. Kerwin, "Cytoprotective effect of caffeic acid phenethyl ester (CAPE) and catechol ring-fluorinated CAPE derivatives against menadione-induced oxidative stress in human endothelial cells," *Bioorganic & medicinal chemistry*, vol. 14, no. 14, pp. 4879–4887, 2006.
- [101] K. Natarajan, S. Singh, T. R. Burke, D. Grunberger, and B. B. Aggarwal, "Caffeic acid phenethyl ester is a potent and specific inhibitor of activation of nuclear transcription factor NF-kappa B," *Proceedings of the National Academy of Sciences*, vol. 93, no. 17, pp. 9090–9095, 1996.
- [102] Y. Kimura, H. Okuda, T. Okuda, T. Hatano, I. Agata, and S. Arichi, "Studies on the activities of tannins and related compounds from medicinal plants and drugs. VII. Effects of extracts of leaves of *Artemisia* species, and caffeic acid and chlorogenic acid on lipid metabolic injury in rats fed peroxidized oil," *Chemical and pharmaceutical bulletin*, vol. 33, no. 5, pp. 2028–2034, 1985.
- [103] J. Laranjinha, Otilia Vieira, Vitor Madeira, and L. Almeida, "Two related phenolic antioxidants with opposite effects on vitamin E content in low density lipoproteins oxidized by ferrylmyoglobin: consumption vs regeneration," *Archives of Biochemistry and Biophysics*, vol. 323, no. 2, pp. 373–381, 1995.
- [104] X. Wei, L. Zhao, Z. Ma et al., "Caffeic acid phenethyl ester prevents neonatal hypoxic-ischaemic brain injury," *Brain*, vol. 127, no. 12, pp. 2629–2635, 2004.
- [105] A. Russo, R. Longo, and A. Vanella, "Antioxidant activity of propolis: role of caffeic acid phenethyl ester and galangin," *Fitoterapia*, vol. 73, pp. S21–S29, 2002.
- [106] M. Ozeren, N. SUCU, L. Tamer et al., "Caffeic acid phenethyl ester (CAPE) supplemented St. Thomas' hospital cardioplegic solution improves the antioxidant defense system of rat myocardium during ischemia-reperfusion injury," *Pharmacological research*, vol. 52, no. 3, pp. 258–263, 2005.
- [107] J.-K. Kim and H.-D. Jang, "Nrf2-mediated HO-1 induction coupled with the ERK signaling pathway contributes to indirect antioxidant capacity of caffeic acid phenethyl ester in HepG2 cells," *International journal of molecular sciences*, vol. 15, no. 7, pp. 12149–12165, 2014.
- [108] H. Kim, W. Kim, S. Yum et al., "Caffeic acid phenethyl ester activation of Nrf2 pathway is enhanced under oxidative state: structural analysis and potential as a pathologically targeted therapeutic agent in treatment of colonic inflammation," *Free Radical Biology and Medicine*, vol. 65, pp. 552–562, 2013.
- [109] C. Gocmez, F. Celik, K. Kamasak et al., "Effects of intrathecal caffeic acid phenethyl ester and methylprednisolone on oxidant/antioxidant status in traumatic spinal cord injuries," *Journal of Neurological Surgery Part A: Central European Neurosurgery*, vol. 76, no. 1, pp. 20–24, 2015.
- [110] B. Akgun, S. Ozturk, G. Artas, and F. S. Erol, "Effects of intrathecal caffeic acid phenethyl ester (CAPE) on IL-6 and TNF- α ; levels and local inflammatory responses in spinal cord injuries," *Turkish neurosurgery*, vol. 28, no. 4, pp. 625–629, 2018.
- [111] H. Ak, İ. Gülşen, T. Karaaslan et al., "The effects of caffeic acid phenethyl ester on inflammatory cytokines after acute spinal cord injury," *Turkish journal of trauma & emergency surgery*, vol. 21, no. 2, pp. 96–101, 2015.
- [112] M. Kasai, H. Fukumitsu, H. Soumiya, and S. Furukawa, "Caffeic acid phenethyl ester reduces spinal cord injury-evoked locomotor dysfunction," *Biomedical research*, vol. 32, no. 1, pp. 1–7, 2011.
- [113] A. Ilhan, U. Koltuksuz, S. Ozen, E. Uz, H. Ciralik, and O. Akyol, "The effects of caffeic acid phenethyl ester (CAPE) on spinal cord ischemia/reperfusion injury in rabbits," *European journal of cardio-thoracic surgery*, vol. 16, no. 4, pp. 458–463, 1999.
- [114] A. Ilhan, O. Akyol, A. Gurel, F. Armutcu, M. Iraz, and E. Oztas, "Protective effects of caffeic acid phenethyl ester against experimental allergic encephalomyelitis-induced oxidative stress in rats," *Free radical biology & medicine*, vol. 37, no. 3, pp. 386–394, 2004.
- [115] A. J. Alonso-Castro, R. Zapata-Bustos, F. Domínguez, A. García-Carrancá, and L. A. Salazar-Olivo, "Magnolia dealbata Zucc and its active principles honokiol and magnolol stimulate glucose uptake in murine and human adipocytes using the insulin-signaling pathway," *Phytomedicine*, vol. 18, no. 11, pp. 926–933, 2011.
- [116] X. Han, Y. Pang, S. Liu et al., "Antidiarrhea and antioxidant activities of honokiol extract from *Magnolia officinalis* cortex in mice," *Tropical Journal of Pharmaceutical Research*, vol. 13, no. 10, pp. 1643–1651, 2014.
- [117] B. H. Kim and J. Y. Cho, "Anti-inflammatory effect of honokiol is mediated by PI3K/Akt pathway suppression," *Acta Pharmacologica Sinica*, vol. 29, no. 1, pp. 113–122, 2008.
- [118] Y.-R. Lin, H.-H. Chen, C.-H. Ko, and M.-H. Chan, "Effects of honokiol and magnolol on acute and inflammatory pain models in mice," *Life Sciences*, vol. 81, no. 13, pp. 1071–1078, 2007.
- [119] Q. Xu, L.-T. Yi, Y. Pan et al., "Antidepressant-like effects of the mixture of honokiol and magnolol from the barks of *Magnolia officinalis* in stressed rodents," *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, vol. 32, no. 3, pp. 715–725, 2008.
- [120] R. Prasad and S. K. Katiyar, "Honokiol, an active compound of magnolia plant, inhibits growth, and progression of cancers of different organs," in *Anti-inflammatory Nutraceuticals and Chronic Diseases*, pp. 245–265, Springer, 2016.
- [121] C. M. Chen, S. H. Liu, and S. Y. Lin-Shiau, "Honokiol, a neuroprotectant against mouse cerebral ischaemia, mediated by preserving Na⁺, K⁺-ATPase activity and mitochondrial functions," *Basic & clinical pharmacology & toxicology*, vol. 101, no. 2, pp. 108–116, 2007.

- [122] Y. Wang, Z. Z. Zhang, Y. Wu, J. Zhan, X. H. He, and Y. L. Wang, "Honokiol protects rat hearts against myocardial ischemia reperfusion injury by reducing oxidative stress and inflammation," *Experimental and therapeutic medicine*, vol. 5, no. 1, pp. 315–319, 2013.
- [123] H.-Q. Yin, Y.-T. Je, Y.-C. Kim et al., "Magnolia officinalis reverses alcoholic fatty liver by inhibiting the maturation of sterol regulatory element-binding protein-1c," *Journal of pharmacological sciences*, vol. 109, no. 4, pp. 486–495, 2009.
- [124] Y. YU, M. LI, N. SU et al., "Honokiol protects against renal ischemia/reperfusion injury via the suppression of oxidative stress, iNOS, inflammation and STAT3 in rats," *Molecular medicine reports*, vol. 13, no. 2, pp. 1353–1360, 2016.
- [125] Y. Hou, S. Peng, X. Li, J. Yao, J. Xu, and J. Fang, "Honokiol alleviates oxidative stress-induced neurotoxicity via activation of Nrf2," *ACS chemical neuroscience*, vol. 9, no. 12, pp. 3108–3116, 2018.
- [126] D. Y. Chuang, M.-H. Chan, Y. Zong et al., "Magnolia polyphenols attenuate oxidative and inflammatory responses in neurons and microglial cells," *Journal of neuroinflammation*, vol. 10, no. 1, p. 786, 2013.
- [127] A. Rajgopal, S. R. Missler, and J. D. Scholten, "Magnolia officinalis (Hou Po) bark extract stimulates the Nrf2-pathway in hepatocytes and protects against oxidative stress," *Journal of ethnopharmacology*, vol. 193, pp. 657–662, 2016.
- [128] W. Chang, Y. Chang, F. Lu, and H.-C. Chiang, "Inhibitory effects of phenolics on xanthine oxidase," *Anticancer research*, vol. 14, no. 2A, pp. 501–506, 1994.
- [129] J. P. Wang, E. F. Hsu, S. L. Raung et al., "Inhibition by magnolol of formylmethionyl-leucyl-phenyl alanine-induced respiratory burst in rat neutrophils," *Journal of pharmacy and pharmacology*, vol. 51, no. 3, pp. 285–294, 1999.
- [130] L. K. Chao, P.-C. Liao, C.-L. Ho et al., "Anti-inflammatory bioactivities of honokiol through inhibition of protein kinase C, mitogen-activated protein kinase, and the NF- κ B pathway to reduce LPS-induced TNF α and NO expression," *Journal of agricultural and food chemistry*, vol. 58, no. 6, pp. 3472–3478, 2010.
- [131] J. Liu, C. Zhang, Z. Liu, J. Zhang, Z. Xiang, and T. Sun, "Honokiol downregulates Kruppel-like factor 4 expression, attenuates inflammation, and reduces histopathology after spinal cord injury in rats," *Spine*, vol. 40, no. 6, pp. 363–368, 2015.
- [132] K.-T. Liou, Y.-C. Shen, C.-F. Chen, C.-M. Tsao, and S.-K. Tsai, "Honokiol protects rat brain from focal cerebral ischemia-reperfusion injury by inhibiting neutrophil infiltration and reactive oxygen species production," *Brain research*, vol. 992, no. 2, pp. 159–166, 2003.
- [133] C.-H. Chang, C.-C. Chyau, C.-L. Hsieh et al., "Relevance of phenolic diterpene constituents to antioxidant activity of supercritical CO₂ extract from the leaves of rosemary," *Natural Product Research*, vol. 22, no. 1, pp. 76–90, 2008.
- [134] M. Nadeem, M. Imran, T. Aslam Gondal et al., "Therapeutic potential of rosmarinic acid: a comprehensive review," *Applied Sciences*, vol. 9, no. 15, p. 3139, 2019.
- [135] S. K. Amoah, L. P. Sandjo, J. M. Kratz, and M. W. Biavatti, "Rosmarinic acid—pharmaceutical and clinical aspects," *Planta medica*, vol. 82, no. 5, pp. 388–406, 2016.
- [136] C. F. Lima, M. Fernandes-Ferreira, and C. Pereira-Wilson, "Phenolic compounds protect HepG2 cells from oxidative damage: relevance of glutathione levels," *Life sciences*, vol. 79, no. 21, pp. 2056–2068, 2006.
- [137] J. Govindaraj and S. Sorimuthu Pillai, "Rosmarinic acid modulates the antioxidant status and protects pancreatic tissues from glucolipotoxicity mediated oxidative stress in high-fat diet: streptozotocin-induced diabetic rats," *Molecular and cellular biochemistry*, vol. 404, no. 1–2, pp. 143–159, 2015.
- [138] C. Lu, Y. Zou, Y. Liu, and Y. Niu, "Rosmarinic acid counteracts activation of hepatic stellate cells via inhibiting the ROS-dependent MMP-2 activity: Involvement of Nrf2 antioxidant system," *Toxicology and applied pharmacology*, vol. 318, pp. 69–78, 2017.
- [139] A. Y. Lee, T. T. Wu, B. R. Hwang et al., "The neuro-protective effect of the methanolic extract of Perilla frutescens var. japonica and rosmarinic acid against H₂O₂-Induced oxidative stress in C6 glial cells," *Biomolecules & therapeutics*, vol. 24, no. 3, pp. 338–345, 2016.
- [140] M. G. Rahbardar, B. Amin, S. Mehri, S. J. Mirnajafi-Zadeh, and H. Hosseinzadeh, "Rosmarinic acid attenuates development and existing pain in a rat model of neuropathic pain: an evidence of anti-oxidative and anti-inflammatory effects," *Phytomedicine*, vol. 40, pp. 59–67, 2018.
- [141] A. Areti, P. Komirishetty, A. K. Kalvala, K. Nellaiappan, and A. Kumar, "Rosmarinic acid mitigates mitochondrial dysfunction and spinal glial activation in oxaliplatin-induced peripheral neuropathy," *Molecular neurobiology*, vol. 55, no. 9, pp. 7463–7475, 2018.
- [142] H. Rong, Y. Liang, and Y. Niu, "Rosmarinic acid attenuates β -amyloid-induced oxidative stress via Akt/GSK-3 β /Fyn-mediated Nrf2 activation in PC12 cells," *Free radical biology & medicine*, vol. 120, pp. 114–123, 2018.
- [143] F. H. C. Melo, E. T. Venâncio, D. P. de Sousa et al., "Anxiolytic-like effect of carvacrol (5-isopropyl-2-methylphenol) in mice: involvement with GABAergic transmission," *Fundamental & clinical pharmacology*, vol. 24, no. 4, pp. 437–443, 2010.
- [144] F. H. C. Melo, B. A. Moura, D. P. de Sousa et al., "Antidepressant-like effect of carvacrol (5-isopropyl-2-methylphenol) in mice: involvement of dopaminergic system," *Fundamental & clinical pharmacology*, vol. 25, no. 3, pp. 362–367, 2011.
- [145] F. Chen, Z. Shi, K. Neoh, and E. Kang, "Antioxidant and antibacterial activities of eugenol and carvacrol-grafted chitosan nanoparticles," *Biotechnology and bioengineering*, vol. 104, no. 1, pp. 30–39, 2009.
- [146] N. Kumar, A. Yadav, S. Gulati, Kanupriya, N. Aggarwal, and R. Gupta, "Antigenotoxic effect of curcumin and carvacrol against parathion induced DNA damage in cultured human peripheral blood lymphocytes and its relation to GSTM1 and GSTT1 polymorphism," *Journal of toxicology*, vol. 2014, Article ID 404236, 7 pages, 2014.
- [147] A. G. Guimarães, M. A. Xavier, M. T. de Santana et al., "Carvacrol attenuates mechanical hypernociception and inflammatory response," *Naunyn-Schmiedeberg's archives of pharmacology*, vol. 385, no. 3, pp. 253–263, 2012.
- [148] B. K. Can, "Biological and pharmacological activities of carvacrol and carvacrol bearing essential oils," *Current pharmaceutical design*, vol. 14, no. 29, pp. 3106–3119, 2008.
- [149] S. Samarghandian, T. Farkhondeh, F. Samini, and A. Borji, "Protective effects of carvacrol against oxidative stress induced by chronic stress in rat's brain, liver, and kidney," *Biochemistry research international*, vol. 2016, Article ID 2645237, 7 pages, 2016.

- [150] S. Banik, M. Akter, S. E. Corpus Bondad, T. Saito, T. Hosokawa, and M. Kurasaki, "Carvacrol inhibits cadmium toxicity through combating against caspase dependent/independent apoptosis in PC12 cells," *Food and Chemical Toxicology*, vol. 134, article 110835, 2019.
- [151] A. L. Chenet, A. R. Duarte, F. J. S. de Almeida, C. M. B. Andrade, and M. R. de Oliveira, "Carvacrol depends on heme oxygenase-1 (HO-1) to exert antioxidant, anti-inflammatory, and mitochondria-related protection in the human neuroblastoma SH-SY5Y cells line exposed to hydrogen peroxide," *Neurochemical research*, vol. 44, no. 4, pp. 884–896, 2019.
- [152] Z.-w. Cui, Z.-x. Xie, B.-f. Wang et al., "Carvacrol protects neuroblastoma SH-SY5Y cells against Fe²⁺-induced apoptosis by suppressing activation of MAPK/JNK-NF- κ B signaling pathway," *Acta Pharmacologica Sinica*, vol. 36, no. 12, pp. 1426–1436, 2015.
- [153] R. R. Crichton, S. Wilmet, R. Legssyer, and R. J. Ward, "Molecular and cellular mechanisms of iron homeostasis and toxicity in mammalian cells," *Journal of inorganic biochemistry*, vol. 91, no. 1, pp. 9–18, 2002.
- [154] M. Mahmoodi, H. Amiri, F. Ayooobi et al., "Carvacrol ameliorates experimental autoimmune encephalomyelitis through modulating pro- and anti-inflammatory cytokines," *Life sciences*, vol. 219, pp. 257–263, 2019.
- [155] E. M. Alvarenga, L. K. Souza, T. S. Araújo et al., "Carvacrol reduces irinotecan-induced intestinal mucositis through inhibition of inflammation and oxidative damage via TRPA1 receptor activation," *Chemico-biological interactions*, vol. 260, pp. 129–140, 2016.
- [156] A. Ganeshpurkar and A. K. Saluja, "The pharmacological potential of rutin," *Saudi Pharmaceutical Journal*, vol. 25, no. 2, pp. 149–164, 2017.
- [157] A. Potapovich and V. Kostyuk, "Comparative study of antioxidant properties and cytoprotective activity of flavonoids," *Biochemistry*, vol. 68, no. 5, pp. 514–519, 2003.
- [158] Z. Nouri, S. Fakhri, K. Nouri, C. E. Wallace, M. H. Farzaei, and A. Bishayee, "Targeting multiple signaling pathways in cancer: the rutin therapeutic approach," *Cancers*, vol. 12, no. 8, p. 2276, 2020.
- [159] F. Mellou, H. Loutrari, H. Stamatis, C. Roussos, and F. N. Kolisis, "Enzymatic esterification of flavonoids with unsaturated fatty acids: effect of the novel esters on vascular endothelial growth factor release from K562 cells," *Process Biochemistry*, vol. 41, no. 9, pp. 2029–2034, 2006.
- [160] H. Javed, M. Khan, A. Ahmad et al., "Rutin prevents cognitive impairments by ameliorating oxidative stress and neuroinflammation in rat model of sporadic dementia of Alzheimer type," *Neuroscience*, vol. 210, pp. 340–352, 2012.
- [161] J. Wu, L. Maoqiang, H. Fan et al., "Rutin attenuates neuroinflammation in spinal cord injury rats," *Journal of Surgical Research*, vol. 203, no. 2, pp. 331–337, 2016.
- [162] P.-H. Pan, S.-Y. Lin, Y.-Y. Wang et al., "Protective effects of rutin on liver injury induced by biliary obstruction in rats," *Free Radical Biology and Medicine*, vol. 73, pp. 106–116, 2014.
- [163] S. Singh, D. K. Singh, A. Meena, V. Dubey, N. Masood, and S. Luqman, "Rutin protects t-butyl hydroperoxide-induced oxidative impairment via modulating the Nrf2 and iNOS activity," *Phytomedicine*, vol. 55, pp. 92–104, 2019.
- [164] J. Almeida, D. Benvegnú, N. Boufleuer et al., "Hydrogels containing rutin intended for cutaneous administration: efficacy in wound healing in rats," *Drug development and industrial pharmacy*, vol. 38, no. 7, pp. 792–799, 2012.
- [165] A. Gęgotek, E. Ambrożewicz, A. Jastrząb, I. Jaročka-Karpowicz, and E. Skrzydlewska, "Rutin and ascorbic acid cooperation in antioxidant and antiapoptotic effect on human skin keratinocytes and fibroblasts exposed to UVA and UVB radiation," *Archives of dermatological research*, vol. 311, no. 3, pp. 203–219, 2019.
- [166] R. Tian, W. Yang, Q. Xue et al., "Rutin ameliorates diabetic neuropathy by lowering plasma glucose and decreasing oxidative stress via Nrf2 signaling pathway in rats," *European journal of pharmacology*, vol. 771, pp. 84–92, 2016.
- [167] S. Moghbelinejad, M. Nassiri-Asl, T. N. Farivar et al., "Rutin activates the MAPK pathway and BDNF gene expression on beta-amyloid induced neurotoxicity in rats," *Toxicology letters*, vol. 224, no. 1, pp. 108–113, 2014.
- [168] R. Wang, Y. Sun, H. Huang, L. Wang, J. Chen, and W. Shen, "Rutin, a natural flavonoid protects PC12 cells against sodium nitroprusside-induced neurotoxicity through activating PI3K/Akt/mTOR and ERK1/2 pathway," *Neurochemical research*, vol. 40, no. 9, pp. 1945–1953, 2015.
- [169] N. O. al-Harbi, F. Imam, M. M. al-Harbi et al., "Rutin inhibits carfilzomib-induced oxidative stress and inflammation via the NOS-mediated NF- κ B signaling pathway," *Inflammopharmacology*, vol. 27, no. 4, pp. 817–827, 2019.
- [170] P. Zhang and X. Ma, "Effect of rutin on spinal cord injury through inhibition of the expression of MIP-2 and activation of MMP-9, and downregulation of Akt phosphorylation," *Molecular medicine reports*, vol. 12, no. 5, pp. 7554–7560, 2015.
- [171] H.-I. Song, X. Zhang, W.-z. Wang et al., "Neuroprotective mechanisms of rutin for spinal cord injury through anti-oxidation and anti-inflammation and inhibition of p38 mitogen activated protein kinase pathway," *Neural regeneration research*, vol. 13, no. 1, pp. 128–134, 2018.
- [172] Y. F. Zhou, B. Guo, M. J. Ye, R. F. Liao, and S. L. Li, "Protective effect of rutin against H₂O₂-induced oxidative stress and apoptosis in human lens epithelial cells," *Current eye research*, vol. 41, no. 7, pp. 933–942, 2016.
- [173] M. Nassiri-Asl, A. Ghorbani, S. Salehisar, E. Asadpour, and H. R. Sadeghnia, "Effect of rutin on oxidative DNA damage in PC12 neurons cultured in nutrients deprivation condition," *Iranian journal of basic medical sciences*, vol. 23, no. 3, pp. 390–395, 2020.
- [174] S. Yao, L. Wang, Q. Chen, T. Lu, X. Pu, and C. Luo, "The effect of mild hypothermia plus rutin on the treatment of spinal cord injury and inflammatory factors by repressing TGF- β /Smad pathway," *Acta Cirúrgica Brasileira*, vol. 36, no. 3, p. e360307, 2021.
- [175] M. Jain and H. S. Parmar, "Evaluation of antioxidative and anti-inflammatory potential of hesperidin and naringin on the rat air pouch model of inflammation," *Inflammation research*, vol. 60, no. 5, pp. 483–491, 2011.
- [176] K. P. Devi, T. Rajavel, S. F. Nabavi et al., "Hesperidin: a promising anticancer agent from nature," *Industrial crops and products*, vol. 76, pp. 582–589, 2015.
- [177] A. Siddiqi, B. Saidullah, and S. Sultana, "Anti-carcinogenic effect of hesperidin against renal cell carcinoma by targeting COX-2/PGE2 pathway in Wistar rats," *Environmental toxicology*, vol. 33, no. 10, pp. 1069–1077, 2018.
- [178] P. Bhargava, V. K. Verma, S. Malik, S. I. Khan, J. Bhatia, and D. S. Arya, "Hesperidin regresses cardiac hypertrophy by

- virtue of PPAR- γ agonistic, anti-inflammatory, antiapoptotic, and antioxidant properties,” *Journal of biochemical and molecular toxicology*, vol. 33, no. 5, article e22283, 2019.
- [179] A. M. Mahmoud, H. M. Mohammed, S. M. Khadrawy, and S. R. Galaly, “Hesperidin protects against chemically induced hepatocarcinogenesis via modulation of Nrf2/ARE/HO-1, PPAR γ and TGF- β 1/Smad3 signaling, and amelioration of oxidative stress and inflammation,” *Chemico-Biological Interactions*, vol. 277, pp. 146–158, 2017.
- [180] H. R. A. Salem, A. El-Raouf, E. M. Saleh, and K. A. Shalaby, “Influence of hesperidin combined with Sinemet on genetical and biochemical abnormalities in rats suffering from Parkinson’s disease,” *Life Sciences*, vol. 9, no. 4, pp. 930–945, 2012.
- [181] D. Wang, L. Liu, X. Zhu, W. Wu, and Y. Wang, “Hesperidin alleviates cognitive impairment, mitochondrial dysfunction and oxidative stress in a mouse model of Alzheimer’s disease,” *Cellular and molecular neurobiology*, vol. 34, no. 8, pp. 1209–1221, 2014.
- [182] M. Ashafaq, L. Varshney, M. H. A. Khan et al., “Neuromodulatory effects of hesperidin in mitigating oxidative stress in streptozotocin induced diabetes,” *BioMed research international*, vol. 2014, pp. 1–9, 2014.
- [183] M. Chen, H. Gu, Y. Ye et al., “Protective effects of hesperidin against oxidative stress of tert-butyl hydroperoxide in human hepatocytes,” *Food and Chemical Toxicology*, vol. 48, no. 10, pp. 2980–2987, 2010.
- [184] C. Zhu, Y. Dong, H. Liu, H. Ren, and Z. Cui, “Hesperetin protects against H₂O₂-triggered oxidative damage via upregulation of the Keap1-Nrf2/HO-1 signal pathway in ARPE-19 cells,” *Biomedicine & Pharmacotherapy*, vol. 88, pp. 124–133, 2017.
- [185] M. S. Aly, S. R. Galaly, N. Moustafa, H. M. Mohammed, S. M. Khadrawy, and A. M. Mahmoud, “Hesperidin protects against diethylnitrosamine/carbon tetrachloride-induced renal repercussions via up-regulation of Nrf2/HO-1 signaling and attenuation of oxidative stress,” *Journal of Applied Pharmaceutical Science*, vol. 7, no. 11, pp. 7–14, 2017.
- [186] S.-D. Heo, J. Kim, Y. Choi, P. Ekanayake, M. Ahn, and T. Shin, “Hesperidin improves motor disability in rat spinal cord injury through anti-inflammatory and antioxidant mechanism via Nrf-2/HO-1 pathway,” *Neuroscience letters*, vol. 715, article 134619, 2020.
- [187] A. I. Carballo-Villalobos, M. E. González-Trujano, N. Alvarado-Vázquez, and F. J. López-Muñoz, “Pro-inflammatory cytokines involvement in the hesperidin antihyperalgesic effects at peripheral and central levels in a neuropathic pain model,” *Inflammopharmacology*, vol. 25, no. 2, pp. 265–269, 2017.
- [188] B. Salehi, A. P. Mishra, M. Nigam et al., “Resveratrol: a double-edged sword in health benefits,” *Biomedicines*, vol. 6, no. 3, p. 91, 2018.
- [189] H. Zhao, S. Chen, K. Gao et al., “Resveratrol protects against spinal cord injury by activating autophagy and inhibiting apoptosis mediated by the SIRT1/AMPK signaling pathway,” *Neuroscience*, vol. 348, pp. 241–251, 2017.
- [190] M. Xu, Z. Cheng, Z. Ding, Y. Wang, Q. Guo, and C. Huang, “Resveratrol enhances IL-4 receptor-mediated anti-inflammatory effects in spinal cord and attenuates neuropathic pain following sciatic nerve injury,” *Molecular pain*, vol. 14, p. 174480691876754, 2018.
- [191] A. Rauf, M. Imran, M. S. Butt, M. Nadeem, D. G. Peters, and M. S. Mubarak, “Resveratrol as an anti-cancer agent: a review,” *Critical reviews in food science and nutrition*, vol. 58, no. 9, pp. 1428–1447, 2018.
- [192] C. A. De La Lastra and I. Villegas, *Resveratrol as an antioxidant and pro-oxidant agent: mechanisms and clinical implications*, Portland Press Ltd., 2007.
- [193] S. Fu, R. Lv, L. Wang, H. Hou, H. Liu, and S. Shao, “Resveratrol, an antioxidant, protects spinal cord injury in rats by suppressing MAPK pathway,” *Saudi journal of biological sciences*, vol. 25, no. 2, pp. 259–266, 2018.
- [194] V. Kesharwani, F. Atif, S. Yousuf, and S. K. Agrawal, “Resveratrol protects spinal cord dorsal column from hypoxic injury by activating Nrf-2,” *Neuroscience*, vol. 241, pp. 80–88, 2013.
- [195] S. W. Tsang, Y.-F. Guan, J. Wang, Z.-X. Bian, and H.-J. Zhang, “Inhibition of pancreatic oxidative damage by stilbene derivative dihydro-resveratrol: implication for treatment of acute pancreatitis,” *Scientific reports*, vol. 6, no. 1, p. 22859, 2016.
- [196] A. K. Singh and M. Vinayak, “Resveratrol alleviates inflammatory hyperalgesia by modulation of reactive oxygen species (ROS), antioxidant enzymes and ERK activation,” *Inflammation Research*, vol. 66, no. 10, pp. 911–921, 2017.
- [197] L. Cheng, Z. Jin, R. Zhao, K. Ren, C. Deng, and S. Yu, “Resveratrol attenuates inflammation and oxidative stress induced by myocardial ischemia-reperfusion injury: role of Nrf2/ARE pathway,” *International journal of clinical and experimental medicine*, vol. 8, no. 7, pp. 10420–10428, 2015.
- [198] N. Tamaki, R. Cristina Orihuela-Campos, Y. Inagaki, M. Fukui, T. Nagata, and H.-O. Ito, “Resveratrol improves oxidative stress and prevents the progression of periodontitis via the activation of the Sirt1/AMPK and the Nrf2/antioxidant defense pathways in a rat periodontitis model,” *Free Radical Biology and Medicine*, vol. 75, pp. 222–229, 2014.
- [199] P. Palsamy and S. Subramanian, “Resveratrol protects diabetic kidney by attenuating hyperglycemia-mediated oxidative stress and renal inflammatory cytokines via Nrf2-Keap1 signaling,” *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, vol. 1812, no. 7, pp. 719–731, 2011.
- [200] A. Daverey and S. K. Agrawal, “Pre and post treatment with curcumin and resveratrol protects astrocytes after oxidative stress,” *Brain Research*, vol. 1692, pp. 45–55, 2018.
- [201] O. ATEŞ, S. CAYLI, E. ALTINOZ et al., “Effects of resveratrol and methylprednisolone on biochemical, neurobehavioral and histopathological recovery after experimental spinal cord injury,” *Acta Pharmacologica Sinica*, vol. 27, no. 10, pp. 1317–1325, 2006.
- [202] J. Liu, L. Yi, Z. Xiang, J. Zhong, H. Zhang, and T. Sun, “Resveratrol attenuates spinal cord injury-induced inflammatory damage in rat lungs,” *International journal of clinical and experimental pathology*, vol. 8, no. 2, pp. 1237–1246, 2015.
- [203] B. P. Xu, M. Yao, Z. J. Li et al., “Neurological recovery and antioxidant effects of resveratrol in rats with spinal cord injury: a meta-analysis,” *Neural regeneration research*, vol. 15, no. 3, pp. 482–490, 2020.
- [204] M. D. Recalde, C. A. Miguel, M. V. Noya-Riobó, S. L. González, M. J. Villar, and M. F. Coronel, “Resveratrol exerts anti-oxidant and anti-inflammatory actions and prevents oxaliplatin-induced mechanical and thermal allodynia,” *Brain Research*, vol. 1748, article 147079, 2020.
- [205] L. Xu, C. Liu, W. Xiang, H. Chen, X. Qin, and X. Huang, “Advances in the study of oxyresveratrol,” *International Journal of Pharmacology*, vol. 10, no. 1, pp. 44–54, 2013.

- [206] H. Du, L. Ma, G. Chen, and S. Li, "The effects of oxyresveratrol abrogates inflammation and oxidative stress in rat model of spinal cord injury," *Molecular medicine reports*, vol. 17, no. 3, pp. 4067–4073, 2018.
- [207] Q.-H. Du, C. Peng, and H. Zhang, "Polydatin: a review of pharmacology and pharmacokinetics," *Pharmaceutical biology*, vol. 51, no. 11, pp. 1347–1354, 2013.
- [208] R. Lv, L. Du, L. Zhang, and Z. Zhang, "Polydatin attenuates spinal cord injury in rats by inhibiting oxidative stress and microglia apoptosis via Nrf2/HO-1 pathway," *Life sciences*, vol. 217, pp. 119–127, 2019.
- [209] R. Lv, L. Du, X. Liu, F. Zhou, Z. Zhang, and L. Zhang, "Polydatin alleviates traumatic spinal cord injury by reducing microglial inflammation via regulation of iNOS and NLRP3 inflammasome pathway," *International immunopharmacology*, vol. 70, pp. 28–36, 2019.
- [210] M. Chen, Y. Hou, and D. Lin, "Polydatin protects bone marrow stem cells against oxidative injury: involvement of Nrf 2/ARE pathways," *Stem cells international*, vol. 2016, 10 pages, 2016.
- [211] Y. Fan, Y. Li, S. Huang, H. Xu, H. Li, and B. Liu, "Resveratrol-primed exosomes strongly promote the recovery of motor function in SCI rats by activating autophagy and inhibiting apoptosis via the PI3K signaling pathway," *Neuroscience Letters*, vol. 736, article 135262, 2020.
- [212] S. Zhang, B. O. Botchway, Y. Zhang, and X. Liu, "Resveratrol can inhibit Notch signaling pathway to improve spinal cord injury," *Annals of Anatomy-Anatomischer Anzeiger*, vol. 223, pp. 100–107, 2019.
- [213] X. Liu, B. O. Botchway, X. Tan, Y. Zhang, and M. Fang, "Resveratrol treatment of spinal cord injury in rat model," *Microscopy research and technique*, vol. 82, no. 3, pp. 296–303, 2019.
- [214] W. Chen, Z. Zhao, Z. Zhao, L. Zhang, and Q. Song, "Resveratrol and puerarin loaded polymeric nanoparticles to enhance the chemotherapeutic efficacy in spinal cord injury," *Biomedical Microdevices*, vol. 22, no. 4, 2020.
- [215] U. Kiziltepe, N. N. D. Turan, U. Han, A. T. Ulus, and F. Akar, "Resveratrol, a red wine polyphenol, protects spinal cord from ischemia-reperfusion injury," *Journal of vascular surgery*, vol. 40, no. 1, pp. 138–145, 2004.
- [216] H. D. Wang, Y. M. Shi, L. Li, J. D. Guo, Y. P. Zhang, and S. X. Hou, "Treatment with resveratrol attenuates sublesional bone loss in spinal cord-injured rats," *British journal of pharmacology*, vol. 170, no. 4, pp. 796–806, 2013.
- [217] Y.-B. Yang and Y.-J. Piao, "Effects of resveratrol on secondary damages after acute spinal cord injury in rats," *Acta Pharmacologica Sinica*, vol. 24, no. 7, pp. 703–710, 2003.
- [218] B. B. Aggarwal, A. Kumar, and A. C. Bharti, "Anticancer potential of curcumin: preclinical and clinical studies," *Anti-cancer research*, vol. 23, no. 1A, pp. 363–398, 2003.
- [219] V. P. Menon and A. R. Sudheer, *Antioxidant and anti-inflammatory properties of curcumin. The molecular targets and therapeutic uses of curcumin in health and disease*, Springer, 2007.
- [220] K. I. Priyadarsini, "Free radical reactions of curcumin in membrane models," *Free Radical Biology and Medicine*, vol. 23, no. 6, pp. 838–843, 1997.
- [221] Z. Xie, B. Wu, G. Shen, X. Li, and Q. Wu, "Curcumin alleviates liver oxidative stress in type 1 diabetic rats," *Molecular medicine reports*, vol. 17, no. 1, pp. 103–108, 2018.
- [222] İ. Akar, İ. İnce, A. Arici et al., "The protective effect of curcumin on a spinal cord ischemia-reperfusion injury model," *Annals of vascular surgery*, vol. 42, pp. 285–292, 2017.
- [223] Z. Q. Liu, S. S. Xing, and W. Zhang, "Neuroprotective effect of curcumin on spinal cord in rabbit model with ischemia/reperfusion," *The journal of spinal cord medicine*, vol. 36, no. 2, pp. 147–152, 2013.
- [224] J. Wu, Q. Li, X. Wang et al., "Neuroprotection by curcumin in ischemic brain injury involves the Akt/Nrf2 pathway," *PLoS one*, vol. 8, no. 3, 2013.
- [225] W. Dai, H. Wang, J. Fang et al., "Curcumin provides neuroprotection in model of traumatic brain injury via the Nrf2-ARE signaling pathway," *Brain research bulletin*, vol. 140, pp. 65–71, 2018.
- [226] W. Yin, X. Zhang, X. Shi, and Y. Li, *Curcumin protects SH-SY5Y cells from oxidative stress by up-regulating HO-1 via phosphatidylinositol 3 kinase/Akt/Nrf-2 and down-regulating HO-2. Molecular neurodegeneration*, Springer, 2012.
- [227] A. Daverey and S. K. Agrawal, "Neuroprotective effects of riluzole and curcumin in human astrocytes and spinal cord white matter hypoxia," *Neuroscience letters*, vol. 738, article 135351, 2020.
- [228] H. Jiang, X. Tian, Y. Guo, W. Duan, H. Bu, and C. Li, "Activation of nuclear factor erythroid 2-related factor 2 cytoprotective signaling by curcumin protect primary spinal cord astrocytes against oxidative toxicity," *Biological & pharmaceutical bulletin*, vol. 34, no. 8, pp. 1194–1197, 2011.
- [229] N. Zhang, G. Wei, J. Ye et al., "Effect of curcumin on acute spinal cord injury in mice via inhibition of inflammation and TAK1 pathway," *Pharmacological Reports*, vol. 69, no. 5, pp. 1001–1006, 2017.
- [230] W. Jin, J. Wang, T. Zhu et al., "Anti-inflammatory effects of curcumin in experimental spinal cord injury in rats," *Inflammation Research*, vol. 63, no. 5, pp. 381–387, 2014.
- [231] B. G. Alvarado-Sanchez, H. Salgado-Ceballos, S. Torres-Castillo et al., "Electroacupuncture and curcumin promote oxidative balance and motor function recovery in rats following traumatic spinal cord injury," *Neurochemical research*, vol. 44, no. 2, pp. 498–506, 2019.
- [232] B. Cemil, K. Topuz, M. N. Demircan et al., "Curcumin improves early functional results after experimental spinal cord injury," *Acta neurochirurgica*, vol. 152, no. 9, pp. 1583–1590, 2010.
- [233] A. M. Sanli, E. Turkoglu, G. Serbes et al., "Effect of curcumin on lipid peroxidation, early ultrastructural findings and neurological recovery after experimental spinal cord contusion injury in rats," *Turkish neurosurgery*, vol. 22, no. 2, pp. 189–195, 2012.
- [234] Y.-S. Lee, D.-C. Cho, C. H. Kim, I. Han, E. Y. Gil, and K.-T. Kim, "Effect of curcumin on the inflammatory reaction and functional recovery after spinal cord injury in a hyperglycemic rat model," *The Spine Journal*, vol. 19, no. 12, pp. 2025–2039, 2019.
- [235] J. Zhang, H. Wei, M. Lin, C. Chen, C. Wang, and M. Liu, "Curcumin protects against ischemic spinal cord injury: the pathway effect," *Neural regeneration research*, vol. 8, no. 36, pp. 3391–3400, 2013.
- [236] Y.-F. Wang, J.-N. Zu, J. Li, C. Chen, C.-Y. Xi, and J.-L. Yan, "Curcumin promotes the spinal cord repair via inhibition of

- glial scar formation and inflammation,” *Neuroscience letters*, vol. 560, pp. 51–56, 2014.
- [237] M.-S. Lin, Y.-Y. Sun, W.-T. Chiu et al., “Curcumin attenuates the expression and secretion of RANTES after spinal cord injury in vivo and lipopolysaccharide-induced astrocyte reactivation in vitro,” *Journal of neurotrauma*, vol. 28, no. 7, pp. 1259–1269, 2011.
- [238] J. Yuan, M. Zou, X. Xiang et al., “Curcumin improves neural function after spinal cord injury by the joint inhibition of the intracellular and extracellular components of glial scar,” *Journal of Surgical Research*, vol. 195, no. 1, pp. 235–245, 2015.
- [239] C.-C. Lin, T.-H. Chiang, W.-J. Chen, Y.-Y. Sun, Y.-H. Lee, and M.-S. Lin, “CISD2 serves a novel role as a suppressor of nitric oxide signalling and curcumin increases CISD2 expression in spinal cord injuries,” *Injury*, vol. 46, no. 12, pp. 2341–2350, 2015.
- [240] J. Yuan, W. Liu, H. Zhu et al., “Curcumin inhibits glial scar formation by suppressing astrocyte-induced inflammation and fibrosis in vitro and in vivo,” *Brain research*, vol. 1655, pp. 90–103, 2017.
- [241] L. Machova Urdzikova, K. Karova, J. Ruzicka et al., “The anti-inflammatory compound curcumin enhances locomotor and sensory recovery after spinal cord injury in rats by immunomodulation,” *International journal of molecular sciences*, vol. 17, no. 1, p. 49, 2016.
- [242] J. Zu, Y. Wang, G. Xu, J. Zhuang, H. Gong, and J. Yan, “Curcumin improves the recovery of motor function and reduces spinal cord edema in a rat acute spinal cord injury model by inhibiting the JAK/STAT signaling pathway,” *Acta histochemica*, vol. 116, no. 8, pp. 1331–1336, 2014.
- [243] D.-s. Yu, Y. Cao, X.-f. Mei et al., “Curcumin improves the integrity of blood-spinal cord barrier after compressive spinal cord injury in rats,” *Journal of the neurological sciences*, vol. 346, no. 1–2, pp. 51–59, 2014.
- [244] H. Ni, W. Jin, T. Zhu et al., “Curcumin modulates TLR4/NF- κ B inflammatory signaling pathway following traumatic spinal cord injury in rats,” *The journal of spinal cord medicine*, vol. 38, no. 2, pp. 199–206, 2015.
- [245] A. Matkowski, P. Kus, E. Goralska, and D. Wozniak, “Mangiferin—a bioactive xanthonoid, not only from mango and not just antioxidant,” *Mini reviews in medicinal chemistry*, vol. 13, no. 3, pp. 439–455, 2013.
- [246] A. J. Núñez Selles, M. Daglia, and L. Rastrelli, “The potential role of mangiferin in cancer treatment through its immunomodulatory, anti-angiogenic, apoptotic, and gene regulatory effects,” *Biofactors*, vol. 42, no. 5, pp. 475–491, 2016.
- [247] M. Zheng and Z. Lu, “Antiviral effect of mangiferin and isomangiferin on herpes simplex virus,” *Chinese Medical Journal*, vol. 103, no. 2, pp. 160–165, 1990.
- [248] S. Saha, P. Sadhukhan, and P. C. Sil, “Mangiferin: a xanthonoid with multipotent anti-inflammatory potential,” *Biofactors*, vol. 42, no. 5, pp. 459–474, 2016.
- [249] T. Miura, H. Ichiki, I. Hashimoto et al., “Antidiabetic activity of a xanthone compound, mangiferin,” *Phytomedicine*, vol. 8, no. 2, pp. 85–87, 2001.
- [250] R. K. Khurana, R. Kaur, S. Lohan, K. K. Singh, and B. Singh, “Mangiferin: a promising anticancer bioactive,” *Pharmaceutical patent analyst*, vol. 5, no. 3, pp. 169–181, 2016.
- [251] A. Dar, S. Faizi, S. Naqvi et al., “Analgesic and antioxidant activity of mangiferin and its derivatives: the structure activity relationship,” *Biological and Pharmaceutical Bulletin*, vol. 28, no. 4, pp. 596–600, 2005.
- [252] E. Kasi Viswanadh, B. Nageshwar Rao, and B. S. Satish Rao, “Antigenotoxic effect of mangiferin and changes in antioxidant enzyme levels of Swiss albino mice treated with cadmium chloride,” *Human & experimental toxicology*, vol. 29, no. 5, pp. 409–418, 2010.
- [253] B.-p. Zhang, J. Zhao, S.-s. Li et al., “Mangiferin activates Nrf2-antioxidant response element signaling without reducing the sensitivity to etoposide of human myeloid leukemia cells in vitro,” *Acta Pharmacologica Sinica*, vol. 35, no. 2, pp. 257–266, 2014.
- [254] S. Saha, P. Sadhukhan, K. Sinha, N. Agarwal, and P. C. Sil, “Mangiferin attenuates oxidative stress induced renal cell damage through activation of PI3K induced Akt and Nrf-2 mediated signaling pathways,” *Biochemistry and biophysics reports*, vol. 5, pp. 313–327, 2016.
- [255] Z. Wang, S. Guo, J. Wang, Y. Shen, J. Zhang, and Q. Wu, “Nrf2/HO-1 mediates the neuroprotective effect of mangiferin on early brain injury after subarachnoid hemorrhage by attenuating mitochondria-related apoptosis and neuroinflammation,” *Scientific reports*, vol. 7, no. 1, pp. 1–12, 2017.
- [256] L. Xu, J. Liang, T. Jin, and F. Zhou, “Neuroprotective effects of mangiferin on acute spinal cord injury in rats and its mechanism,” *Chinese journal of reparative and reconstructive surgery*, vol. 30, no. 8, pp. 1019–1025, 2016.
- [257] P. K. Leong, J. Chen, and K. M. Ko, “Development of Chinese herbal health products for the prevention of aging-associated diseases,” in *Natural Products and Drug Discovery*, pp. 73–104, Elsevier, 2018.
- [258] N. Chen, P. Y. Chiu, and K. M. Ko, “Schisandrin B enhances cerebral mitochondrial antioxidant status and structural integrity, and protects against cerebral ischemia/reperfusion injury in rats,” *Biological and Pharmaceutical Bulletin*, vol. 31, no. 7, pp. 1387–1391, 2008.
- [259] R. Checker, R. S. Patwardhan, D. Sharma et al., “Schisandrin B exhibits anti-inflammatory activity through modulation of the redox-sensitive transcription factors Nrf2 and NF- κ B,” *Free Radical Biology and Medicine*, vol. 53, no. 7, pp. 1421–1430, 2012.
- [260] S.-P. Ip, C.-T. Che, Y.-C. Kong, and K.-M. Ko, “Effects of Schisandrin B pretreatment on tumor necrosis factor- α induced apoptosis and Hsp70 expression in mouse liver,” *Cell stress & chaperones*, vol. 6, no. 1, pp. 44–48, 2001.
- [261] T. H. Lee, C. H. Jung, and D.-H. Lee, “Neuroprotective effects of Schisandrin B against transient focal cerebral ischemia in Sprague-Dawley rats,” *Food and chemical toxicology*, vol. 50, no. 12, pp. 4239–4245, 2012.
- [262] R. A. Thandavarayan, V. V. Giridharan, S. Arumugam et al., “Schisandrin B prevents doxorubicin induced cardiac dysfunction by modulation of DNA damage, oxidative stress and inflammation through inhibition of MAPK/p53 signaling,” *PLoS One*, vol. 10, no. 3, article e0119214, 2015.
- [263] Q. Chen, H. Zhang, Y. Cao et al., “Schisandrin B attenuates CCl₄-induced liver fibrosis in rats by regulation of Nrf2-ARE and TGF- β /Smad signaling pathways,” *Drug design, development and therapy*, vol. 11, pp. 2179–2191, 2017.
- [264] Y. Wu, Z.-c. Li, L.-q. Yao, M. Li, and M. Tang, “Schisandrin B alleviates acute oxidative stress via modulation of the Nrf2/Keap1-mediated antioxidant pathway,” *Applied Physiology, Nutrition, and Metabolism*, vol. 44, no. 1, pp. 1–6, 2019.

- [265] T. Qian, P. K. Leong, K. M. Ko, and W. Chan, "Investigation of in vitro and in vivo metabolism of Schisandrin B from *Schisandrae fructus* by liquid chromatography coupled electrospray ionization tandem mass spectrometry," *Pharmacology & Pharmacy*, vol. 6, no. 8, pp. 363–373, 2015.
- [266] D. Xin, Z. Hu, H. Huo et al., "Schisandrin B attenuates the inflammatory response, oxidative stress and apoptosis induced by traumatic spinal cord injury via inhibition of p53 signaling in adult rats," *Molecular medicine reports*, vol. 16, no. 1, pp. 533–538, 2017.
- [267] N. Zhu, C. Cai, A. Zhou, X. Zhao, Y. Xiang, and C. Zeng, "Schisandrin B prevents hind limb from ischemia-reperfusion-induced oxidative stress and inflammation via MAPK/NF- κ B pathways in rats," *BioMed research international*, vol. 2017, Article ID 4237973, 2017.
- [268] Y. Sun, F. Gong, J. Yin et al., "Therapeutic effect of apocynin through antioxidant activity and suppression of apoptosis and inflammation after spinal cord injury," *Experimental and therapeutic medicine*, vol. 13, no. 3, pp. 952–960, 2017.
- [269] D. Impellizzeri, E. Mazzon, E. Esposito, I. Paterniti, P. Bramanti, and S. Cuzzocrea, "Effect of apocynin, an inhibitor of NADPH oxidase, in the inflammatory process induced by an experimental model of spinal cord injury," *Free radical research*, vol. 45, no. 2, pp. 221–236, 2011.
- [270] S. Kang, S. Liu, H. Li, D. Wang, and X. Qi, "Baicalin effects on rats with spinal cord injury by anti-inflammatory and regulating the serum metabolic disorder," *Journal of cellular biochemistry*, vol. 119, no. 9, pp. 7767–7779, 2018.
- [271] Y. Cao, G. Li, Y. F. Wang et al., "Neuroprotective effect of baicalin on compression spinal cord injury in rats," *Brain research*, vol. 1357, pp. 115–123, 2010.
- [272] F. du, X. Wang, B. Shang et al., "Gastrodin ameliorates spinal cord injury via antioxidant and anti-inflammatory effects," *Acta Biochimica Polonica*, vol. 63, no. 3, pp. 589–593, 2016.
- [273] P. Xia, X. Gao, L. Duan, W. Zhang, and Y.-F. Sun, "Mulberrin (Mul) reduces spinal cord injury (SCI)-induced apoptosis, inflammation and oxidative stress in rats via miRNA-337 by targeting Nrf-2," *Biomedicine & Pharmacotherapy*, vol. 107, pp. 1480–1487, 2018.
- [274] D.-s. Yu, Y.-s. Wang, Y.-l. Bi et al., "Salvianolic acid A ameliorates the integrity of blood-spinal cord barrier via miR-101/Cul3/Nrf2/HO-1 signaling pathway," *Brain research*, vol. 1657, pp. 279–287, 2017.
- [275] Z.-k. Fan, G. Lv, Y.-f. Wang et al., "The protective effect of salvianolic acid B on blood-spinal cord barrier after compression spinal cord injury in rats," *Journal of Molecular Neuroscience*, vol. 51, no. 3, pp. 986–993, 2013.
- [276] C. S. Park, J. Y. Lee, H. Y. Choi et al., "Gallic acid attenuates blood-spinal cord barrier disruption by inhibiting Jmjd3 expression and activation after spinal cord injury," *Neurobiology of Disease*, vol. 145, article 105077, 2020.
- [277] Y. H. Yang, Z. Wang, J. Zheng, and R. Wang, "Protective effects of gallic acid against spinal cord injury-induced oxidative stress," *Molecular medicine reports*, vol. 12, no. 2, pp. 3017–3024, 2015.
- [278] Z. Liu, X. Yao, B. Sun et al., "Pretreatment with kaempferol attenuates microglia-mediate neuroinflammation by inhibiting MAPKs-NF- κ B signaling pathway and pyroptosis after secondary spinal cord injury," *Free Radical Biology and Medicine*, vol. 168, pp. 142–154, 2021.
- [279] I. Paterniti, D. Impellizzeri, R. Di Paola, M. Navarra, S. Cuzzocrea, and E. Esposito, "A new co-ultramicrosized composite including palmitoylethanolamide and luteolin to prevent neuroinflammation in spinal cord injury," *Journal of neuroinflammation*, vol. 10, no. 1, pp. 1–11, 2013.
- [280] L.-B. Shi, P.-F. Tang, W. Zhang, Y.-P. Zhao, L.-C. Zhang, and H. Zhang, "Naringenin inhibits spinal cord injury-induced activation of neutrophils through miR-223," *Gene*, vol. 592, no. 1, pp. 128–133, 2016.
- [281] A. R. Khalatbary and H. Ahmadvand, "Neuroprotective effect of oleuropein following spinal cord injury in rats," *Neurological research*, vol. 34, no. 1, pp. 44–51, 2012.
- [282] D. Impellizzeri, E. Esposito, E. Mazzon et al., "The effects of a polyphenol present in olive oil, oleuropein aglycone, in an experimental model of spinal cord injury in mice," *Biochemical pharmacology*, vol. 83, no. 10, pp. 1413–1426, 2012.
- [283] L. Tao, X. Liu, W. Da, Z. Tao, and Y. Zhu, "Pycnogenol achieves neuroprotective effects in rats with spinal cord injury by stabilizing the mitochondrial membrane potential," *Neurological Research*, vol. 42, no. 7, pp. 597–604, 2020.
- [284] M.-J. Tsai, J.-F. Liao, D.-Y. Lin et al., "Silymarin protects spinal cord and cortical cells against oxidative stress and lipopolysaccharide stimulation," *Neurochemistry international*, vol. 57, no. 8, pp. 867–875, 2010.
- [285] C. Liu, Z. Shi, L. Fan, C. Zhang, K. Wang, and B. Wang, "Resveratrol improves neuron protection and functional recovery in rat model of spinal cord injury," *Brain Research*, vol. 1374, pp. 100–109, 2011.
- [286] H. S. Kavakli, C. Koca, and O. Alici, "Antioxidant effects of curcumin in spinal cord injury in rats," *Turkish Journal of Trauma and Emergency Surgery*, vol. 17, no. 1, pp. 14–18, 2011.
- [287] K. T. Kim, M. J. Kim, D. C. Cho et al., "The neuroprotective effect of treatment with curcumin in acute spinal cord injury: Laboratory Investigation," *Neurologia medico-chirurgica*, vol. 54, no. 5, pp. 387–394, 2014.
- [288] W. Li, S. Yao, H. Li, Z. Meng, and X. Sun, "Curcumin promotes functional recovery and inhibits neuronal apoptosis after spinal cord injury through the modulation of autophagy," *The journal of spinal cord medicine*, vol. 1-9, 2021.
- [289] X. Liu, Y. Zhang, Y. Yang et al., "Therapeutic effect of curcumin and methylprednisolone in the rat spinal cord injury," *The Anatomical Record*, vol. 301, no. 4, pp. 686–696, 2018.
- [290] Z.-H. Wang, Y.-X. Xie, J.-W. Zhang et al., "Carnosol protects against spinal cord injury through Nrf-2 upregulation," *Journal of Receptors and Signal Transduction*, vol. 36, no. 1, pp. 72–78, 2016.
- [291] J. Xi, X. Luo, Y. Wang et al., "Tetrahydrocurcumin protects against spinal cord injury and inhibits the oxidative stress response by regulating FOXO4 in model rats," *Experimental and therapeutic medicine*, vol. 18, no. 5, pp. 3681–3687, 2019.
- [292] E. Collyer, A. Catenaccio, D. Lemaitre et al., "Sprouting of axonal collaterals after spinal cord injury is prevented by delayed axonal degeneration," *Experimental neurology*, vol. 261, pp. 451–461, 2014.
- [293] H. Katoh, K. Yokota, and M. G. Fehlings, "Regeneration of spinal cord connectivity through stem cell transplantation and biomaterial scaffolds," *Frontiers in cellular neuroscience*, vol. 13, p. 248, 2019.
- [294] M. S. Chen, A. B. Huber, M. E. van der Haar et al., "Nogo-A is a myelin-associated neurite outgrowth inhibitor and an

- antigen for monoclonal antibody IN-1," *Nature*, vol. 403, no. 6768, pp. 434–439, 2000.
- [295] A. Josephson, J. Widenfalk, H. Widmer, L. Olson, and C. Spenger, "NOGO mRNA expression in adult and fetal human and rat nervous tissue and in weight drop injury," *Experimental neurology*, vol. 169, no. 2, pp. 319–328, 2001.
- [296] W. A. Barton, B. P. Liu, D. Tzvetkova et al., "Structure and axon outgrowth inhibitor binding of the Nogo-66 receptor and related proteins," *The EMBO journal*, vol. 22, no. 13, pp. 3291–3302, 2003.
- [297] S. Li, J.-E. Kim, S. Budel, T. G. Hampton, and S. M. Strittmatter, "Transgenic inhibition of Nogo-66 receptor function allows axonal sprouting and improved locomotion after spinal injury," *Molecular and Cellular Neuroscience*, vol. 29, no. 1, pp. 26–39, 2005.
- [298] T. Liebscher, L. Schnell, D. Schnell et al., "Nogo-A antibody improves regeneration and locomotion of spinal cord-injured rats," *Annals of Neurology*, vol. 58, no. 5, pp. 706–719, 2005.
- [299] T. GrandPré, S. Li, and S. M. Strittmatter, "Nogo-66 receptor antagonist peptide promotes axonal regeneration," *Nature*, vol. 417, no. 6888, pp. 547–551, 2002.
- [300] U. Gundimeda, T. H. McNeill, B. A. Barseghian et al., "Polyphenols from green tea prevent antineurotogenic action of Nogo-A via 67-kDa laminin receptor and hydrogen peroxide," *Journal of neurochemistry*, vol. 132, no. 1, pp. 70–84, 2015.
- [301] Takekazu Kubo, Katsuhiko Hata, Atsushi Yamaguchi, and Toshihide Yamashita, "Rho-ROCK inhibitors as emerging strategies to promote nerve regeneration," *Current pharmaceutical design*, vol. 13, no. 24, pp. 2493–2499, 2007.
- [302] W. Young, "Spinal cord regeneration," *Cell transplantation*, vol. 23, no. 4-5, pp. 573–611, 2014.
- [303] S. A. Benitah, P. F. Valerón, and J. C. Lacal, "ROCK and nuclear factor- κ B-dependent activation of cyclooxygenase-2 by Rho GTPases: effects on tumor growth and therapeutic consequences," *Molecular Biology of the Cell*, vol. 14, no. 7, pp. 3041–3054, 2003.
- [304] H. Tokuda, S. Takai, R. Matsushima-Nishiwaki et al., "Function of Rho-kinase in prostaglandin D₂-induced interleukin-6 synthesis in osteoblasts," *Prostaglandins, leukotrienes and essential fatty acids*, vol. 79, no. 1-2, pp. 41–46, 2008.
- [305] M. Leibinger, A. Andreadaki, and D. Fischer, "Role of mTOR in neuroprotection and axon regeneration after inflammatory stimulation," *Neurobiology of disease*, vol. 46, no. 2, pp. 314–324, 2012.
- [306] A. Hervera, F. de Virgiliis, I. Palmisano et al., "Reactive oxygen species regulate axonal regeneration through the release of exosomal NADPH oxidase 2 complexes into injured axons," *Nature cell biology*, vol. 20, no. 3, pp. 307–319, 2018.
- [307] Q. Han, Y. Xie, J. D. Ordaz et al., "Restoring cellular energetics promotes axonal regeneration and functional recovery after spinal cord injury," *Cell metabolism*, vol. 31, no. 3, pp. 623–641.e8, 2020.
- [308] H.-J. Lin, C.-C. Su, H.-F. Lu et al., "Curcumin blocks migration and invasion of mouse-rat hybrid retina ganglion cells (N18) through the inhibition of MMP-2,-9, FAK, Rho A and ROCK-1 gene expression," *Oncology reports*, vol. 23, no. 3, pp. 665–670, 2010.
- [309] R. Requejo-Aguilar, A. Alastrue-Agudo, M. Cases-Villar et al., "Combined polymer-curcumin conjugate and ependymal progenitor/stem cell treatment enhances spinal cord injury functional recovery," *Biomaterials*, vol. 113, pp. 18–30, 2017.
- [310] S. Okada, M. Hara, K. Kobayakawa, Y. Matsumoto, and Y. Nakashima, "Astrocyte reactivity and astrogliosis after spinal cord injury," *Neuroscience research*, vol. 126, pp. 39–43, 2018.
- [311] C. C. Stichel, K. Lips, G. Wunderlich, and H. W. Müller, "Reconstruction of transected postcommissural fornix in adult rat by Schwann cell suspension grafts," *Experimental neurology*, vol. 140, no. 1, pp. 21–36, 1996.
- [312] J. Frisé, C. B. Johansson, C. Török, M. Risling, and U. Lendahl, "Rapid, widespread, and longlasting induction of nestin contributes to the generation of glial scar tissue after CNS injury," *The Journal of cell biology*, vol. 131, no. 2, pp. 453–464, 1995.
- [313] J. S. Rudge, Y. Li, E. M. Pasnikowski et al., "Neurotrophic factor receptors and their signal transduction capabilities in rat astrocytes," *European Journal of Neuroscience*, vol. 6, no. 5, pp. 693–705, 1994.
- [314] A. Nadal, E. Fuentes, J. Pastor, and P. A. McNaughton, "Plasma albumin is a potent trigger of calcium signals and DNA synthesis in astrocytes," *Proceedings of the National Academy of Sciences*, vol. 92, no. 5, pp. 1426–1430, 1995.
- [315] R. Wang, X. Zhang, J. Zhang et al., "Oxygen-glucose deprivation induced glial scar-like change in astrocytes," *PLoS One*, vol. 7, no. 5, article e37574, 2012.
- [316] R. Shechter, O. Miller, G. Yovel et al., "Recruitment of beneficial M2 macrophages to injured spinal cord is orchestrated by remote brain choroid plexus," *Immunity*, vol. 38, no. 3, pp. 555–569, 2013.
- [317] D. J. Donnelly, E. E. Longbrake, T. M. Shawler et al., "Deficient CX3CR1 signaling promotes recovery after mouse spinal cord injury by limiting the recruitment and activation of Ly6C^{lo}/iNOS⁺ macrophages," *Journal of Neuroscience*, vol. 31, no. 27, pp. 9910–9922, 2011.
- [318] A. Badner, R. Vawda, A. Laliberte et al., "Early intravenous delivery of human brain stromal cells modulates systemic inflammation and leads to vasoprotection in traumatic spinal cord injury," *Stem cells translational medicine*, vol. 5, no. 8, pp. 991–1003, 2016.
- [319] E. V. Voß, J. Škuljec, V. Gudi et al., "Characterisation of microglia during de- and remyelination: Can they create a repair promoting environment?," *Neurobiology of disease*, vol. 45, no. 1, pp. 519–528, 2012.
- [320] J. E. Herrmann, T. Imura, B. Song et al., "STAT3 is a critical regulator of astrogliosis and scar formation after spinal cord injury," *Journal of Neuroscience*, vol. 28, no. 28, pp. 7231–7243, 2008.
- [321] GUO, JIN, ISHIKAWA et al., "Regulation of beta-chemokine mRNA expression in adult rat astrocytes by lipopolysaccharide, proinflammatory and immunoregulatory cytokines," *Scandinavian journal of immunology*, vol. 48, no. 5, pp. 502–508, 1998.
- [322] P. Krupa, B. Svobodova, J. Dubisova, S. Kubinova, P. Jendelova, and L. Machova Urdzikova, "Nano-formulated curcumin (Lipodisq™) modulates the local inflammatory response, reduces glial scar and preserves the white matter after spinal cord injury in rats," *Neuropharmacology*, vol. 155, pp. 54–64, 2019.
- [323] J. Ruzicka, L. M. Urdzikova, A. Kloudova et al., "Anti-inflammatory compound curcumin and mesenchymal stem cells in

- the treatment of spinal cord injury in rats," *Acta Neurobiologiae Experimentalis*, vol. 78, no. 4, pp. 358–374, 2018.
- [324] J. Yuan, B. O. Botchway, Y. Zhang, X. Tan, X. Wang, and X. Liu, "Curcumin can improve spinal cord injury by inhibiting TGF- β -SOX9 signaling pathway," *Cellular and molecular neurobiology*, vol. 39, no. 5, pp. 569–575, 2019.
- [325] L. Xu, B. O. Botchway, S. Zhang, J. Zhou, and X. Liu, "Inhibition of NF- κ B signaling pathway by resveratrol improves spinal cord injury," *Frontiers in neuroscience*, vol. 12, p. 690, 2018.
- [326] W. Rong, Y.-w. Pan, X. Cai et al., "The mechanism of naringin-enhanced remyelination after spinal cord injury," *Neural regeneration research*, vol. 12, no. 3, pp. 470–477, 2017.
- [327] R. Subbarayan, R. Barathidasan, S. T. Raja et al., "Human gingival derived neuronal cells in the optimized caffeic acid hydrogel for hemitranssection spinal cord injury model," *Journal of cellular biochemistry*, vol. 121, no. 3, pp. 2077–2088, 2020.
- [328] M. Karus, A. Ulc, M. Ehrlich et al., "Regulation of oligodendrocyte precursor maintenance by chondroitin sulphate glycosaminoglycans," *Glia*, vol. 64, no. 2, pp. 270–286, 2016.
- [329] E. J. Bradbury, L. D. Moon, R. J. Popat et al., "Chondroitinase ABC promotes functional recovery after spinal cord injury," *Nature*, vol. 416, no. 6881, pp. 636–640, 2002.
- [330] H. Suzuki, C. S. Ahuja, R. P. Salewski et al., "Neural stem cell mediated recovery is enhanced by chondroitinase ABC pretreatment in chronic cervical spinal cord injury," *PLoS One*, vol. 12, no. 8, article e0182339, 2017.
- [331] P. Lu, L. L. Jones, and M. H. Tuszynski, "Axon regeneration through scars and into sites of chronic spinal cord injury," *Experimental neurology*, vol. 203, no. 1, pp. 8–21, 2007.
- [332] R. R. Williams and M. B. Bunge, "Schwann cell transplantation: a repair strategy for spinal cord injury?," *Progress in brain research*, vol. 201, pp. 295–312, 2012.
- [333] R. Quertainmont, D. Cantinieaux, O. Botman, S. Sid, J. Schoenen, and R. Franzen, "Mesenchymal stem cell graft improves recovery after spinal cord injury in adult rats through neurotrophic and pro-angiogenic actions," *PloS one*, vol. 7, no. 6, article e39500, 2012.
- [334] P. Tabakow, W. Jarmundowicz, B. Czapiga et al., "Transplantation of autologous olfactory ensheathing cells in complete human spinal cord injury," *Cell transplantation*, vol. 22, no. 9, pp. 1591–1612, 2013.
- [335] G. Kumagai, P. Tsoulfas, S. Toh, I. McNiece, H. M. Bramlett, and W. D. Dietrich, "Genetically modified mesenchymal stem cells (MSCs) promote axonal regeneration and prevent hypersensitivity after spinal cord injury," *Experimental neurology*, vol. 248, pp. 369–380, 2013.
- [336] D. D. Pearse, F. C. Pereira, A. E. Marcillo et al., "cAMP and Schwann cells promote axonal growth and functional recovery after spinal cord injury," *Nature medicine*, vol. 10, no. 6, pp. 610–616, 2004.
- [337] K. Zukor, S. Belin, C. Wang, N. Keelan, X. Wang, and Z. He, "Short hairpin RNA against PTEN enhances regenerative growth of corticospinal tract axons after spinal cord injury," *Journal of Neuroscience*, vol. 33, no. 39, pp. 15350–15361, 2013.
- [338] A. Tandon, S. J. Singh, and R. K. Chaturvedi, "Stem cells as potential targets of polyphenols in multiple sclerosis and Alzheimer's disease," *BioMed research international*, vol. 2018, Article ID 1483791, 30 pages, 2018.
- [339] S. K. Tiwari, S. Agarwal, B. Seth et al., "Curcumin-loaded nanoparticles potently induce adult neurogenesis and reverse cognitive deficits in Alzheimer's disease model via canonical Wnt/ β -catenin pathway," *ACS nano*, vol. 8, no. 1, pp. 76–103, 2014.
- [340] D. R. Ormond, C. Shannon, J. Oppenheim et al., "Stem cell therapy and curcumin synergistically enhance recovery from spinal cord injury," *PLoS One*, vol. 9, no. 2, article e88916, 2014.
- [341] W. Wanjiang, C. Xin, C. Yaxing et al., "Curcumin improves human umbilical cord-derived mesenchymal stem cell survival via ERK1/2 signaling and promotes motor outcomes after spinal cord injury," *Cellular and Molecular Neurobiology*, vol. 1–12, 2020.
- [342] W.-S. Bang, K.-T. Kim, Y. J. Seo, D.-C. Cho, J.-K. Sung, and C. H. Kim, "Curcumin increase the expression of neural stem/progenitor cells and improves functional recovery after spinal cord injury," *Journal of Korean Neurosurgical Society*, vol. 61, no. 1, pp. 10–18, 2018.
- [343] P. Bonilla, J. Hernandez, E. Giraldo et al., "Human-induced neural and mesenchymal stem cell therapy combined with a curcumin nanoconjugate as a spinal cord injury treatment," *International journal of molecular sciences*, vol. 22, no. 11, p. 5966, 2021.
- [344] D. S. Yu, L. B. Liu, Y. Cao et al., "Combining bone marrow stromal cells with green tea polyphenols attenuates the blood-spinal cord barrier permeability in rats with compression spinal cord injury," *Journal of Molecular Neuroscience*, vol. 56, no. 2, pp. 388–396, 2015.
- [345] S. J. Hewlings and D. S. Kalman, "Curcumin: a review of its effects on human health," *Foods*, vol. 6, no. 10, p. 92, 2017.
- [346] V. Krup, H. Prakash L, and H. A., "Pharmacological activities of turmeric (*Curcuma longa* Linn): a review," *Journal of Homeopathy & Ayurvedic Medicine*, vol. 2, no. 4, p. 2167, 2013.
- [347] J.-J. Yu, L.-B. Pei, Y. Zhang, Z.-Y. Wen, and J.-L. Yang, "Chronic supplementation of curcumin enhances the efficacy of antidepressants in major depressive disorder: a randomized, double-blind, placebo-controlled pilot study," *Journal of clinical psychopharmacology*, vol. 35, no. 4, pp. 406–410, 2015.
- [348] G. Belcaro, M. R. Cesarone, M. Dugall et al., "Efficacy and safety of Meriva[®], a curcumin-phosphatidylcholine complex, during extended administration in osteoarthritis patients," *Alternative Medicine Review*, vol. 15, no. 4, pp. 337–344, 2010.
- [349] N. Kertia, A. H. Asdie, W. Rochmah, and Marsetyawan, "Ability of curcuminoid compared to diclofenac sodium in reducing the secretion of cyclooxygenase-2 enzyme by synovial fluid's monocytes of patients with osteoarthritis," *Acta Medica Indonesiana*, vol. 44, no. 2, pp. 105–113, 2012.
- [350] Y. Panahi, A. R. Rahimnia, M. Sharafi, G. Alishiri, A. Saburi, and A. Sahebkar, "Curcuminoid treatment for knee osteoarthritis: a randomized double-blind placebo-controlled trial," *Phytotherapy Research*, vol. 28, no. 11, pp. 1625–1631, 2014.
- [351] R. Satoskar, S. Shah, and S. Shenoy, "Evaluation of anti-inflammatory property of curcumin (diferuloyl methane) in patients with postoperative inflammation," *International journal of clinical pharmacology, therapy, and toxicology*, vol. 24, no. 12, pp. 651–654, 1986.

- [352] M. Hatefi, M. R. H. Ahmadi, A. Rahmani, M. M. Dastjerdi, and K. Asadollahi, "Effects of curcumin on bone loss and biochemical markers of bone turnover in patients with spinal cord injury," *World neurosurgery*, vol. 114, pp. e785–e791, 2018.
- [353] D. J. Allison and D. S. Ditor, "Targeting inflammation to influence mood following spinal cord injury: a randomized clinical trial," *Journal of neuroinflammation*, vol. 12, no. 1, pp. 1–10, 2015.
- [354] S. Dolati, Z. Babaloo, H. Ayromlou et al., "Nanocurcumin improves regulatory T-cell frequency and function in patients with multiple sclerosis," *Journal of neuroimmunology*, vol. 327, pp. 15–21, 2019.
- [355] S. Asadi, M. S. Gholami, F. Siassi, M. Qorbani, K. Khamoshian, and G. Sotoudeh, "Nano curcumin supplementation reduced the severity of diabetic sensorimotor polyneuropathy in patients with type 2 diabetes mellitus: A randomized double-blind placebo-controlled clinical trial," *Complementary therapies in medicine*, vol. 43, pp. 253–260, 2019.
- [356] C. Moussa, M. Hebron, X. Huang et al., "Resveratrol regulates neuro-inflammation and induces adaptive immunity in Alzheimer's disease," *Journal of neuroinflammation*, vol. 14, no. 1, pp. 1–10, 2017.
- [357] S. Seyyedebrahimi, H. Khodabandehloo, E. Nasli Esfahani, and R. Meshkani, "The effects of resveratrol on markers of oxidative stress in patients with type 2 diabetes: a randomized, double-blind, placebo-controlled clinical trial," *Acta diabetologica*, vol. 55, no. 4, pp. 341–353, 2018.
- [358] J. Most, S. Timmers, I. Warnke et al., "Combined epigallocatechin-3-gallate and resveratrol supplementation for 12 wk increases mitochondrial capacity and fat oxidation, but not insulin sensitivity, in obese humans: a randomized controlled trial," *The American journal of clinical nutrition*, vol. 104, no. 1, pp. 215–227, 2016.
- [359] F. Hendouei, H. Sanjari Moghaddam, M. R. Mohammadi, N. Taslimi, F. Rezaei, and S. Akhondzadeh, "Resveratrol as adjunctive therapy in treatment of irritability in children with autism: a double-blind and placebo-controlled randomized trial," *Journal of clinical pharmacy and therapeutics*, vol. 45, no. 2, pp. 324–334, 2020.
- [360] E. M. Yiu, G. Tai, R. E. Peverill et al., "An open-label trial in Friedreich ataxia suggests clinical benefit with high-dose resveratrol, without effect on frataxin levels," *Journal of neurology*, vol. 262, no. 5, pp. 1344–1353, 2015.
- [361] J. Morillas-Ruiz, J. Rubio-Perez, M. Albaladejo, P. Zafrilla, S. Parra, and M. Vidal-Guevara, "Effect of an antioxidant drink on homocysteine levels in Alzheimer's patients," *Journal of the neurological sciences*, vol. 299, no. 1–2, pp. 175–178, 2010.
- [362] P. Valensi, C. le Devehat, J.-L. Richard et al., "A multicenter, double-blind, safety study of QR-333 for the treatment of symptomatic diabetic peripheral neuropathy: a preliminary report," *Journal of Diabetes and its Complications*, vol. 19, no. 5, pp. 247–253, 2005.
- [363] A. A. Verlaet, B. Ceulemans, H. Verhelst et al., "Effect of Pycnogenol® on attention-deficit hyperactivity disorder (ADHD): study protocol for a randomised controlled trial," *Trials*, vol. 18, no. 1, pp. 1–9, 2017.
- [364] D. O. Clark, H. Xu, L. Moser et al., "MIND food and speed of processing training in older adults with low education, the MINDSpeed Alzheimer's disease prevention pilot trial," *Contemporary clinical trials*, vol. 84, article 105814, 2019.
- [365] T. Traustadóttir, S. S. Davies, A. A. Stock et al., "Tart cherry juice decreases oxidative stress in healthy older men and women," *The Journal of nutrition*, vol. 139, no. 10, pp. 1896–1900, 2009.
- [366] V. Ferianec, M. Fülöp, M. Ježovičová et al., "The oak-wood extract Robuvit® improves recovery and oxidative stress after hysterectomy: a randomized, double-blind, placebo-controlled pilot study," *Nutrients*, vol. 12, no. 4, p. 913, 2020.
- [367] Y. X. You, S. Shahar, N. F. Rajab et al., "Effects of 12 weeks Cosmos caudatus supplement among older adults with mild cognitive impairment: a randomized, double-blind and placebo-controlled trial," *Nutrients*, vol. 13, no. 2, p. 434, 2021.
- [368] D. di Pierro, C. Ciaccio, D. Sbardella et al., "Effects of oral administration of common antioxidant supplements on the energy metabolism of red blood cells. Attenuation of oxidative stress-induced changes in Rett syndrome erythrocytes by CoQ10," *Molecular and cellular biochemistry*, vol. 463, no. 1–2, pp. 101–113, 2020.
- [369] M. Dvořáková, M. Sivoňová, J. Trebatická et al., "The effect of polyphenolic extract from pine bark, Pycnogenol® on the level of glutathione in children suffering from attention deficit hyperactivity disorder (ADHD)," *Redox Report*, vol. 11, no. 4, pp. 163–172, 2006.
- [370] Y. Panahi, M. S. Hosseini, N. Khalili et al., "Effects of curcumin on serum cytokine concentrations in subjects with metabolic syndrome: a post-hoc analysis of a randomized controlled trial," *Biomedicine & pharmacotherapy*, vol. 82, pp. 578–582, 2016.
- [371] Y. Panahi, M. S. Hosseini, N. Khalili, E. Naimi, M. Majeed, and A. Sahebkar, "Antioxidant and anti-inflammatory effects of curcuminoid-piperine combination in subjects with metabolic syndrome: a randomized controlled trial and an updated meta-analysis," *Clinical nutrition*, vol. 34, no. 6, pp. 1101–1108, 2015.
- [372] G. Patki, N. Solanki, F. Atrooz, F. Allam, and S. Salim, "Depression, anxiety-like behavior and memory impairment are associated with increased oxidative stress and inflammation in a rat model of social stress," *Brain research*, vol. 1539, pp. 73–86, 2013.
- [373] F. Allam, A. T. Dao, G. Chugh et al., "Grape powder supplementation prevents oxidative stress-induced anxiety-like behavior, memory impairment, and high blood pressure in rats," *The Journal of nutrition*, vol. 143, no. 6, pp. 835–842, 2013.
- [374] N. Solanki, A. Salvi, G. Patki, and S. Salim, "Modulating oxidative stress relieves stress-induced behavioral and cognitive impairments in rats," *International Journal of Neuropsychopharmacology*, vol. 20, no. 7, pp. 550–561, 2017.
- [375] J. M. Zuidema, R. J. Gilbert, and D. J. Osterhout, "Nanoparticle technologies in the spinal cord," *Cells Tissues Organs*, vol. 202, no. 1–2, pp. 102–115, 2016.
- [376] Y. Peng, L. Chen, S. Ye et al., "Research and development of drug delivery systems based on drug transporter and nanoformulation," *Asian Journal of Pharmaceutical Sciences*, vol. 15, no. 2, pp. 220–236, 2020.